Published in final edited form as:

Biochem Biophys Res Commun. 2013 September 27; 439(3): . doi:10.1016/j.bbrc.2013.08.056.

MicroRNA-133a-1 regulates inflammasome activation through uncoupling protein-2

Sayantani Bandyopadhyay[#], Troy Lane[#], Rajanbabu Venugopal, Prasanna Tamarapu Parthasarathy, Young Cho, Lakshmi Galam, Richard Lockey, and Narasaiah Kolliputi^{*}

Division of Allergy and Immunology, Department of Internal Medicine, Morsani College of Medicine, University of South Florida, 12901 Bruce B. Downs Blvd., Tampa, FL 33612

Abstract

Inflammasomes are multimeric protein complexes involved in the processing of IL-1 β through Caspase-1 cleavage. NLRP3 is the most widely studied inflammasome, which has been shown to respond to a large number of both endogenous and exogenous stimuli. Although studies have begun to define basic pathways for the activation of inflammasome and have been instrumental in identifying therapeutics for inflammasome related disorders; understanding the inflammasome activation at the molecular level is still incomplete. Recent functional studies indicate that microRNAs (miRs) regulate molecular pathways and can lead to diseased states *when hampered or overexpressed*. Mechanisms *involving the* miRNA regulatory network in the activation of inflammasome and IL-1 β processing is yet unknown. This report *investigates* the involvement of miR-133a-1 in the activation of inflammasome (NLRP3) and IL-1 β production. miR-133a-1 is known to target the mitochondrial uncoupling protein 2 (UCP2). The role of UCP2 in inflammasome activation has remained elusive. To understand the role of miR-133a-1 in regulating inflammasome activation, we either overexpressed or suppressed miR-133a-1 in differentiated THP1 cells that express *the* NLRP3 inflammasome. Levels of Caspase-1 and IL-1 β were analyzed by *Western* blot analysis. For the first time, we showed that overexpression of miR-133a-1 *increases* Caspase-1 p10 and IL-1 β p17 cleavage, concurrently suppressing mitochondrial uncoupling protein 2 (UCP2). Surprisingly, our results demonstrated that miR-133a-1 controls inflammasome activation without affecting the basal expression of the individual inflammasome components NLRP3 and ASC or its immediate downstream targets proIL-1 β and pro-Caspase-1. *To confirm the involvement of UCP2 in the regulation of inflammasome activation, Caspase-1 p10 and IL-1 β p17 cleavage in UCP2 of overexpressed and silenced THP1 cells were studied.* Suppression of UCP2 by siRNA enhanced the inflammasome activity stimulated by H₂O₂ and, conversely, overexpression of UCP2 decreased the inflammasome activation. *Collectively*, these studies suggest that miR-133a-1 *suppresses* inflammasome activation via the suppression of UCP2.

© 2013 Elsevier Inc. All rights reserved.

^{*}Address correspondence to: Narasaiah Kolliputi, Division of Allergy and Immunology, Department of Internal Medicine, Morsani College of Medicine, University of South Florida, 12901 Bruce B. Downs Blvd., MDC 19, Tampa, FL 33612, Phone: 813-974-5419, Fax: 813-974-8575, nkollipu@health.usf.edu.

[#]Equally contributed

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

Inflammasomes are multi-protein structures that regulate the activation of Caspase-1 and the maturation of pro-inflammatory cytokines like IL-1 β , IL-18, and IL-33 [1]. Inflammasome activation is a two-step process; the first signal is *enabled* through the activation of pathogen response receptors (PRRs). Activated PRRs stimulate NF- κ B and prime *the formation of the* inflammasome complex. The second signal comes from a range of stimuli *such as* ATP, uric acid crystals, hydrogen peroxide, reactive oxygen species (ROS), or intracellular stimuli such as sterile inflammation [2]. Among the wide variety of inflammasomes, the NLRP3 inflammasome complex is well studied [3].

Although the precise mechanisms of activation are not known, studies demonstrate that NLRP3 is activated by a wide range of compounds: *they include* both exogenous *stimuli* as well as host ligands including bacterial RNA, ATP, uric acid crystals, antiviral imidazoquinoline compounds, ceramide, and oxygen toxicity [4,5,6,7]. So far, based on these findings, three key mechanisms have been described to account for NLRP3 activation [3]. One *mechanism for* NLRP3 activation is potassium efflux [8]. External ATP is recognized by the P₂X₇ receptor, a cation channel, *which facilitates the activation of* potassium efflux that, in turn, triggers NLRP3 activation [8]. The generation of mitochondria-derived ROS plays a critical role *in* the activation of NLRP3 [9]. Phagolysosomal destabilization also activates NLRP3; *it is* caused by large crystals and particulates such as monosodium urate (MSU), adjuvant alum, asbestos, and silica [10]. Upon activation of NLRP3, it oligomerizes and recruits the ASC domain which, in turn, recruits pro-Caspase-1. This event leads to auto-proteolytic cleavage of pro-Caspase-1 and formation of active Caspase-1. Active Caspase-1 cleaves pro-IL-1 β *and leads to* secretion of active IL-1 β [11].

One defense *mechanism* of the innate immune system *involves* inflammasomes *which* combat invading microbes via activation of Caspase-1 and the production of mature pyrogenic cytokine IL-1 β [3]. IL-1 β is an essential mediator of the inflammatory response causing fever, hypotension, and production of other pro-inflammatory cytokines [12]. Inflammasomes also take part in a variety of cellular activities including cell proliferation, differentiation, and apoptosis [13]. The synthesis of IL-1 β is very tightly regulated by several mechanisms; however, mutations in the NLRP3 gene are associated with a spectrum of auto-inflammatory diseases characterized by excessive production of IL-1 β , *they include diseases such as* cryopyrin-associated periodic syndrome (CAPS), familial cold auto-inflammatory syndrome, Muckle-Wells syndrome, and chronic infantile cutaneous neurological articular syndrome [14,15,16]. *Inflammasomes* also *aggravat* gout [17], asbestosis, silicosis [10,18], and Alzheimer's disease [19].

Recently, microRNAs (miRs) have emerged as a discrete class of biological regulators with conserved functions that have the potential to control IL-1 β generation [20]. miRs play a crucial role in cell proliferation, cell differentiation, apoptosis, developmental timing, and other regulatory pathways [21,22,23,24,25,26]. Increasing evidence *in* the involvement of miRs in clinical disease models *demonstrate that* miRs are considered promising agents in *clinically-associated* research [22,23,24,25,26]. In general, miRs are short, single-stranded, non-protein coding RNA molecules [26,27]. miRs bind to the complementary untranslated region (UTR) of the target mRNA and cleaves the segment: resulting in instability and subsequent degradation of the mRNA [22,28].

In the present study, we *investigate* the role of miR-133A in inflammasome activation and IL-1 β production. miR-133-a-1 was first characterized in mice; it is homologous to some other species, including invertebrates [27]. There are three miR-133 genes identified in the

human genome: miR-133a-1, miR-133a-2, and miR-133b [28]. Some in vitro studies suggest that *upregulation* of miR-133a-1 suppresses UCP2 expression [29]. UCP2 is an integral membrane protein and a member of the large UCP family; it plays a role in ATP synthesis and ROS production [30,31]. UCP2 is widely expressed in several tissues compared to the other UCP family members including the spleen, kidney, pancreas, and immune system. [32]. A previous study shows UCP2 negatively regulates ROS and induces an anti-inflammatory response [31]. It plays a critical role in UCP2-mediated mitochondrial function in phagocytosis [33].

In this report, we *demonstrate* the activation of Caspase-1 p10 and IL-1 β p17 by *miR-133a-1-induced* suppression of UCP2 in *inflammasome-stimulated* THP1 cells.

2. Materials and Methods

2.1. Cells and reagents

THP1 cells derived from human lung adenocarcinoma were obtained from American Type Culture Collection (ATCC, Manassas, VA) and were maintained in RPMI-1640 medium as per vendor instructions. PMA (Sigma-Aldrich, St. Louis, MO, USA), anti-mouse (Cell Signaling Technology, Danvers, MA), anti-goat (Santa Cruz, Inc., CA, USA), or anti-rabbit (Cell signaling) HRP conjugated secondary antibodies were bought from *its listed companies*. Primary antibodies specific to NLRP3 (Enzo Life Sciences, Inc. Farmingdale, NY, USA), ASC (Enzo) Caspase-1 (Santa Cruz), UCP2 (R&D systems, MN, USA), IL-1 β (Cell signaling), β -Actin (Cell signaling), and GAPDH (Cell signaling) were used in *Western blot*.

2.2. Inflammasome activation

Inflammasome was stimulated in THP1 cells *as previously* mentioned [34]. In brief, THP1 cells were differentiated by treating *it* with 0.5 μ g/ml of PMA for three hours, and the medium was replaced with complete RPMI 1640 overnight. *After 24 hours*, cells were transfected with miR-133-a1 or miR vector control. *After 24 hours*, cells were briefly washed with PBS and stimulated with 5mM H₂O₂ for 3 hours or 2.5 mM ATP for 1 hour. The activation of inflammasome was confirmed by measuring the IL-1 β secretion and comparing the expression of mature IL-1 β and Caspase-1 expression by ELISA and *Western blot* analysis.

2.3. Micro RNA and SiRNA Transfection

THP1 cells were seeded at 0.5 million cells per well in a six-well plate, and differentiated with 0.5 μ g/ml of PMA for three hours. To remove the residual effect of PMA, the medium was replaced with complete RPMI 1640 overnight. *After 24 hours*, the medium was replaced by serum-free RPMI-1640 and cells were transfected with 2.5 μ g/well DNA of pCMV-miR control (OriGene, Rockville, MD) and miR-133a-1 (Origene) plasmid using Lipofectamine® LTX plus Reagent (Invitrogen, CA, USA) as per manufacturer instructions. After 4 hours of transfection, the medium was replaced with complete RPMI-1640. *After 24 hours*, cells were washed with PBS and medium was replaced by serum-free RPMI-1640 for two hours, then cells were treated with 5mM H₂O₂ for 3 hours.

2.4. Western blot

Cells were lysed with 150 μ l of RIPA buffer and 10 μ l/ml of protease and phosphatase inhibitors; protein quantification was performed using the Pierce BCA Protein Assay Kit (Thermo Scientific, Rockford, IL). Equal quantities of protein *were* loaded into SDS-PAGE, and protein was transferred to PVDF membrane. The membrane was blocked in 5% nonfat dry milk, and probed with primary and secondary antibodies diluted in blocking buffer. A

Pierce ECL *Western* blotting substrate kit (Thermo scientific Rockford, IL, USA) was used to detect the protein under ChemiDoc XRS (Bio-Rad, Hercules, CA, USA). Bands were quantified and densitometric analysis was performed using ImageJ (Data not shown).

2.5. Immunofluorescence and immunocytochemistry

Differentiated THP1 cells in 35mm, 4 *compartment*, and *glass-bottom* dish (Greiner bio-one, Maybachstr, Germany) were transfected with pCMV-miR control and pCMV-miR-133 plasmid containing IRES-tGFP for 24 *hours*. Localization of GFP was observed under the Olympus IX81 inverted microscope equipped with the 3i Yokogawa spinning disk scanner and CCD cameras with revolving laser arrays. *Cy3-labeled siRNA* scramble was transfected into differentiated THP1 cells. At 48 *hour* post-transfection, the cells were fixed with 4% paraformaldehyde; the nuclei were stained with DAPI, and examined for Cy3 localization under the microscope.

2.6 Bone marrow macrophage isolation from knockout mice

Approval of the study protocol was obtained from the University of South Florida Institutional Animal Care and Use Committee. All mice were maintained in a specific-pathogen-free animal facility at the University of South Florida and animal experiments were carried out according to the provisions of the Animal Welfare Act, PHS Animal Welfare Policy, and the principles of the NIH Guide for the Care and Use of Laboratory Animals. Bone marrow cells were obtained from ASC^{-/-}, NLRP3^{-/-} and wild type (WT) mice (The Jackson Laboratory, Bar Harbor, ME) by dissecting the *leg bones at the thigh bone*, then flushing the marrow from the leg bones with a 23-gauge needle and L-conditioned DMEM medium (M. A. Bioproducts, Walkersville, MD) supplemented with penicillin/streptomycin (P/S: 100 IU/ml and 100 pg/ml, respectively). The cells were dissociated by gentle vortexing, centrifuged at 1000 rpm, and resuspended in medium.

The cell suspension was seeded at 1 million cells per well in a 12-well plate. Medium was replaced once in every 3 days and cells were used for further experiments.

2.7 Statistical analysis

All experiments were performed in triplicate. A paired t-test was used to determine the statistical significance. A p-value of $p < 0.05$ was accepted as statistically significant.

3. Results

3.1 miR-133a-1 enhances Caspase-1 activation and IL-1 β processing

To understand the role of miR-133a-1 in the activation of NLRP3 inflammasome, THP1 cells were transfected with pCMV-miR (Vector control) or pCMV-miR-133a-1, followed by stimulation with *an* inflammasome activator, H₂O₂, as mentioned in the *Materials and Methods* section. After transfection and post-inflammasome stimulation, supernatants were used to check *Caspase-1* p10 and cleaved IL-1 β p17 expression by *Western* blot. Our results indicate that in pCMV-miR-133a-1, the transfection of miR-133a-1 itself increases *Caspase-1* p10 compared to miR vector control (Fig. 1A). In addition, inflammasome stimulation by H₂O₂ also further increases *Caspase-1* p10 release in supernatants. This data demonstrates that miR-133a-1 enhances *Caspase-1* p10 activation. Since inflammasome stimulation and *Caspase-1* activation process mature IL-1 β , secretion of mature IL-1 β p17 was analyzed in vector control and miR-133a-1 transfected THP1 cells. In miR vector control-transfected cells, the activation of IL-1 β p17 was higher in H₂O₂ stimulated cells (Fig. 1A). This confirms the inflammasome activation by H₂O₂. In miR-133a-1 transfected cells, the transfection of miR-133a-1 increase IL-1 β p17 release when compared to miR control vector. In addition, inflammasome stimulation by H₂O₂ also further increases IL-1 β

p17. This data demonstrates that miR-133a-1 enhances IL-1 β p17 processing and secretion. Since the plasmids encoding precursor microRNA also possess internal ribosome entry site (IRES)-driven tGFP the transfection was confirmed by checking the tGFP expression in a fluorescence microscope at 24 *hour* post-transfection (Fig. 1B). In order to understand whether miR-133a-1 only controls inflammasome activation or if it affects the basal expression of inflammasome components and its effector proteins, the expression of NLRP3, ASC, pro IL-1 β and *pro-Caspase-1* were analyzed by *Western* blot. The expression pattern of inflammasome components in miR-vector control and miR-133a-1 transfected cells under untreated and *H₂O₂-treated conditions* remain unchanged (Fig. 1C). These results suggest that *miR-133a-1* is involved in inflammasome activation that can lead to Caspase-1 activation *as well as* IL-1 β processing and secretion.

3.2 miR-133a-1 suppress UCP2 expression

The miRNA database shows that the miR-133a-1 targets the untranslated region (UTR) of UCP2. To confirm that miR-133a-1 controls UCP2, the expression of UCP2 was checked by *Western* blot in *miRNA-overexpressed* THP1 cells. The expression of UCP2 was *downregulated* in *miR-133a-1-transfected* THP1 cells (Fig. 1D). This data demonstrated *miR-133a-1-mediated* down-regulation of UCP2.

3.3 Silencing UCP2 enhances inflammasome activation

To understand the direct influence of UCP2 on inflammasome activation, Caspase-1 and IL-1 β secretion levels were examined in the UCP2 silenced cells. Human UCP2 *in siRNA's* were transfected into THP1 cells by using Lipofectamine as per manufacturer instructions. The transfection efficiency in 48 *hour* post-transfected cells were confirmed by examining the *Cy3- labeled siRNA-transfected* cells under microscope and *Western* blot analysis (Fig. 2A,2B). Compared to the scrambled siRNA control, UCP2 *siRNA-transfected* cells *exhibited* a significant increase in *Caspase-1* and IL-1 β secretion levels in response to *H₂O₂* (Fig. 2C). These results indicate that UCP2 suppression leads to activation of *the* inflammasome.

3.4 UCP2 overexpression limits Caspase-1 and IL-1 β production

Since *UCP2-silenced* cells enhanced inflammasome activation, *Caspase-1* and IL-1 β secretion *were* compared in *UCP2-overexpressed* THP1 cells. Human UCP2 cDNA tagged with GFP was transfected into THP1 cells; the transfection efficiency was confirmed by examining the GFP expression under microscope and by checking UCP2 levels by *Western* blot analysis (Fig. 3A, 3B). The *H₂O₂* mediated activation of *Caspase-1* and IL-1 β *decreased more significantly* in *UCP2-overexpressed* cells than the vector control (Fig. 3C). This demonstrated that UCP2 may act as a negative regulator of inflammasome activation.

3.5 UCP2 is upregulated in the absence of NLRP3 and ASC

To define the role of inflammasome components, NLRP3 and ASC, in the regulation of UCP2, bone marrow macrophages were derived from NLRP3 and ASC knockout mice (NLRP3^{-/-}, ASC^{-/-}), and the expression of UCP2 was compared with wild type controls (WT). UCP2 levels were increased in NLRP3^{-/-} and ASC^{-/-} cells when compared to *the* wild type *controls* (Fig. 4). This showed that UCP2 is negatively regulated by NLRP3 and ASC.

4. Discussion

In this study, we demonstrated the involvement of miR-133a-1 in the regulation of inflammasome effector proteins Caspase-1p10 and IL-1 β p17 by suppressing its target UCP2. Inflammasomes are multiprotein complexes that are involved in IL-1 β processing and secretion through a tightly regulated process. Since IL-1 β is directly correlated with

various disorders that include autoinflammatory, autoimmune, and infectious diseases; approaches to inhibit IL-1 β production *should* gain priority in clinical research. Recent studies are focusing on cellular mechanisms regulating IL-1 β processing and secretion. Since the discovery of *inflammasomes*, many molecular mechanisms have been explained to *present* the activation and processing of IL-1 β . However, there *is* little evidence that link *microRNA-mediated* regulation of inflammasome activation and IL-1 β processing. Recent studies with miR-155 unlocked the possibilities of *IL-1 β* regulation through the modulation of miRNA expression [35]. For the first time, we have *presented* the involvement of miR-133a-1 in modulating the activation of NLRP3 inflammasome.

Our results *indicate* that overexpression of miR-133a-1 increases *Caspase* and IL-1 β p17 levels in response to inflammasome stimuli: H₂O₂ (Fig. 3). In addition, *miR-133a-1-mediated* suppression of UCP2 (Fig. 1D) observed in this report was in accordance with previous studies [29]. Studies with UCP2 overexpression and siRNA *show* a direct relationship between UCP2 and the activation of Caspase-1 and processing of IL-1 β (Fig. 2, 3). Inhibition of UCP2 has been previously reported to elevate ROS and *not complete* phagocytosis. UCP2 *may regulate* inflammasome activation by modulating cellular ROS and phagocytosis. To further check the role of inflammasome components in regulating UCP2, we checked UCP2 levels in inflammasome component knockout mice NLRP3^{-/-} and ASC^{-/-}. Our results *reveal* that UCP2 is negatively regulated by NLRP3 and ASC (Fig. 4).

The inflammasome activation is often a two-step process, and the basal expression levels of individual inflammasome components are not necessarily modulated to regulate inflammasome activation. Our present data demonstrated that miR-133a-1 did not alter the basal expression of individual components of the NLRP3 complex (Fig. 1C), *however*, it regulated IL-1 β processing and expression levels of its effector protein Caspase-1 (Fig. 1A). Hence, miR-133a-1 *may regulate* inflammasome activation by disturbing NLRP3 complex formation. Additional studies comparing the co-localization of ASC, NLRP3, and Caspase-1 will address this phenomenon. In conclusion, our results *demonstrate the* novel role of miR-133a-1 in enhancing inflammasome activation and secretion of *Caspase-1* and IL-1 β in human macrophages.

Acknowledgments

This work was funded by the American Heart Association National Scientist Development Grant 09SDG2260957 and National Institutes of Health R01 HL105932 to NK and the Joy McCann Culverhouse Endowment to the Division of Allergy and Immunology. Authors thank Dr. Brenda Flam for critical reading and editing this manuscript.

References

1. Johnson KE, Chikoti L, Chandran B. Herpes simplex virus 1 infection induces activation and subsequent inhibition of the IFI16 and NLRP3 inflammasomes. *J Virol.* 2013; 87:5005–5018. [PubMed: 23427152]
2. Land WG. Transfusion-Related Acute Lung Injury: The Work of DAMPs. *Transfus Med Hemother.* 2013; 40:3–13. [PubMed: 23637644]
3. Rathinam VA, Vanaja SK, Fitzgerald KA. Regulation of inflammasome signaling. *Nat Immunol.* 2012; 13:333–332. [PubMed: 22430786]
4. Kanneganti TD, Ozoren N, Body-Malapel M, Amer A, Park JH, Franchi L, Whitfield J, Barchet W, Colonna M, Vandenabeele P, Bertin J, Coyle A, Grant EP, Akira S, Nunez G. Bacterial RNA and small antiviral compounds activate caspase-1 through cryopyrin/Nalp3. *Nature.* 2006; 440:233–236. [PubMed: 16407888]

5. Mariathasan S, Weiss DS, Newton K, McBride J, O'Rourke K, Roose-Girma M, Lee WP, Weinrauch Y, Monack DM, Dixit VM. Cryopyrin activates the inflammasome in response to toxins and ATP. *Nature*. 2006; 440:228–232. [PubMed: 16407890]
6. Kolliputi N, Galam L, Parthasarathy PT, Tipparaju SM, Lockey RF. NALP-3 inflammasome silencing attenuates ceramide-induced transepithelial permeability. *J Cell Physiol*. 2012; 227:3310–3316. [PubMed: 22169929]
7. Kolliputi N, Shaik RS, Waxman AB. The inflammasome mediates hyperoxia-induced alveolar cell permeability. *J Immunol*. 2010; 184:5819–5826. [PubMed: 20375306]
8. Petrilli V, Papin S, Dostert C, Mayor A, Martinon F, Tschopp J. Activation of the NALP3 inflammasome is triggered by low intracellular potassium concentration. *Cell Death Differ*. 2007; 14:1583–1589. [PubMed: 17599094]
9. Jin C, Flavell RA. Molecular mechanism of NLRP3 inflammasome activation. *J Clin Immunol*. 2010; 30:628–631. [PubMed: 20589420]
10. Dostert C, Petrilli V, Van Bruggen R, Steele C, Mossman BT, Tschopp J. Innate immune activation through Nalp3 inflammasome sensing of asbestos and silica. *Science*. 2008; 320:674–677. [PubMed: 18403674]
11. Bryant C, Fitzgerald KA. Molecular mechanisms involved in inflammasome activation. *Trends Cell Biol*. 2009; 19:455–464. [PubMed: 19716304]
12. Dinarello CA. Interleukin-1 beta, interleukin-18, and the interleukin-1 beta converting enzyme. *Ann N Y Acad Sci*. 1998; 856:1–11. [PubMed: 9917859]
13. Simsa-Maziel S, Monsonago-Ornan E. Interleukin-1beta promotes proliferation and inhibits differentiation of chondrocytes through a mechanism involving down-regulation of FGFR-3 and p21. *Endocrinology*. 2012; 153:2296–2310. [PubMed: 22492305]
14. Dowds TA, Masumoto J, Zhu L, Inohara N, Nunez G. Cryopyrin-induced interleukin 1beta secretion in monocytic cells: enhanced activity of disease-associated mutants and requirement for ASC. *J Biol Chem*. 2004; 279:21924–21928. [PubMed: 15020601]
15. Agostini L, Martinon F, Burns K, McDermott MF, Hawkins PN, Tschopp J. NALP3 forms an IL-1beta-processing inflammasome with increased activity in Muckle-Wells autoinflammatory disorder. *Immunity*. 2004; 20:319–325. [PubMed: 15030775]
16. Church LD, Cook GP, McDermott MF. Primer: inflammasomes and interleukin 1beta in inflammatory disorders. *Nat Clin Pract Rheumatol*. 2008; 4:34–42. [PubMed: 18172447]
17. Martinon F, Petrilli V, Mayor A, Tardivel A, Tschopp J. Gout-associated uric acid crystals activate the NALP3 inflammasome. *Nature*. 2006; 440:237–241. [PubMed: 16407889]
18. Cassel SL, Eisenbarth SC, Iyer SS, Sadler JJ, Colegio OR, Tephly LA, Carter AB, Rothman PB, Flavell RA, Sutterwala FS. The Nalp3 inflammasome is essential for the development of silicosis. *Proc Natl Acad Sci U S A*. 2008; 105:9035–9040. [PubMed: 18577586]
19. Tan MS, Yu JT, Jiang T, Zhu XC, Tan L. The NLRP3 Inflammasome in Alzheimer's Disease. *Mol Neurobiol*. 2013
20. Perry MM, Moschos SA, Williams AE, Shepherd NJ, Larner-Svensson HM, Lindsay MA. Rapid changes in microRNA-146a expression negatively regulate the IL-1beta-induced inflammatory response in human lung alveolar epithelial cells. *J Immunol*. 2008; 180:5689–5698. [PubMed: 18390754]
21. Cogswell JP, Ward J, Taylor IA, Waters M, Shi Y, Cannon B, Kelnar K, Kemppainen J, Brown D, Chen C, Prinjha RK, Richardson JC, Saunders AM, Roses AD, Richards CA. Identification of miRNA changes in Alzheimer's disease brain and CSF yields putative biomarkers and insights into disease pathways. *J Alzheimers Dis*. 2008; 14:27–41. [PubMed: 18525125]
22. Ameres SL, Zamore PD. Diversifying microRNA sequence and function. *Nat Rev Mol Cell Biol*. 2013; 14:475–488. [PubMed: 23800994]
23. Sun K, Lai EC. Adult-specific functions of animal microRNAs. *Nat Rev Genet*. 2013; 14:535–548. [PubMed: 23817310]
24. Yoshino H, Seki N, Itesako T, Chiyomaru T, Nakagawa M, Enokida H. Aberrant expression of microRNAs in bladder cancer. *Nat Rev Urol*. 2013; 10:396–404. [PubMed: 23712207]
25. Szabo G, Bala S. MicroRNAs in liver disease. *Nat Rev Gastroenterol Hepatol*. 2013

26. Allegra A, Alonci A, Campo S, Penna G, Petrunaro A, Gerace D, Musolino C. Circulating microRNAs: new biomarkers in diagnosis, prognosis and treatment of cancer (review). *Int J Oncol.* 2012; 41:1897–1912. [PubMed: 23026890]
27. Lagos-Quintana M, Rauhut R, Yalcin A, Meyer J, Lendeckel W, Tuschl T. Identification of tissue-specific microRNAs from mouse. *Curr Biol.* 2002; 12:735–739. [PubMed: 12007417]
28. Liu N, Bezprozvannaya S, Williams AH, Qi X, Richardson JA, Bassel-Duby R, Olson EN. microRNA-133a regulates cardiomyocyte proliferation and suppresses smooth muscle gene expression in the heart. *Genes Dev.* 2008; 22:3242–3254. [PubMed: 19015276]
29. Chen X, Wang K, Chen J, Guo J, Yin Y, Cai X, Guo X, Wang G, Yang R, Zhu L, Zhang Y, Wang J, Xiang Y, Weng C, Zen K, Zhang J, Zhang CY. In vitro evidence suggests that miR-133a-mediated regulation of uncoupling protein 2 (UCP2) is an indispensable step in myogenic differentiation. *J Biol Chem.* 2009; 284:5362–5369. [PubMed: 19073597]
30. Krauss S, Zhang CY, Lowell BB. The mitochondrial uncoupling-protein homologues. *Nat Rev Mol Cell Biol.* 2005; 6:248–261. [PubMed: 15738989]
31. Basu Ball W, Kar S, Mukherjee M, Chande AG, Mukhopadhyaya R, Das PK. Uncoupling protein 2 negatively regulates mitochondrial reactive oxygen species generation and induces phosphatase-mediated anti-inflammatory response in experimental visceral leishmaniasis. *J Immunol.* 2011; 187:1322–1332. [PubMed: 21705615]
32. Echtay KS. Mitochondrial uncoupling proteins--what is their physiological role? *Free Radic Biol Med.* 2007; 43:1351–1371. [PubMed: 17936181]
33. Park D, Han CZ, Elliott MR, Kinchen JM, Trampont PC, Das S, Collins S, Lysiak JJ, Hoehn KL, Ravichandran KS. Continued clearance of apoptotic cells critically depends on the phagocyte Ucp2 protein. *Nature.* 2011; 477:220–224. [PubMed: 21857682]
34. Taxman DJ, Lei Y, Zhang S, Holley-Guthrie E, Offenbacher S, Ting JP. ASC-dependent RIP2 kinase regulates reduced PGE2 production in chronic periodontitis. *J Dent Res.* 2012; 91:877–882. [PubMed: 22828789]
35. Ceppi M, Pereira PM, Dunand-Sauthier I, Barras E, Reith W, Santos MA, Pierre P. MicroRNA-155 modulates the interleukin-1 signaling pathway in activated human monocyte-derived dendritic cells. *Proc Natl Acad Sci U S A.* 2009; 106:2735–2740. [PubMed: 19193853]

Abbreviations

miR	micro RNA
UCP2	Uncoupling protein 2
siRNA	Silencing RNA
IL-1β	Interleukin 1 beta
THP1	human monocyte cell line

Highlights

1. We identified a novel mechanism for inflammasome activation by miR-133a-1
2. Overexpression of miR-133a-1 lead to activation of inflammasome by suppression of UCP2.
3. Suppression of UCP2 by siRNA enhanced the activity.

Figure 1. miR-133-a1 enhances *H₂O₂*-mediated inflammasome activation and IL-1β production in THP1 cells and suppresses its target UCP2

Differentiated THP1 cells were transfected with miR-133-a1 or *vector* control (miR control). At 48 hour post-transfection, inflammasomes were stimulated by adding H₂O₂ at 5mM for four hours. (A) Expression levels of *Caspase-1* p10 and cleaved IL-1β p17 in the supernatants were measured by *Western* blot analysis. (B) Transfection efficiency of micro RNA plasmids (*vector* control and miR-133-a1) was confirmed by examining the localization of IRES-GFP under bright field and fluorescence microscopy. (C) Expression levels of NLRP3, ASC, pro-IL1β, *pro-Caspase-1*, and GAPDH in lysates were observed under immunoblot. (D) Differentiated THP1 cells were transfected with miR-133-a1 or *vector* control (miR control) plasmids. Expression levels of UCP2 at 48 hour post-transfection were analyzed by *Western* blot.

Figure 2. UCP2 suppression augments inflammasome activation

siRNA specific to UCP2 (siRNA1, 2, 3), Cy3-labeled siRNA control, and scrambled siRNA control were transfected into differentiated THP1 cells. (A) At 48 hour post-transfection, expression of Cy3 was examined under fluorescence microscope. siRNA (in red) localization with nucleus, in blue, can be seen in the transfected cells. The left image shows THP1 cells exposed to siRNA without transfection reagent. (B) Western blot analysis of UCP2 was utilized to confirm the silencing of UCP2 in transfected THP1 cells. (C) After transfection with siRNAs and controls, THP1 cells were stimulated with 5mM H₂O₂ and secretion of mature IL-β p17 and Caspase-1 p10 in the supernatants was compared by Western blot analysis.

Figure 3. UCP2 overexpression suppressed IL-1 β and Caspase-1p10 production in THP1 cells pCMV-UCP2-GFP or pCMV-GFP (vector) was transfected into differentiated THP1 cells seeded at 0.5 million cells/well in a six-well plate with coverslip. Untransfected cells acted as controls. (A) 48 *hour* post-transfection, cells were fixed in 4% paraformaldehyde, and the expression of GFP was examined under fluorescence microscope. (B) Expression of UCP2 in cell lysates transfected with UCP2-GFP or *vector* was analyzed by *Western* blot to confirm transfection. (C) After post-transfection with plasmids overexpressing UCP2 and controls, THP1 cells were stimulated with 5mM H₂O₂ and secretion of mature IL- β p17 and Caspase-1 p10 in the supernatants was compared by *Western* blot analysis.

Figure 4. UCP2 is upregulated in bone marrow macrophages of NLRP3 and ASC deficient mice
 Bone marrow macrophages from wild type, NLRP3^{-/-}, and ASC^{-/-} knockout mice were isolated, inflammasome was stimulated using H₂O₂, and the expression levels of UCP2, NLRP3, and GAPDH were compared by *Western* blot analysis.