Published in final edited form as:

J Control Release. 2013 December 28; 172(3): . doi:10.1016/j.jconrel.2013.09.005.

HPMA copolymer-based combination therapy toxic to both prostate cancer stem/progenitor cells and differentiated cells induces durable anti-tumor effects

Yan Zhou¹, Jiyuan Yang¹, Johng S. Rhim², and Jindřich Kopeček^{1,3,*}

¹Department of Pharmaceutics and Pharmaceutical Chemistry/CCCD, University of Utah, Salt Lake City, UT 84112, USA

²Uniformed Services University of the Health Sciences, Bethesda, MD 20814

³Department of Bioengineering, University of Utah, Salt Lake City, UT 84112, USA

Abstract

Current treatments for prostate cancer are still not satisfactory, often resulting in tumor regrowth and metastasis. One of the main reasons for the ineffective anti-prostate cancer treatments is the failure to deplete cancer stem-like cells (CSCs) - a subset of cancer cells with enhanced tumorigenic capacity. Thus, combination of agents against both CSCs and bulk tumor cells may offer better therapeutic benefits. Several molecules with anti-cancer stem/progenitor cell activities have been under preclinical evaluations. However, their low solubility and nonspecific toxicity limit their clinical translation. Herein, we designed a combination macromolecular therapy containing two drug conjugates: HPMA copolymer-cyclopamine conjugate (P-CYP) preferentially toxic to cancer stem/progenitor cells, and HPMA copolymer-docetaxel conjugate (P-DTX) effective in debulking the tumor mass. Both conjugates were synthesized using RAFT (reversible addition-fragmentation chain transfer) polymerization resulting in narrow molecular weight distribution. The killing effect of the two conjugates against bulk tumor cells and CSCs were evaluated in vitro and in vivo. In PC-3 or RC-92a/hTERT prostate cancer cells, P-CYP preferentially kills and impairs the function of CD133+ prostate cancer stem/progenitor cells; P-DTX was able to kill bulk tumor cells instead of CSCs. In PC-3 xenograft mice model, combination of P-DTX and P-CYP showed the most effective and persistent tumor growth inhibitory effect. In addition, residual tumors contained less CD133+ cancer cells following combination or P-CYP treatments, indicating selective killing of cancer cells with stem/progenitor cell properties.

Keywords

N-(2-hydroxypropyl)methacrylamide (HPMA); Combination therapy; Cancer stem/progenitor cells; Cyclopamine; Docetaxel

© 2013 Elsevier B.V. All rights reserved.

*Corresponding author: Jindřich Kopeček, Center for Controlled Chemical Delivery, 20 S 2030 E, BPRB 205B, University of Utah, Salt Lake City, UT 84112-9452, USA. Phone: (801) 581-7211; Fax: (801) 581-7848. jindrich.kopecek@utah.edu (J. Kopeček).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

Despite advances in anti-prostate cancer therapies, patients with either localized or progressive disease frequently suffer from relapse, progression and metastasis [1, 2]. This phenomenon can be well explained by an important dimension of tumor cell heterogeneity; tumor cells reside in several levels of differentiation statuses, including cancer stem cells (CSCs) and progenitor cells with infinite or certain degree of self-renew and differentiation abilities, and differentiated cancer cells depleted of these abilities [3–5]. The failure of traditional anti-cancer therapies is largely attributed to the existence of CSCs that are drug-resistant and survive after treatment, despite the efficient killing of differentiated tumor cells [6–8]. Indeed, these drug-resistant cancer cells with enhanced tumor-reinitiating capacity were found to survive and even expand following standard chemo- or radio-therapy [5, 9, 10]. Thus, therapies targeting both proliferating bulk tumor cells and self-renewable cancer stem/progenitor cells that give rise to differentiated progenies may offer superior therapeutic benefit by preventing tumor growth, regrowth and metastasis [11].

CSCs have been identified in several hematopoietic and solid tumors [12, 13]. In prostate cancer, the cell compartment enriched with prostate cancer stem/progenitor cells has been identified by the expression of biomarker profiles similar to normal prostate stem cells, or by growing cells under anchorage-independent condition [5, 14, 15]. Importantly, these cells exhibit much more enhanced stem cell properties and in vivo tumorigenicity than the bulk tumor cell counterparts [5, 14–17].

Noticeably, critical signaling pathways governing the stem cell self-renewal and differentiation were found to be active in the putative prostate CSCs [18, 19]. Among these, the hedgehog (Hh) pathway is not only involved in prostate cancer growth and progression, but also is critical in promoting prostate CSC self-renewal and growth [19–21]. Activation of a central Hh pathway element GLI1 can transform normal prostate epithelial cells into tumorigenic cancer cells; knockdown of Hh pathway by small molecule inhibitor cyclopamine down-regulates CSC self-renewal gene expression and leads to long-term prostate cancer regression [19]. However, cyclopamine is poorly water soluble and possesses unfavorable in vivo behavior and toxicity [22]. To be therapeutically effective, cyclopamine must be administered frequently at high doses via i.p. or s.c. route.

HPMA copolymers as drug carriers offer superior solubility, enhanced tumor accumulation and reduced nonspecific toxicity of the small molecule drugs in the application for cancer treatments [23, 24]. We have previously showed that attaching cyclopamine to the water-soluble and biocompatible HPMA copolymer retained its anti-prostate CSC activity in vitro [25]. HPMA copolymer-cyclopamine conjugate (P-CYP), although not potent in inhibiting bulk tumor growth, preferentially decreased the CD133+ prostate cancer stem/progenitor cell population. On the other hand, docetaxel, a traditional first line chemotherapeutic agent for prostate cancer, is effective in debulking the tumor mass; similarly, HPMA copolymer-docetaxel conjugate (P-DTX) showed growth inhibition in prostate cancer xenografts [26, 27]. Therefore, combination macromolecular therapeutics by intravenous administration of both P-DTX and P-CYP may effectively kill both bulk tumor cells and cancer stem/progenitor cells [28].

In this study, we evaluated the differential effects of P-CYP and P-DTX on cancer stem and differentiated cells in PC-3 and RC-92a/hTERT prostate cancer cells. Moreover, we evaluated the long-term tumor inhibition effects of single agent and combination of P-CYP and P-DTX in nude mice bearing PC-3 xenografts. Our results demonstrate that the combination macromolecular therapy induces more effective and durable tumor inhibitory effect and anti-CSC effect than either P-CYP or P-DTX single treatment.

2. Materials and methods

2.1. Materials

Cyclopamine (free base) was purchased from LC Laboratories (Woburn, MA). Docetaxel was purchased from AK Scientific (Union City, CA). 2,2'-azobis[2-(2-imidazolin-2-yl)propane]dihydrochloride (VA-044) and 2,2'-azobis(2,4-dimethyl valeronitrile) (V-65) were from Wako Chemicals (Richmond, VA). Papain and cathepsin B were purchased from Sigma (St. Louis, MO). 4-cyanopentanoic acid dithiobenzoate [29], *N*-(2-hydroxypropyl)methacrylamide (HPMA) [30], *N*-methacryloylglycylphenylalanylleucylglycine (MA-GFLG-OH) [31], and *N*-methacryloylglycylphenylalanylleucylglycyl thiazolidine-2-thione (MA-GFLG-TT) [32] were synthesized as previously described.

2.2. Synthesis and characterization of HPMA copolymer-cyclopamine conjugate (P-CYP)

The synthesis of P-GFLG-CYP was similar as previously described [25]. Firstly, HPMA copolymer precursor with enzymatically degradable side chains containing thiazolidine-2-thione (TT) reactive groups (P-GFLG-TT) was prepared; then CYP was conjugated to the copolymer side chains to yield P-CYP. Briefly, for the synthesis of P-GFLG-TT, reversible addition-fragmentation chain-transfer (RAFT) copolymerization of HPMA (1288.8 mg, 9 mmol, 90 mol%) and MA-GFLG-TT (561.7 mg, 1 mmol, 10 mol%) were carried out in a degassed ampoule in the presence of VA 044 (0.0143 mmol) as initiator and 4-cyanopentanoic acid dithiobenzoate (3.6×10^{-3} mmol) as chain transfer agent. The copolymerization was performed in methanol at 40°C for 20 h. The reaction mixture was precipitated in acetone/ether and purified by dissolution-precipitation in methanol-acetone to yield the product as yellow solid (1005 mg). This copolymer was further end-group modified to remove the dithiobenzoate group from the chain ends as previously described [25]. Briefly, P-GFLG-TT (1005 mg, Mn 37.4 kDa, 26.7 μ mol) and V65 (40x excess, 265 mg, 1068 μ mol) were reacted in methanol at 50°C for 4 h. The end-modified copolymer was purified by dissolution-precipitation in methanol-acetone twice and dried under vacuum (approximately 800 mg). The TT content in the copolymer (4.46×10^{-4} mmol/mg, 7.8 mol%) was determined by UV spectrophotometry in methanol ($\epsilon_{305} = 10,800 \text{ L mol}^{-1} \text{ cm}^{-1}$). The molecular weight and molecular weight distribution (Mw 39.9 kDa, PDI 1.07) was determined by size-exclusion chromatography (SEC) (Superose 6 HR10/30 column, Pharmacia) on an AKTA FPLC system with miniDAWN and OptilabEX detectors using acetate/30% acetonitrile as mobile phase.

HPMA copolymer-cyclopamine conjugate was obtained by the reaction of end-modified P-GFLG-TT (230 mg) with cyclopamine (Mw 411.6, 3x excess of TT mol content, 127 mg, 307.8 μ mol) in pyridine at 50°C for 28 h. After reaction, the unreacted TT groups were removed by adding excess of 1-amino-2-propanol. The conjugate was precipitated into ether/acetone and purified by dissolution-precipitation in methanol-acetone. The dried product was dissolved in deionized water, purified through PD-10 column and freeze-dried. Yield: 230 mg. To estimate the CYP content, the conjugate was exposed to excess of papain for cleavage of the GFLG linker. A calibration curve of the AUC of the free CYP peak eluted on high-performance liquid chromatography (HPLC) was constructed. The amount of CYP released from the conjugate was calculated based on its AUC (6.6 wt%).

2.3. Synthesis of polymerizable derivative of docetaxel (MA-GFLG-DTX)

MA-GFLG-DTX was synthesized as previously described [26]. Briefly, MA-GFLG-OH (248 mg, 0.54 mmol), DTX (396mg, 0.49 mmol) and DMAP (4-dimethylaminopyridine, 60 mg, 0.49 mmol) were dissolved in 2 mL *N,N*-dimethylformamide (DMF) and cooled to 0°C. DMF solution of DCC (*N,N*'-dicyclohexylcarbodiimide, 131 mg, 0.64 mmol) was added

dropwise. The reaction mixture was stirred at 4°C for 24 h. After reaction, dicyclohexylurea (DCU) was removed by filtration. The filtrate was washed with $\text{KHSO}_4\text{NaHCO}_3$ and NaCl solutions sequentially and dried over MgSO_4 . The yield was 450 mg (73.4%). The crude product was dissolved in ethyl acetate and purified on silica gel column using ethyl acetate:actone 8:1 as the running buffer. The mass of the purified monomer (158 mg) was confirmed by MALDI-TOF mass spectrometry ($[\text{M}+\text{Na}^+ 1272.62]$).

2.4. Synthesis of HPMA copolymer-docetaxel conjugate (P-DTX)

P-DTX was synthesized by RAFT copolymerization of HPMA (277.8 mg, 1.94 mmol, 97 mol%) and MA-GFLG-DTX (75 mg, 0.06 mmol, 3 mol%) using V65 (8×10^{-4} mmol) as initiator and 4-cyanopentanoic acid dithiobenzoate (3.2×10^{-3} mmol) as chain transfer agent. The mixture in methanol was bubbled with nitrogen for 30 min, sealed and polymerized at 50°C for 35 h. After reaction, it was precipitated in acetone/ether, dissolved in methanol and re-precipitated into acetone to yield white solid 178 mg (50.5%). The product was end-modified in the presence of 40× molar excess of V65 (29.3 mg) in methanol at 50°C for 4 h. The yield of the final end-modified P-DTX (Mw 38.6 kDa, PDI 1.06) was 165 mg. DTX content in P-DTX was determined by enzymatic cleavage of GFLG in the presence of excess of papain [26]. The cleaved sample was run on HPLC and the amount of DTX was calculated based on the AUC of the DTX elution peak and the calibration curve of free DTX with known concentrations.

2.5. Cell culture

PC-3 prostate cancer cell line was purchased from American Type Culture Collection (ATCC, Manassas, VA). The hTERT-immortalized primary human prostate tumor-derived cell line (RC-92a/hTERT) was from Dr. Rhim's group [15, 33]. PC-3 and RC-92a/hTERT cells were grown and maintained at 37°C, 5% CO_2 in a keratinocyte serum-free medium (K-SFM) supplemented with bovine pituitary extract (BPE) and recombinant epidermal growth factor (EGF) (Life Technologies, Gaithersburg, MD). The cells were routinely trypsinized and subcultured when reaching 90% confluency.

2.6. Prostatosphere culture

Single cell suspensions of RC-92a/hTERT cells or PC-3 cells were prepared in K-SFM supplemented with 10 ng/mL EGF, 10 ng/mL basic fibroblast growth factor and B27 (all from Invitrogen, Carlsbad, CA). PC-3 cells were plated in 96-well ultra low attachment plate at the density of 500 cells/well. Cells were exposed to 100 μL drug or conjugate containing sphere growth medium at the final concentrations as follows: P-CYP (CYP equivalent 5 μM , 10 μM), CYP (5 μM , 10 μM), P-DTX (DTX equivalent 2 nM, 10 nM), or DTX (2 nM, 10 nM) for three days, followed by addition of 200 μL fresh sphere growth medium. RC-92a/hTERT cells were plated at the density of 500 cells/well in 6-well ultra low attachment plate. Cells were exposed to 1 mL solution of P-CYP (CYP equivalent 5 μM , 10 μM), CYP (5 μM , 10 μM), P-DTX (DTX equivalent 1 nM, 2 nM) or DTX (1 nM, 2 nM) for three days, followed by addition of 2 mL sphere growth medium. The concentrations of DTX or DTX equivalent in the conjugate that inhibit the cell growth of each cell line by around 20% (moderate inhibition) and 50% (significant inhibition) were chosen. After eight-day culture, the numbers of spheres formed with diameter between 50 to 200 μm were counted under the microscope [34].

2.7. Cell surface marker expression by flow cytometry and fluorescent activated cell sorting (FACS)

To determine CD133 expression in PC-3 cells with or without treatments, cells were seeded in 6-well plates at the density of 4×10^4 cells/well and grew for 24 h. Then cells were

exposed to different concentrations of drug/conjugate-containing medium for 72 h, with fresh medium and vehicle (medium containing <0.5% vol. DMSO) treated groups as control. After treatments, cells were harvested, washed and resuspended in cold staining buffer (phosphate buffered saline pH 7.2, 0.5% bovine serum albumin, 2 mM EDTA) at the density of up to 1×10^7 /mL. Cells were pre-blocked with 3% bovine serum albumin and stained with mouse anti-human CD133/2 (293C3; allophycocyanin (APC) labeled; Miltenyi Biotec, Auburn, CA) in the dark at 4°C for 20 min. After staining, cells were thoroughly washed twice. Hoechst 33258 was added 10 min before analysis to exclude dead cells. Samples were run on FACScan flow cytometer (Becton Dickinson) and data were analyzed by FlowJo (Tree Star). The data were collected from three independent experiments. For each batch of experiment, the CD133 expression level in untreated control cells were set as 100%; CD133 expression following different treatments were normalized to the untreated control.

For the isolation of CD133+ and CD133-RC-92a/hTERT cells, the cells were stained using the same procedure as described above. Live cells were gated and sorted using FACS Aria cell sorter (Becton Dickinson) as previously described [25].

2.8. Cytotoxicity assay

Cytotoxic effects of free and HPMa copolymer-conjugated cyclopamine and docetaxel at different drug concentrations were determined using Cell Counting Kit-8 (CCK-8, Dojindo, Japan). Briefly, PC-3 cells were seeded in 96-well plates in triplicate, grew for 24 h and were incubated with CYP, DTX, P-CYP and P-DTX, respectively for another 72 h. After treatments, cells were washed and incubated with CCK-8 solution (15× dilution in fresh growth medium) at 37°C, 5% CO₂ for 1 h. The absorbance in each well was measured by a microplate reader at 450 nm (630 nm as reference). Untreated cells were set as 100% viable.

2.9. Apoptosis assay

Annexin V assay was performed to evaluate the induction of apoptosis by P-CYP in sorted CD133+ and CD133-RC-92a/hTERT cells. Immediately after sorting, CD133+ cells, CD133- cells and unsorted cells were seeded in 6-well plates at the same density and treated with drug/conjugate for 72 h. Following treatments, cells were harvested and stained with Annexin V kit according to the manufacturer's protocol (Oncogene Research Products, Boston, MA). Vehicle treated cells served as controls.

2.10. Cell growth inhibition in vitro

For short-term evaluation, PC-3 cells were seeded in 96-well plates at the density of 5000/well and allowed to grow for 24 h. Cells were then exposed to single conjugate or drug (P-CYP at 10 μM CYP equivalent, P-DTX at 2 nM DTX equivalent, CYP 10 μM, or DTX 2 nM), as well as the simultaneous combination of P-CYP (CYP equivalent 10 μM) + P-DTX (DTX equivalent 2 nM), and combination of CYP 10 μM + DTX 2 nM for 72 h. Immediately after treatments, CCK-8 assay was performed. For the evaluation of long-term cell growth inhibition, cells were exposed to DTX or P-DTX (10 nM drug equivalent) for 24 h and then CYP or P-CYP (10 μM) for another 48 h. Vehicle-treated cells and cells exposed to single drug/conjugate were controls. After 72 h, cells were washed and incubated in fresh growth medium continued for up to two weeks. Media were replaced every three days. CCK-8 assay was performed on 0, 3, 7, 11, and 14 days post treatment.

2.11. In vivo anti-tumor activity

The anti-tumor activities of P-DTX and P-CYP, individually and in combination, were evaluated in nude mice bearing PC-3 xenografts. Animal studies were carried out according to the University of Utah IACUC guidelines under approved protocols. Viable PC-3 cells

(2.5×10^6) suspended in 200 μL phosphate buffered saline were inoculated subcutaneously into the right flank region of each 6–8 week male nude mouse. Two-three weeks after tumor cell inoculation when the tumor sizes reached an average of 50 mm^3 treatments were initiated. P-DTX and P-CYP at the doses indicated below were both dissolved in sterile water:saline (1:1) solution (total volume about 250 μL). Mice in P-DTX treated group received a single intravenous injection of P-DTX at the dose of DTX equivalent 10 mg/kg on day 0 ($n=4$); mice in P-CYP treated group received a total of six doses of P-CYP (CYP equivalent 40 mg/kg) twice per week via intravenous injections ($n=4$); For combination treatment group, mice received a single dose of P-DTX (DTX equivalent 10 mg/kg) on day 0 and six doses of P-CYP (CYP equivalent 40 mg/kg) twice per week starting on day 1 ($n=5$). Saline treated group served as control ($n=4$). Tumor growth and body weight were monitored every other day from day 0 for as long as 100 days, unless the body weight loss exceeded 20% due to tumor growth or the tumor weight exceeded 10% of the body weight. Tumor volumes were calculated as $V = \text{width}^2 \times \text{length} / 2$. At the end of experiments, mice were sacrificed. Tumors and major organs (liver, spleen, kidney and intestine) were harvested. Tissues were fixed in 10% formalin, paraffin embedded, sectioned and stained with hematoxylin and eosin (H&E). The tissue sections were examined under the microscope.

2.12. Characterizations of residual tumor cells

Immediately after sacrificing the mice, tumor single cell suspensions were prepared [35]. Briefly, the tumor mass was chopped into small pieces using #10 surgical blade and digested in collagenase IV solution 3 mg/mL for 30 min, followed by filtration through 70 μm mesh. Dead cells and red blood cells were discarded from the cell suspension by histopaque-1077 gradient centrifugation. For CD133 expression analysis by flow cytometry, tumor cells were sampled randomly from the harvested single cell suspensions and were stained with APC-conjugated anti-CD133/2 (293C3) monoclonal antibody and Hoechst 33258 as described in 2.7. For sphere forming assay, dissociated tumor cells from each mouse were seeded in 4–5 wells of a 96-well plate at the density of 500 cells/well and grew for 10 days before counting the number of spheres formed.

2.13. Statistical analysis

One-way analysis of variance coupled with bonferroni post-hoc analysis were used to compare three or more groups, with p values of <0.05 indicating statistically significant differences.

3. Results and discussion

3.1. Synthesis and characterization of P-CYP and P-DTX

The synthesis of both P-CYP and P-DTX employed controlled RAFT polymerization chemistry (Fig. 1). For P-CYP, HPMA copolymer precursor containing reactive functional groups in the side chains (P-GFLG-TT) was prepared first, followed by the conjugation of free CYP. The CYP content in the P-CYP conjugate was 6.6 wt%. Since the sterical environment and electron density of the nitrogen atom affect the rate of TT aminolysis, the reactivity of the secondary amine in CYP with TT group may not be as effective as that of the primary amines [36]. Herein, CYP was added only at the last step of P-CYP synthesis, thus saving the amount of CYP consumed. P-DTX was synthesized by direct polymerization of HPMA with polymerizable monomer containing DTX (MA-GFLG-DTX). The DTX content in the P-DTX could be controlled by adjusting the feed ratio of HPMA and MA-GFLG-DTX; this also permitted to control the solubility of P-DTX in aqueous environment. Both synthetic routes yielded HPMA copolymer-drug conjugates with well-controlled molecular weight and narrow molecular weight distribution (Table 1). In this combination

therapy design, the two different drugs were conjugated to two HPMA copolymer backbones respectively; this allows the easy control of drug incorporation. In addition, both the simultaneous and sequential administration of two conjugates can be achieved, in order to investigate the combinational effects of different dosing regimens.

3.2. In vitro evaluation of stem cell-like properties following drug/conjugate treatments

Prostasphere forming ability under the anchorage-independent culture condition has been utilized to characterize stem cell self-renewal behavior of prostate cancer cells [5, 34]. We observed that there was a significant reduction in the number of spheres formed from both RC-92a/hTERT and PC-3 cells, when exposed to 10 μ M CYP or CYP equivalent (in P-CYP) (Fig. 2A,B). In contrast, DTX and P-DTX treatments did not impair the prostasphere forming capacity (Fig. 2A,B), when PC-3 or RC-92a/hTERT cells were exposed to DTX or P-DTX at the concentrations that inhibit the cell growth by about 20% (DTX equivalent concentrations of 2 nM for PC-3 and 1 nM for RC-92a/hTERT cells) and 50% (DTX equivalent concentrations of 10 nM for PC-3 and 2 nM for RC-92a/hTERT cells). In particular, our and other's studies support that CD133+ prostate cancer stem/progenitor cell subpopulations in both cell lines possess much more enhanced sphere forming ability than CD133-cells [25, 37, 38]. Taken together, our results indicate that CYP or P-CYP, in contrast to DTX or P-DTX, inhibits the self-renewal ability of prostate CSCs.

We next investigated whether P-CYP inhibits CSC self-renewal ability via reducing the CD133+ subpopulation within the whole cell population. Following P-CYP (CYP equivalent 10 μ M) or CYP treatments, the proportion of CD133+ PC-3 cells significantly decreased, while the whole cell population proliferation remained above 95% relative to controls for the P-CYP group and above 90% for the CYP group (Fig. 3). In contrast, DTX or P-DTX treatments suppressed whole cell population proliferation significantly at DTX equivalent concentration of 10 nM (around 50% compared to controls), but the percentages of CD133+ cells still remained similar or slightly increased. These results agree with those obtained in sphere forming assay: P-CYP or CYP more specifically decreased the population of CD133+ cancer stem/progenitor cells, leading to fewer sphere formation; instead, P-DTX or DTX affected mainly the CD133-bulk tumor cells, thus could not inhibit the sphere formation. Based on the differential therapeutic effects of P-CYP and P-DTX towards prostate CSCs and non-CSCs respectively, we speculate that combination of P-CYP and P-DTX may yield better anti-tumor outcome than either of the single agents.

In the above experiments, DTX and P-DTX showed similar levels of cell growth inhibition, which could be due to the gradual hydrolytic release of free DTX from P-DTX over the 72 h incubation period. For P-CYP, the amide bond between GFLG linker and cycloamine was stable in the growth medium over 72 h incubation. P-CYP and CYP inhibit the CD133+ cells at concentrations of the same magnitude; this could be attributed to the mechanism of action of CYP. CYP inhibits the Hh pathway by directly binding to a cell surface receptor SMO (smoothened), which is an important regulatory element along the Hh pathway [39]. It was indicated that the steric hindrance of the HPMA copolymer did not significantly affect the interaction of (bound) CYP with SMO.

The main mechanism of cycloamine's inhibitory effect on CSCs is through the inhibition of Hh pathway [19, 39]. Cycloamine inhibits the activity of SMO receptor, thus down-regulating the GLI1 transcriptional factor and its downstream gene products, eventually inducing apoptosis of the affected cancer cells [39]. We tested the mRNA levels of GLI1 in prostate cancer cells with and without P-CYP (or CYP) treatments at the effective dose used in our experiments. GLI1 mRNA expression was significantly lower following both CYP and P-CYP treatments relative to the vehicle control (Fig. 4). The results indicate that CYP or P-CYP exert their therapeutic effect, if not all, at least partially through down-regulating

the Hh pathway activity. Other mechanisms, beyond the scope of this study, may be operative. For example, CYP could down-regulate the expression of MDR receptors on cancer cells, potentially sensitizing drug-resistant CSCs to traditional anti-cancer therapies [40]; in some cancer models, CYP could modulate the tumor stromal cells [41].

To evaluate the cell killing effect by P-CYP toward CD133+ prostate cancer stem/progenitor cells and CD133-cells, the Annexin V assay was performed on CD133+ and CD133-RC-92a/hTERT cells immediately following isolation by FACS. P-CYP treatment induced a significantly higher level of apoptosis in CD133+ cells (23.8% normalized to untreated CD133+ cells) than in CD133-cells (1.1% normalized to untreated CD133-cells) and the whole cell population (2.8% normalized to untreated cells) (Fig. 5). Similar trends were observed in CYP treated cells. It may suggest that the impairment of the sphere forming ability and the reduction in the proportion of CD133+ cells were due to the killing of CD133+ cells by CYP or P-CYP.

3.3. Growth inhibition of prostate cancer cells in vitro by combination therapy

The inhibition of prostate cancer cell growth by P-CYP and P-DTX, individually and in combination, was evaluated immediately after treatments and in long-term. Consistent with data in Fig. 3, after 72 h treatment, CYP and P-CYP (CYP equivalent 10 μ M) resulted in little or very low level of growth inhibition (Fig. 6). DTX and P-DTX (DTX equivalent 2 nM) inhibit prostate cancer cell growth more effectively (around 20% inhibition) (Fig. 6). For cells exposed to the P-CYP + P-DTX combination, the level of growth inhibition (around 24% of control) was slightly greater than that induced by either of the single agent (around less than 1% for P-CYP and 20% for P-DTX). However, this difference was not statistically significant. For CYP + DTX combination, a more significant cell growth inhibition (around 43%) was observed compared to single CYP (around 10%) or DTX (around 16%) treatment (Fig. 6). This observation agrees with previous reports on free CYP and DTX combination using other prostate cancer cell lines [42]. CYP could also be cytotoxic to bulk prostate cancer cells at high doses [19, 42]. Nevertheless, based on our results in both RC-92a/hTERT cells and PC-3 cells, CYP and P-CYP is more effective in killing the relatively rare CD133+ prostate CSCs.

When studying the long-term cytotoxic effects, the combination of P-CYP and P-DTX showed more definite therapeutic benefit over the single P-CYP or P-DTX treatment (Fig. 7). Immediately after 72 h sequential treatments, the differences in cell growth inhibition between DTX and DTX+CYP group or P-DTX and P-DTX+P-CYP group were indistinguishable. However, starting from 10 days after treatments, the conjugate or drug combination group showed significantly stronger growth inhibition than P-DTX or DTX alone group respectively (Fig. 7). When comparing the recovery rate of residual PC-3 cells, the cells exposed to only P-DTX or DTX treatments grew back faster than those subjected to the combination treatments.

3.4 In vivo anti-tumor efficacy

One important end point for evaluating in vivo anti-tumor efficacy is the tumor growth inhibitory effect in long-term. In all three treatment groups, P-CYP, P-DTX and combination of P-DTX + P-CYP, the tumors grew slower than the tumors in mice subjected to saline treatment. At the end of the treatment on day 22, the average tumor volumes in all three-treatment groups were significantly smaller than that in saline group (Fig. 8A). However, within 40 days of observation, there was no statistically significant difference in tumor volumes among the P-CYP, P-DTX and combination groups. When monitoring the tumor growth for longer periods, the difference in tumor volumes between P-DTX (or P-CYP group) and combination group became progressively larger, reaching statistical

significance starting from day 42 ($p < 0.05$) (Fig. 8A). The tumor growth rate in the combination group (the slope of the growth curve) did not increase, or even started to decrease after the stop of the treatment; on the contrary, the tumors in P-DTX treated group started to regrow faster. In addition, mice in control, P-DTX and P-CYP groups were kept up to 26 days, 58 days and 64 days, respectively from the beginning of treatment. These mice had to be sacrificed due to big tumor sizes or tumor growth-related weight loss. In contrast, four out of five mice in the combination group could live without lethal tumor burden and weight loss beyond 64 days and two of them lived over 100 days.

Consistent with the in vitro evaluation, P-CYP was less potent in inhibiting the bulk tumor cell growth than P-DTX. P-CYP alone led to PC-3 tumor growth retardation compared to saline treated group, although tumors eventually grew to about 30 times of the original size on average. Thus, P-CYP or combination therapy did not show immediate benefit in tumor inhibition over the P-DTX treatment. However, combination therapy significantly suppressed the tumor growth for a longer period compared to P-DTX therapy, demonstrating the long-term anti-tumor benefit by inhibiting both tumor stem/progenitor and differentiated cells.

3.5 Ex vivo evaluation of stem cell-like properties in residual tumor cells

To verify whether in vivo administration of P-DTX, P-CYP or their combination affected the cancer stem-like cell subpopulation, the percentage of CD133+ cancer cells and the sphere-forming ability of tumor cells harvested from the residual tumor masses were evaluated. As shown in Fig. 9A, both P-CYP and combination treatments resulted in decreased proportion of CD133+ cells compared to saline group; on the contrary, P-DTX failed to reduce the CD133+ cell population. According to another report, the CD133+ cancer cell population in prostate cancer xenografts after therapeutic interventions still represents stem cell properties [35]; therefore, decrease in CD133+ cells demonstrated the attenuation of CSC activities in the residual tumors. Similarly, the numbers of prostaspheres formed by the tumor cells isolated from P-CYP and combination group were significantly fewer than that isolated from P-DTX treated group (Fig. 9B). Taken together, P-CYP alone or in combination with P-DTX indeed decreased CD133+ cancer stem/progenitor cells, which may contribute to more effective long-term tumor growth inhibition. In future studies, we shall evaluate the tumorigenicity of residual tumor cells when the dosing regimen of the combination therapy was further optimized.

3.6. Toxicity

As shown in Fig. 8B, the body weights of the mice in all four groups did not drop after saline or conjugate administration. The body weights increased steadily at similar rates in all groups, except when tumors grew too big and affected the mice body weights. This indicates that P-DTX, P-CYP and their combination, at the doses administered, are well tolerated. In addition, the chronic toxicities of each dosing regimen were evaluated by histological analysis of the major organs (Fig. 10). No obvious pathological changes were observed in livers, spleens and kidneys of the mice from all treatment groups. Within the experiment duration, we did not observe histological changes in intestines, which is a rapidly self-renewing organ with robust stem cell activities.

4. Conclusions

Most of the anti-cancer combination therapies engage two or more drugs with independent mechanisms of action on cancer cells, in order to achieve synergistic therapeutic effects. As CSC hypothesis revived and cancer cell heterogeneity was taken into consideration, another modality of combination therapy which combines therapeutic agents targeting different

cancer cell populations have shown promising efficacy inhibiting tumor growth and preventing tumor regrowth [37]. In this study, we showed that HPMA copolymer-cyclopamine conjugate preferentially targets the cancer stem/progenitor cell population at doses that do not reduce bulk tumor cells. P-CYP inhibited the self-renew ability and induced apoptosis of CD133+ prostate cancer stem/progenitor cells. The combination of P-CYP and an effective bulk cancer cell-killing conjugate P-DTX showed more effective long-term tumor growth inhibition in vitro and in PC-3 prostate cancer xenografts in mice than single agent therapies.

Acknowledgments

The research was supported in part by the NIH grant CA132831 from the National Cancer Institute. We thank Dr. Rui Zhang and Monika Sima for the help on animal studies and Professor Bryan E. Welm for the consultation on in vitro and ex vivo biological experiments.

References

1. MacVicar GR, Hussain MH. Emerging therapies in metastatic castration-sensitive and castration-resistant prostate cancer. *Curr. Opin. Oncol.* 2013; 25:252–260. [PubMed: 23511665]
2. Grubb RL, Kibel AS. Prostate cancer: screening, diagnosis and management in 2007. *Mol. Med.* 2007; 104:408–412.
3. Medema JP. Cancer stem cells: the challenges ahead. *Nat. Cell Biol.* 2013; 15:338–344. [PubMed: 23548926]
4. Chaffer CL, Weinberg RA. A perspective on cancer cell metastasis. *Science.* 2011; 331:1559–1564. [PubMed: 21436443]
5. Tang DG, Patrawala L, Calhoun T, Bhatia B, Choy G, Schneider-Broussard R, Jeter C. Prostate cancer stem/progenitor cells: identification, characterization and implications. *Mol. Carcinogenesis.* 2007; 46:1–14.
6. Shackleton M, Quintana E, Fearon ER, Morrison SJ. Heterogeneity in cancer: cancer stem cells versus clonal evolution. *Cell.* 2009; 138:822–829. [PubMed: 19737509]
7. Rosen JM, Jordan CT. The increasing complexity of the cancer stem cell paradigm. *Science.* 2009; 234:1670–1677. [PubMed: 19556499]
8. Dick JE. Looking ahead in cancer stem cell research. *Nat. Biotechnol.* 2009; 27:44–46. [PubMed: 19131997]
9. Yu F, Yao H, Zhu P, Zhang X, Pan Q, Gong C, Huang Y, Hu X, Su F, Lieberman J, Song E. Let-7 regulates self-renewal and tumorigenicity of breast cancer cells. *Cell.* 2007; 131:1109–1123. [PubMed: 18083101]
10. Dubrovskaya A, Elliott J, Salamone RJ, Kim S, Aimone LJ, Walker JR, Watson J, Sauveur-Michel M, Garcia-Echeverria C, Cho CY, Reddy VA, Schultz PG. Combination therapy targeting both tumor-initiating and differentiated cell populations in prostate carcinoma. *Clin. Cancer Res.* 2010; 16:5692–5702. [PubMed: 21138868]
11. Gupta PB, Fillmore CM, Jiang G, Shapira SD, Tao K, Kuperwasser C, Lander ES. Stochastic state transitions give rise to phenotypic equilibrium in populations of cancer cells. *Cell.* 2011; 146:633–644. [PubMed: 21854987]
12. Lapidot T, Sirard C, Vormoor J, Murdoch B, Hoang T, Caceres-Cortes J, Minden M, Paterson B, Caligiuri MA, Dick JE. A cell initiating human acute myeloid leukaemia after transplantation into SCID mice. *Nature.* 1994; 367:645–648. [PubMed: 7509044]
13. Al-Hajj M, Wicha MS, Benito-Hernandez A, Morrison SJ, Clarke MF. Prospective identification of tumorigenic breast cancer cells. *Proc. Natl. Acad. Sci. USA.* 2003; 100:3983–3988. [PubMed: 12629218]
14. Collins AT, Berry PA, Hyde C. Prospective identification of tumorigenic prostate cancer stem cells. *Cancer Res.* 2005; 65:10946–10951. [PubMed: 16322242]
15. Miki J, Furusato B, Li H, Gu Y, Takahashi H, Egawa S, Sesterhenn IA, McLeod DG, Srivastava S, Rhim JS. Identification of putative stem cell markers, CD133 and CXCR4, in hTERT-

- immortalized primary nonmalignant and malignant tumor-derived human prostate epithelial cell lines and in prostate cancer specimens. *Cancer Res.* 2007; 67:3153–3161. [PubMed: 17409422]
16. Kelly K, Yin J. Prostate cancer and metastasis initiating stem cells. *Cell Res.* 2008; 18:528–537. [PubMed: 18414448]
 17. Wang ZA, Shen MM. Revisiting the concept of cancer stem cells in prostate cancer. *Oncogene.* 2010; 30:1261–1271. [PubMed: 21119602]
 18. Teng Y, Wang X, Wang Y, Ma D. Wnt/ β -catenin signaling regulates cancer stem cells in lung cancer A549 cells. *Biochem. Biophys. Res. Commun.* 2010; 392:373–379. [PubMed: 20074550]
 19. Karhadkar SS, Bova GS, Abdallah N, Dhara S, Gardner D, Maitra A, Isaacs JT, Berman DM, Beachy PA. Hedgehog signaling in prostate regeneration, neoplasia and metastasis. *Nature.* 2004; 431:707–712. [PubMed: 15361885]
 20. Sanchez P, Hernandez AM, Stecca B, Kahler AJ, DeGueme AM, Barrett A, Beyna M, Datta MW, Datta S, Ruiz I, Altaba A. Inhibition of prostate cancer proliferation by interference with SONIC HEDGEHOG-GLI1 signaling. *Proc. Natl. Acad. Sci. USA.* 2004; 101:12561–12566. [PubMed: 15314219]
 21. Bar EE, Chaudhry A, Lin A, Fan X, Schreck K, Matsui W, Piccirillo S, Vescovi AL, DiMeco F, Olivi A, Eberhart CG. Cyclopamine-mediated hedgehog pathway inhibition depletes stem-like cancer cells in glioblastoma. *Stem Cells.* 2007; 25:2524–2533. [PubMed: 17628016]
 22. Lipinski RJ, Hutson PR, Hannam PW, Nydza RJ, Washington IM, Moore RW, Girdaukas GG, Peterson RE, Bushman W. Dose- and route-dependent teratogenicity, toxicity, and pharmacokinetic profiles of the hedgehog signaling antagonist cyclopamine in the Mouse. *Toxicol. Sci.* 2008; 104:189–197. [PubMed: 18411234]
 23. Kopeček J, Kopecková P. HPMA copolymers: origins, early developments, present, and future. *Adv. Drug Deliv. Rev.* 2010; 62:122–149. [PubMed: 19919846]
 24. Kopeček J. Polymer-drug conjugates: origins, progress to date and future directions. *Adv. Drug Deliv. Rev.* 2013; 65:49–59. [PubMed: 23123294]
 25. Zhou Y, Yang J, Kopeček J. Selective inhibitory effect of HPMA copolymer-cyclopamine conjugate on prostate cancer stem cells. *Biomaterials.* 2012; 33:1863–1872. [PubMed: 22138033]
 26. Liu J, Kopecková P, Pan H, Sima M, Böhler P, Wolf P, Elsässer-Beile U, Kopeček J. Prostate cancer-targeted *N*-(2-hydroxypropyl)methacrylamide copolymer/docetaxel conjugates. *Macromol. Biosci.* 2012; 12:412–422. [PubMed: 22493797]
 27. Peng Z-H, Sima M, Salama ME, Kopecková P, Kopeček J. Spacer length impacts the efficacy of targeted docetaxel conjugates in prostate specific membrane antigen expressing prostate cancer. *J. Drug Targeting.* in press.
 28. Zhou Y, Kopeček J. Biological rationale for the design of polymeric anti-cancer nanomedicines. *J. Drug Targeting.* 2013; 21:1–26.
 29. Mitsukami Y, Donovan MS, Lowe AB, McCormick CL. Water-soluble polymers. 81. Direct synthesis of hydrophilic styrenic-based homopolymers and block copolymers in aqueous solution via RAFT. *Macromolecules.* 2001; 34:2248–2256.
 30. Kopeček J, Bažilová H. Poly[*N*-(2-hydroxypropyl)methacrylamide]. 1. Radical polymerization and copolymerization. *Eur. Polym. J.* 1973; 9:7–14.
 31. Rejmanová P, Pohl J, Baudyš M, Kostka V, Kopeček J. Polymers containing enzymatically degradable bonds. 8. Degradation of oligopeptide sequences in *N*-(2-hydroxypropyl)methacrylamide copolymers by bovine spleen cathepsin B. *Makromol. Chem.* 1983; 184:2009–2020.
 32. Šubr V, Ulbrich K. Synthesis and properties of new *N*-(2-hydroxypropyl)methacrylamide copolymers containing thiazolidine-2-thione reactive groups. *React. Funct. Polym.* 2006; 66:1525–1538.
 33. Gu Y, Li H, Miki J, Kim KH, Furusato B, Sesterhenn IA, Chu WS, McLeod DG, Srivastava S, Ewing CM, Isaacs WB, Rhim JS. Phenotypic characterization of telomerase-immortalized primary nonmalignant and malignant tumor-derived human prostate epithelial cell lines. *Exp. Cell Res.* 2006; 312:831–843. [PubMed: 16413016]
 34. Wang S. Anchorage-independent growth of prostate cancer stem cells. *Methods Mol. Biol.* 2009; 568:151–160. [PubMed: 19582425]

35. Patrawala L, Calhoun T, Schneider-Broussard R, Li H, Bhatia B, Tang S, Reilly JG, Chandra D, Zhou J, Claypool K, Coghlan L, Tang DG. Highly purified CD44+ prostate cancer cells from xenograft human tumors are enriched in tumorigenic and metastatic progenitor cells. *Oncogene*. 2006; 25:1696–1708. [PubMed: 16449977]
36. Nagao Y, Seno K, Kawabata K, Miyasaka T, Takao S, Fujita E. Monitored aminolysis of 3-acylthiazolidine-2-thione: a new convenient synthesis of amide. *Tetrahedron Lett*. 1980; 21:841–844.
37. Dubrovskaya A, Elliott J, Salamone RJ, Kim S, Aimone LJ, Walker JR, Watson J, Sauveur-Michel M, García-Echeverría C, Cho CY, Reddy VA, Schultz PG. Combination therapy targeting both tumor-initiating and differentiated cell populations in prostate carcinoma. *Clin. Cancer Res*. 2010; 16:5692–5702. [PubMed: 21138868]
38. Dubrovskaya A, Kim S, Salamone RJ, Walker JR, Maira S-M, García-Echeverría C, Schultz PG, Reddy VA. The role of PTEN/Akt/PI3K signaling in the maintenance and viability of prostate cancer stem-like cell populations. *Proc. Natl. Acad. Sci. USA*. 2009; 106:268–273. [PubMed: 19116269]
39. Altaba AR, Sánchez P, Dahmane N. Hedgehog signaling in cancer formation and maintenance. *Nat. Rev. Cancer*. 2003; 3:903–911. [PubMed: 14737121]
40. Steg AD, Katre AA, Bevis KS, Ziebarth A, Dobbin ZC, Shah MM, Alvarez RD, Landen CN. Smoothed antagonists reverse taxane resistance in ovarian cancer. *Mol. Cancer Ther*. 2012; 11:1587–1597. [PubMed: 22553355]
41. Olive KP, Jacobetz MA, Davidson CJ, Gopinathan A, McIntyre D, Honess D, Madhu B, Goldgraben MA, Caldwell ME, Allard D, Frese KK, Denicola G, Feig C, Combs C, Winter SP, Ireland-Zecchini H, Reichelt S, Howat WJ, Chang A, Dhara M, Wang L, Rückert F, Grützmann R, Pilarsky C, Izeradjene K, Hingorani SR, Huang P, Davies SE, Plunkett W, Egorin M, Hruban RH, Whitebread N, McGovern K, Adams J, Iacobuzio-Donahue C, Griffiths J, Tuveson DA. Inhibition of hedgehog signaling enhances delivery of chemotherapy in a mouse model of pancreatic cancer. *Science*. 2009; 324:1457–1461. [PubMed: 19460966]
42. Mimeault M, Johansson SL, Henichart J-P, Depreux P, Batra SK. Cytotoxic effects induced by docetaxel, gefitinib, and cyclopamine on side population and nonside population cell fractions from human invasive prostate cancer cells. *Mol. Cancer Ther*. 2010; 9:617–630. [PubMed: 20179163]

Figure 1.
Synthetic scheme of A) HPMA copolymer-cyclopamine conjugate (P-CYP) and B) HPMA copolymer-docetaxel conjugate (P-DTX).

Figure 2. Prostasphere formation by A) PC-3 cells and B) RC-92a/hTERT cells following free and HPMA copolymer conjugated cyclopamine and docetaxel treatments. Data represents mean $\pm$ SD. *, $p < 0.05$.

Figure 3. CD133 expression and cytotoxicity in PC-3 cells following CYP, DTX, P-CYP and P-DTX treatments. Black columns: CD133 expression level (%); gray columns: cell viability (%). *, $p < 0.05$.

Figure 4. GLI1 mRNA expression levels in PC-3 cells following CYP and P-CYP treatments. *, $p < 0.05$.

Figure 5. Cell death induced by CYP and P-CYP in RC-92a/hTERT cells and sorted CD133+/- cells. A) Flow cytometry profiles of cells treated with P-CYP; B) Summary of the apoptosis levels in the whole cell line (white bars), CD133-cells (gray bars) and CD133+ cells (red bars).

Figure 6. Immediate PC-3 cell growth inhibition relative to untreated cells by P-CYP, P-DTX, and simultaneous combination of P-CYP and P-DTX.

Figure 7. Long-term PC-3 cell growth inhibition by single agent and the combination of free drugs (A) and conjugates (B). // - data were not recorded due to the over-confluent cell growth; *, $p < 0.05$; **, $p < 0.01$.

Figure 8.

A) Tumor growth inhibition by P-CYP (CYP equivalent 40 mg/kg), P-DTX (DTX equivalent 10 mg/kg), and combination of P-DTX (DTX equivalent 10 mg/kg) and P-CYP (CYP equivalent 40 mg/kg) in PC-3 tumor-bearing nude mice. * represents statistically significant difference between P-DTX or P-CYP single treatment and combination treatment groups ($p < 0.05$); B) Percentage of mean body weight change in mice treated with P-CYP, P-DTX, and combination of P-CYP and P-DTX. Data are presented as mean $\pm$ SD.

Figure 9. Characterization of the residual tumor cells harvested from tumor bearing mice following in vivo treatments. A) Percentage of CD133 expression level changes in residual tumor cells following in vivo treatments by saline, P-DTX (DTX equivalent 10 mg/kg, single dose), P-CYP (CYP equivalent 40 mg/kg, 6 doses), and combination of P-DTX (DTX 10 mg/kg, single dose) and P-CYP (CYP 40 mg/kg, 6 doses); B) Changes in prostasphere forming ability of the residual tumor cells following in vivo treatments (the same dosing regimens as described in A). *, $p < 0.05$.

Figure 10.

Histological evaluation of organs harvested from PC-3 tumor-bearing mice treated with saline, P-DTX, P-CYP and combination agents. The tissues were sliced and stained with hematoxylin and eosin (H&E).

Table 1

Characterization of HPMA copolymer-cycloamine conjugate (P-CYP) and HPMA copolymer-docetaxel conjugate (P-DTX)

Conjugate	Mn _{theory} , kDa	Mn, kDa	Mw, kDa	Mw/Mn	Drug content (wt%)
P-CYP	45	37.4	39.9	1.07	6.6
P-DTX	45	36.4	38.6	1.06	7.2