Published in final edited form as:

Free Radic Biol Med. 2013 December ; 65: . doi:10.1016/j.freeradbiomed.2013.10.006.

Independent roles of methionine sulfoxide reductase A in mitochondrial ATP synthesis and as antioxidant in retinal pigment epithelial cells

Ying Dun^a, Jade Vargas^a, Nathan Brot^b, and Silvia C. Finnemann^{a,*}

^aDepartment of Biological Sciences, Center for Cancer, Genetic Diseases, and Gene Regulation, Fordham University, Bronx, NY 10458, USA

^bCenter for Molecular Biology and Biotechnology, Florida Atlantic University, Boca Raton, FL 33431, USA, and Department of Microbiology and Immunology, Weill Medical College of Cornell University, New York, NY 10065, USA

Abstract

The antioxidant enzyme methionine sulfoxide reductase A (MsrA) is highly expressed in the retinal pigment epithelium (RPE), a support tissue for neighboring photoreceptors. MsrA protein levels correlate with sensitivity of RPE in culture to experimental oxidative stress. To investigate whether and how MsrA affects RPE functionality regardless of oxidative stress, we tested the effects of acute silencing or overexpression of MsrA on the phagocytosis of photoreceptor outer segment fragments (POS), a demanding, daily function of the RPE that is essential for vision. Endogenous MsrA localized to mitochondria and cytosol of rat RPE in culture. RPE cells manipulated to express higher or lower levels of MsrA than control cells showed no signs of cell death but increased or decreased, respectively, POS binding as well as engulfment. These effects of altered MsrA protein concentration on phagocytosis were independent of the levels of oxidative stress. However, altering MsrA expression had no effect on phagocytosis when mitochondrial respiration was inhibited. Furthermore, ATP content directly correlated with MsrA protein levels in RPE cells that used mitochondrial oxidative phosphorylation for ATP synthesis but not in RPE cells that relied on glycolysis alone. Overexpressing MsrA was sufficient to increase specifically the activity of complex-IV of the respiratory chain, while activity of complex-II and mitochondrial content were unaffected. Thus, MsrA likely enhances ATP synthesis by increasing complex-IV activity. Such contribution of MsrA to energy metabolism is independent of its function in protection from elevated oxidative stress but contributes to routine but vital photoreceptor support by RPE cells.

Keywords

mitochondria; MsrA; oxidative stress; phagocytosis; respiratory chain; retinal pigment epithelium

© 2013 Elsevier Inc. All rights reserved.

*Corresponding author. Phone: +1 718 817 3630; fax: +1 718 817 3514; finnemann@fordham.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Introduction

The enzyme methionine sulfoxide reductase A (MsrA) is widely distributed among mammalian tissues and localizes to mitochondria, the cytosol, and/or the nucleus depending on cell type [1, 2]. Different transcripts of the MSRA gene yield MsrA isoforms that assume distinct subcellular distributions [3, 4]. MsrA acts enzymatically to reduce *S*-methionine sulfoxide residues in proteins thereby repairing protein modifications induced by oxidative stress [1, 5]. Notably, MsrA has recently been found to promote both the oxidation of methionine and reduction of methionine sulfoxide [6]. Intriguing examples of how MsrA may oxidize and reduce methionine residues in proteins to reversibly regulate functionality of specific proteins have also been reported. Methionine residue oxidation slows down the inactivation of voltage-dependent potassium channels expressed in *Xenopus* oocytes, a process reversed by MsrA overexpression [7]. Methionine oxidation contributes to the activation of calcium/calmodulin-dependent protein kinase II suggesting a possible role for reversible oxidation in signal transduction pathways [8]. Identification of specific MsrA substrates and cellular processes controlled by MsrA remains an active area of investigation.

Survival and functionality in vision of photoreceptor neurons in the retina require continuous support by the neighboring retinal pigment epithelium (RPE)¹ (reviewed in [9]). Like photoreceptors, mammalian RPE cells are post-mitotic and subjected to a lifetime of photo-oxidative stress. Most RPE functions are dependent on sufficient availability of ATP generated by oxidative phosphorylation in mitochondria. Mitochondrial defects severely impair the functions of the RPE *in vivo* and in cell culture [10, 11]. Decline in mitochondrial activity is associated with aging of the human RPE and the development of age-related macular degeneration (AMD) [12]. The molecular mechanisms controlling mitochondrial ATP synthesis efficiency in RPE cells have not yet been extensively studied.

Earlier reports have shown a role for MsrA in protection of RPE cells from excess oxidative stress (reviewed in [13]). In rat retina, MsrA is abundant in the RPE [1]. In monkey retina, MsrA levels are highest in the RPE in the macular region of the retina where RPE cells must support a particularly high number of tightly packed cone photoreceptors [3]. In human retina, MsrA localizes to the RPE and in part to drusen deposits beneath the RPE that are associated with AMD [14]. RPE cells in culture respond to moderate levels of experimental oxidative stress by increasing MsrA expression. Acutely reducing MsrA of RPE cells by gene silencing enhances cytotoxicity of oxidative stress [3, 14].

We hypothesized that MsrA may support the routine functions of unstressed RPE cells. Whether or not MsrA fulfills functions in RPE cells other than protection from acute oxidative damage has not yet been directly investigated. The continuous clearance of shed photoreceptor outer segment fragments (POS) by phagocytosis and their prompt and complete digestion are among critical RPE responsibilities. POS phagocytosis employs the RPE F-actin cytoskeleton and its phago-lysosomal organelles all of which must be intact and dynamic [15, 16]. POS phagocytosis is a costly process that requires ATP synthesis by RPE mitochondria [10]. Sensitive experimental uptake assays can accurately and with high sensitivity quantify phagocytic binding and engulfment of purified POS by RPE cells in culture. In this study, we characterized the effects of specifically decreasing or increasing MsrA on the phagocytic function of RPE cells in culture reasoning that even moderate changes in RPE function will affect RPE phagocytosis. We compared the effects of altered MsrA expression on phagocytic activity and cell viability in the presence of hydrogen peroxide, trolox antioxidant, or mitochondrial respiratory chain inhibitors. We determined

¹**Abbreviations:** used AMD, age-related macular degeneration; β -gal, β -galactosidase; PBS, phosphate buffered saline; PFA, paraformaldehyde; POS, shed photoreceptor outer segment tips; RPE, retinal pigment epithelium; siRNA, silencing RNAs.

that MsrA promotes phagocytic function by increasing the activity of complex-IV of the respiratory chain and as a result mitochondrial ATP synthesis, regardless of the levels of oxidative stress. Conversely, MsrA protection from damage by hydrogen peroxide was unaffected by mitochondrial inhibition. Thus, MsrA supports RPE function by independently supporting mitochondrial ATP synthesis and counteracting oxidative damage.

Materials and methods

All reagents were purchased from Sigma (St. Louis, MO) or Invitrogen (Carlsbad, CA) unless otherwise specified.

Antibodies

Opsin clone B6-30 (a gift from Dr. Paul Hargrave [17]), cyclophilin D, MsrA, tubulin, (all Abcam, Cambridge, MA), complex-I subunit NDUF8 (clone 20E9DH10C12), complex-II subunit 30 kDa (clone 21A11AE7), complex-III subunit Core 2 (clone 13G12AF12BB11), complex-IV subunit I (clone 1D6E1A8), complex-V -subunit (clone 15H4C4), (all Invitrogen), -galactosidase (-gal) (Santa Cruz Biotechnologies, Santa Cruz, CA), cytochrome C, porin (Cell Signaling, Cambridge, MA).

Cell culture, silencing RNA (siRNA) transfections, and adenoviral infections

RPE cells for primary culture were isolated from 9 - 11 day-old Long Evans rats housed under cyclic 12 h light: 12 h dark light conditions and fed *ad libitum* using a previously published procedure [18, 19]. Briefly, cornea, lens, iris and vitreous body were removed from eyes that had been enucleated immediately after animal asphyxiation with CO₂. Eyecups were incubated with 1 mg/ml hyaluronidase in Hank's balanced saline solution without Ca²⁺ and Mg²⁺ for 45 min at 37°C. The neural retina was removed and eyecups were incubated with 2 mg/ml trypsin in Hank's balanced saline solution with Ca²⁺ and Mg²⁺ for 30 to 45 min at 37°C. RPE sheets were manually collected from the underlying choroid. Purified RPE patches were grown in 96-well plates on collagen IV-coated glass coverslips in DMEM 10% FBS at 37°C, 5% CO₂ for 5-7 days without passaging before experiments. Only primary cell samples that retained the obvious cobblestone appearance of polarized RPE were used for experiments. All procedures involving animals were performed according to the ARVO Statement for the Use of Animals in Ophthalmic and Vision Research and the National Institutes of Health *Guide for the Care and Use of Laboratory Animals* and were reviewed and approved by the Fordham University Institutional Animal Care and Use Committee.

Immortalized rat RPE-J cells (ATCC, Manassas, VA) were maintained at 32°C and 8% CO₂ in DMEM supplemented with 4% FBS (CELlect Gold, ICN, Irvine, CA) and were sub-cultured every seven days. Post-confluent, differentiated cells were used six days after plating.

To silence MsrA, RPE-J cells received a mixture of four different 21-nucleotide RNAs specifically targeting rat MsrA (Accell SMARTpool siRNA, Dharmacon/Thermo Fisher Scientific, Lafayette, CO). Accell non-targeting siRNA pool (Dharmacon) confirmed to have minimal targeting of known genes in human, mouse and rat cells was used as a negative control. Cells in 96-well plates were incubated with 100 nmoles siRNA mix per well. Cells in 24-well plates were incubated with 250 nmoles siRNA mix per well. For experiments shown in figure 9, cells in 96-well plates were incubated with 50 nmoles siRNA mix per well. Uptake of siRNA was enhanced using Metafectene Pro (Biontex, San Diego, CA) following the manufacturer's protocol.

Established replication-defective adenoviruses were used to induce expression of -gal (Cell Biolabs, San Diego, CA) or MsrA [20]. Adenovirus was applied at a concentration of 1×10^9 particles/ml in serum-free DMEM. Post-confluent RPE-J cells were infected four days after plating by incubating with this adenovirus dilution for 16 hours. Primary RPE cells were infected 5 days after plating by incubating with the same adenovirus dilution for 2 hours.

Following either siRNA or adenovirus treatment, cells were incubated in regular growth medium for 40 to 48 hours before use. As appropriate, 20 M trolox, 50 or 200 M hydrogen peroxide, 0.25 M antimycin, or 0.1 M oligomycin were included in media during experiments.

POS phagocytosis assays

Published protocols using sucrose density gradient centrifugation were followed to purify POS from pig eyes obtained fresh from a local slaughterhouse and to covalently label POS with FITC isomer I [21, 22]. RPE-J cells received FITC-POS suspended in DMEM with 5% FBS at a ratio of ~10 to 1 for 3.5 hours before three washes with PBS containing 0.2 mM CaCl_2 and 1 mM MgCl_2 (PBS-CM). For subsequent immunoblotting or cell viability analysis, cells were processed as described in sections below. To quantify total (bound plus internalized) FITC-POS, cells were fixed with ice-cold methanol followed by rehydration in PBS-CM. To quench FITC emission of surface-bound POS, cells were incubated with 0.25% trypan blue in PBS-CM for 10 min and washed twice in PBS-CM before fixation and rehydration. FITC emission of total or internal POS were quantified by fluorescence scanning with a Typhoon Trio Scanner (GE Healthcare) as described in detail previously [23].

Quantification of cell viability

RPE-J cells were treated with siRNA followed by phagocytosis assays, peroxide, and/or drug treatments before labeling live and dead cells with different fluorescent dyes using the LIVE/DEAD® Viability/Cytotoxicity Kit for mammalian cells (Invitrogen) according to the manufacturer's protocol. The test principle was validated first by fluorescence microscopy, which confirmed green fluorescence of live cells and red fluorescence of dead cells. A Spectramax M2e plate reader (Molecular Devices, Sunnyvale, CA) was then used to quantify fluorescence emission at 530 nm after excitation at 485 nm of cells in flat bottom, black wall 96-well plates, which directly corresponds to numbers of live cells per well.

Cell lysis, cell fractionation, SDS-PAGE and immunoblotting

RPE-J or primary RPE cells were chilled on ice before lysis in 50 mM HEPES, pH 7.4, 150 mM NaCl, 10% glycerol, 1.5 mM MgCl_2 , 1% Triton X-100 freshly supplemented with 1% protease and phosphatase inhibitor cocktails (HNTG buffer). Clarified whole cell lysates representing equal numbers of cells were boiled for 3 min in reducing SDS sample buffer and separated on SDS-polyacrylamide gels. Fractions of cells enriched in mitochondria were obtained by differential centrifugation [24]. Briefly, cells were homogenized in buffer containing 225 mM mannitol, 75 mM sucrose, 5 mM Hepes, 1 mM EGTA, pH 7.4, with a Dounce homogenizer. Homogenate was centrifuged at 2,000 g for 5 min at 4°C, and the pellet was discarded. The supernatant was then centrifuged at 14,000 g for 10 min at 4°C yielding a pellet enriched in mitochondria and a supernatant depleted from mitochondria. Both fractions were supplemented with SDS-sample buffer and separated side-by-side with total extracts on SDS-polyacrylamide gels. Proteins were transferred to nitrocellulose membranes, which were blocked with non-fat milk and incubated sequentially with primary antibodies and appropriate horseradish peroxidase-conjugated secondary antibodies (Jackson ImmunoResearch Laboratories, West Grove, PA) followed by ECL chemiluminescence

detection (Perkin-Elmer, Waltham, MA). X-ray films were scanned and processed using Photoshop CS4 (Adobe, San Jose, CA). Bands were quantified by densitometry using ImageQuant™ TL 7.0.

Immunofluorescence staining and microscopy

To label live mitochondria using a fluorescent probe, primary rat RPE or RPE-J cells were incubated at growth temperature for 15 min with 0.5 M MitoTracker Red FM in serum-free DMEM followed by a 15 min incubation in serum-free DMEM and fixation with 4% PFA. Fixed cells were permeabilized with 0.2% Triton-X100 in PBS-CM for 5 min before sequential incubation with rabbit antibody to MsrA and with donkey anti rabbit IgG conjugated to AlexaFluor488. For double immunolabeling, cells were fixed with ice-cold methanol for 5 min before sequential incubation with MsrA antibody and complex-IV protein I antibody raised in mouse followed by a mix of donkey-anti-rabbit IgG conjugated to AlexaFluor488 and donkey-anti-mouse IgG conjugated to AlexaFluor594. Nuclei were counterstained with DAPI. Images of single x-y focal planes were acquired on a Leica TSP5 laser scanning confocal microscopy system (Leica, Mannheim, Germany) using sequential scanning mode and compiled using Photoshop CS4.

Quantification of cellular ATP content

ATP was extracted from RPE-J cells in 24-well plates or primary rat RPE in 96-well plates with ice-cold 0.5% trichloroacetic acid before neutralization with 0.05 M Tris-acetate. ATP content in serial dilutions of whole cell extracts was measured using the ENLITEN ATP Luminescence Assay Kit and measured with a Glomax Luminometer (both Promega, Madison, WI). A concentration gradient of ATP (10^{-11} to 10^{-7} M) (Promega) was also measured for each experiment.

Quantification of cellular mitochondrial complex-II and complex-IV activities

Cells in 24-well plates were resuspended and extracted with detergent according to the protocols and with the materials provided in the Complex-II Enzyme Activity Microplate Assay Kit or the Complex-IV Rodent Enzyme Activity Microplate Assay Kit (both Abcam). These assays quantify the specific enzymatic activity of individual respiratory chain complexes that are captured from sample extracts by antibodies immobilized in 96-well plates. Each assay yielded relative complex activity in whole cell extracts of test cells as compared to activity in whole cell extracts of control cells. Cell counting and comparative immunoblotting were used to confirm that samples represented equal cell numbers.

Statistical analysis

All experiments were performed at least three times independently. In each phagocytosis assay, triplicate samples were analyzed. In mitochondrial assays, duplicate samples were analyzed. Samples were first analyzed using one-way ANOVA to determine significant variance among groups. If significance was established, difference between select sample groups was determined using Bonferroni's multiple comparison post-hoc test. Student's *t* test was used to determine the difference between two samples where appropriate. Statistics used is indicated in figure legends. *P* values < 0.05 were considered a statistically significant difference for all analyses.

Results

To characterize the functions of MsrA in RPE cells, we first sought to identify an appropriate RPE model system. Earlier immunohistological experiments demonstrated that rat RPE cells *in situ* express MsrA but those experiments did not reveal the subcellular

distribution of MsrA [1]. Here, we immunolabeled MsrA in cells that had either been live-labeled with MitoTracker dye before fixation (Fig. 1, A and E) or that were also immunolabeled after fixation with antibody to mitochondrial complex-IV (Fig. 1, D and H). Arrows point out colocalization in magnified areas. Both co-staining methods detected MsrA mainly in mitochondria and to a lesser extent in the cytoplasm of unpassaged rat RPE cells in primary culture similar to the distribution observed in human and monkey primary RPE cells [3, 14] (Figure 1, A to D). The rat RPE-derived RPE-J cell line retains expression of many proteins characteristic of mammalian RPE *in situ* and provides a reliable and robust model for the RPE's phagocytic function [23]. Here, we found that RPE-J cells possess mitochondrial and cytoplasmic MsrA similar to primary rat RPE (Fig. 1, E to H). We therefore chose this cell line as the principal RPE model for our investigation.

Silencing MsrA impairs RPE phagocytosis and increases toxicity of oxidative stress

To reduce MsrA protein levels, we applied a mixture of silencing RNAs (siRNA) specific to MsrA to confluent RPE-J cells. This treatment reduced MsrA protein levels two days later by 77% on average (Fig. 2, A and B). Incubation of cells with POS for 3.5 hours did not change MsrA levels (Fig. 2A, compare lanes m and POS). We found that cells expressing less MsrA bound and engulfed POS only poorly compared to untreated cells (Fig. 2, C and D, compare white and black bars). Treatment of RPE cells with a mixture of control siRNA that do not specifically target any mammalian transcript did not significantly alter MsrA protein, POS binding or POS engulfment indicating that the effects observed were MsrA specific (Fig. 2, B to D, compare white and gray bars). Importantly, MsrA silencing did not significantly affect RPE cell viability regardless of phagocytic challenge (Fig. 2E).

MsrA overexpression increases the phagocytic activity of RPE cells

To study the effects of elevated MsrA protein levels on RPE phagocytosis, we infected RPE cells with recombinant adenovirus encoding full-length human MsrA (MsrA virus) or control virus encoding -galactosidase (-gal), a protein that does not affect RPE phagocytosis [15]. We used immunoblotting to compare MsrA content in total cell homogenates and separated fractions of cells obtained by differential centrifugation yielding supernatant fractions that were depleted from mitochondria (sup) and pelleted fractions that were enriched in mitochondria (mito). Infection with MsrA virus increased total cellular MsrA protein levels 3.7-fold on average compared to untreated cells while -gal virus promoted -gal expression but did not alter MsrA (Fig. 3, A and B). Both cell fractions contained significantly more MsrA in cells infected with MsrA virus than in uninfected or -gal virus infected cells (Fig. 3, A and B). However, the increase in MsrA content was somewhat higher in the mitochondria-enriched cell fraction than in the mitochondria-depleted fraction (Fig. 3B, compare black bars, sup, and black bars, mito). We then compared POS phagocytosis by cells expressing -gal or MsrA with phagocytosis by uninfected cells. MsrA expression was unaffected by POS challenge (Fig. 3C). Cells overexpressing MsrA contained elevated levels of total POS opsin representing bound plus engulfed POS (Fig. 3C). Separate quantification of POS binding and POS engulfment showed that cells overexpressing MsrA bound and internalized significantly more POS than control cells (Fig. 3, D and E). Internalization was enhanced slightly more than binding. Thus, MsrA overexpression likely promotes both steps of the phagocytic process.

MsrA-mediated changes of RPE phagocytic activity require mitochondrial respiration

We next sought to determine the cellular mechanisms through which MsrA affects RPE phagocytosis. We first challenged the cells with POS in the presence of 20 M trolox, a water-soluble, vitamin E based antioxidant or with 50 M hydrogen peroxide. We had previously established that these reagents at the concentrations used did not by themselves

affect phagocytosis by RPE-J cells [10, 16]. Here, we found that MsrA silencing specifically decreased POS uptake and excess MsrA specifically increased POS uptake even in the presence of trolox or hydrogen peroxide (Fig. 4, A and B). Thus, neither antioxidant nor increased oxidative stress changed the alterations in phagocytosis caused by lower or higher levels of MsrA. Next, we challenged the cells with POS in the presence of two different inhibitors of mitochondrial respiration. Strikingly, altering MsrA expression lost its effect on RPE phagocytosis if the mitochondrial respiratory chain was inhibited by antimycin, which targets complex-III (Fig. 4C) or oligomycin, which targets complex-V (Fig. 4D). None of the compounds used affected MsrA expression levels (Fig. 4E). Taken together, these results demonstrate that MsrA does not alter RPE phagocytosis by counteracting oxidative stress. However, the effect of MsrA on RPE phagocytic activity requires mitochondrial respiration.

Susceptibility to oxidative stress of RPE cells lacking MsrA is unrelated to mitochondrial respiration

Earlier studies documented that lack of MsrA increases the levels of cell death in response to experimental oxidative stress [3, 14]. Here, we found that treatment with 50 M hydrogen peroxide for 3.5 hours did not significantly affect RPE cell viability regardless of MsrA protein level (Fig. 5A). However, incubation with 200 M hydrogen peroxide proved lethal to 40% of cells treated with MsrA siRNA but had no significant effect on the survival of cells treated with control siRNA (Fig. 5B). We wondered whether this increased susceptibility to oxidative stress of RPE lacking MsrA was related to mitochondrial respiration like the decreased phagocytic function (Fig. 4). We thus quantified survival of RPE cells after treatment with 200 M hydrogen peroxide in the presence of inhibitors of mitochondrial respiration or antioxidant. At the end of a 3.5-hour incubation with 200 M hydrogen peroxide plus antimycin or oligomycin, viability of cells with silenced MsrA remained significantly lower than viability of cells treated with control siRNA (Fig. 5C, antimycin, oligomycin). In contrast, co-incubation with antioxidant largely prevented hydrogen peroxide-induced cell death of cells lacking MsrA (Fig. 5C, trolox). Thus, providing antioxidant alleviates the increased susceptibility of RPE cells with decreased levels of MsrA to the cytotoxic effects of oxidative stress but inhibiting mitochondrial respiration does not.

MsrA concentration affects mitochondrial respiration and cellular ATP content

Because MsrA affected phagocytosis only if the respiratory chain was active, we suspected that MsrA might alter the energy metabolism of RPE cells. Therefore, we quantified cellular ATP content of RPE cells with altered MsrA protein levels. Indeed, 24 hours after being supplied with regular growth medium containing 25 mM glucose (high glucose), the ATP content of RPE cells with elevated levels of MsrA was increased by 29% compared to cells expressing -gal. The ATP content of RPE cells with silenced MsrA was decreased by 25% compared to cells treated with control siRNA and by 42% compared to untreated control cells (Fig. 6A). -gal control adenovirus had no effect on ATP content but cells receiving control siRNA alone possessed 22% less ATP than untreated control cells (Fig. 6A, compare white and gray bars).

To more specifically examine the causal relationship of MsrA and ATP synthesis, we next depleted ATP from cells with two consecutive 30 min incubations with carbon-source free medium before supplying cells with medium containing either physiological levels of glucose (5.6 mM) or pyruvate (1 mM) for 1 hour in the presence or absence of respiratory chain inhibitor oligomycin. Pyruvate as energy source depends on an active mitochondrial respiratory chain to yield ATP while glucose may yield ATP through glycolysis regardless of mitochondrial activity. We observed that RPE cells treated with either control or MsrA siRNA responded to the energy depletion-repletion regimen with frequent cell detachment.

Therefore, we continued these experiments using only adenovirus-infected cells. Supplying these cells with glucose at a physiological concentration yielded a small but significant increase in ATP content in cells overexpressing MsrA over cells expressing -gal unless mitochondrial activity was inhibited (Fig. 6B). If cells were supplied with glucose in the presence of oligomycin, there was no difference in ATP content between cells differing in MsrA levels (Fig. 6B, oligomycin). If cells were supplied with pyruvate as the sole carbon source, the ATP content of cells overexpressing MsrA was 34% higher than the ATP content of control cells (Fig. 6C). As expected, cells contained very little ATP if they received pyruvate in the presence of oligomycin (Fig. 6C, oligomycin). Adding trolox or sublethal amounts of hydrogen peroxide had no effect on ATP content of pyruvate-fed cells (Fig. 6D). Taken together, these data indicate that mitochondrial synthesis of ATP is more efficient in RPE cells with higher levels of MsrA. In contrast, MsrA levels do not affect the ATP content of cells whose respiration is inhibited by oligomycin.

MsrA controls the specific activity of the respiratory chain rather than mitochondrial load

RPE cells overexpressing MsrA could increase mitochondrial ATP synthesis by generating more mitochondria or by increasing the specific activity of mitochondrial ATP synthesis. To distinguish between these mechanisms we first quantified mitochondrial protein content of RPE cells. Figure 7 shows representative immunoblots of one protein of each of the five functional protein complexes of the mitochondrial respiratory chain, the soluble mitochondrial matrix proteins cyclophilin D and cytochrome C, and the outer mitochondrial membrane protein porin. Tubulin, MsrA and -gal were also detected to confirm total cell load and MsrA silencing or viral overexpression (Fig. 7). Repeated similar immunoblotting studies revealed that there was no significant change in any of the mitochondrial proteins tested with treatment. We also noted a decrease of about 20% in cell yield in samples treated with siRNA. However, this change was not specific to silencing MsrA.

Finally, quantification of the enzymatic activity of complex-IV of the respiratory chain (also known as cytochrome C oxidase) showed a robust and specific increase if MsrA was overexpressed and a significant decrease if MsrA was silenced (Fig. 8A). Similar assays revealed only small changes in complex-II activity in response to MsrA up- or downregulation that did not reach statistical significance (Fig. 8B). Thus, RPE cells respond to MsrA overexpression by increasing the specific activity of complex-IV, which in turn increases mitochondrial respiration.

Our data thus far showed that cells with silenced MsrA showed significant decreases in POS phagocytosis (Fig. 2 and 4), resistance to oxidative stress (Fig. 5), ATP content (Fig. 6) and complex-IV activity in comparison to cells treated with control siRNA and thus as response specifically to loss of MsrA. However, an additional decrease in ATP (Fig. 6) and in cell yield (Fig. 7) in comparison to untreated cells was shared by cells treated with control and MsrA specific siRNA suggesting that our siRNA treatment affected cell functionality non-specifically as well. To experimentally separate specific and non-specific effects of the silencing treatments, we repeated silencing assays applying only half as much siRNA as previously. We found that this treatment decreased MsrA protein levels by ~50% (Fig. 9, A and B) and was sufficient to significantly impair RPE phagocytic activity (Fig. 9, A and C) and decrease ATP content (Fig. 9D). Importantly, comparable control siRNA treatment had no effect on ATP levels (Fig. 9D, compare gray to white bar), and immunoblotting analysis confirmed normal cell yield of cells subjected to control or MsrA siRNA (Fig. 9A). Taken together, these results further support the conclusion that decreasing protein levels of MsrA impairs RPE mitochondrial function and phagocytosis.

Unpassaged primary rat RPE cells increase mitochondrial function and ATP synthesis in response to extra MsrA like the RPE-J cell line

To determine whether our results were representative for RPE cells in general, we repeated the key experiments of our study testing the response by rat primary RPE cells. We isolate these cells from rat eyes for culture for up to one week only as epithelial explants, in which the majority of cells do not proliferate and retains many characteristics of its epithelial and RPE phenotype. Adenoviral infection that increased MsrA protein level of rat primary RPE by 3.2-fold on average had no effect on expression of any of the mitochondrial or cell marker proteins tested (Fig. 10A). Yet, primary RPE with elevated MsrA increased their ATP content by 39% on average over cells expressing -gal during a 1-hour incubation in medium with pyruvate as sole carbon source (Fig. 10B). Like RPE-J cells, primary rat RPE responded to extra MsrA with an increase in activity of complex-IV (Fig. 10C) but not of complex-II (Fig. 10D) of their respiratory chain. Finally, MsrA overexpression was also sufficient to promote increased uptake of POS particles by primary rat RPE (Fig. 10E). Taken together, these results suggest that MsrA improves mitochondrial energy metabolism of rat primary RPE, which enhances their function.

Discussion

This study identifies a role for the antioxidant enzyme MsrA in controlling specifically the activity of complex-IV of the mitochondrial respiratory chain. Increasing or decreasing levels of MsrA protein altered respiratory chain complex-IV activity, ATP levels, and the ATP-dependent phagocytic function of RPE cells. MsrA changes had no effect on ATP content or phagocytosis if cells used glycolysis alone for ATP synthesis indicating that these effects require active mitochondrial respiration. In contrast, MsrA effects were independent of levels of oxidative stress. Taken together, our study exploring RPE cells in culture reveals a novel role for MsrA in supporting mitochondrial ATP synthesis that is important for the functionality of RPE cells. Future experiments comparing energy metabolism and POS phagocytosis by RPE cells *in situ* of wild-type and MsrA null mice will show how this activity of MsrA affects retinal health and functionality.

Mitochondria serve critical functions in cells maintaining calcium homeostasis, dissipating reactive oxidative species and synthesizing ATP [25]. Earlier studies have linked MsrA expression levels to resistance to oxidative stress in RPE and numerous other cell types [3, 13, 26-28]. Furthermore, Nan and colleagues reported enlarged mitochondria in heart muscle of MsrA^{-/-} mice [29]. The same study also showed enhanced susceptibility of MsrA^{-/-} cardiomyocytes in culture to oxidative stress but the mitochondrial energy metabolism of these cells was not assessed. In the present study, we are expanding our knowledge of MsrA further showing that MsrA protein levels also regulate efficiency of mitochondrial respiration and thus ATP synthesis independently of oxidative stress.

In this study we experimentally manipulated levels of MsrA protein using gene expression silencing and adenoviral overexpression. Viral overexpression of MsrA increased MsrA content preferentially but not solely in mitochondrial organelles. Interestingly, cells may use endogenous mechanisms to similarly increase mitochondrial MsrA in response to oxidative stress [14]. Our experiments did not reveal changes in MsrA following hydrogen peroxide treatment likely because our challenge was restricted to a 3.5-hour time period. In part cellular adaptation of MsrA content takes place at the level of transcription [14]. Its specificity for mitochondrial MsrA suggests the intriguing possibility that RPE cells under oxidative stress may produce certain splice variants of MsrA transcripts, may re-distribute MsrA to mitochondria, or may regulate specific activity of MsrA to increase mitochondrial ATP synthesis, which in turn increases efficiency of antioxidant defense mechanisms.

Deciphering which mechanism/s cells employ to regulate MsrA in response to oxidative insult will require future studies.

Our experiments reveal that modifying MsrA protein levels is sufficient to change ATP content and that MsrA does not increase ATP synthesis by increasing cellular mitochondrial load. Rather, elevating levels of MsrA increases the rate of mitochondrial respiration by increasing enzymatic activity of complex-IV (but not of complex-II) of the respiratory chain. Although we did not test the activities of complexes-I, -III, and -V, which may also be affected, these results suggest that MsrA acts specifically on complex-IV.

The *in vitro* assays we performed quantified activities of individual respiratory chain complexes immunocaptured from whole cell extracts. These experiments revealed increased activity of complex-IV but not of complex-II. Intact mitochondria in live cells should yield higher ATP content only if the respiratory chain as a whole increases activity. Our experiments showed that increasing MsrA elevates ATP content in RPE cells that rely solely on mitochondrial respiration for ATP synthesis. We thus speculate that the complex-IV activity increase caused by extra MsrA indirectly upregulates activities of the remaining components of the respiratory chain. Alternatively, complex-IV activity may be the rate-limiting step of respiration by RPE cells such that its increase will yield more ATP even if activities of complex-II and possibly other aspects of the respiratory chain remain unchanged. Indeed, it has been proposed earlier that cells may alter specifically complex-IV efficiency to adjust ATP synthesis to different signaling and stress conditions [30].

Given the presence of MsrA in mitochondria it is possible that MsrA directly interacts with complex-IV proteins. Oxidized methionine residues of complex-IV protein/s may thus be substrates for MsrA. Complex-IV is a non-covalent assembly of thirteen different protein subunits, three of which are encoded by the mitochondrial genome. To our knowledge, there have been no reports yet on the oxidation of methionine residues in complex-IV or other respiratory chain complex proteins. MsrA may act to reverse methionine oxidation in complex-IV proteins that occurred non-specifically. Conceptually, altering respiratory chain activity in response to changes in overall mitochondrial oxidative burden would be a physiologically relevant pathway of adaptation. However, our results show that altering the overall oxidative burden of cells by either adding hydrogen peroxide or trolox antioxidant did not affect the ability of MsrA to change ATP synthesis and cell functions that rely on it. This result supports the conclusion that regulation of respiratory chain activity by MsrA is largely independent of levels of oxidative stress. We speculate that MsrA may modify specific methionine residues by both oxidizing and reducing them given its innate dual enzymatic function [6]. Future analysis of oxidation state of methionine residues in specific respiratory chain proteins will be needed to investigate this intriguing possibility.

Acknowledgments

Paul Hargrave kindly provided opsin antibody clone B6-30 for our study. We thank Mark P. Connor for excellent technical assistance. This work was supported by National Institutes of Health Grant EY013295 from the National Eye Institute and a Fordham Faculty Fellowship (to S.C.F.). The authors declare no conflict of interest.

References

- [1]. Moskovitz J, Jenkins NA, Gilbert DJ, Copeland NG, Jursky F, Weissbach H, Brot N. Chromosomal localization of the mammalian peptide-methionine sulfoxide reductase gene and its differential expression in various tissues. *Proc. Natl. Acad. Sci. U. S. A.* 1996; 93:3205–3208. [PubMed: 8622914]

- [2]. Kim HY, Gladyshev VN. Role of structural and functional elements of mouse methionine-S-sulfoxide reductase in its subcellular distribution. *Biochemistry*. 2005; 44:8059–8067. [PubMed: 15924425]
- [3]. Lee JW, Gordiyenko NV, Marchetti M, Tserentsoodol N, Sagher D, Alam S, Weissbach H, Kantorow M, Rodriguez IR. Gene structure, localization and role in oxidative stress of methionine sulfoxide reductase A (MSRA) in the monkey retina. *Exp. Eye Res.* 2006; 82:816–827. [PubMed: 16364291]
- [4]. Kim HY, Gladyshev VN. Alternative first exon splicing regulates subcellular distribution of methionine sulfoxide reductases. *BMC Mol. Biol.* 2006; 7:11. [PubMed: 16542431]
- [5]. Moskovitz J, Weissbach H, Brot N. Cloning the expression of a mammalian gene involved in the reduction of methionine sulfoxide residues in proteins. *Proc. Natl. Acad. Sci. U. S. A.* 1996; 93:2095–2099. [PubMed: 8700890]
- [6]. Lim JC, You Z, Kim G, Levine RL. Methionine sulfoxide reductase A is a stereospecific methionine oxidase. *Proc. Natl. Acad. Sci. U. S. A.* 2011; 108:10472–10477. [PubMed: 21670260]
- [7]. Ciorba MA, Heinemann SH, Weissbach H, Brot N, Hoshi T. Modulation of potassium channel function by methionine oxidation and reduction. *Proc. Natl. Acad. Sci. U. S. A.* 1997; 94:9932–9937. [PubMed: 9275229]
- [8]. Erickson JR, Joiner ML, Guan X, Kutschke W, Yang J, Oddis CV, Bartlett RK, Lowe JS, O'Donnell SE, Aykin-Burns N, Zimmerman MC, Zimmerman K, Ham AJ, Weiss RM, Spitz DR, Shea MA, Colbran RJ, Mohler PJ, Anderson ME. A dynamic pathway for calcium-independent activation of CaMKII by methionine oxidation. *Cell*. 2008; 133:462–474. [PubMed: 18455987]
- [9]. Strauss O. The retinal pigment epithelium in visual function. *Physiol. Rev.* 2005; 85:845–881. [PubMed: 15987797]
- [10]. Vives-Bauza C, Anand M, Shirazi AK, Magrane J, Gao J, Vollmer-Snarr HR, Manfredi G, Finnemann SC. The age lipid A2E and mitochondrial dysfunction synergistically impair phagocytosis by retinal pigment epithelial cells. *J. Biol. Chem.* 2008; 283:24770–24780. [PubMed: 18621729]
- [11]. Zhao C, Yasumura D, Li X, Matthes M, Lloyd M, Nielsen G, Ahern K, Snyder M, Bok D, Dunaief JL, LaVail MM, Vollrath D. mTOR-mediated dedifferentiation of the retinal pigment epithelium initiates photoreceptor degeneration in mice. *J. Clin. Invest.* 2011; 121:369–383. [PubMed: 21135502]
- [12]. Jarrett SG, Lewin AS, Boulton ME. The importance of mitochondria in age-related and inherited eye disorders. *Ophthalmic Res.* 2010; 44:179–190. [PubMed: 20829642]
- [13]. Sreekumar PG, Hinton DR, Kannan R. Methionine sulfoxide reductase A: Structure, function and role in ocular pathology. *World J. Biol. Chem.* 2011; 2:184–192. [PubMed: 21909460]
- [14]. Sreekumar PG, Kannan R, Yaung J, Spee CK, Ryan SJ, Hinton DR. Protection from oxidative stress by methionine sulfoxide reductases in RPE cells. *Biochem. Biophys. Res. Commun.* 2005; 334:245–253. [PubMed: 15993845]
- [15]. Mao Y, Finnemann SC. Essential diurnal Rac1 activation during retinal phagocytosis requires v 5 integrin but not tyrosine kinases FAK or MerTK. *Mol. Biol. Cell.* 2012; 23:1104–1114. [PubMed: 22262456]
- [16]. Yu CC, Nandrot EF, Dun Y, Finnemann SC. Dietary antioxidants prevent age-related retinal pigment epithelium actin damage and blindness in mice lacking v 5 integrin. *Free. Radic. Biol. Med.* 2012; 52:660–670. [PubMed: 22178979]
- [17]. Adamus G, Zam ZS, Arendt A, Palczewski K, McDowell JH, Hargrave PA. Anti-rhodopsin monoclonal antibodies of defined specificity: characterization and application. *Vision Res.* 1991; 31:17–31. [PubMed: 2006550]
- [18]. Finnemann SC. Focal adhesion kinase signaling promotes phagocytosis of integrin-bound photoreceptors. *EMBO J.* 2003; 22:4143–4154. [PubMed: 12912913]
- [19]. Nandrot EF, Kim Y, Brodie SE, Huang X, Sheppard D, Finnemann SC. Loss of synchronized retinal phagocytosis and age-related blindness in mice lacking v 5 integrin. *J. Exp. Med.* 2004; 200:1539–1545. [PubMed: 15596525]

- [20]. Yermolaieva O, Xu R, Schinstock C, Brot N, Weissbach H, Heinemann SH, Hoshi T. Methionine sulfoxide reductase A protects neuronal cells against brief hypoxia/reoxygenation. *Proc. Natl. Acad. Sci. U. S. A.* 2004; 101:1159–1164. [PubMed: 14745014]
- [21]. Molday RS, Hicks D, Molday L. Peripherin. A rim-specific membrane protein of rod outer segment discs. *Invest. Ophthalmol. Vis. Sci.* 1987; 28:50–61. [PubMed: 2433249]
- [22]. Finnemann SC, Bonilha VL, Marmorstein AD, Rodriguez-Boulan E. Phagocytosis of rod outer segments by retinal pigment epithelial cells requires v 5 integrin for binding but not for internalization. *Proc. Natl. Acad. Sci. U. S. A.* 1997; 94:12932–12937. [PubMed: 9371778]
- [23]. Finnemann SC, Rodriguez-Boulan E. Macrophage and retinal pigment epithelium phagocytosis: apoptotic cells and photoreceptors compete for v 3 and v 5 integrins, and protein kinase C regulates v 5 binding and cytoskeletal linkage. *J. Exp. Med.* 1999; 190:861–874. [PubMed: 10499924]
- [24]. Kwong JQ, Henning MS, Starkov AA, Manfredi G. The mitochondrial respiratory chain is a modulator of apoptosis. *J. Cell Biol.* 2007; 179:1163–1177. [PubMed: 18086914]
- [25]. Nunnari J, Suomalainen A. Mitochondria: in sickness and in health. *Cell.* 2012; 148:1145–1159. [PubMed: 22424226]
- [26]. Marchetti MA, Lee W, Cowell TL, Wells TM, Weissbach H, Kantorow M. Silencing of the methionine sulfoxide reductase A gene results in loss of mitochondrial membrane potential and increased ROS production in human lens cells. *Exp. Eye Res.* 2006; 83:1281–1286. [PubMed: 16934804]
- [27]. Moskovitz J, Bar-Noy S, Williams WM, Requena J, Berlett BS, Stadtman ER. Methionine sulfoxide reductase (MsrA) is a regulator of antioxidant defense and lifespan in mammals. *Proc. Natl. Acad. Sci. U. S. A.* 2001; 98:12920–12925. [PubMed: 11606777]
- [28]. Zhao H, Kim G, Liu C, Levine RL. Transgenic mice overexpressing methionine sulfoxide reductase A: Characterization of embryonic fibroblasts. *Free. Radic. Biol. Med.* 49:641–648. [PubMed: 20510353]
- [29]. Nan C, Li Y, Jean-Charles PY, Chen G, Kreymerman A, Prentice H, Weissbach H, Huang X. Deficiency of methionine sulfoxide reductase A causes cellular dysfunction and mitochondrial damage in cardiac myocytes under physical and oxidative stresses. *Biochem. Biophys. Res. Commun.* 2010; 402:608–613. [PubMed: 20971073]
- [30]. Arnold S. The power of life--cytochrome c oxidase takes center stage in metabolic control, cell signalling and survival. *Mitochondrion.* 2012; 12:46–56. [PubMed: 21640202]

- Levels of MsrA protein regulate ATP content of RPE cells.
- MsrA dependent ATP changes in turn alter the phagocytic activity of RPE cells.
- MsrA's effect on ATP content and phagocytosis is independent of oxidative stress.
- MsrA's effect on ATP content and phagocytosis requires mitochondrial respiration.
- MsrA regulates activity of complex IV but not complex II of the respiratory chain.

Figure 1. MsrA localizes to cytosol and mitochondria in rat RPE in primary culture and the RPE-J cell line

Unpassaged rat RPE in primary culture (A to D) and post-confluent, polarized RPE-J cells (E to H) were stained and analyzed side-by-side. Images show representative single x-y plane confocal microscopy sections acquired using the same acquisition settings for all fields. For each field, the upper row shows MsrA or control image in green, the center row shows mitochondrial marker or control in red, and the lower row shows the merged image. Insets in A, D, E, and H show areas marked by white rectangles in the main fields with additional 2.5-fold magnification. Arrows point out examples of colocalization, which appears yellow. DAPI nuclei stain in blue is also shown in all fields. A and E show cells

incubated live with MitoTracker Red for 15 min followed by fixation in paraformaldehyde and labeling with MsrA antibodies. B and F show control cells incubated with medium instead of MitoTracker. C and G show control cells incubated with MitoTracker Red followed by fixation and labeling only with secondary antibody conjugated to AlexaFluor488. D and H show cells fixed in ice-cold methanol and co-stained with antibodies to MsrA and mitochondrial protein complex-IV subunit I. Scale bars, 10 μ m.

Figure 2. Silencing MsrA decreases POS binding and engulfment by RPE cells

RPE-J cells were treated with control siRNA or MsrA siRNA for 2 days before POS challenge for 3.5 hours and analysis by immunoblotting or viability determination. A shows a representative immunoblot comparing MsrA protein levels in cells treated with control or MsrA siRNA as indicated and incubated with medium (m) or POS (POS). The same blot membrane is shown probed sequentially for MsrA, tubulin as loading control, and POS-opsin as marker for phagocytosed POS as indicated. B shows MsrA protein levels in cells after POS challenge, C and D show quantification of surface-bound POS and internalized POS, respectively. E shows relative cell viability as compared to untreated cells incubated with assay medium (set as 100%) of cells after incubation for 3.5 hours with assay medium

or POS as indicated. (B to E) Untreated RPE-J cells (white bars, /, set as 100% control) were compared to cells treated with control siRNA (gray bars, si-c) or MsrA siRNA (black bars, si-M). All bars show mean $\pm$ S. D. of three independent experiments with triplicate samples each. One-way ANOVA revealed significant differences among samples ($P < 0.05$) for data in B to D but not for E. For B to D, Bonferroni post-hoc comparison showed significant difference of cells treated with MsrA siRNA to both untreated and control siRNA treated cells. Control siRNA treated cells did not differ from untreated cells. Asterisks indicate significant difference to untreated cells.

Figure 3. MsrA overexpression increases POS binding and engulfment by RPE cells

RPE-J cells were infected with recombinant adenovirus encoding either -galactosidase (-gal) as control or MsrA two days before cell fractionation (A and B) or challenge with medium (m) or POS for 3.5 hours (C to E). A shows a representative immunoblot of whole cell lysates (lanes total), mitochondria-depleted supernatant after mitochondrial pelleting (lanes sup) and pelleted cell content enriched in mitochondria (lanes mito). All lanes represent the same number of cells. The same blot membrane is shown probed sequentially for MsrA, -gal, cytoplasmic marker tubulin, and mitochondrial marker porin, as indicated. Bars in B show fold changes in MsrA content in the different fractions analyzed in A harvested from cells overexpressing -gal (gray bars, -g) or MsrA (black bars, M) as compared to MsrA

content of such fractions in uninfected cells, which were set as 1. C shows a representative immunoblot comparing MsrA protein levels in whole cell lysates prepared from cells infected with -gal or MsrA virus as indicated and incubated with medium (m) or POS (POS). The same blot membrane is shown probed sequentially for MsrA, -gal, tubulin as loading control, and POS-opsin as marker for phagocytosed POS as indicated. Bars in D and E show quantification of surface-bound POS and internalized POS, respectively. Untreated RPE-J cells (white bars, /, set as 100% control) were compared to cells infected with -gal virus (gray bars, -g) or MsrA virus (black bars, M). (B, D and E) Bars show mean $\pm$ S. D. of three independent experiments with triplicate samples each. One-way ANOVA revealed significant differences among samples for B, D, and E ($P < 0.05$). Bonferroni post-hoc comparison showed significant difference of cells treated with MsrA siRNA to both untreated and control siRNA treated cells. Control siRNA treated cells did not differ from untreated cells. Asterisks indicate significant difference to untreated cells.

Figure 4. Effects of MsrA silencing or overexpression on POS phagocytosis require an active mitochondrial respiratory chain

RPE-J cells that were untreated (white bars, /), treated with control siRNA (gray bars, si-c), infected with control -gal virus (gray bars, -g), treated with MsrA siRNA (black bars, si-M) or infected with MsrA virus (black bars, M) were challenged for 3.5 hours with POS in the presence of the water-soluble antioxidant trolox (A), hydrogen peroxide (B), mitochondrial respiratory chain inhibitors antimycin (C) or oligomycin (D) at concentrations as indicated before quantification of internalized POS by fluorescence scanning. Bars show relative POS internalization of cells compared to POS internalization by untreated cells, which was set as 1. Bars show mean $\pm$ S. D. of three independent experiments with triplicate samples each.

For A to D, one-way ANOVA revealed significant differences among samples ($P < 0.05$). For A and B, Bonferroni post-hoc comparison of select samples showed significant difference of cells treated with MsrA siRNA to both untreated and control siRNA treated cells, and of MsrA virus treated cells to both control virus and untreated cells. Control siRNA treated cells and control virus treated cells did not differ from untreated cells. For C and D, Bonferroni post-hoc test of select samples showed significant difference between untreated cells and MsrA siRNA treated cells. However, this difference was not directly related to MsrA reduction as MsrA siRNA treated cells did not differ significantly from control siRNA treated cells. There was no significant difference of untreated cells to control siRNA treated cells, to control virus treated cells, or to MsrA virus treated cells. For A to D, individual asterisks indicate significant difference to untreated cells. Arrows above bars connected by black lines show significant difference between control and MsrA specific treatment. n.s. above bars connected by black lines shows lack of difference between control and MsrA specific treatment. E shows representative immunoblots of cells treated as above (lanes marked as for bar graphs), and lysed after POS challenge for 3.5 hours in the presence of trolox, hydrogen peroxide, antimycin, or oligomycin, as indicated. Solvent control for the upper panel was 1% water, solvent control for the lower panel was 0.5% ethanol. Upper and lower set of panels each shows an individual blot membrane probed sequentially for MsrA, -gal, and tubulin as loading control, as indicated. Band densitometry did not reveal differences in MsrA content depending on treatment (data not shown).

Figure 5. MsrA silencing decreases RPE cell viability in response to oxidative stress and this increased sensitivity is not altered by inhibitors of the mitochondrial respiratory chain but is alleviated by antioxidant

RPE-J cells were not pre-treated (white bars, /) or were treated with control siRNA (gray bars, si-c) or MsrA siRNA (black bars, si-M) for 2 days. Numbers of live cells in each sample were determined by fluorescence spectrometry after challenge for 3.5 hours with 50 M (A) or 200 M (B) hydrogen peroxide. In C, cells treated with control siRNA or with MsrA siRNA as indicated received mitochondrial respiration inhibitors antimycin or oligomycin or the water-soluble antioxidant trolox as indicated during the 3.5-hour challenge with 200 M hydrogen peroxide before viability quantification. Bars show relative cell viability compared to viability of cells not treated with siRNA and not challenged with hydrogen peroxide, which was set as 100%. Bars show mean $\pm$ S. D. of three independent experiments with triplicate samples each. One-way ANOVA revealed significant differences among samples ($P < 0.05$) for data in B and C but not for A. For B and C, Bonferroni post-hoc comparison showed significant difference of cells treated with MsrA siRNA to both untreated and control siRNA treated cells. Control siRNA treated cells did not differ from untreated cells. Asterisks indicate significant differences of MsrA siRNA treated cells to control siRNA treated cells.

Figure 6. Altering MsrA protein levels affects ATP content in RPE cells that use mitochondrial respiration

RPE-J cells that were either untreated (white bar, /), treated with control siRNA (gray bars, si-c), infected with control -gal virus (gray bars, -g), treated with MsrA siRNA (black bars, si-M) or infected with MsrA virus (black bars, M) were fed with growth medium containing high (25 mM) glucose 24 hours before harvest (A), or were starved during two 30 minute incubations with carbon source-free medium followed by 1 hour supply with physiological concentrations of glucose (5.6 mM, B) or pyruvate (1 mM, C and D). Select samples were fed with medium in the presence of oligomycin, trolox, or 50 M hydrogen peroxide as indicated. ATP was extracted from harvested cells and quantified. Bars show mean $\pm$ S. D. of three to four independent experiments with triplicate samples each. For A, one-way ANOVA revealed significant differences among samples ($P < 0.05$). Bonferroni post-hoc comparison between select samples showed significant difference of cells treated with MsrA siRNA to both untreated and control siRNA treated cells. Control siRNA treated cells also differed significantly from untreated cells. MsrA virus treated cells differed significantly from both untreated and control virus treated cells. Control virus treated cells did not differ from untreated cells. Individual asterisks mark treated cells that are different from untreated cells, while asterisks above lines indicate significant difference between two treated groups as connected by lines. In B to D, asterisks indicate significant difference of cells overexpressing MsrA compared to cells expressing -gal as determined by Student's *t* test.

Figure 7. Altering MsrA protein levels has no effect on mitochondrial protein content

Mitochondrial protein levels were compared among whole cell lysates of RPE-J cells that were either untreated (/), had received control siRNA or MsrA siRNA (siRNA, si-c, si-M), or had been infected with control -gal or MsrA virus (ADV, -g, M). Immunoblots show detection of proteins as indicated in one representative set of samples, of four independent sets tested.

Figure 8. Altering MsrA protein levels specifically affects activity of complex-IV of the mitochondrial respiratory chain

Enzymatic activity of respiratory chain complex-IV (A) or complex-II (B) was quantified in extracts of RPE-J cells that were either untreated (white bars, /), received control siRNA (gray bars, si-c), control -gal virus (gray bars, -g), MsrA siRNA (black bars, si-M) or MsrA virus (black bars, M). Bars show mean $\pm$ S. D. of four (complex-IV) or three (complex-II) independent experiments with triplicate samples each. For data in A and B, one-way ANOVA revealed significant differences among samples ($P < 0.05$). For A, Bonferroni post-hoc comparison between select samples showed significant difference of cells treated with MsrA siRNA to both untreated and control siRNA treated cells. Control siRNA treated cells also differed significantly from untreated cells. MsrA virus treated cells differed significantly from both untreated and control virus treated cells. Control virus treated cells did not differ from untreated cells. Individual asterisks mark treated cells that are different from untreated cells (/), while asterisks above lines indicate significant difference between two treated groups as connected by lines. For B, Bonferroni post-hoc comparison of select samples showed that none of the treated cells differed from untreated cells and cells with more or less MsrA did not differ in activity from their respective controls, as indicated by

lines stating not significant (n. s.). Significant differences were found between cells treated with MsrA siRNA and cells treated with MsrA virus. As significant differences were also found between cells treated with control siRNA and cells treated with MsrA virus, these differences were not related to MsrA expression.

Figure 9. Halving MsrA protein levels in RPE cells is sufficient to decrease POS phagocytosis and ATP content under conditions that maintain normal cell yield

RPE-J cells were untreated (white bars, /) or treated with 50 nmoles (half the amount used in our previous experiments) control siRNA (gray bars, si-c) or MsrA siRNA (black bars, si-M) and analyzed 2 days later. (A to C) Cells were challenged with medium (m) or with POS particles in medium (POS) for 3.5 hours before lysis and immunoblotting analysis of protein overexpression, POS-opsin marker, and mitochondrial proteins as indicated. A shows a representative blot result. (B to D) Bar graphs compare results from cells treated with control siRNA or with MsrA specific siRNA to untreated control cells, whose results were set as 100%. (B) MsrA protein content was quantified in cells after POS challenge by densitometry. MsrA levels were normalized to tubulin content in each sample. (C) total POS uptake (indicating bound plus internal POS) was established by quantifying POS-opsin marker protein content in cell lysates as shown in A by densitometry. POS content was normalized to tubulin content. (D) ATP content of cells was quantified 24 hours after feeding with growth medium containing 25 mM glucose. All experiments were performed three times independently with duplicate samples per experiment. Bars show mean $\pm$ S. D. (B to D) One-way ANOVA revealed significant differences among samples ($P < 0.05$). Bonferroni post-hoc tests established difference of MsrA siRNA treated cells to control siRNA treated cells and to untreated cells. Asterisks indicate difference to untreated cells.

Figure 10. Unpassaged rat RPE in primary culture respond to MsrA overexpression by increasing ATP content, complex-IV activity, and POS uptake

Primary RPE cells were infected with adenovirus encoding β -gal or MsrA and analyzed 2 days after infection. (A) Cells were challenged with medium (m) or with POS particles in medium (POS) for 1.5 hours before lysis and immunoblotting analysis of protein overexpression, POS-opsin marker, and mitochondrial proteins as indicated. (B to E) All bar graphs compare results from cells expressing β -gal (gray bars, -g) with results from cell overexpressing MsrA (black bars, M). (B) ATP content was quantified in cells that had been depleted from carbon sources during two 30-minute incubations with glucose-free medium before incubation with 1 mM pyruvate in glucose-free medium for 1 hour. (C and D)

complex-IV and complex-II activity was quantified in whole cell extracts. Bars show activity of MsrA overexpressing cells relative to activity of cells expressing -gal, which is set as 1. (E) total POS uptake (indicating bound plus internal POS) was established by quantifying POS-opsin marker protein content in cell lysates as shown in A by densitometry. POS content was normalized to tubulin content and set as 1 for control cells expressing -gal. All experiments were performed three times independently with duplicate samples per experiment. Bars show mean $\pm$ S. D. Asterisks indicate statistical difference of cells overexpressing MsrA from cells expressing -gal as determined by Student's *t* test.