Concomitant medications:

- All the patients were required to be receiving all the current guideline directed medications. Consequently 99.3% were on aspirin, and 96% on clopidogrel. Amongst the adjuvant anti-thrombotic therapy 33% received unfractionated heparin, 3% received bivalirudin, and 10.9% received fondaparinux; GpIIb-IIIa inhibitors were given to 25% of patients.

Principle findings:

- A total of 7021 patients were enrolled, 3507 to radial access and 3514 to femoral access.
- About 28% presented with ST-segment elevation myocardial infarction (STEMI), 22% had diabetes, and 67% underwent PCI, while 8% underwent CABG.
- The success rates were similar and high in both the arms (95.3%). The operators were fairly high volume and trained with a median annual PCI rate of 300 out of which approximately 40% were radial PCIs.
- The first important finding was a relatively high cross-over rate. The rate was 7.6% in the radial arm compared to 2% in the femoral arm and this difference was statistically significant ($P < 0.0001$).
- There was no difference in the primary quadruple end point of death, MI, stroke, and non-CABG related major bleeding (3.7% for radial vs 4% for femoral access).
- The secondary end point of death, MI, or stroke were similar in the two arms (3.2% vs 3.2%).
- As there was a very high interest in the safety end point of bleeding it was analysed in detail. There was no difference in bleeding if non-CABG major bleeding was taken as a whole (0.7% for radial vs 0.9% for femoral). However, as expected major vascular access site complications (1.4% vs 3.7%; $P < 0.0001$) and ACUTY non-CABG major bleeding (1.9% vs 4.5%, $P < 0.0001$) were significantly lower in the radial access arm.
- Another area of concern with the radial access is the speed. The study showed no significant difference in the overall PCI time between the two arms (35 minutes vs 34 minutes). However, the total fluoroscopy time was significantly different between the two (9.3 minutes vs 8 minutes, $P < 0.0001$).
- The most significant parameter where radial access scored impressively over femoral access was patient satisfaction. Patient preferred access site for next procedure was almost twice as high with radial access (90% vs 49%, $P < 0.0001$).

Interpretation

Where do the results of this study leave us? The results are fairly predictable. In experienced hands both sites lead to high and equal success rates. There is no increase in the total procedure time with radial access although there is a significant increase in total fluoroscopy time making cumulative radiation exposure an important safety issue for operators who opt for a very high or exclusive radial practice over a prolonged period of time. Although overall bleeding is same but vascular site bleeding is an important issue and radial access clearly establishes its superiority over femoral access. The issue which probably will drive the future increase in use of radial access as shown in this study is patient satisfaction. There is a substantial difference of having to lie down for hours with pressure and sheaths in areas which may be discomforting and embarrassing for many patients and being able to move about immediately with a small band tied to the arm.

Perspective

Like many other “this or that” situations the message here is also “this and that rather than this or that”. Interventional cardiologists should become proficient in both the accesses and select the access on an individual basis keeping in mind the speed, bleeding risk and of course the patient preference. Contrary to the expectation that radial route will be safer for patients, this study failed to show increased safety for radial route. The main reason for this is that the non-access bleeds were nearly equal in both the arms. This shows that our search for further means of reducing bleeding complications should continue.

Contributed by
Abid Hussain, Upendra Kaul
 Fortis Escorts Heart Institute,
 New Delhi, India.

1. **Rivaroxaban in patients with a recent acute coronary syndrome.** Mega JL, Braunwald E, Wiviott SD, ATLAS ACS 2–TIMI 51 Investigators. *N Engl J Med* 2012;366:9–19. doi: 10.1016/S0019-4832(12)60037-6

Background: Acute coronary syndromes (ACS) arise from coronary atherosclerosis with superimposed thrombosis. Since factor Xa plays a central role in thrombosis, the inhibition of factor Xa with low-dose rivaroxaban might improve cardiovascular outcomes in patients with a recent ACS.

Methods: In this double-blind, placebo-controlled trial, we randomly assigned 15,526 patients with a recent ACS to receive twice-daily doses of either 2.5 mg or 5 mg of rivaroxaban or placebo for a mean of 13 months and up to 31 months. The primary efficacy end point was a composite of death from cardiovascular causes, myocardial infarction (MI), or stroke.

Results: Rivaroxaban significantly reduced the primary efficacy end point, as compared with placebo, with respective rates of 8.9% and 10.7% (hazard ratio in the rivaroxaban group, 0.84; 95% confidence interval [CI], 0.74–0.96; $P=0.008$), with significant improvement for both the twice-daily 2.5-mg dose (9.1% vs 10.7%, $P=0.02$) and the twice-daily 5-mg dose (8.8% vs 10.7%, $P=0.03$). The twice-daily 2.5-mg dose of rivaroxaban reduced the rates of death from cardiovascular causes (2.7% vs 4.1%, $P=0.002$) and from any cause (2.9% vs 4.5%, $P=0.002$), a survival benefit that was not seen with the twice-daily 5-mg dose. As compared with placebo, rivaroxaban increased the rates of major bleeding not related to coronary artery bypass grafting (2.1% vs 0.6%, $P<0.001$) and intracranial haemorrhage (0.6% vs 0.2%, $P=0.009$), without a significant increase in fatal bleeding (0.3% vs 0.2%, $P=0.66$) or other adverse events. The twice-daily 2.5-mg dose resulted in fewer fatal bleeding events than the twice-daily 5-mg dose (0.1% vs 0.4%, $P=0.04$).

Conclusion: In patients with a recent ACS, rivaroxaban reduced the risk of the composite end point of death from cardiovascular causes, MI, or stroke. Rivaroxaban increased the risk of major bleeding and intracranial haemorrhage but not the risk of fatal bleeding.

Perspective

Since coronary artery disease especially in the form of ACS is a prothrombotic state, it makes great logic to use long-term oral anti-coagulation. However, in the presence of dual antiplatelet therapy, the so called triple therapy it is fraught with high-risk of bleeding complications but in settings like a patient with atrial fibrillation who undergoes stenting, it becomes mandatory to use triple therapy for variable periods of time. Earlier meta-analysis revealed improved cardiovascular outcomes with addition of warfarin to aspirin in ACS patients but with increased bleeding complications. The ATLAS ACS TIMI 51 trial is a further extension to use triple therapy in patients with recent ACS in general. Rivaroxaban is an oral anti-coagulant that directly and selectively inhibits factor Xa. Aspirin was used in 98% of the patients and thienopyridin in 93% of patients. Rivaroxaban added to dual antiplatelet therapy following ACS significantly reduced the primary end point of cardiovascular death, myocardial infarction and stroke at a mean follow-up of 13 months. The lower dose (2.5 mg twice daily) used in one arm also showed a survival benefit. Although there was increased incidence of major bleeding including intracranial haemorrhage, there was no fatal bleeding with use of add-on rivaroxaban. The result was irrespective of the mode of presentation (ST-segment elevation myocardial infarction [STEMI], non-STEMI [NSTEMI] or unstable angina). Index event was a STEMI in 50.3% of the patients, an NSTEMI in 25.6% and unstable angina in 24.0%. The mean time from the index event to randomisation was 4.7 days. The mean duration of treatment was 13.1 months. Rivaroxaban reduced the risk of stent thrombosis as well. The benefit with rivaroxaban was consistent among the sub-groups except for patients with a history of stroke or transient ischaemic attack. There was no evidence of increased liver toxicity. Earlier in the ATLAS ACS TIMI 46 and APPRAISE 1 trials, rivaroxaban and apixaban showed trends towards a reduction in cardiovascular events. However, further studies are required to have this reproducible result in varied subset of ACS patients with other co-morbid conditions. We need a large trial which compares rivaroxaban to warfarin in the presence of dual antiplatelet therapy in post ACS patients before this new idea is accepted.

2. **Effect of two intensive statin regimens on progression of coronary disease.** Nicholls SJ, Ballantyne CM, Barter PJ. *N Engl J Med* 2011;365:2078–87. doi: 10.1016/S0019-4832(12)60038-8

Background: Statins reduce adverse cardiovascular outcomes and slow the progression of coronary atherosclerosis in proportion to their ability to reduce low-density lipoprotein (LDL) cholesterol. However, few studies have either assessed the ability of intensive statin treatments to achieve disease regression or compared alternative approaches to maximal statin administration.

Methods: We performed serial intravascular ultrasonography in 1039 patients with coronary disease, at baseline and after 104 weeks of treatment with either atorvastatin, 80 mg daily, or rosuvastatin, 40 mg daily, to compare the effect of these two intensive statin regimens on the progression of coronary atherosclerosis, as well as to assess their safety and side-effect profiles.

Results: After 104 weeks of therapy, the rosuvastatin group had lower levels of LDL cholesterol than the atorvastatin group (62.6 mg/dL vs 70.2 mg/dL [1.62 mmol/L vs 1.82 mmol/L], $P<0.001$), and higher levels of high-density lipoprotein (HDL) cholesterol (50.4 mg/dL vs 48.6 mg/dL [1.30 mmol/L vs 1.26 mmol/L], $P=0.01$). The primary efficacy end point, percent atheroma volume (PAV), decreased by 0.99% (95% confidence interval [CI], –1.19 to –0.63) with atorvastatin and by 1.22% (95% CI, –1.52 to –0.90) with rosuvastatin ($P=0.17$). The effect on the secondary efficacy end point, normalised total atheroma volume (TAV), was more favourable with rosuvastatin than with atorvastatin: –6.39 mm³ (95% CI, –7.52 to –5.12), as compared with –4.42 mm³ (95% CI, –5.98 to –3.26) ($P=0.01$). Both agents induced regression in the majority of patients: 63.2% with atorvastatin and 68.5% with rosuvastatin for PAV ($P=0.07$) and 64.7% and 71.3%, respectively, for TAV ($P=0.02$). Both agents had acceptable side-effect profiles, with a low incidence of laboratory abnormalities and cardiovascular events.

Conclusion: Maximal doses of rosuvastatin and atorvastatin resulted in significant regression of coronary atherosclerosis. Despite the lower level of LDL cholesterol and the higher level of HDL cholesterol achieved with rosuvastatin, a similar degree of regression of PAV was observed in the two treatment groups.