Letters to the Editor

Robert J. Snyder, DPM, CWS^{a,*}

^a*Medical Director, Wound Healing Center at University Hospital, Tamarac, Florida, USA*

Question:

Is there any place for whirlpool in wound care nowadays?

*Milton Fechter
New York, NY*

Answer: Whirlpool (WP) or hydrotherapy has seen a gradual but significant decline in wound management over the past decade. Some of the reasons for the decline are related to costs associated with the following factors:

1. Purchase cost of the equipment
2. Cost of maintaining the equipment
3. Cost of water
4. Cost of disinfecting the whirlpool tanks
5. Costs of getting the patient to and from the therapy
6. Cost of time it takes to treat the patient

Other reasons for decline of whirlpool therapy for wounds include the following:

1. Risk of cross-contamination between patients who use the same whirlpool equipment
2. Less time efficient and wound treatment effective than other methods (eg pulsed lavage with vacuum) used to irrigate, cleanse, reduce microorganisms, reduce solubilized slough and thick exudates
3. Very difficult to take WP bedside compared with pulsed lavage with vacuum
4. No vacuum with WP
5. Can't control WP jet pressure to safe level As mentioned in my first article on the role of physical therapy in wound management, I will be addressing many other

wound interventions (including but not limited to WP) in subsequent issues of the JACCWS.

*Luther C. Kloth, PT, MS, FAPTA, CWS, FACCWS
Emeritus Professor in Physical Therapy
Department of Physical Therapy
Marquette University*

Question:

In patients with mixed arterial and venous pathology, what are your guidelines for percutaneous perforator treatment? Do you use a scale like the ankle-brachial index, with a cutoff at a certain level to decide whether the patient is a candidate for the VNUS procedure? How do you decide whether the procedure is warranted in the presence of arterial disease in the patient with an incompetent perforator?

*Robert Pachecho, MD
Miami, FL*

Answer: Thanks for your question. The treatment of perforator disease uses the general indications of the presence of incompetent perforator veins (usually defined as vein closure time >0.5 seconds and size >3 mm or size >4 mm regardless of flow dynamics) and a clinical-etiology-anatomy-pathophysiology (CEAP) class 4, 5, or 6. Other indications include high suspicion of pain secondary to the perforator vein or if the perforator vein is the likely source of varicose veins with a CEAP class of at least 2. In regard to mixed disease, arterial disease should be considered if an ulcer is present and has poor healing or if the patient is unable to tolerate compression because of suspected ischemic pain. Clinical suspicion should prioritize the more pressing problem, ie, address the arterial disease first if limb threat exists. Certainly, mixed disease can be more difficult to treat,

* Corresponding author.

E-mail address: DrWound@aol.com

but a systemic approach should yield the best results. Hope this helps.

Honesto Poblete, MD
Center for Vein Disease
Mt. Sinai School of Medicine

Question:

In a patient with mixed venous-arterial disease and who has incompetent perforators causing gravity-dependent swelling in both ankles, at what point would an intermittent pneumatic compression (IPC) pump become a challenge or a problem for the 'patient's arterial condition? Would you suggest a specific ankle-brachial index cutoff? In my practice, we use a lot of IPC pumps for gravity-dependent swelling when the patient cannot use compression bandages for various reasons. The question comes up with mixed disease.

Brenda Jackson, MD
Newark, NJ

Answer: This is a great question as mixed cases can be very challenging to manage! To begin with, the patient would need to have been thoroughly evaluated with vascular testing. At what level are the incompetent perforators? What has been done (if anything is possible) to correct them? I would also want to make sure the patient does not have any grafts or stents prior to the application of IPC. As far as an ankle-brachial index cutoff, I would recommend 0.8. Caution always needs to be employed when dealing with arterial insufficiency (AI). If IPC is being used, I recommend starting with low pressure, 50 mm Hg, then slowly stepping it up based on the patient's tolerance to no more than 125 mm Hg. I am more inclined to recommend the use of low compression systems (elasticated bandages such as Tubigrip or custom low compression stockings) supplemented with walking and exercise.

Heather Hettrick, PT, PhD, CWS, MLT, FACCWS
American Medical Technologies
Director of Clinical Education

Question:

CMS mandated that effective November 2008, products that contain papain/urea and papain/urea/chlorophyllin complex sodium could no longer be manufactured, and effective January 21, 2009, these products could no longer be shipped in interstate commerce. Now that these products have been removed from the market, what other chemical debridement options exist?

Leslie Cohen
Queens, NY

Answer: Many options for debridement exist. To address chemical debridement specifically, topical collagenase ointments are enzymes that are indicated for the debridement of chronic dermal ulcers and severely burned areas. Collagenase works by digesting collagen in necrotic tissue. Hypertonic

saline gels soften and debride dry and moist necrotic tissue to create an environment for wound healing. The high-concentration sodium gel creates a hypertonic environment and draws drainage from the wound to naturally soften eschar.

Kari Bechert, MPT
ACCWS

Question:

I work as a physician assistant (PA) in orthopedics. Please tell me what roles PAs play in chronic wound care settings?

Lisa Abraham, PA-C
Long Island, NY

Answer: Currently, PAs are not often involved in wound care. However, there is plenty of opportunity. Wound care clinics need practitioners who can debride wounds, prescribe treatments and medications, and make diagnostic and therapeutic decisions. The PA is ideally suited to these tasks, especially working under a certified wound care physician. Local specialists (vascular surgeons, orthopedists) often have an interest in wound care but are unable to devote their full time to its practice. A PA working with one of these specialists would allow the specialist to participate in the practice but maintain a full surgical schedule. You can begin by taking a wound care course (such as the American College of Certified Wound Specialists Back-to-Basics course), attending a national wound care meeting, and looking into wound care certification (such as the certified wound specialist certification by the American Academy of Wound Management). Then find a local physician who does wound care and talk to him or her about the possibilities.

Wayne C. Stuart, MD, CWS
Chair Elect, ACCWS
Assistant Professor
Physician Assistant Program, DeSales University

Question:

In your article on "Wound Closure and the Reconstructive Ladder in Plastic Surgery," you discuss the need to minimize tension on the closure sight and the fact that swelling can be a factor in flap or skin graft failure. I have seen this occur especially in patients who have procedures for wound closure on the lower extremities where edema is more likely to be problematic and elevation often difficult and inadequate. Can you discuss the methods of edema management that you have found successful in preventing edema in lower extremities? Are you comfortable with compression bandages, such as short stretch or multilayer systems, for edema management after a flap or skin graft, provided that the patient has adequate perfusion? Congratulations on an informative article in the first edition of the *Journal of the American College of Certified Wound Specialists!*

Rose Hamm, DPT, CWS
Los Angeles, CA

Answer: Thank you for your question on edema management in postoperative flap surgery. Edema occurs after any surgery to a certain extent. However, in traumatic injury to the lower extremity with extensive soft tissue damage, a significant edema is more likely to take place. Immediate flap reconstructive surgery is not recommended and should be postponed until partial or complete resolution of the edema is observed. Edema is harmful to healing, not only by the increased tension on the closure site, but also by the decrease of oxygen and nutrient supply to the wound. Adding compression therapy to the flap site to treat edema may decrease blood perfusion, which may lead to necrosis. Increased tension on any surgical closure may lead to skin necrosis, infection, and wound dehiscence. To treat postoperative flap edema, I would assure good fluid resuscitation, electrolyte balance, good nutrition, and if possible, elevation. Applying pressure on skin grafts, on the other hand, is not harmful to the graft and is even encouraged to allow graft take.

Richard Simman, MD, CWS, FACCWS

Plastic and Reconstructive Surgery

Associate Professor, Wright State School of Medicine

Question:

A patient recently came in after walking into her wheelchair and gashing open her leg. It was a jagged but more or less linear cut that was pretty deep. She had done this the night before and came to our office about 11:00 AM. That would make the time from trauma about 13 hours. My question is, how long after this type of trauma can the wound be closed primarily by suturing?

Linda Blackburn, RN

Newark, NJ

Answer: Thank you for your question. The window for acute wound closure depends on the degree of contamination and the mechanism of injury of the wound. Human and dog bites, for example, are considered highly contaminated. When located on the torso and extremities, they are cleaned vigorously and approximated loosely, and the patients are placed on oral antibiotics and dressing changes with frequent follow-ups. When located on the face, these wounds are cleaned and closed nicely. Because of the increased vascularity, facial wounds heal better. Scarring disfigurement is more significant on the face, and normal closure for better aesthetic outcome warrants the risk of infection that may result from tight closure. In plastic surgery, Mohs defects referred by dermatologists after basal and squamous cell carcinomas may be closed days later without high risks of infection. These wounds are kept clean and covered with topical antibiotics and dressing, and the patients are kept on oral antibiotics until wound closure.

In your case, you describe a clean wound with no crush injury to wound edges 13 hours after the accident. I would anesthetize the wound, vigorously clean it with soap and

saline or other antiseptic solution, and then close it nicely. Keep the patient on oral antibiotics with frequent office visits to detect any early sign of infection.

Richard Simman, MD, CWS, FACCWS

Plastic and Reconstructive Surgery

Associate Professor, Wright State School of Medicine

Question:

The article on wound bed prep combines sharp debridement and surgical debridement under the same category. Some older journals have them classified as separate categories. Why have they been grouped together?

Simon Young

Nyack NY

Answer 1: When I read Chuck Gokoo's article, it is hard to say whether he separated or combined them (sharp and surgical debridement), and he actually differentiated between sharp debridement and surgical debridement, but I do not know that they are really separate or not. All of the debridement codes are surgical procedures, but more extensive for the surgical, OR codes. I think that it has to do with the magnitude of the procedure, but I think that they are both surgical debridement. The bedside debridement, in which there may be still some slough or eschar remaining, a thin rim as he says, is more like what is done on a regular basis at the wound care center. This may leave some tissue behind requiring autolytic debridement but also tends to be a more selective form of debridement because it leaves living tissue behind at the expense of also possibly leaving some nonviable tissue behind. They are grouped together because they are all surgical codes, but the goals represented by the different types of surgical codes are different.

Jane Fore

Answer 2: *Sharp and surgical debridement* are usually used synonymously.

However, surgical debridement can be viewed as excisional while sharp could include scraping the wound surface partially, fully, or subcutaneously without excising the wound edges.

Medicare has different codes for each.

Robert Snyder

Comment:

A recent article¹ in this publication perpetuates two longstanding myths about the action of Unna's boot (UB). This device is very effective at eliminating leg edema, but not by "compression" and not by action of a "muscle pump." About 30 years ago, Dr. H.I. Lippman of our clinic supervised unpublished measurements demonstrating that the cross-sectional area of the ankle diminishes when it is flexed (as when walking). This is analogous to what happens to a full

toothpaste tube when the cap is removed and the tube is rapidly bent. The result of bending the tube is rapid ejection of toothpaste. Anyone can replicate Dr. Lippman's measurements by using a simple tape measure.

If the ankle and adjacent foot and leg from the base of the toes to the popliteal fossa are wrapped in the flexible but nonstretchable UB, we have an analogue to the toothpaste tube. Whenever the ankle is briskly flexed (as by ordinary walking), fluid will be rapidly ejected from the leg, thereby effectively reducing edema. Thus it would be appropriate to refer to this device as an *ankle pump*, not a *muscle pump* or *compression system*. Compression systems include stretchable devices such as surgical hose and stretch bandages—not the UB, which, like toothpaste tubes, works by virtue of its *nonstretchability*.

The UB will not remove edema if the ankle is not flexed. Thus, for patients who cannot walk or flex their ankles, the UB will be useless unless an aide passively flexes the affected ankle for several hours every day.

We use the UB as the major treatment for venous stasis ulcers of the legs and find it to be virtually 100% effective in curing these wounds.

1. Hettrick H. The science of compression therapy for chronic venous insufficiency edema. *J Am Coll Cert Wound*

Care Specialists. 2009;1(1):20-27.

Richard K. Bernstein, MD, FACE, FACN, FCCWS

Peripheral Vascular Disease Clinic

Jacobi Medical Center

Albert Einstein College of Medicine