CASE REPORT

Nodular kidney involvement in a patient with idiopathic Sweet's syndrome

Reem Daloul,¹ Sagar Gupta,¹ Maria Marin²

¹Department of Internal Medicine, William Beaumont Hospital, Royal Oak, Michigan, USA

²Department of Nephrology, William Beaumont Hospital, Royal Oak, Michigan, USA

Correspondence to

Dr Reem Daloul,
Reem.Daloul@beumont.edu

SUMMARY

A 40-year-old previously healthy woman presented with a history of painful maculopapular rash, fever and diffuse arthralgia. The skin biopsy was consistent with Sweet's syndrome. Further workup with abdominal CT revealed bilateral hypoenhancing renal lesions involving the cortex and medulla, as well as terminal ileal thickening. Her renal function was preserved. The urinalysis revealed microscopic haematuria and sterile pyuria. Upper and lower gastrointestinal endoscopies were unremarkable and terminal ileal biopsy was normal. The evaluation for secondary causes of Sweet's syndrome was negative. She responded to corticosteroid therapy. A repeat abdominal CT scan two months later showed complete resolution of renal and ileal lesions. One year after the initial presentation the patient is doing well with no recurrent symptoms. We concluded that the abnormal renal findings were most likely due to sterile abscesses in the course of idiopathic Sweet's syndrome.

BACKGROUND

Renal involvement in Sweet's syndrome is uncommon. It usually manifests as proteinuria. Less common findings are haematuria, renal insufficiency and mesangiocapillary glomerulonephritis. We believe this might be the first reported case of mass-like involvement of the kidneys in the course of Sweet's syndrome.

CASE PRESENTATION

A 40-year-old woman was in her usual state of health until 2 weeks prior to her hospital admission. She initially noticed a non-pruritic, painful, vesiculopapular eruption at the posterior aspect of her left shoulder which then became maculopapular before it spread to the upper part of her back and both of the upper and lower extremities. The rash did not involve her palms or soles. She denied insect bites or any sick contacts. Two days following the rash onset, she noticed a high fever up to 102° F and severe diffuse arthralgia that was limiting her daily activities.

The patient was empirically treated with doxycycline for a possible Lyme disease but fever and arthralgia continued to worsen. Acyclovir was subsequently added and biopsy of her shoulder and calf lesions was performed. The patient noted no improvement in her symptoms and ibuprofen provided minimal relief. She presented to the emergency department.

Her medical and surgical histories were negative. Family history was significant for type 2 diabetes mellitus in her mother and both grandmothers.

There was no history of tobacco use and she consumed alcohol only occasionally. She had two normal pregnancies and no medication allergies. Prior to the current disease she was only taking a multivitamin per day. At the time of admission, she was on ibuprofen, doxycycline and acyclovir.

On physical examination, she was alert and oriented, well-nourished and in no acute distress. Her blood pressure was 128/75 mm Hg with no orthostatic drop, pulse 84/min, temperature 96.8° F, respiratory rate 20 and oxygen saturation 100% on room air. Skin examination revealed multiple 1.5–10 cm areas of erythematous maculopapular skin lesions that were tender to touch. There was mild erythema and warmth over both ankles with no effusion or limitation in the range of motion. She had tenderness over lateral epicondyles bilaterally, knee joint margins and anserine bursae. There was no lymphadenopathy or hepatosplenomegaly. The rest of her physical examination was unremarkable.

INVESTIGATIONS

The complete blood count showed white cell count of 9.2 billion/L (neutrophils 6.2, lymphocytes 2.0, monocytes 0.8, eosinophils, 0.1 and immature granulocyte 0.02). The haemoglobin level was 12.2 g/dL and platelets 313 billion/L. Electrolytes were normal. Her creatinine measure was 0.59 mg/dL with blood urea nitrogen 8 mg/dL. Liver function testing was normal. The erythrocyte sedimentation rate was 66 mm/h (0–18). Immunoglobulin G and immunoglobulin M (IgM) antibody for Lyme disease, Venereal Disease Research Laboratory, hepatitis B core antibody IgM and hepatitis B virus (HBV) surface antibody and antigen, hepatitis C virus (HCV) antibody, parvovirus B19 IgM antibody and antistreptolysin O returned negative or normal. Antinuclear antibody, double-stranded DNA, antineutrophil cytoplasmic antibody, Sjogren SSA and SSB antibody, rheumatoid factor, cyclic citrullinated peptide, complements C3 and C4 and ACE were within normal range. The urinalysis revealed 2+ blood, negative for protein, with 4 red blood cells/hpf and 9 white cell count/hpf and 35 squamous epithelial cells/lpf. Urine culture was negative. Urine protein to creatinine ratio was 0.1.

The skin biopsy revealed a moderately dense mixed inflammatory cell infiltrate displaying a perivascular and interstitial distribution in the papillary and superficial reticular dermis with a smaller amount of inflammation noted in the deeper reticular dermis in a predominantly perivascular distribution.

CrossMark

To cite: Daloul R, Gupta S, Marin M. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2013-010075

The infiltrate was composed mostly of neutrophils, histiocytes and lymphocytes suggestive of Sweet's syndrome.

Imaging data included chest X-ray, which was normal, and chest, abdomen and pelvis CT scan, which demonstrated multiple hypoenhancing lesions involving renal cortex and medulla bilaterally (figure 1A–C) and terminal ileal wall thickening. The colonoscopy was normal except for non-bleeding internal haemorrhoids. Terminal ileum was grossly normal and biopsy was negative, including evaluation by flow cytometry. A kidney biopsy was initially planned to investigate the nature of the hypoenhancing lesions. Given the skin biopsy findings consistent with Sweet's syndrome and negative infectious workup, decision was made to defer renal biopsy and start treatment.

DIFFERENTIAL DIAGNOSIS

Sweet's syndrome is characterised by a constellation of painful well demarcated skin lesions, fever and leukocytosis. A wide spectrum of mucocutaneous and systemic disorders may present with similar dermatological and systemic manifestations. Possibilities include autoimmune diseases, vasculitis, infections (Lyme disease, Syphilis, HBV, HCV, HIV, parvovirus B19 and streptococcal infection), drug eruptions, pyoderma gangrenosum, erythema multiforme and erythema nodosum.

Differential diagnosis of hypoenhancing renal nodules on CT include metastatic lesions, lymphoma, infiltrative diseases, infectious aetiologies such as tuberculosis and inflammatory/vasculitis processes.

TREATMENT

The patient was treated with intravenous methylprednisone 40 mg daily for 4 days, followed by oral prednisone 40 mg/day. This was tapered and discontinued after 1 month. She had immediate response to the treatment, with resolution of her fever, joint pain and rash.

OUTCOME AND FOLLOW-UP

The patient was seen in a follow-up visit 3 weeks after hospital discharge. At that time she was clinically doing well on prednisone 20 mg/day. Her renal function was normal and the urinalysis was negative for blood, protein and leucocytes. A repeat abdominal CT scan 2 months after the initial presentation demonstrated a complete resolution of renal nodules (figure 2A,B) and normal terminal ileum. Given complete resolution of the hypoenhancing kidney lesions, the invasive procedure of a renal biopsy was no longer indicated. One year after the initial presentation, the patient has stayed in remission with no recurrent symptoms.

DISCUSSION

Sweet's syndrome falls under the spectrum of clinical entity commonly referred to as 'neutrophilic dermatoses' which comprises Sweet's syndrome, pyoderma gangrenosum, subcorneal pustular dermatosis and erythema elevatum diutinum. Originally described by Dr Robert Douglas Sweet in 1964, it is characterised by fever, neutrophilia and tender well-demarcated erythematous skin lesions (papules, nodules and plaques). When biopsied, the skin lesions reveal a mature neutrophilic infiltrate in the upper dermis and papillary dermal oedema.

Sweet's syndrome can present in several clinical settings: classical (or idiopathic), malignancy associated or drug-induced. The pathophysiology is not well established and many theories abound, including immune complex reaction, superantigen activation, abnormal function of neutrophils or role of cytokines-interleukin (IL-1, IL-3, IL-8) or granulocyte/macrophage colony stimulating factor.¹

Figure 1 (A–C) Abdominal CT scan showing hypoenhancing renal lesions.

The diagnosis of this dermatosis warrants a detailed investigation to look for a possible underlying malignancy as it can sometimes be the cutaneous harbinger of either an undiagnosed visceral

Figure 2 (A and B) Follow-up abdominal CT scans showing a complete resolution of the hypoenhancing lesions.

malignancy or cancer recurrence. Most common malignancies are acute myelogenous leukaemia and solid tumours arising in the genitourinary organs, breast and gastrointestinal tract.

Although mainly characterised by mature neutrophil cutaneous infiltration, Sweet's syndrome and other neutrophilic dermatoses represent more of a systemic disease as neutrophils may infiltrate almost any organ in the body. Extracutaneous manifestations are most commonly reported in the eyes, mouth, bone, intestine, muscles, spleen and lungs.

Renal involvement is uncommon and has been reported in the context of both idiopathic and malignancy-associated Sweet's syndrome. It most commonly presents as proteinuria, and less often as haematuria^{2–4} and renal insufficiency.^{4–5} Biopsy-diagnosed mesangiocapillary glomerulonephritis has also been described.^{5–8} Renal involvement usually parallels the clinical course of skin lesions.⁹ Steroid therapy results in complete resolution of proteinuria and restoration of normal kidney function in most patients.¹⁰ Some patients may continue to have worsening kidney function requiring a long-term dialysis.¹¹

We believe that nodular or mass-like involvement of the kidneys has not been reported previously. Nodular involvement of other intra-abdominal viscera, most commonly the spleen,^{12–14} but also the liver, pancreas, lymph nodes and digestive tract has been

observed.¹⁴ These lesions were found to be sterile neutrophilic abscesses as biopsies showed neutrophilic infiltration without vasculitis and negative microbiological cultures. Such abscesses may occur during the acute phase of Sweet's syndrome, months to years prior to cutaneous manifestations or during episodes of recurrence. A challenging entity of systemic sterile neutrophilic abscess in the absence of cutaneous involvement has also been reported.¹⁴

Similar to cutaneous lesions, sterile neutrophilic abscess have dramatic response to steroid and may recur following steroid taper with or without recurrence of skin lesions. Though no histological diagnosis was made for the kidney lesions encountered in our patient, the clinical course and resolution with steroids treatment are highly suggestive of sterile neutrophilic abscess as the underlying pathology. Other possible differential diagnoses including infectious and inflammatory aetiologies were excluded prior to the initiation of steroid treatment.

Learning points

- ▶ Sweet's syndrome is characterised by fever, neutrophilia and tender erythematous skin lesions. It can be idiopathic, malignancy-associated or drug-induced.
- ▶ Extracutaneous manifestations are common and can potentially affect every organ in the body. Renal involvement most commonly presents as proteinuria. Less common presentations include haematuria, renal failure, biopsy-diagnosed mesangiocapillary glomerulonephritis and mass-like lesions.
- ▶ Recognition of the dermatological and/or the extracutaneous manifestations can save the patient from unnecessary testing and delays in treatment.
- ▶ Systemic corticosteroids remain the mainstay treatment and often lead to dramatic improvement in signs and symptoms of the disease.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Kluger N, Gil-Bistes D, Guillot B, *et al.* Inflammatory cells, cytokines and matrix metalloproteinases in amicrobial pustulosis of the folds and other neutrophilic dermatoses. *Dermatology* 2011;222:123–7.
- 2 Gunawardena DA, Gunawardena KA, Ratnayaka RMRS, *et al.* The clinical spectrum of Sweet's syndrome (acute febrile neutrophilic dermatosis): a report of eighteen cases. *Br J Dermatol* 1975;92:363–73.
- 3 Sitjas D, Puig L, Cuatrecasas M, *et al.* Acute febrile neutrophilic dermatosis (Sweet's syndrome). *Int J Dermatol* 1993;32:261–8.
- 4 Matta M, Kurban AK. Sweet's syndrome: systemic association. *Cutis* 1973;12:561–5.
- 5 Von Den Driesch P. Sweet's syndrome (acute febrile neutrophilic dermatosis). *J Am Acad Dermatol* 1994;31:535–56.
- 6 Kemmett D, Hunter JAA. Sweet's syndrome: a clinicopathologic review of twenty-nine cases. *J Am Acad Dermatol* 1990;23:503–7.
- 7 Frayha R, Matta M, Kurban A. Sweet's syndrome simulating systemic lupus erythematosus. *Dermatologica* 1972;144:321–4.
- 8 Esteves C, Goncalves M, Barba SG, *et al.* Sweet's syndrome. Report of a case with renal involvement and ANCA positivity. *Eur J Intern Med* 1995;6:195–7.
- 9 Cohen PR, Talpaz M, Kurzrock R. Malignancy-associated Sweet's syndrome: review of the world literature. *J Clin Oncol* 1988;6:1887–97.
- 10 Christ E, Linka A, Jacky E, *et al.* Sweet's syndrome involving the musculoskeletal system during treatment of promyelocytic leukemia with all-trans retinoic acid. *Leukemia* 1996;10:731–4.

- 11 Unis ME, Hill GS. Sweet's syndrome associated with acute renal failure. *Cutis* 1987;40:139–42.
- 12 Vignon-Pennanmen MD. A case of sweet syndrome with spleen and lymph node involvement preceded by parvovirus B19 infection, and a review of literature on extracutaneous sweet syndrome. *Am J Dermatopathol* 2010;32:621–7.
- 13 Klinger S, Mathis N, Jackson S. Bullous Sweet syndrome associated with an aseptic splenic abscess. *Cutis* 2009;84:255–8.
- 14 Vignon-Pennanmen MD. The extracutaneous involvement in the neutrophilic dermatoses. *Clin Dermatol* 2000;18:339–47.

Copyright 2013 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.

BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow