Published in final edited form as:

Schizophr Res. 2007 August ; 94(0): . doi:10.1016/j.schres.2007.04.008.

Cognition and functional outcome among deaf and hearing people with schizophrenia

Heather K. Horton^{a,*}

^aSchool of Social Welfare, University at Albany, Richardson Hall 280, 135 Western Avenue, Albany, NY 12203, United States

1. Introduction

Specific cognitive abilities have been shown to predict functional outcomes among people with schizophrenia (PWS) (Goldberg and Green, 2002; Green, 1996; Green et al., 2000). However, previous studies have been limited to hearing PWS; cognitive functioning in deaf PWS, and its relationship to outcomes, has only rarely been studied. Viggo C. Hansen (1929) conducted the earliest work, and fewer than twenty empirical articles have been published since that date (for a reviews see Horton, 2005; Vernon, 1999; Vernon, 1995).

On the other hand, a sizeable body of knowledge related to *nonclinical* samples of deaf people has been generated. Natural experiments created by deaf people born to hearing parents (90% of the deaf population) have lead to extensive research regarding the effect of delayed language acquisition (Emmorey, 1993; Mayberry, 1991, 1992, 1995; Neville et al., 1997; Singleton and Newport, 2004); the relationship between language and spatial cognition (Emmorey, 1993; Emmorey, 2002; Mayberry, 1992; Neville, 1995; Poizner et al., 1987); the attainment of cognitive and linguistic milestones (Bellugi et al., 1990; Chamberlain et al., 2000; Church and Goldin-Meadow, 1986; Goldin-Meadow et al., 1996; Goldin-Meadow and Mylander, 1991; Newport and Meier, 1985); the neural organization of American Sign Language (ASL) (Bellugi et al., 1989; Emmorey et al., 2002; Emmorey et al., 1998; Neville, 1988; Neville, 1995; Neville et al., 1998); and the relationship between language and affect (Corina, 1989; Corina et al., 1999; Reilly et al., 1990a; Reilly et al., 1990b). There is evidence that deaf children who have been exposed to a complete language model from birth, as is the case with deaf children born to deaf parents (i.e., native signers or "early language learners"), perform the same as hearing children of hearing parents on many cognitive measures (Marschark, 1993; Marschark and Clark, 1993; Mayberry, 1992).

There are important differences as well as similarities in neural processing between deaf and hearing people. Similarities between signed and spoken language in interhemispheric organization reveal that the left hemisphere (specifically frontal as well as temporoparietal areas of the brain) are responsible for language *independent of the language modality* (Neville et al., 1997; Neville et al., 1998; Petitto et al., 2000). For example, when Broca's area is damaged in a hearing person, orofacial movement deficits related to speech production as well as deficits in written output are evidenced (Broca, 1861; Broca, 1865; Mohr et al., 1978). Similarly, deaf subjects with Broca's area damage have movement

© 2007 Elsevier B.V. All rights reserved.

*Corresponding author. Tel.: +1 518 442 5331; Fax: 518 442 5380. hhorton@albany.edu (H.K. Horton).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

deficits—in their limbs rather than their mouths—related to the production of ASL (Hickok et al., 1999; Neville et al., 1997). In short, Broca's area controls language motor output, whether orofacial or limb movements are involved in that output.

The differences in neural processing between deaf and hearing groups can be found in specific aspects of spatial cognition. Deaf people have displayed superior mental rotation abilities (McKee, 1987; Talbot and Haude, 1993), greater accuracy in remembering object orientation (Emmorey et al., 1998), and superior copying abilities compared to hearing subjects who do not sign (Emmorey, 1993; Emmorey et al., 1993). Taken together, cognitive data from nonclinical samples of deaf people point to the idea that native signers—whether deaf or hearing—display enhanced visual-spatial processing abilities compared to hearing people who do not know sign language or who learn it later in childhood (Bellugi et al., 1990; Bettger et al., 1997; McCullough and Emmorey, 1997). Native signers include deaf and hearing persons whose first language is ASL. Typically, these are people born to deaf parents and thus exposed to sign language from birth. Rather than deafness per se, it appears that enhancements among native signers are due to experience with a visual-spatial language and are tied to specific linguistic-processing requirements (e.g., interpretation of grammatical facial expressions and perspective transformations) (Emmorey, 1993, p. 153).

The current study tested two hypotheses. The first is that particular domains of cognition (e.g., early visual processing, verbal and visual-spatial memory) will predict levels of functional outcome among deaf PWS just as they have been proven to do among hearing PWS. There are little or no empirical data regarding the relationship between cognition and outcome among deaf PWS; however, a priori there is no reason to assume that the relationship will not manifest. A within-group (deaf) analysis is employed with the goal of replicating data from the (hearing) schizophrenia research literature.

The second hypothesis is that differences between deaf and hearing participants will manifest in levels of cognitive ability. In particular, deaf PWS are expected to outperform their hearing counterparts on measures of early visual processing and visual-spatial memory. A between-group (deaf, hearing) analysis is employed in an attempt to replicate findings from the nonclinical and deafness-oriented psychological literature that deaf people have superior visual processing abilities compared to hearing people. The finding has proven to be particularly strong among deaf compared to hearing people *without* mental illness, and just one study supports the finding that visual processing skills of deaf PWS are superior to those of hearing people *with* schizophrenia (Kubota et al., 2002).

2. Materials and Methods

2.1 Subjects

The study was approved by the University of Chicago Institutional Review Board. PWS were recruited from all deaf and hearing consumers at a psychiatric rehabilitation agency in Chicago (approximately 3,000 consumers). In total, 87 people agreed to participate in the study (46 deaf, 41 hearing). A diagnosis of schizophrenia or schizoaffective disorder was determined by the Structured Clinical Interview for DSM-IV Axis I Disorders-Patient Edition (SCID) (First et al., 1997). The SCID was translated into ASL and back-translated into English by the author and a native signer. The author was trained to administer the SCID by an experienced, doctoral-level trainer at the University of Chicago, Department of Psychiatry (the author achieved an interclass correlation of .78 against the gold standard trainer). Twenty people did not meet the SCID-based diagnostic inclusion criteria (11 deaf, 9 hearing). In addition, two people, each with a history of head injury, were excluded (1 deaf, 1 hearing).

The final sample included 65 subjects (34 deaf, 31 hearing). Among the 34 deaf subjects, 32 completed all three testing sessions. Of the two who did not participate in the final session, one gave no reason, and the other was not available until after the data collection period ended. For these two deaf subjects, cognitive data are missing.

2.2 Sample Characteristics

Table 1 presents demographic information for both deaf and hearing subjects. The groups were similar with respect to gender, race, diagnosis, housing status, employment status, and psychiatric history variables. The mean age of illness onset was 20 years old; the average total number of lifetime hospitalizations was 8; subjects experienced their most recent hospitalizations 6 years prior to the test date; and the average illness duration was 26 years.

In comparing the two groups, the same number of subjects attended and completed college. One meaningful difference was the number within each group who graduated from high school. Of 16 hearing subjects who entered high school, only 2 graduated. Among deaf subjects, 21 entered high school and 17 graduated. The strength of the relationship between education and the dependent variable (functional outcome) was moderate ($r = .442$) and thus was statistically controlled in all between-group comparisons.

2.3 Measures

Measures were based on the Measurement and Treatment Research to Improve Cognition in Schizophrenia (MATRICS) initiative (Green et al., 2004). Of the six cognitive domains suggested by the MATRICS consensus battery, four are evaluated in the current project: early visual processing (EVP), vigilance, visual-spatial memory, and verbal memory.

Tests that represent the independent measurement of early visual and attentional abilities—with no working memory demands—are employed in this project (Asarnow et al., 1991; Asarnow and Nuechterlein, 1994; Silverstein et al., 1998). Further, measures used to assess visual-spatial processing that do not overlap with linguistic ability were chosen in order to avoid the possible confounds arising for deaf people who are “late language learners.”

Test instructions and stimuli were administered in sign language for deaf subjects (either ASL or Pidgin Signed English) and spoken English for hearing subjects. A random-number generator was used to order the presentation of measures for each subject.

2.3.1 Early Visual Processing—Early visual processing (EVP) was measured with the Partial Report Span of Apprehension test ([Span] Asarnow and Nuechterlein, 1994). The Span is a computer-based measure requiring subjects to identify which of two target letters (T or F) appear on the monitor by pressing one of two response buttons as quickly as possible. Because the task uses alphabetic letters as stimuli it is conceptualized as a linguistic-based cognitive task. A measure of reaction time—in milliseconds—represents the length of time it took subjects to respond to the presentation of the stimuli. Two different series of stimuli are displayed randomly: a 3-letter matrix and a 12-letter matrix.

Span-of-apprehension performance among people with schizophrenia increasingly diverges from that of non-mentally ill and psychiatric controls as the number of items to be recalled increases (Suslow, 1998; Asarnow et al., 1991). The testing room specifications established by Asarnow and Nuechterlein (1994) were followed and were also followed for measures of vigilance.

2.3.2 Vigilance—Vigilance was measured with the Degraded Stimulus Continuous Performance Test ([DS-CPT] Nuechterlein and Asarnow, 1999). The DS-CPT has been

widely used to measure sustained attentional deficits in schizophrenia. Subjects monitor a random series of single numbers, which are presented continuously at a rate of approximately one per second. Subjects are asked to indicate that they have detected a target event by pressing a response button, and they are to avoid responding to nontarget stimuli. The target stimulus—the number zero—is presented 480 times. Sensitivity A' (i.e., A -prime) is a nonparametric measure of the subject's ability (sensitivity) to discriminate between target and nontarget stimuli; the score is reflected as a percentage.

Clear deficits in attention, as measured by continuous performance tests, are found among people with schizophrenia when compared to nonclinical groups for both verbal as well as visual-spatial material (Nuechterlein, et al., 1992; Chen et al., 1997; Elvevag et al., 2001).

2.3.3 Visual-Spatial Memory—Visual-spatial memory was measured by the Complex Figure Test ([CFT] Osterrieth, 1944; Visser, 1970–1973). The CFT uses an intricate stimulus that is asymmetrical in design. Subjects are asked to copy the figure and then draw it again from memory. Three indices are measured: the “strategy” employed to complete the task, perceptual organization (VSP-copy), and immediate recall (VSP-recall). A 36-point scoring system refers to specific lines and details of the figure to be recalled; points are awarded for each line properly placed. Among people with schizophrenia, retrieval has been shown to be affected more than encoding (McClain, 1983; Paulsen et al., 1995).

2.3.4 Verbal Memory—Verbal memory, referred to as word memory, was measured with the Rey Auditory Verbal Learning Test ([AVLT] Rey, 1964), which consists of several lists containing 15 words each. Seven trials test subjects on number of words correctly recalled; the total short-term word-memory score is the sum of words recalled correctly on trials 1 through 5 (high score = 75). Long-term word memory is evaluated by asking subjects to recall the original list of 15 words after a 20-minute delay. A recognition scale reveals the ability to distinguish between target and nontarget words.

When schizophrenic subjects are asked to recall a list of words after a short delay (i.e., short-term verbal memory), with or without cues, a deficit is almost always observed (Calev et al., 1983; Heinrichs & Zakzanis, 1998; Stirling et al., 1997; Strauss et al., 1990). This is true of immediate and delayed-recall tasks for verbal material.

2.3.5 Functional Outcome—The Multnomah Community Ability Scale ([MCAS]: Barker et al., 1994a; 1994b) was used to operationalize outcome. The MCAS is a rating scale developed for use with people with chronic mental illness living in the community. The score across four subscales is summed to create an overall functional outcome rating, hereafter referred to simply as “outcome.” Three levels of disability are described by the final score: low (17–47), medium (48–62), and high (63–85). The researcher met with caseworkers individually and completed an MCAS for each participant. Reliability statistics were calculated with Cronbach's alpha ($\alpha = .90$, $F = 10.11$, $p = .000$).

2.3.6 Illness Severity—Illness severity was operationalized by the total score on the Brief Psychiatric Rating Scale ([BPRS]: Ventura et al., 1993). After training, the author achieved an interclass correlation of .82 against a gold standard and administered the BPRS to all subjects. Positive and negative symptom clusters were determined by conducting a restricted confirmatory factor analysis (CFA) on the 24 BPRS items. The anergia factor (blunted affect, emotional withdrawal and motor retardation, self-neglect) and the thought disorder factor (unusual thought content, grandiosity, and suspiciousness) were used to represent positive and negative symptom clusters.

2.4 Statistical Analyses

To test the first hypothesis, a series of bivariate regressions was employed. The tests evaluated the relationship between cognition and outcome among deaf subjects ($n = 32$). Post hoc, analyses of the hearing sample ($n = 31$) and the total sample ($n=63$), were conducted. To test the second hypothesis, ANCOVAs were used to compare the deaf and hearing subjects' levels of cognitive ability ($n = 63$).

Bonferroni corrections were used to account for sequential testing on the same set of subjects. The alpha level was corrected for each set of hypotheses. The first hypothesis ($n = 31$) entailed five comparisons ($p = .0102$, t -value for 2-tails: 2.744 ; 1 tail: ≥ 2.452). The second hypothesis ($n = 61$) also entailed five comparisons ($p = .0102$, t -value for 2-tails 2.657 ; 1 tail: ≥ 2.388). Results presented represent 2-tailed tests of significance unless otherwise noted.

Level of skew and kurtosis with their standard errors (SE) were analyzed in all variables. An a priori decision was made to transform variables if skew/SE and or kurtosis/SE was greater than ± 2 . Three variables were transformed: VSP-copy, and the BPRS anergia and thought disorder factors.

Outliers were removed according to a specific guideline: No more than 5% of cases were removed ($n=3$) from any one analysis (BMDP, 1994; Cohen and Cohen, 1983). A correlation matrix depicting the strength of associations among the primary study variables is presented in Table 2.

3. Results

3.1 Cognition and Outcome: Deaf Subjects

The results in Table 3 are ordered from the strongest to the weakest predictor of outcome for deaf subjects. The hypothesis that cognition predicts outcome among deaf PWS was supported.

3.2 Cognition and Outcome: Hearing Subjects

For informative purposes, Table 4 presents the cognitive predictors of outcome for hearing subjects. Table 5 presents a side-by-side analysis of the strongest predictors of outcome for the deaf subjects, hearing subjects, and the total sample.

3.3 Differences in Cognitive Performance Between Deaf and Hearing PWS

The results of the second research question are displayed in Table 6. An analysis of covariance was used to control for the difference in high school graduation rates among a subgroup of deaf and hearing subjects (see section 2.2). The hypothesis that differences exist across the deaf and hearing samples on indices of EVP and visual-spatial memory was not supported.

Post hoc, the remaining domains of cognition were compared across deaf and hearing subjects, and one domain revealed a significant difference (italicized in Table 6). Deaf subjects scored higher than hearing subjects on the word memory/recognition task. This difference was not hypothesized and should be interpreted cautiously.

4. Discussion

This study tested two hypotheses developed from separate domains of inquiry. The research questions are linked by the importance of visual and visual-spatial processing among deaf

people. The first question addressed the relationship between cognition and outcome. It is a common finding in the hearing-oriented schizophrenia literature that cognitive processing is a reliable predictor of outcome. This study replicated that finding among deaf PWS. The strongest predictor of outcome for deaf subjects was VSM-copy followed by VSM-recall, EVP, and vigilance. The final domain, word memory, was not significant once corrected. The most potent predictors of outcome for the deaf sample include abilities that place less of a demand on linguistic processing and more on visual-spatial processing.

Only two of the five predictors (EVP and word memory) were statistically significant for hearing subjects. VSM-recall and VSM-copy were weak trends in relation to outcome, while vigilance was not related to outcome in the hearing sample. Overall, and for unknown reasons, cognition was a more potent predictor of outcome for deaf than for hearing subjects. Nonetheless, it was the linguistic-based cognitive abilities that best predicted outcome for the hearing subjects.

The importance of attending to the linguistic and nonlinguistic demands of cognitive tasks was borne out in the current study by the finding that nonlinguistic visual processing mechanisms were the most potent predictors of outcome for deaf subjects (VSM-recall and copy) while linguistic-based visual mechanisms (EVP and word memory) were among the weakest predictors of outcome for deaf subjects. For hearing subjects, the opposite was empirically supported. That is, nonlinguistic visual-spatial abilities (VSM-recall and copy) had only weak relationships to outcome, and the linguistic-based cognitive tasks (EVP and word memory) were the strongest predictors of outcome.

The relevance of *nonlinguistic* visual-spatial processing for deaf subjects and, likewise, linguistic-based cognitive processing for hearing subjects is in line with empirical work emphasizing the way in which a lifelong reliance on visual rather than auditory mechanisms for information and linguistic processing influences cognition. The standard method of separating *verbal* mechanisms from those that are *visual-spatial* is challenged when considering deaf people because their “verbal” communication domain *is* visual-spatial. The linguistic and nonlinguistic components of visual-spatial processing must be specifically dissociated from one another, as can be seen by the difference between tasks that require the processing of language (e.g., EVP and word memory) and tasks that require the processing of geometric shapes (e.g., VSM-recall and copy). These differences may have implications for rehabilitation efforts for deaf and hearing PWS.

The second question addressed directly the differences in cognitive performance between deaf and hearing subjects. Empirical evidence from the deafness-oriented psychological literature has long revealed that deaf signers, and especially native signers, have visual-spatial abilities that are superior to those of hearing and deaf people who either learned sign language “late” or had never been exposed to sign language (the definition of “late” varies in the literature). In the current project, differences in certain abilities were found between subjects who learned sign language before, compared to after, their fifth birthdays (results presented in a forthcoming paper).

While the raw data reveals that deaf subjects in the current study performed better than hearing subjects on the VSM-copy and recall task, differences were not significant, and there were essentially no differences between deaf and hearing subjects on the measure of EVP. It is perhaps not surprising that differences were not found on the measure of EVP because it represents a linguistic-based processing skill. Future research should compare deaf and hearing subjects on measures of cognition that fall squarely within the domain of nonlinguistic visual and/or visual-spatial processing.

Unexpected significant differences arose with regard to word memory/recognition. Deaf subjects correctly chose target from nontarget words at a higher rate than hearing subjects. Without further hypothesis testing, it is difficult to determine the reasons for such performance. The finding does not correspond to the well-documented delay in other domains of memory among deaf people (especially children). Results in this area of research, however, have been inconsistent (Bebko, 1984; Harris and Moreno, 2004; Arnold and Mills, 2001; Arnold and Murray, 1998).

It is clear that cognition varies with important changes in outcome and the quality of life (Green et al., 2000; Addington et al., 1998). To contribute to the expansion of theories of social and functional dysfunction among PWS, we must explain the nature of that relationship. That is, *how* do below average scores on cognitive tasks relate to social and functional competence? The presence of cognitive deficits by themselves is an insufficient explanation of the profound social disabilities associated with schizophrenia (Carter and Flesher, 1995, p. 215). This is especially apparent among PWS who display impoverished social competence along with relatively intact cognition (Heinrichs and Zakzanis, 1998; Heinrichs, 2007).

A recent suggestion is that the relationship between cognition and outcome may be mediated by *social* cognition (e.g., Theory of Mind). Targeting social cognition as a potential mediator addresses concerns regarding the gap between microprocesses such as attention or memory and real-world social and functional behavior. A mediating hypothesis has been suggested in the literature (e.g., Green, et al., 2000; Kee et al., 1998) and is tested by the author in a forthcoming paper (see Horton & Silverstein, 2007).

The goals of this study were twofold. The first was to establish that cognition predicts outcomes among deaf people, as we know it does among hearing PWS. Support for the hypothesis was provided and moves towards an understanding of whether pathways to social impairment are distinct across the populations. The second goal was to address the lack of empirical research among deaf people with serious mental illness by providing an empirical foundation for future work. As part of this goal, special attention was paid to visual and visual-spatial processing among deaf people because of their reliance on such mechanisms for information and linguistic processing.

Acknowledgments

I would like to thank the members and staff of Thresholds Psychiatric Rehabilitation Center for their contribution to this research. I am also grateful to R. F. Asarnow and K. H. Nuechterlein for putting at my disposal software packages for neuropsychological testing.

References

- Addington J, McCleary L, Munroe-Blum H. Relationship between cognitive and social dysfunction in schizophrenia. *Schizophr Res.* 1998; 34(1–2):59–66. [PubMed: 9824877]
- Arnold P, Mills M. Memory for faces, shoes, and objects by deaf and hearing signers and hearing nonsigners. *J Psycholinguist Res.* 2001; 30(2):185–195. [PubMed: 11385825]
- Arnold P, Murray C. Memory for faces and objects by deaf and hearing signers and hearing nonsigners. *J Psycholinguist Res.* 1998; 27(4):481–497. [PubMed: 9691334]
- Asarnow, RF.; Granholm, E.; Sherman, T. Span of apprehension in schizophrenia. In: Steinhauer, SR.; Gruzeli, JH.; Zubin, J., editors. *Handbook of Schizophrenia: Neuropsychology, Psychophysiology, and Information Processing.* Vol. 5. Amsterdam: Elsevier; 1991. p. 335–370.
- Asarnow, RF.; Nuechterlein, KH. *Directions for Use of the UCLA Span of Apprehension Program, Version 3.5 for Arrays 3 and 12 Letters, on IBM AT and Fully Compatible Microcomputers.* UCLA Department of Psychiatry and Biobehavioral Sciences; Los Angeles, CA: 1994.

- Barker S, Barron N, McFarland BH, Bigelow DA. A community ability scale for chronically mentally ill consumers: Part I. Reliability and validity. *Community Ment Health J.* 1994a; 30(5):363–383. [PubMed: 7956112]
- Barker S, Barron N, McFarland BH, Bigelow DA, Carnahan T. A community ability scale for chronically mentally ill consumers: Part II. Applications *Community Ment Health J.* 1994b; 30:459–472.
- Bebko JM. Memory and rehearsal characteristics of profoundly deaf children. *J Exp Child Psychol.* 1984; 38(3):415–28. [PubMed: 6520582]
- Bellugi U, Poizner H, Klima ES. Language, modality and the brain. *Trends Neurosc.* 1989; 12(10): 380–388.
- Bellugi, U.; O'Grady, L.; Lillo-Martin, L.; O'Grady, M.; van Hoek, K.; Corina, D. Enhancement of spatial cognition in deaf children. In: Volterra, V.; Erting, C., editors. *From Gesture to Language in Hearing and Deaf Children.* Springer-Verlag; New York: 1990. p. 278-298.
- Bettger JG, Emmorey K, McCullough S, Bellugi U. Enhanced facial discrimination: Effects of experience with American Sign Language. *J Deaf Stud Deaf Educ.* 1997; 2(4):223–233. [PubMed: 15579850]
- BMDP Statistical Software . BMDP New System for Windows Version 1.0. Los Angeles, CA: 1994.
- Broca MP. Remarques sur le siege de la faculté du langage articulé, suivies d'une observation d'aphemie (Perte de la Parole). *Bull Mem Soc Anat Paris.* 1861; 36:330–357.
- Broca PC. Sur le siege de la faculté du langage articule. *Bulle la Soc Anthropol Paris.* 1865; 6:377–393.
- Calev A, Venables PH, Monk AE. Evidence for distinct verbal memory pathologies in severely and mildly disturbed schizophrenics. *Schizophr Bull.* 1983; 9(2):247–264. [PubMed: 6867632]
- Carter M, Flesher S. The neurosociology of schizophrenia: vulnerability and functional disability. *Psychiatry.* 1995; 58(3):209–224. [PubMed: 8539302]
- Chamberlain, C.; Morford, JP.; Mayberry, RI. *Language Acquisition by Eye.* Hillsdale, NJ: Lawrence Erlbaum; 2000.
- Chen EYH, Lam LCW, Chen RYL, Nguyen DGH, Chan CKY, Wilkins AJ. Neuropsychological correlates of sustained attention in schizophrenia. *Schizophr Res.* 1997; 24(3):299–310. [PubMed: 9134590]
- Church RB, Goldin-Meadow S. The mismatch between gesture and speech as an index of transitional knowledge. *Cogn.* 1986; 23(1):43–71.
- Cohen, J.; Cohen, P. *Applied Multiple Regression/Correlation Analysis for the Behavioral Sciences.* 2. Hillsdale, NJ: Lawrence Erlbaum; 1983.
- Corina DP. Recognition of affective and noncanonical linguistic facial expressions in hearing and deaf subjects. *Brain Cognit.* 1989; 9(2):227–237. [PubMed: 2923713]
- Corina DP, Bellugi U, Reilly J. Neuropsychological studies of linguistic and affective facial expressions in deaf signers. *Lang Speech.* 1999; 42(2–3):307–331. [PubMed: 10767992]
- Dickinson D, Coursey RD. Independence and overlap among neurocognitive correlates of community functioning in schizophrenia. *Schizophr Res.* 2002; 56(1–2):161–170. [PubMed: 12084430]
- Dickinson D, Iannone VN, Wilk CM, Gold JM. General and specific cognitive deficits in schizophrenia. *Biol Psychiatry.* 2004; 55(8):826–833. [PubMed: 15050864]
- Elvevag B, Weinstock DM, Akil M, Kleinman JE, Goldberg TE. A comparison of verbal fluency tasks in schizophrenic patients and normal controls. *Schizophr Res.* 2001; 51(2–3):119–126. [PubMed: 11518632]
- Emmorey K. Processing a dynamic visual-spatial language: psycholinguistic studies of American Sign Language. *J Psycholinguist Res.* 1993; 22(2):153–187. [PubMed: 8366475]
- Emmorey, K. *Language, Cognition, and the Brain: Insights from Sign Language Research.* Erlbaum; Mahwah, NJ: 2002.
- Emmorey K, Damasio H, McCullough S, Grabowski T, Ponto LLB, Hichwa RD, Bellugi U. Neural systems underlying spatial language in American Sign Language. *NeuroImage.* 2002; 17(2):812–824. [PubMed: 12377156]

- Emmorey K, Kosslyn SM, Bellugi U. Visual imagery and visual-spatial language: enhanced imagery abilities in deaf and hearing ASL signers. *Cogn.* 1993; 46(2):139–181.
- Emmorey K, Klima ES, Hickok G. Mental rotation within linguistic and non-linguistic domains in users of American Sign Language. *Cogn.* 1998; 68(3):221–246.
- First, MB.; Spitzer, RL.; Gibbon, M.; Williams, JBW. Structured Clinical Interview for DSM-IV Axis I Disorders—Patient Edition (SCID-I/P, Version 2.0, 4/97 revision). Biometrics Research Department New York State Psychiatric Institute; New York: 1997.
- Goldberg, T.; Green, M. Neurocognitive functioning in patients with schizophrenia: an overview. In: Davis, KL.; Charney, D.; Coyle, JT.; Nemeroff, C., editors. *Neuropsychopharmacology: The Fifth Generation of Progress*. Williams & Wilkins, Lippincott; Philadelphia: 2002. p. 657–669.
- Goldin-Meadow S, McNeil D, Singleton J. Grammatical expression in the manual modality. *Psychol Rev.* 1996; 103(1):34–55. [PubMed: 8650298]
- Goldin-Meadow, S.; Mylander, C. Home sign systems in deaf children: the development of morphology without a conventional language model. In: Siple, P.; Fischer, SD., editors. *Theoretical Issues in Sign Language Research*. University of Chicago Press; Chicago: 1991. p. 41–63. (Vol. 2: Psychology)
- Green MF. What are the functional consequences of neurocognitive deficits in schizophrenia? *Am. J Psychiatry.* 1996; 153(3):321–330.
- Green MF, Kern RS, Braff DL, Mintz J. Neurocognitive deficits and functional outcome in schizophrenia: are we measuring the “right stuff”? *Schizophr. Bull.* 2000; 26(1):119–136.
- Green MF, Nuechterlein KH, Gold JM, Barch DM, Cohen J, Essock S, Fenton WS, Frese F, Goldberg TE, Heaton RK, Keefe RSE, Kern RS, Kraemer H, Stover E, Weinberger DR, Zalcman S, Marder SR. Approaching a consensus cognitive battery for clinical trials in schizophrenia: The NIMH-MATRICES conference to select cognitive domains and test criteria. *Biol Psychiatry.* 2004; 56:301–307. [PubMed: 15336511]
- Green MF, Kern RS, Braff DL, Mintz J. Neurocognitive deficits and functional outcome in schizophrenia: Are we measuring the “right stuff”? *Schizophr Bull.* 2000; 26(1):119–136. [PubMed: 10755673]
- Hansen, VC. Beretning om Sindslidelse blandt Damarks Døvtumme. Johansen; Copenhagen, Denmark: 1929.
- Harris M, Moreno C. Deaf children’s use of phonological coding: evidence from reading, spelling, and working memory. *J Deaf StudDeaf Educ.* 2004; 9(3):253–268.
- Heinrichs W, Miles A, McDermid Vaz S, Zargarian T, Miki A, Smith D. Verbally gifted schizophrenia patients and healthy controls: Differences in functional status, but not cognition? *Schizophr Bull.* 2007; 33(2):560.
- Heinrichs RW, Zakzanis K. Neurocognitive deficit in schizophrenia: a quantitative review of the evidence. *Neuropsychol.* 1998; 12(3):426–445.
- Hickok G, Wilson M, Clark K, Klima ES, Kritchinsky M, Bellugi U. Discourse deficits following right hemisphere damage in deaf signers. *Brain Lang.* 1999; 66(2):233–248. [PubMed: 10190988]
- Horton, HK. Unpublished doctoral dissertation. University of Chicago; Chicago: 2005. Schizophrenia, deafness and functional outcome: the role of neurocognition and social cognition.
- Horton HK, Silverstein SM. Social cognition as a mediator of neurocognition and functional outcome among deaf people with schizophrenia. *Schizophr Bull.* 2007; 33(2):591.
- Kee KS, Kern RS, Green MF. Perception of emotion and neurocognitive functioning in schizophrenia: what’s the link? *Psychiatry Res.* 1998; 81(1):57–65. [PubMed: 9829651]
- Kubota Y, Querel C, Pelion F, Laborit J, Laborit MFM, Gorog F, Okada T, Murai T, Sato W, Yoshikawa S, Toichi M, Hayashi T. Facial affect recognition in pre-lingually deaf people with schizophrenia. *Schizophr Res.* 2002; 63(1):265–270.
- Marschark, M. *Psychological Development of Deaf Children*. Oxford University Press; New York: 1993.
- Marschark, M.; Clark, MD. *Psychological Perspectives on Deafness*. Lawrence Erlbaum; Hillsdale NJ: 1993.

- Mayberry, RI. Children's memory for sign and fingerspelling in relation to production rate and sign language input. In: Siple; Fischer, SD., editors. *Theoretical Issues in Sign Language Research*. University of Chicago Press; Chicago: 1991. p. 211-229. (Vol. 2: Psychology)
- Mayberry, RI. The cognitive development of deaf children: recent Insights. In: Segalowitz, SJ.; Rapin, I., editors. *Handbook of Neuropsychology*. Vol. 7. Elsevier Science; Oxford: 1992. p. 51-68.
- Mayberry, RI. Mental phonology and language comprehension, or What does that sign mistake mean?. In: Emmorey, K.; Reilly, JS., editors. *Language, Gesture and Space*. Hillsdale, NJ: Lawrence Erlbaum; 1995. p. 355-370.
- McClain L. Encoding and retrieval in schizophrenics' free recall. *J Nerv Ment Dis*. 1983; 171(8):471-479. [PubMed: 6875531]
- McCullough S, Emmorey K. Face processing by deaf ASL signers: evidence for expertise in distinguishing local features. *J Deaf Stud Deaf Educ*. 1997; (4):212-222. [PubMed: 15579849]
- McKee, D. Unpublished doctoral dissertation. University of Pittsburgh; Pittsburgh, PA: 1987. An analysis of specialized cognitive functions in deaf and hearing signers.
- Mohr JP, Pessin MS, Finkelstein S, Funkenstein HH, Duncan GW, Davis KR. Broca aphasia: pathologic and clinical. *Neurology*. 1978; 28(4):311-324. [PubMed: 565019]
- Neville, H. Cerebral organization for spatial attention. In: Stiles-Davis, J.; Kritchevsky, M.; Bellugi, U., editors. *Spatial Cognition: Brain Bases and Development*. Lawrence Erlbaum; Hillsdale, NJ: 1988. p. 327-341.
- Neville, HJ. Developmental specificity in neurocognitive development in humans. In: Gazzaniga, M., editor. *The Cognitive Neuroscience*. MIT Press; Cambridge, MA: 1995.
- Neville HJ, Bavelier D, Corina D, Rauschecker J, Karni A, Lalwani A, Braun A, Clark V, Jezzard P, Turner R. Cerebral organization for language in deaf and hearing subjects: Biological constraints and effects of experience. *Proc Natl Acad Sci*. 1998; 95:922-929. [PubMed: 9448260]
- Neville HJ, Coffey SA, Lawson DS, Fischer A, Emmorey K, Bellugi U. Neural systems mediating American Sign Language: effects of sensory experience and age of acquisition. *Brain Lang*. 1997; 57(3):285-308. [PubMed: 9126418]
- Newport, EL.; Meier, RP. The acquisition of American Sign Language. In: Slobin, DI., editor. *The crosslinguistic study of language acquisition*. Vol. 1. Lawrence Erlbaum; Hillsdale, NJ: 1985.
- Nuechterlein, KH.; Asarnow, RF. *Degraded Stimulus Continuous Performance Test (DS-CPT): Program for IBM-Compatible Micro-Computers, Version 8.12* [Computer software]. Los Angeles: 1999.
- Nuechterlein KH, Dawson ME, Gitlin M, Ventura J, Goldstein MJ, Snyder KS, Yee CM, Mintz J. Developmental processes in schizophrenic disorders: longitudinal studies of vulnerability and stress. *Schizophr Bull*. 1992; 18(3):387-425. [PubMed: 1411329]
- Osterrieth PA. Le test de copie d'une figure complexe: contribution à l'étude de la perception et de la memoire. *Arch Psychol*. 1944; 30:286-350.
- Paulsen JS, Heaton RK, Sadek JR, Perry W, Delis DC, Braff D, Kuck J, Zisook S, Jeste DV. The nature of learning and memory impairments in schizophrenia. *J Int Neuropsychol Soc*. 1995; 1:88-99. [PubMed: 9375213]
- Petitto LA, Zatorre RJ, Gauna K, Nikelski EJ, Dostie D, Evans AC. Speech-like cerebral activity in profoundly deaf people processing signed language: implications for the neural basis of human language. *PNAS (Proceedings of the National Academy of Sciences of the United States of America)*. 2000; 97(25):13961-13966.
- Poizner, H.; Klima, ES.; Bellugi, U. *What the Hands Reveal About the Brain*. MIT Press; Cambridge, MA: 1987.
- Reilly, JS.; McIntire, M.; Bellugi, U. Faces: the relationship between language and affect. In: Volterra, V.; Erting, C., editors. *From Gesture to Language in Hearing and Deaf Children*. Springer-Verlag; New York: 1990a. p. 128-141.
- Reilly JS, McIntire M, Bellugi U. The acquisition of conditionals in American Sign Language: grammaticized facial expressions. *Appl Psycholinguist*. 1990b; 11(1):369-392.
- Rey, A. *L'examen clinique en psychologie*. Presses Universitaires de France; Paris: 1964.

- Silverstein SM, Osborn LM, Palumbo DR. Rey-Osterrieth complex figure test performance in acute, chronic, and remitted schizophrenia patients. *J Clin Psychol.* 1998; 54(7):985–994. [PubMed: 9811135]
- Singleton JL, Newport EL. When learners surpass their models: The acquisition of American Sign Language from inconsistent input. *Cogn Psychol.* 2004; 49(4):370–407. [PubMed: 15342259]
- Steinberg AG, Sullivan VJ, Loew RC. Cultural and linguistic barriers to mental health service access: the deaf consumer's perspective. *Amer J Psychiatry.* 1998; 155(7):982–984. [PubMed: 9659872]
- Stirling JD, Hellewell JSE, Hewitt J. Verbal memory impairment in schizophrenia: No sparing of short term recall. *Schizophr Res.* 1997; 25(2):85–95. [PubMed: 9187007]
- Strauss ME, Reynolds KS, Jayaram G, Tune LE. Effects of anticholinergic medication on memory in schizophrenia. *Schizophr Res.* 1990; 3(2):127–129. [PubMed: 1980611]
- Suslow T. Attentional scanpath on the span of apprehension task in paranoid schizophrenia. *Ger J Psychiatry.* 1998; 1(1):1–5.
- Talbot KF, Haude RH. The relationship between sign language skill and spatial visualization ability: mental rotation of three-dimensional objects. *Percept Mot Skills.* 1993; 77(3, pt. 2):1387–1391. [PubMed: 8170794]
- Ventura J, Green MF, Shaner A, Liberman RP. Training and quality assurance with the Brief Psychiatric Rating Scale: "The drift busters." *Int. J Methods Psychiatr Res.* 1993; 3(4):221–244.
- Vernon M. An historical perspective on psychology and deafness. *J Am Deaf Rehabil Assoc.* 1995; 29(2):8–13.
- Vernon M. Historical overview of inpatient care of mental patients who are deaf. *Am Ann Deaf.* 1999; 144 (1):51–61. [PubMed: 10230083]
- Visser, RSH. Complex Figure Test. Swets Test Publishers; Lisse, The Netherlands: 1970–1973.

Table 1

Demographics and Clinical Variables of Study Participants (n = 65)

Variable	Deaf n = 34	Hearing n = 31	Sig.	Total n = 65
Mean age (sd)	45 (8.9)	47 (9.2)		46 (9.1)
Sex				
Female	13	9		22
Male	21	22		43
Race				
White	18	12		30
Black	12	15		27
Asian	1	1		2
Latino	3	1		4
Biracial	0	2		2
Education				
No or some HS	4	14		18
HS grad	17	2	.000*	19
Any college	11	11		22
College grad.	2	4		6
Housing status				
Residential or nursing home	15	11		26
Semi-independent	5	10		15
SRO or hotel	0	3		3
Own apt.	14	7		21
Diagnosis				
Schizophrenia	26	24		50
Schizoaffective	8	7		15
Illness severity				
Mild	7	3		10
Moderate	6	10		16
Moderate-Severe	6	12		18
Severe	15	6		21
Employment				
Not employed	25	23		48
Noncompetitive	5	7		12
Competitive	4	1		5
Mean negative symptom rating^a	1.90	1.92		1.91
Range 1–7 (sd)	(1.16)	(1.16)		(1.16)
Mean positive symptom rating^b	2.13	2.17		2.15
Range 1–7 (sd)	(1.51)	(1.65)		(1.58)
Mean functional outcome score (sd)	59/85 (14)	61/85 (10)		-

SRO – Single room occupancy

* Scheffé p = .000.

^a BPRS Anergia factor including blunted affect, emotional withdrawal, motor retardation & self-neglect.

^b BPRS thought disorder factor including suspiciousness, unusual thought content and grandiosity.

Table 2

Correlations (Pearson) Between Study Variables.

	Func. out.	Illness sever.	Neg. symp.	Pos. symp.	VM	VSM copy	VSM recall	Vig. ^a	EVP	Educ.
Func. out.	1.000									
Illness sev.	-.586** .000	1.000								
Neg. symp.	-.253* .043	.345** .005	1.000							
Pos. symp.	-.256* .041	.449** .000	.000 1.000	1.000						
VM	.380** .002	-.265* .036	-.229 .071	.019 .884	1.000					
VSM copy	.278* .027	-.055 .666	.061 .633	-.043 .739	.395** .001	1.000				
VSM recall	.254* .045	-.276* .029	-.015 .909	-.047 .714	.415** .001	.454** .000	1.000			
Vig.^a	.180 .157	-.346** .007	-.282* .029	.051 .694	.354** .004	.197 .121	.283* .024	1.000		
EVP	.367** .003	-.301* .017	-.221 .081	-.123 .335	.325** .009	.360** .004	.378** .002	.193 .131	1.000	
Educ.	.442** .000	-.113 .371	.001 .992	.113 .373	.241 .058	.441** .000	.182 .154	.161 .219	.027 .835	1.000

**
p < .01;

*
p < .05.

Func. Out. = MCAS functional outcome; Illness sev. = BPRS illness severity; VM = Visual-spatial memory apprehension; VSM Recall = Visual-spatial memory recall; Vig. = Vigilance (DS-of education)

^aThree outliers were removed to improve fit.

Table 3
Functional Outcome Regressed on to Cognition: Bivariate Associations (Deaf Subjects, n = 32)

Predictors	Beta (Standardized)	Degrees of freedom	Functional outcome t-stat	p-value	r ² (adj.)
VSM (copy) ^a	.505	28	3.04**	.005	.227
VSM (recall) ^b	.466	29	2.79**	.009	.189
EVP ^a	.465	28	2.73*	.011	.187
Vigilance (A') ^a	.433	28	2.50*	.019	.157
Word memory	.386	31	2.29	.029	.121

** p < .05, corrected.

* p < .05, corrected, 1-tailed.

EVP – Early visual processing. VSM – Visual spatial memory.

^aThree outliers were removed to gain better fit.

^bOne outlier was removed to gain better fit.

Functional Outcome Regressed on to Cognition: Bivariate Associations (Hearing Subjects, n = 31)

Table 4

Predictors	Beta standardized	Degrees of freedom	Functional outcome t-stat	p-value	r ² (adj.)
EVP	.449	30	2.71 *	.011	.174
Word memory	.399	30	2.34 *	.026	.130
VSM-recall ^a	.334	29	1.87	.071	.080
VSM-copy	.283	30	1.59	.123	.048
Vigilance (A')	.256	30	1.43	.164	.033

* p < .05

Note: These relationships were analyzed post hoc and were not subjected to a Bonferroni correction.

EVP – Early visual processing. VSM – Visual spatial memory.

^aTwo outliers were removed to gain better fit.

Table 5

Strongest Predictors of Functional Outcome by Group (Deaf, Hearing)

Deaf (n = 32) ^a	Hearing (n = 31) ^a	Total (n = 63)
VSM-copy p = .005	Early visual processing p = .011	Word memory p = .002
		VSM-recall p = .002
VSM-recall p = .009	Word memory p = .026	Early visual processing p = .003
Early visual processing p = .011	VSM-recall p = .071	Vigilance p = .006
Vigilance p = .019	VSM-copy p = .123	VSM-copy p = .016
Word memory p = .029	Vigilance p = .164	

^aThe 2 left hand columns (Deaf and Hearing) are reproductions of Tables 3 and 4 respectively.

Table 6

ANCOVAs: Levels of Cognitive Ability (Education as Covariate), Deaf and Hearing Comparisons

Measures of cognition	Deaf mean (sd)	Hearing mean (sd)	F	p-value
Early visual processing	.70 (.09)	.68 (.12)	.445	.507
Early visual processing: Reaction time in milliseconds	997 (348)	1066 (357)	.365	.548
Visual-spatial memory: Strategy ^a	2.22 (1.1)	2.35 (1.0)	.062	.804
Visual-spatial memory: Copy score	.80 (.20)	.75 (.19)	.342	.561
Visual-spatial memory: Recall	.39 (.18)	.32 (.14)	2.02	.160
Verbal memory: Recognition	.18 (.03)	.16 (.03)	5.420	.023*

* p < .05 (uncorrected).

^a Visual-spatial memory strategy scores range from 1–3; lower scores represent a higher level of ability.