Published in final edited form as:

FEBS Lett. 2013 April 17; 587(8): . doi:10.1016/j.febslet.2013.01.021.

One size doesn't fit all: the oligomeric states of α B crystallin

Scott P. Delbecq and Rachel E. Klevit*

Department of Biochemistry, Box 357350, University of Washington, Seattle, WA 98195-7350, USA

Abstract

Small Heat Shock Proteins (sHSPs) are a diverse family of molecular chaperones that delay protein aggregation through interactions with nonnative and aggregate-prone protein states. This function has been shown to be important to cellular viability and sHSP function/dysfunction is implicated in many diseases, including Alzheimer's and Alexander disease. Though their gene products are small, many sHSPs assemble into a distribution of large oligomeric states that undergo dynamic subunit exchange. These inherent properties present significant experimental challenges for characterizing sHSP oligomers. Of the ten mammalian sHSPs, the human sHSP 03B1;B crystallin is a paradigm example of sHSP oligomeric properties. Advances in our understanding of sHSP structure, oligomeric distribution, and dynamics have prompted the proposal of several models for the oligomeric states of B. The aim of this review is to highlight characteristics of α B crystallin (α B) that are key to understanding its structure and function. The current state of knowledge, existing models, and outstanding questions that remain to be addressed are presented.

Keywords

small heat shock protein; α B crystallin; oligomers; pseudoatomic models

Small Heat Shock Proteins (sHSPs) are a diverse class of protein chaperones observed across all kingdoms of life [1]. Though the list of specific sHSPs functions continues to grow, sHSPs share an ability to interact with partly unfolded and aggregate-prone proteins to delay the formation of insoluble protein aggregates *in vitro* [2]. This so-called "holdase" function is held to be responsible for the increased viability sHSPs impart on cells under stress conditions and may explain the association of sHSPs with diseases tied to protein aggregation [3–5]. While the ability of sHSPs to delay protein aggregation has been known for decades, our current understanding of how sHSPs work remains limited due in large part to the fact that neither the sHSPs nor their aggregating "clients" are amenable to conventional structural and biochemical methods for characterizing protein structure and protein-protein interactions in detail. Recent developments in nuclear magnetic resonance spectroscopy (NMR), electron microscopy (EM), native mass spectrometry (MS), small angle x-ray scattering (SAXS), and other techniques are beginning to allow a more detailed view into the sHSPs and their interactions [6–10]. Among metazoan sHSPs, much of the recent effort has focused on human α B crystallin (HSPB5), which is highly expressed in eye

© 2013 Federation of European Biochemical Societies. Published by Elsevier B.V. All rights reserved.

*Corresponding Author klevit@uw.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

lens, muscle, and brain [5, 11, 12]. The aim of this review is to highlight characteristics of α B crystallin (α B) that are key to understanding its structure and function. We present the current state of knowledge, inadequacies of current models, and identify outstanding questions that remain to be addressed and challenges to be overcome. The sub-class of heat shock proteins known as the *small* heat shock proteins are defined by their shared architecture, composed of a conserved central domain called the “alpha crystallin domain” (“ACD”) which is flanked by a highly variable N-terminal region (“NTR”) and a variable C-terminal region (“CTR”) that contains a small conserved motif known as the “IXI/V” (Figure 1). Despite their moniker of “small” (based on the size of the gene products), sHSPs are functionally very large because they assemble into oligomers. The diversity and properties of sHSP oligomers are highly unusual and are believed to be key to their broad holdase function. In particular, their tendency to form large polydisperse (variable number of subunits) oligomeric species that undergo dynamic subunit exchange is a distinguishing feature of this family of proteins. The properties of sHSPs and their cellular function involve two types of protein-protein interactions: 1) homomeric interactions responsible for the assembly of and dynamic equilibria among sHSP oligomers and 2) heteromeric interactions between sHSP species and their client proteins. An additional layer of complexity arises from the fact that metazoan sHSPs are thought to form hetero oligomers (i.e., oligomers composed of more than one type of sHSP subunit), thereby allowing for hetero-homomeric interactions (i.e., sHSPsHSP interactions). Furthermore, there is mounting evidence that some sHSPs including α B and HSP27 (HSPB1) interact with specific folded cellular proteins in addition to partly unfolded proteins, indicating that further parsing may be required in the heteromeric category of interactions [13, 14]. An ultimate goal is a unifying model for the myriad interactions that define sHSP function.

Homomeric Interactions: A distribution of oligomeric states

There are two distinguishing features of sHSP oligomers. First, different sHSPs form oligomers with different numbers of subunits, with oligomers ranging from dimers to 40mers across all sHSPs. For example, the sHSP from the thermophile *Methanococcus jannaschii*, Hsp16.5, forms a spherical 24mer with octahedral symmetry and the sHSP from wheat, WHSP16.9, forms a dodecameric double disc with 3_2 symmetry [15, 16]. Second, metazoan sHSPs such as α B form a polydisperse population of oligomers containing different numbers of subunits [17]. Species within the population are in a highly dynamic equilibrium, making it almost impossible to isolate or capture a single discrete species for detailed investigation. Furthermore, the species and the equilibria among them are strongly influenced by small changes in pH or by phosphorylation [18, 19]. Thus, a realistic model for α B and its metazoan relatives must take into account the nature of all oligomeric species present. This requires answers to the following questions: Are there a finite number of states, i.e., are there upper and/or lower bounds to the number of subunits that exist in the oligomeric state? Is there a continuum of N, N+1, N+2, etc. oligomeric states? and What governs the interconversion among states? In addition, questions related to α B function include: Are there certain states responsible for holdase activity? Are there (inactive) storage forms? and Do different forms interact with different clients? Although the field is far from having answers to all these questions, progress is being made at a growing pace.

In general, proteins belong to one of three categories, defined by their overall structural behavior: globular, fibrous, or intrinsically disordered. The first two categories include proteins that self-associate to form larger structures. Self-associating globular proteins adopt discrete tertiary and quaternary structures; fibrous proteins adopt repeating structures of indefinite size that are usually insoluble. α B is unusual in its tendency *not* to favor oligomers of a specific size while remaining one of the most highly soluble proteins known. In self-assembling systems, protein concentration is predicted to alter the population and/or

size of oligomeric species observed. Notably, there are large differences in α B cellular concentrations: it is a major component in the eye lens but exists at much lower concentrations in other tissues [12]. A number of studies have been directed towards understanding the size and distribution of α B oligomers. B behaves as a very large species in the range of 600 kDa (corresponding to ca. 30 subunits) when analyzed using conventional methods such as size exclusion chromatography (SEC)[20]. Simple thermodynamic considerations would predict that larger species should be favored as the concentration of α B increases. It is therefore surprising that the distribution of α B oligomers has been shown to be concentration independent over a concentration range spanning three orders of magnitude by analytical SEC [20]. However, approaches such as SEC may not detect changes in 600 kDa species that correspond to differences of one or two 20 kDa subunits. Unlike SEC, native Mass Spectrometry can provide fine-grained information regarding α B subunit distribution. α B oligomers ranging from 10-40mers have been detected by these methods [7]. The relative abundance of oligomers ranging from 22mers to 34mers was determined and used to model the full distribution of oligomeric states present under a specific condition [19]. Of particular note, the data appear to fit a Poisson distribution of species centered around 26mers to 28mers, in concordance with earlier SEC observations. Such a distribution is consistent with a model in which interactions responsible for each of the oligomeric species are isoenergetic [19, 21]. It is notable that at physiological pH the mass distribution observed by native MS is biased to oligomers containing even numbers of subunits (i.e., 24mer, 26mer, 28mer. etc.)[19]. The bias for even numbered oligomers is lost under acidic conditions (pH 5)[19]. Global changes in the oligomeric state as a function of pH have also been observed by SAXS, where the ensemble of particles at low pH conditions have a larger radius of gyration and maximum diameter than under physiological conditions[21]. The MS observations are consistent with a dimeric building block being largely responsible for oligomer assembly under physiological conditions. The data suggest that monomeric units exist, at least transiently, at lower pH and can be incorporated into and/or dissociate from α B oligomers. Thus, the homomeric interactions that define and determine α B dimers are fundamental to the structure and properties of α B oligomers. The dimer-monomer transition implied by the MS analysis suggests these interactions are pH dependent. Therefore, a model for α B at physiological pH should be based mainly on dimeric units whose dimer interface is sensitive to changes in pH.

MS analysis suggests that a continuum of oligomers exist simultaneously but provides no direct information about the existence of smaller species (<10mers) that might represent assembly intermediates. However, sedimentation velocity (SV) measurements by analytical ultracentrifugation offer additional insights. Concordant with the native MS, α B sediments as a broad distribution with sedimentation coefficients that correspond to a range between 18mers and 36mers [10]. Under slight destabilizing conditions (1.0 M guanidine), these large species are no longer detected and discrete species corresponding to dimers, hexamers, dodecamers, and 18mers are detected. Such species may represent structural intermediates from which larger species are assembled. These species are quite evanescent, as under an only slightly higher guanidine concentration of 1.5 M α B sediments entirely as monomer. Altogether the data suggest that α B exists as a distribution of oligomeric species that are built of units that include dimers, hexamers, and multiples of hexamers, but that the lifetime and population of the smaller species limit their detection under native conditions.

Homomeric Interactions: Structural Features

The behavior of α B described poses a serious challenge to structural biology. Specifically, how much can we ever know about the structures that make up the distribution of species observed? What features do they have in common and how do they interconvert? In this section, we will present current knowledge regarding α B structure and highlight the

homomeric interactions that determine structure as well as those that appear to be responsible for the polydisperse nature.

THE ALPHA CRYSTALLIN DOMAIN (ACD)

Of the three regions that comprise a sHSP (Figure 1), the central ACD is the best characterized. As the only conserved element of sHSPs, the properties of ACDs are likely fundamental to function [15, 22]. When expressed on their own, ACDs fold autonomously into a discrete structure, allowing atomic-level structures of many ACDs to be solved [23–26]. The fact that ACDs exhibit behavior more like a “normal” globular protein identifies the NTR and/or the CTR as determinants of polydispersity. In all (wild-type) sHSPs that have been structurally characterized, the ACD forms a homodimer and is therefore considered responsible for the lowest level of self-assembly, i.e., between monomer and dimer [23–26]. Protomer structures of the ~90-residue ACDs are highly similar from bacterial to human examples, while there is diversity at the level of dimer construction. All determined structures exhibit a 6-to-8-stranded -sandwich structure for the ACD protomer, (Figure 2). The available structures of sHSP oligomers from non-metazoan organisms contain ACD homodimers built via domainswapping. All metazoan ACD structures solved to date show an antiparallel homodimer with a dimer interface predominantly consisting of a long 6+7 strand [23–26](Figure 2). Structural differences among ACD structures have been previously discussed in detail [25]

Relevant to the topic of protein-protein interactions, a notable feature in available ACD structures is the remarkable plasticity they exhibit with respect to the register of the dimer interface. Three different strand registrations have been observed, one of which has been observed under neutral pH conditions [25]. Importantly, ACD crystals obtained at higher or lower pH have registers shifted by one residue in either direction. While it is not clear what functional significance the observed dimers of differing strand register have, the plasticity that such behavior implies is likely relevant. The ability to form not just a single discrete anti-parallel intermolecular β -sheet but rather numerous ones (at least as captured in crystals) suggests that even the ACD region has evolved to allow for dynamic behavior that could contribute to polydispersity of oligomers. Whether the different dimer interfaces can interconvert directly via a strandsliding mechanism or whether the conversion goes via a dimer-to-monomer dissociation is an open question. Relevant to this point, studies of the isolated α B ACD reveal that monomeric ACD becomes more favorable at lower pH [6]. Altogether, the structures and properties of the α B ACD homodimer discussed here are consistent with the pH-dependent dimermonomer exchange proposed on the basis of distributions observed by native MS for oligomeric α B [19]. Thus, intrinsic properties that contribute to the special behavior of α B oligomers are recapitulated in isolated ACDs, allowing for their more detailed parsing using the more tractable ACD dimers for certain investigations.

THE C-TERMINAL REGION (CTR)

Unlike the ACD, the CTR does not fold autonomously. Available structural information regarding CTRs comes from crystal structures of ACDs that also contain a portion of the CTR and from NMR studies of α B oligomers. CTRs of metazoan sHSPs are diverse both in length and sequence, with the only identifiable conserved element being the so-called IXI/V motif, a threeresidue sequence of alternating isoleucines (or valine)[22]. The IXI/V motif tends to be closer to the ACD than to the extreme C-terminal end of sHSPs, but the exact spacing between the end of an ACD and the IXI/V is not conserved among sHSPs. Existing information regarding the CTR of α B and its relatives highlights two main properties: 1) binding of the IXI/V motif into a groove at the edge of an ACD called the β 4/ β 8 groove (Figure 2) and 2) high flexibility of the extreme C-terminus.

The C-terminal IXI/V motif has been observed bound to the $\beta 4/\beta 8$ groove of an ACD in many experimentally determined structures of sHSPs. In all crystal structures where such an interaction is possible (i.e., the IXI/V region is contained in the protein constructs whose structures were determined) an interaction is observed [15, 16, 26]. Indeed, the bound state of this motif is observed across multiple kingdoms of life. In all but one example, (a truncated form of αA crystallin), the interaction is intermolecular, i.e., an IXI/V of one dimeric unit is bound to the ACD of another dimer (Figure 2). Although crystal packing could enable or enhance such “intermolecular” contacts, the inter-dimer contact is also observed in solid-state NMR studies of αB oligomer, making such an explanation less likely [27]. Furthermore, intermolecular binding of a peptide corresponding to the αB CTR to αB ACD homodimers has been observed in solution-state NMR studies [28]. That the interaction between the two isolated αB regions closely recapitulates what is observed in oligomers supports the idea that the ACD/CTR interaction contributes to the structure and properties of αB . Solid-state NMR spectra of αB oligomer reveal a pH-dependent alteration in the $\beta 4/\beta 8$ groove and IXI/V motif [27]. Specifically, single resonances are observed for residues in these structural elements under neutral conditions, while multiple resonances are detected at pH 6.5. The resonances at pH 7.5 correspond to the IXI-bound state, while the heterogeneity implied by multiple resonances at pH 6.5 points to a plasticity in the interaction. The observation of distinct multiple resonances as opposed to a single averaged resonance indicates that different structural copies of the $\beta 4/\beta 8$ groove and the IXI/V motif exist within oligomers, although the differences could be fairly subtle. As will be discussed in a later section, the ACD/CTR interaction serves to define the second level of protein-protein interaction in αB oligomers, namely hexameric structures comprised of three αB dimers, each linked via their IXI/V motif to their neighbor (Figure 3b). Whether the interaction contributes thermodynamically to oligomer assembly is less clear. Notably, in many but not all sHSPs, mutation of the IXI/V motif results in reduced mass oligomers [29–32].

The CTR/ACD protein-protein interaction must be understood and interpreted not only in the context of structural properties but also in terms of the prevailing dynamics. Solution-based NMR methods on oligomeric sHSPs are able to detect resonances derived from the last 10-15 residues of the protein indicating this region is disordered and flexible relative to the rest of the protein [33]. Resonances detected by this method start at the extreme C-terminus and end in close proximity to the IXI/V motif. For example, the most N-terminal resonance detected in human αB is three residues after the second Ile in the IXI/V motif (Figure 1)[33]. However, in αA crystallin, the first detectable residue precedes the first Ile of its IXI/V motif by one residue. The observations suggest that small differences in sequence and possibly solution conditions can affect the binding of the IXI/V motif and the flexibility of the CTR [33].

Solution NMR methods that specifically detect methyl groups in highly deuterated molecules are able to observe resonances for the Ile residues in αB oligomers. Among the nine resonances observed are two very strong resonances that have been assigned as the two Ile residues of the IXI/V motif. Relaxation data for these peaks are most consistent with the isoleucines existing in a predominantly flexible state [8]. Thus, similar to the observation that the homodimer interaction that defines the ACD homodimer exhibits dynamic and structural plasticity, the CTR/ACD interaction appears also to have a strongly dynamic character that is dependent on conditions including pH and temperature [27, 34].

THE N-TERMINAL REGION (NTR)

The NTR remains the most enigmatic region of αB as its properties render it difficult to study biochemically and structurally. The NTR is the most divergent region among sHSPs both in length and sequence [15, 22].

Nevertheless, it is clear that the NTR is largely responsible for the assembly of oligomers and their distribution. Truncation of the NTR yields an α B that forms oligomers that are both considerably smaller than normal (species ~40-180kDa can be observed) and that show a concentration dependence [20]. This assigns the role of driving the assembly of large α B oligomers and the dynamic distribution of oligomers to the NTR. Such a role hints at some special feature(s) of NTR sequences, but what these might be remains to be uncovered. Intriguingly, insertion of a small portion (14 residues) of the NTR of HSPB1/HSP27, which forms large, polydisperse oligomers like α B, into that of a bacterial sHSP that forms distinct 24mers that can be crystallized generated a chimeric sHSP that forms a 48mer [35]. The finding implies that the peculiar properties of NTRs are both context dependent and that one can evolve sHSPs that form different oligomers simply by altering the NTR. In this regard, we note that the NTR is subject to phosphorylation in α B and other human sHSPs (Figure 1) [36, 37]. Serine phosphorylation or its mimicry by serine-to-aspartate substitutions is reported to shift the distribution of oligomers towards smaller species when analyzed by SEC [18, 37]. As phosphorylation is thought to produce more active α B, the result implies that smaller species may represent holdase-active forms, but this implication has not been confirmed experimentally. Native MS analysis of α B phosphomimics provides a slightly different picture. Here, oligomers within the same range as seen in the native species are observed, but there appears to be an effect on the distribution of species [40]. A single Ser-to-Asp phosphomimic was indistinguishable to the native α B whereas a double substitution (there are three phosphorylation sites recognized on α B) shifts the distribution from the native dimer-biased distribution (i.e., oligomers with even numbers of subunits) to a smoother distribution in which odd numbered oligomers are also populated. The authors of this study suggest that phosphorylation of the NTR is associated with a weakening of the dimer interface but given current knowledge of the elements that define α B dimers it is difficult to propose a structurally sensible model for how this might be accomplished.

Presently, the only experimentally-derived structural information available for the α B NTR comes from ssNMR studies. The first five N-terminal residues are highly flexible and the remainder of the NTR displays significant heterogeneity [21]. While only single resonances are observed for ACD residues in spectra of α B oligomers at pH 7.6, multiple and/or lower intensity resonances are observed for many NTR residues, indicating the presence of heterogeneous states [21]. For example, there are eight distinct resonances apparent for nuclei from Met68 indicating there are at least 8 different environments for the residue [6]. Whether different environments exist simultaneously within a single oligomer species or are specific to different oligomeric species is unknown, but it is likely that both explanations are correct. Despite the heterogeneity, there are restraints observed by ssNMR consistent with helical structure (for residues 14-17 and 27-31) and with an antiparallel -hairpin structure (for residues 48-50 and 61-63) (Figure 1)[21]. Though the NMR observations indicate that such secondary structure exists, the multiple resonances observed for these regions should be taken as an indication that they exist as an ensemble in different environments and possibly different conformations. Again, whether different NTR structures can interconvert within an oligomer or are oligomer-specific cannot be experimentally determined with available technology. Altogether, the data depict an NTR that cannot be described by a single structure or in a single unique environment.

OLIGOMER MODELS

The inherent size and polydispersity of sHSPs make them challenging targets to characterize structurally. The heterogeneity in structure and the distribution of oligomeric states imposes the reality that no single structure can accurately depict the “structure” of α B. Thus it is not (presently) possible to generate atomic-level structures of α B experimentally. However, pseudo-atomic models derived from the merging of data from different methodologies

(listed below), have been proposed for the α B oligomer (Figure 3). The key technological breakthrough was the ability to obtain detailed information at atomic level from solid-state NMR on samples of α B oligomer. The first model, proposed by Jehle and colleagues in 2011, is based predominantly on ssNMR restraints along with SAXS data and previously reported negative-stain EM density [21, 46]. Braun et al. subsequently proposed a model in which they fit the ACD dimer structure determined from the ssNMR data and homology models for the CTR and NTR into higher resolution cryo-EM density [10, 27]. Both groups used cross-linking to verify features of the models. A third model for α B oligomer has been proposed based on native MS measurements [8, 9, 19, 34]. Rather than providing a pseudo-atomic model, it presents a residue-specific model for the kinetics of oligomer subunit exchange and the distribution of oligomers observed. It is important to keep in mind that none of these models propose that a single structure can represent the oligomeric state of α B. The models each aim to provide a structural rationale for known properties of α B. Consistent with the previous sections of this article, models should have the following features: 1) be built of dimeric units in which the ACD β 6+7 strand forms the dimer interface, 2) place dimers in positions relative to each other that allows for the inter-dimer ACD-IXI/V interaction, and 3) provide a structurally reasonable explanation for the observed distribution of and dynamic interchange among oligomer species. Furthermore, realistic models should provide hints or insights regarding the “holdase” function of α B.

Jehle Model—The model proposed by Jehle et al. relies on ssNMR restraints to provide atomic level structural information and previously reported 20Å negative stain-EM density to define the overall architecture [21, 46]. ssNMR restraints were sufficient to define the structure of the ACD homodimer and the interdimer contact between the C-terminal IXI/V motif and the ACD B4/8 Groove within the oligomer (Figure 2). Three copies of ACD dimers, each with its CTR IXI/V bound into the groove of a neighboring ACD fit into the prominent triangular torus feature observed in the EM density, thereby satisfying two of the requirements of a model as outlined above (Figure 3b). This provides an atomic-level model of a 3-fold symmetric hexameric (trimer of dimers) torus for α B residues 60-163 (~60% of the protein sequence). The negative-stain EM density indicates an arrangement of four hexameric rings in tetrahedral symmetry, thereby providing a way to form a 24-subunit oligomer (Figure 3c,d top). The inherent symmetry of the 24mer aided the modeling of the oligomer and therefore a pseudo-atomic model for the 24mer was generated rather than other more abundant species. In the 24mer model, all copies of the ACD dimer and the CTR through the IXI/V sequence are identical, consistent with the single set of ssNMR resonance observed for these regions.

As previously discussed, there are limited NMR restraints for the NTR and multiple conformations and/or environments exist simultaneously. For the sake of simplicity, secondary structure for the NTR was generated using homology modeling biased towards structures containing secondary structure consistent with sparse ssNMR restraints (Figure 1). Placement of these elements in the interior of the 24mer was based on the inability to fit experimental SAXS data of α B oligomer with a hollow particle model. Cysteine cross-linking was used to validate the intermolecular proximity for regions of the NTR within the oligomer, suggesting that the placement of NTRs relative to each other is realistic.

A 24mer represents only ~5% of the oligomeric population of α B and is at the lower end of the distribution observed by native MS [19]. Does the Jehle model provide hints about how 26mers, 28mers, etc. are built and why their interchange is so facile? By placing the NTRs primarily on the inside of their model, Jehle et al., posited that additional α B dimers can pack in a region between hexameric rings to build larger oligomers from the 24mer scaffold. There are six such sites in the model, allowing for oligomers up to 36mers, encompassing the majority of the oligomeric population. Each site within the symmetrical 24mer is

identical, providing a way to isoenergetically form 26mer, 28mer, and so on, consistent with the normal distribution of oligomers.

Braun Model—Braun and colleagues used cryo-EM density that afforded higher resolution (9.4 Å) than their earlier 20 Å negative-stain EM density [10]. Similar to the negative stain data, density corresponding to hexameric rings are a prominent feature of the cryo-EM density and trimers of the ssNMR-determined ACD dimers were placed into this density. New features observed in the cryo-EM density include density on the surface of the particles at the three-fold axes where hexameric rings converge and an absence of density in the core of the particle (Figure 3b–d bottom). To account for the density at the 3-fold axis NTRs were modeled into this region. The Jehle model does not satisfy this density. The resulting 24mer model is a hollow sphere compared to the more compact 24mer particle proposed in the Jehle model (Figure 3c,d). The NTR is where the two models are most divergent (Figure 1 and 3c,d). The NTRs differ between the two models not only in their placement within the oligomer but also in their secondary structure, which Braun et al. chose to generate by homology modeling that does not account for observed sparse ssNMR restraints for the NTR (Figure 1). Thus, the modeled NTR in the Braun model differs from the Jehle model and does not satisfy ssNMR restraints. As the NTR is the most structurally heterogeneous element of a polydisperse oligomer population, it is perhaps not surprising that differences in this region arise in different models.

In an effort to characterize oligomers other than 24mers, single particle images were surveyed for projections consistent with a range of oligomers (biased towards hexamer-based oligomers). Images consistent with a broad range of oligomers were indeed observed, suggesting that though there is a bias towards the 24mer-based assembly and for oligomers with even numbers of subunits, other oligomeric structures are present in an α B oligomeric population at low, but detectable levels.

Baldwin model—A third model for α B oligomer is based on solution NMR relaxation and native MS measurements [8, 9, 19, 34]. Rather than provide a structural picture, the goal of this model is to explain the underlying thermodynamics and kinetics that drive oligomer formation and subunit exchange. The experimental observations on which the model is based are 1) subunit exchange rate between α A and α B oligomers measured by MS and 2) measured relaxation properties of NMR peaks assigned to the C-terminal IXI/V motif in the context of oligomer in solution. Methyl-TROSY NMR data indicate that the IXI/V motif exists in two states, a highly populated dynamic flexible state and a lowly populated (~2% of the population) bound state [34]. The unobserved bound state is inferred to possess properties (i.e., NMR resonances) that would be consistent with bound states observed by ssNMR and crystallographic methods. The proposed predominance of a mobile IXI/V motif contrasts with previously observed ssNMR, crystallographic data, and the observation of CTR peptide binding to α B ACDs in solution by NMR [27, 15, 16, 28, 26]. It is presently unclear whether the discordance among these observations can be rationalized by the variable experimental conditions required for each specialized method and/or differences in what species are actually detected by the various techniques [34].

An exchange rate between the abundant flexible IXI/V and the inferred IXI bound state calculated from NMR relaxation measurements is similar to the rate of subunit exchange between α B and α A oligomers, if one assumes two IXI/V interactions must exist in the bound state for a monomer to leave the oligomer. From this observation, Baldwin et al. suggest that the on/off rate of the CTR IXI/V motif is rate-determining for subunit exchange. The model assumes that all protomers in an oligomer are equivalent and make the maximum number of inter/intramolecular contacts possible, including an intra-dimer interaction (i.e., the ACD dimer interaction) and two CTR interactions per subunit (i.e., accepting and

donating a CTR IXI/V motif). The proposed requirement for bound IXI/V motif in subunit dissociation is novel and somewhat contrary to previous suggestions that the IXI/V motif makes interactions that define and/or stabilize oligomers.

The collision-cross section of α B oligomers derived from ion mobility MS was used to ascertain the overall oligomer architecture. In contrast to the EM density used as a basis for the two pseudo-atomic models, a MS-based 24mer model with octahedral symmetry has been proposed in which an ACD dimer makes each “edge”. The current model does not account for the α B NTR and offers no hypothesis or prediction of how this region contributes to subunit exchange or oligomer formation. Given the inherent heterogeneity and polydispersity of α B oligomers, it should perhaps not be surprising that different models exist. It is likely that features of each realistically represent aspects of α B. Parsing out what those are is critical in generating a more unified picture and is a challenge to be met by future efforts.

Heteromeric Homomeric Interactions: mixed α B oligomers

Oligomers composed of subunits of more than one sHSP have been inferred to exist both in cells and *in vitro* [41–45]. Of the proposed sHSP heterooligomers, the lens sHSPs α A and α B crystallin heterooligomer is the best characterized [47, 48]. However, we still have a poor understanding of how heteromeric species are assembled and the necessity of having to account for more than one type of subunit in what is already an extremely complicated system makes the prospect of generating atomic-level models seem a low probability. Nevertheless, it should be possible to identify which of the protein-protein interactions that define homomeric α B oligomers can also occur between two different types of subunit. At the lowest level, can heteromeric ACD dimers form? Such species have not been observed directly, but given the high degree of conservation among at least some of the human sHSPs (α A and α B, for example), it seems likely that heterodimeric species are possible [22]. That heterooligomer could be composed of homodimers or of heterodimers further increases the possible diversity and complexity. Similarly, the high conservation of both the β 4/ β 8 ACD groove and the IXI/V motifs among at least a subset of sHSPs predicts that it is possible to assemble heteromeric hexameric sub-structures. Indeed, it has been observed that peptides mimicking different IXI/V motifs from other sHSPs are able to bind the β 4/ β 8 ACD groove of α B [28]. The next level of assembly raises the question of interactions involving NTRs. These are the most divergent and the least is known about what the determinants for NTR-directed assembly are. A likely scenario is that the diversity among NTRs could serve as both positive and negative determinants that control the types and compositions of hetero-oligomers that can form [35].

Heteromeric Interactions: Client Binding

The single most outstanding question regarding sHSPs is how they interact with client proteins to delay the formation of insoluble protein aggregates. Fundamental knowledge such as 1) what α B species interact with clients and is this client-specific, 2) what regions within an α B oligomer interact with clients and is this client-specific, 3) what properties of a client are recognized in this interaction and 4) what is the lifetime of an α B-client interaction is currently almost completely lacking. Reports that small sequences from α B on their own display the ability to chaperone aggregation-prone proteins raise the specter that the function derives from as yet undefined properties of interactions between disordered polypeptides [39]. Further, if and how dynamic properties of sHSPs contribute to function remains unclear. Given what is known about interactions involving α B, it is reasonable to expect that those involving client interactions will also be evanescent and highly dynamic. The sheer diversity of potential unfolded client proteins that α B may encounter in a cell under stress

conditions argues that there will not be singular solutions and that once again, an inherent diversity of protein-protein interactions define the system. Although they represent an important step forward in our understanding of the structure of α B, the recent models discussed above have not revealed profound insight into how sHSPs function. Similar to studies that ultimately provided atomic-level information about α B, it is likely that the ability to glean information from simplified systems will allow for progress to be made in the future towards obtaining answers to the questions raised. This will no doubt require the use of model clients that can be characterized at the residue, if not atomic level.

Heteromeric Interactions: Folded Client Binding

There is a growing body of literature that implicates α B in the binding of specific cellular proteins such as VEGF-A, Bax, and caspase-3 that serve to involve α B in cellular processes that include angiogenesis and apoptosis [13, 14]. These proteins are stably structured, at least under native conditions, so the presumption is that α B can also interact with certain folded proteins. Are such heteromeric protein-protein interactions distinct from those involving unfolded clients? Does α B recognize an unstructured region of these otherwise folded proteins or does it capture a transiently unfolded state? These questions just scratch the surface of what promises to invoke even more diversity into α B structure and function.

CONCLUSION

The past several years have seen remarkable progress in information regarding how the sHSP α B is assembled into an array of oligomeric species. This has been made possible by technological developments and by the ability to combine information from multiple experimental details to overcome the intransigence of the system to conventional structural biology approaches. In this review, we have attempted to highlight both the new insights gained from these efforts and the challenges that remain to be met. It is clear that the inherent and highly unusual properties of α B put new demands on how we think about its structural biology. As the functional repertoire of α B and the human diseases and disorders in which it is implicated continue to expand, interest in these fascinating protein assemblies and their highly unusual properties will continue to grow apace.

Acknowledgments

R. E. K. is supported by NIH grant 1R01 EY017370. S. P. D. is supported in part by NIGMS 2T32 GM008268.

References

1. Haslbeck M, Franzmann T, Weinfurter D, Buchner J, Buchner J. Some like it hot: the structure and function of small heat-shock proteins. *Nat Struct Mol Biol.* 2005; 10:842–846. [PubMed: 16205709]
2. Horwitz J. Alpha-crystallin can function as a molecular chaperone. *Proc Natl Acad Sci USA.* 1992; 89:10449–10453. [PubMed: 1438232]
3. Vicart P, Caron A, Guicheney P, Li Z, Prévost M, Faure A, Chateau D, Chapon F, Tomé F, Dupret J, Paulin D, Fardeau M. A missense mutation in the α B-crystallin chaperone gene causes a desmin-related myopathy. *Nat Genet.* 1998; 20:92–95. [PubMed: 9731540]
4. Goldstein LE, Muffat JA, Cherny RA, Moir RD, Ericsson MH, Huang X, Mavros C, Coccia JA, Faget KY, Fitch KA, Masters CL, Tanzi RE Jr, LTC, Bush AI. Cytosolic β -amyloid deposition and supranuclear cataracts in lenses from people with Alzheimer's disease. *Lancet.* 2003; 361:1258–1265. [PubMed: 12699953]
5. Kato K, Inaguma Y, Ito H, Iida K, Iwamoto I, Kamei K, Ochi N, Ohta H, Kishikawa M. Ser-59 is the major phosphorylation site in α B-crystallin accumulated in the brains of patients with Alexander's disease. *J Neurochem.* 2001; 76:730–736. [PubMed: 11158243]

6. Jehle S, van Rossum B, Stout JR, Noguchi SM, Falber K, Rehbein K, Oschkinat H, Klevit RE, Rajagopal P. α B-crystallin: a hybrid solid-state/solution-state NMR investigation reveals structural aspects of the heterogeneous oligomer. *J Mol Biol.* 2009; 385:1481–1497. [PubMed: 19041879]
7. Aquilina JA, Benesch JL, Bateman OA, Slingsby C, Robinson CV. Polydispersity of a mammalian chaperone: mass spectrometry reveals the population of oligomers in α B-crystallin. *Proc Natl Acad Sci USA.* 2003; 100:10611–10616. [PubMed: 12947045]
8. Baldwin AJ, Hilton GR, Lioe H, Bagnieris C, Benesch JLP, Kay LE. Quaternary Dynamics of α B-Crystallin as a Direct Consequence of Localised Tertiary Fluctuations in the C-Terminus. *J Mol Biol.* 2011; 413:310–320. [PubMed: 21839749]
9. Baldwin AJ, Lioe H, Hilton GR, Baker LA, Rubinstein JL, Kay LE, Benesch JLP. The Polydispersity of α B-Crystallin Is Rationalized by an Interconverting Polyhedral Architecture. *Structure.* 2011; 19:1855–1863. [PubMed: 22153508]
10. Braun N, Zacharias M, Peschek J, Kastenmüller A, Zou J, Hanzlik M. Multiple molecular architectures of the eye lens chaperone α B-crystallin elucidated by a triple hybrid approach. *Proc Natl Acad Sci U S A.* 2011; 108:20491–20496. [PubMed: 22143763]
11. Tallot P, Grongnet, David JC. Dual perinatal and developmental expression of the small heat shock proteins [FC12] α B-crystallin and Hsp27 in different tissues of the developing piglet. *Biol Neonate.* 2003; 83:281–288. [PubMed: 12743458]
12. de Jong, WW.; Bloemendal, H., editors. *Molecular and Cellular Biology of the Eye Lens.* New York: Wiley; 1981. p. 221–278.
13. Kase S, He S, Sonoda S, Kitamura M, Spee C, Wawrousek E, Ryan SJ, Kannan R, Hinton DR. α B-crystallin regulation of angiogenesis by modulation of VEGF. *Blood.* 2010; 115:3398–3406. [PubMed: 20023214]
14. Hu WF, Gong L, Cao Z, Ma H, Ji W, Deng M, Liu M, Hu XH, Chen P, Yan Q, Chen HG, Liu J, Sun S, Zhang L, Liu JP, Wawrousek E, Li DW. α A- and α B-crystallins interact with caspase-3 and Bax to guard mouse lens development. *Curr Mol Med.* 2012; 12:177–187. [PubMed: 22280356]
15. Kim KK, Kim R, Kim SH. Crystal structure of a small heatshock protein. *Nature.* 1998; 394:595–599. [PubMed: 9707123]
16. van Montfort RL, Basha E, Friedrich KL, Slingsby C, Vierling E. Crystal structure and assembly of a eukaryotic small heat shock protein. *Nat Struct Biol.* 2001; 8:1025–1030. [PubMed: 11702068]
17. Haley DA, Horwitz J, Stewart PL. The small heat-shock protein, α B-crystallin, has a variable quaternary structure. *J Mol Biol.* 1998; 277:27–35. [PubMed: 9514758]
18. Ito H, Kamei K, Iwamoto I, Inaguma Y, Nohara D, Kato K. Phosphorylation-induced change of the oligomerization state of α B-crystallin. *J Biol Chem.* 2001; 276:5346–5352. [PubMed: 11096101]
19. Baldwin AJ, Lioe H, Robinson CV, Kay LE, Benesch JLP. α B-crystallin polydispersity is a consequence of unbiased quaternary dynamics. *J mol biol.* 2011; 413:297–309. [PubMed: 21839090]
20. Horwitz J. α B-Crystallin: The Quest For A Homogeneous Quaternary Structure. *Exp Eye Res.* 2009; 88:190–194. [PubMed: 18703051]
21. Jehle S, Vollmar BS, Bardiaux B, Dove KK, Rajagopal P, Gonen T, Oschkinat H, Klevit RE. N-terminal domain of α B-crystallin provides a conformational switch for multimerization and structural heterogeneity. *Proc Natl Acad Sci U S A.* 2011; 108:6409–6414. [PubMed: 21464278]
22. Taylor RP, Benjamin IJ. Small heat shock proteins: a new classification scheme in mammals. *J Mol Cellular Cardiol.* 2005; 38:433–444. [PubMed: 15733903]
23. Bagnris C, Bateman OA, Naylor CE, Cronin N, Boelens WC, Keep NH, Slingsby C. Crystal structures of α B-crystallin domain dimers of α B-crystallin and Hsp20. *J Mol Biol.* 2009; 392:1242–1252. [PubMed: 19646995]
24. Baranova EV, Weeks SD, Beelen S, Bukach OV, Gusev NB, Strelkov SV. Three-dimensional structure of α -crystallin domain dimers of human small heat shock proteins HSPB1 and HSPB6. *J Mol Biol.* 2011; 411:110–122. [PubMed: 21641913]

25. Clark AR, Naylor CE, Bagnris C, Keep NH, Slingsby C. Crystal structure of R120G disease mutant of human α B-crystallin domain dimer shows closure of a groove. *J Mol Biol.* 2011; 408:118–134. [PubMed: 21329698]
26. Laganowsky A, Benesch JLP, Landau M, Ding L, Sawaya MR, Cascio D, Huang Q, Robinson CV, Horwitz J, Eisenberg D. Crystal structures of truncated α A and α B crystallins reveal structural mechanisms of polydispersity important for eye lens function. *Protein Sci.* 2010; 19:1031–1043. [PubMed: 20440841]
27. Jehle S, Rajagopal P, Bardiaux B, Markovic S, Khne R, Stout JR, Higman VA, Klevit RE, van Rossum BJ, Oschkinat H. Solid-state NMR and SAXS studies provide a structural basis for the activation of α B-crystallin oligomers. *Nat Struct Mol Biol.* 2010; 17:1037–1042. [PubMed: 20802487]
28. Delbecq SP, Jehle S, Klevit R. Binding determinants of the small heat shock protein, α B-crystallin: recognition of the "IXI" motif. *EMBO J.* 2012 Ahead of print.
29. Saji H, Iizuka R, Yoshida T, Abe T, Kidokoro S-I, Ishii N, Yohda M. Role of the IXI/V motif in oligomer assembly and function of StHsp14.0, a small heat shock protein from the acidothermophilic archaeon, *Sulfolobus tokodaii* strain 7. *Proteins.* 2008; 71:771–782. [PubMed: 17979194]
30. Studer S, Obrist M, Lentze N, Narberhaus F. A critical motif for oligomerization and chaperone activity of bacterial α -heat shock proteins. *Eur J Biochem.* 2002; 269:3578–3586. [PubMed: 12135498]
31. Pasta SY, Raman B, Ramakrishna T, Rao CM. The IXI/V motif in the C-terminal extension of α -crystallins: alternative interactions and oligomeric assemblies. *Mol Vis.* 2004; 10:655–662. [PubMed: 15448619]
32. Giese KC, Vierling E. Mutants in a small heat shock protein that affect the oligomeric state. *J Biol Chem.* 2004; 279:32674–32683. [PubMed: 15152007]
33. Treweek TM, Rekas A, Walker MJ, Carver JA. A quantitative NMR spectroscopic examination of the flexibility of the C-terminal extensions of the molecular chaperones, α A- and α B-crystallin. *Exp Eye Res.* 2010; 91:691–699. [PubMed: 20732317]
34. Baldwin AJ, Walsh P, Hansen DF, Hilton GR, Benesch JLP, Sharpe S, Kay LE. Probing dynamic conformations of the high-molecularweight α B-crystallin heat shock protein ensemble by NMR spectroscopy. *J Am Chem Soc.* 2012; 134:15343–15350. [PubMed: 22916679]
35. McHaourab HS, Lin Y-L, Spiller BW. Crystal structure of an activated variant of small heat shock protein Hsp16.5. *Biochemistry.* 2012; 51:5105–5112. [PubMed: 22670769]
36. Miesbauer LR, Zhou X, Yang Z, Yang Z, Sun Y, Smith DL, Smith JB. Post-translational modifications of water-soluble human lens crystallins from young adults. *J Biol Chem.* 1994; 269:12494–12502. [PubMed: 8175657]
37. Stokoe D, Engel K, Campbell DG, Cohen P, Oaestep M. Identification of MAPKAP kinase 2as a major enzyme responsible for the phosphorylation of the small mammalian heat shock proteins. *FEBS Lett.* 1992; 313:307–313. [PubMed: 1332886]
38. McDonald ET, Bortolus M, Koteiche HA, Mchaourab HS. Sequence, structure, and dynamic determinants of Hsp27 (HspB1) equilibrium dissociation are encoded by the N-terminal domain. *Biochemistry.* 2012; 51:1257–1268. [PubMed: 22264079]
39. Bhattacharyya J, Udupa EGP, Wang J, Sharma KK. Mini- α B Crystallin : A Functional Element of α B-Crystallin with Chaperone-like Activity. *Biochemistry.* 2006; 45:3069–3076. [PubMed: 16503662]
40. Aquilina JA, Benesch JLP, Ding LL, Yaron O, Horwitz J, Robinson CV. Phosphorylation of α B-crystallin alters chaperone function through loss of dimeric substructure. *J Biol Chem.* 2004; 279:28675–28680. [PubMed: 15117944]
41. Zantema A, Verlaan-De Vries M, Maasdam D, Bol S, van der Eb A. Heat shock protein 27 and α B-crystallin can form a complex, which dissociates by heat shock. *J Biol Chem.* 1992; 267:12936–12941. [PubMed: 1618790]
42. Kato K, Goto S, Inaguma Y, Hasegawa K, Morishita R, Asano T. Purification and characterization of a 20-kDa protein that is highly homologous to α B crystallin. *J Biol Chem.* 1994; 269:15302–15309. [PubMed: 8195168]

43. Sugiyama Y, Suzuki A, Kishikawa M, Akutsu R, Hirose T, Waye MMY, Tsui SKW, Yoshida S, Ohno S. Muscle develops a specific form of small heat shock protein complex composed of MKBP/ HSPB2 and HSPB3 during myogenic differentiation. *Biochemistry*. 2000; 275:1095–1104.
44. Bukach OV, Glukhova AE, Seit-Nebi AS, Gusev NB. Heterooligomeric complexes formed by human small heat shock proteins HspB1 (Hsp27) and HspB6 (Hsp20). *Biochim Biophys Acta*. 2009; 1794:486–495. [PubMed: 19100870]
45. Engelsman J, Boros S, Dankers PYW, Kamps B, Vree Egberts WT, Bo de CS, Lane LA, Aquilina JA, Benesch J, Robinson CV, de Jong WW, Boelens WC. The small heat-shock proteins HSPB2 and HSPB3 form well-defined heterooligomers in a unique 3 to 1 subunit ratio. *J Mol Biol*. 2009; 393:1022–1032. [PubMed: 19715703]
46. Peschek J, Braun N, Franzmann TM, Georgalis Y, Haslbeck M, Weinkauff S, Buchner J. The eye lens chaperone alpha-crystallin forms defined globular assemblies. *Proc Natl Acad Sci U S A*. 2009; 106:13272–13277. [PubMed: 19651604]
47. Siezen RJ, Bindels JG, Hoenders HJ. The quaternary structure of bovine alpha-crystallin. Chemical crosslinking with bifunctional imido esters. *Eur J Biochem*. 1980; 107:243–249. [PubMed: 7398636]
48. van den Oetelaar PJ, Van Someren PF, Thomson JA, Siezen RJ, Hoenders HJ. A dynamic quaternary structure of bovine alphacrystallin as indicated from intermolecular exchange of subunits. *Biochemistry*. 1990; 29:3488–3493. [PubMed: 2354148]

Figure 1.

α Bdomain architecture, primary and secondary structure. The three domains/regions that define sHSPs are highlighted on the sequence of α B, with NTR residues in blue text, ACD residues in green, and CTR residues in red. Every tenth residue is underlined as a point of reference. In the NTR, three known phosphorylation sites are shown in cyan. The IXI/V motif in the CTR is shown in the black box. Secondary structure is summarized in three rows above the sequence, as cylinders (α -helix) and arrows (β -strand). TOP (magenta): secondary structure in the Braun model (PDB ID 2ygd). MIDDLE (black): secondary structure in the Jehle model. BOTTOM: experimentally-derived structural information, with secondary structure determined from ssNMR-based ACD structure shown in light blue, structure inferred from sparse ssNMR restraints shown in grey, and flexible regions measured by ssNMR and solution based NMR are shown in open circles.

Figure 2. ssNMR structure of the α B ACD dimer (PDB 2klr) with the CTR IXI/V motif bound. Residues involved in intermolecular contacts are highlighted on both protomers. Regions involved in intra-dimer contacts that define the dimer interface are shown in blue and the β 4/ β 8 groove involved in inter-dimer interactions with the CTR IXI/V motif (from a neighboring dimer) are shown in red. The CTR sequence PERTIPITREEK is shown in black (bound to one protomer for clarity) and the Iles of the IXI/V motif are rendered as sticks. Two views of the ACD are presented, rotated 180° about the x-axis.

Figure 3.

Comparison of the two pseudo-atomic models of an α B 24mer. The NTR (Blue), ACD (Green), and CTR (red) are displayed in different colors in the Jehle model (top row) and the Braun model (bottom row). The two models are aligned in the middle row, with the Jehle model in cyan and the Braun model in black. The hierarchy of assembly for the 24mer models is represented across columns A-D. The dimeric building block defined by intra-dimer contacts (A) are shown assembled into hexameric rings through CTR/ACD inter dimer interactions (B) and four hexamers are assembled through NTR interactions to form the 24mer oligomer (C and D). The models differ significantly in the placement of the NTR. The view in column D (rotated 55° about the y-axis relative to column C) clearly shows the Braun model to be a hollow particle with the NTR on the surface while the core of the particle is filled with the NTR in the Jehle model.