Published in final edited form as:

Biochem Biophys Res Commun. 2013 October 25; 440(3): . doi:10.1016/j.bbrc.2013.09.094.

Developmental *cis*-regulatory analysis of the cyclin D gene in the sea urchin *Strongylocentrotus purpuratus*

Christopher M. McCarty^{a,b,c,e} and James A. Coffman^{a,b,d,e}

^aMount Desert Island Biological Laboratory, PO Box 35, Salisbury Cove, Maine, USA 04672

^bThe Graduate School of Biomedical Science and Engineering, The University of Maine, Orono, ME USA 04469

Abstract

Cyclin D genes regulate the cell cycle, growth and differentiation in response to intercellular signaling. While the promoters of vertebrate cyclin D genes have been analyzed, the *cis*-regulatory sequences across an entire cyclin D locus have not. Doing so would increase understanding of how cyclin D genes respond to the regulatory states established by developmental gene regulatory networks, linking cell cycle and growth control to the ontogenetic program. Therefore, we conducted a *cis*-regulatory analysis on the cyclin D gene, *SpccD*, of the sea urchin, *Strongylocentrotus purpuratus*, during embryogenesis, identifying upstream and intronic sequences, located within six defined regions bearing one or more *cis*-regulatory modules each.

Keywords

Sea urchin embryo; development; Cyclin D; *Strongylocentrotus purpuratus*; *cis*-regulatory analysis

INTRODUCTION

Cyclin D family genes link cell cycle control to the genomically encoded program of multicellular development [1–4], by responding to transcription factors activated by intercellular signals [1, 2] to promote cell cycle transit from G1 to S phase [3–5], cell growth [6], and differentiation in multiple systems including spermatogonia [7], myeloid cells [8], skeletal muscle [9], and the nervous system [10]. Mis-expression is deleterious: cyclin D genes are over-expressed in many cancers [11–15], and in the sea urchin, ectopic expression during cleavage is lethal [16].

The sea urchin *Strongylocentrotus purpuratus* has one cyclin D gene [17], which is expressed in a dynamic pattern during embryogenesis [16]. At blastula stage, expression is global. During gastrulation, expression restricts to gut, oral ectoderm, and ciliary band [16], a pattern displayed by ~40% of genes expressed in the sea urchin embryo, concordant with

© 2013 Elsevier Inc. All rights reserved.

Correspondence to: James A. Coffman.

^ccmccarty@mdibl.org

^djcoffman@mdibl.org

^e12072889880

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

continued growth and proliferation [18]. Cyclin D regulates patterning of the embryo, since its knockdown produces malformed larvae [16].

Many developmentally important genes have now been linked into developmental gene regulatory networks (dGRNs) [19–21]. The linkages between genes are mediated by *cis*-regulatory modules (CRMs) that function by binding transcription factors, many of which are activated by intercellular signaling [19, 21]. Typically, developmentally regulated genes contain multiple CRMs [21]. Although the promoters of various cyclin D genes have been dissected [2, 22], no cyclin D gene has been subjected to a comprehensive *cis*-regulatory analysis to identify CRMs. Such an analysis is a necessary step in elucidating how dGRNs control cell proliferation. We therefore undertook a *cis*-regulatory analysis *SpccycD* in the sea urchin *Strongylocentrotus purpuratus*, which is highly amenable such analyses [23–30]. We used a high throughput method from Nam and colleagues [31] to examine the entire locus.

MATERIALS AND METHODS

Embryo culture

Strongylocentrotus purpuratus obtained from the Pt. Loma Marine Invertebrate Lab (Lakeside, CA) were maintained in a seawater aquarium at 12°C. Gametes were obtained by shaking or injection with 0.55 M KCl [32]. Embryos were cultured at 15°C.

Preparing reporter constructs and BAC DNA

Regions to be assayed were amplified from *SpccycD* BAC or genomic DNA template by polymerase chain reaction (PCR) using high fidelity DNA polymerases (from Roche or New England Biolabs) and the primers in Supplemental Table 1.

‘13-tag’ reporters were obtained from mini-preps of cultures grown with chloramphenicol (12.5 µg/ml) and Copy Control Induction Solution (Epicentre). These reporters were modified from that presented in [31] (J. Nam, personal communication, May, 2011) in that the *Sp-gatae* basal promoter was replaced with an *Sp-nodal* basal promoter using primers new_mNBP and end_core-polyA (Supplemental Table 1) in a reaction employing the cycling shown in Supplemental Table 2.

Amplified regions were ligated to EpGFPII [33], after which the resultant constructs were linearized; or linked as directed [31] to the indicated modified 13-tag reporters (Supplemental Table 3). 13-tag-region-linked reporter products visualized using SYBR Safe and a Safe Imager (Invitrogen) rather than UV light were gel purified (Nucleospin Gel and PCR Cleanup, Clontech). Sequencing showed that only 13-tag-linked region 3 showed significant contamination.

BAC DNA was obtained from stab cultures of BAC 4013 F-18 mCherry (from Sp Genome Research Resource, Caltech) grown under chloramphenicol selection and purified as described [34].

Microinjection

10 µl injection solutions of EpGFPII-linked reporters [33] contained ~10 nmols reporter construct, 165 to 200 ng HindIII digested genomic DNA, 0.12 M KCl. Injection solutions comprising 13-tag-linked regions were made as described [31], except Master Pool volume in the 10 µl injection mix for all experiments but #8 was increased from 0.5 to 1 µl. In Experiment #8, Master Pool concentration was 5× greater than in other presented experiments. Embryos in seawater containing 1 mM PABA salt were injected with the appropriate reporters using standard methods [35]. About 100–150 embryos were injected

with EpGFPII-linked regions, and 200 with 13-tag reporter-linked regions. Hatched embryos were transferred to ~1.5 ml artificial seawater for subsequent culture.

Preparing RNA, DNA and cDNA

To assay endogenous *SpccycD*, embryos were cultured at ~1200 embryos / 4 ml. RNA was obtained (Rneasy Plus mini kit, Qiagen), and DNA was removed as directed. RNA equivalent to 30 ng per 20 μ l was converted to cDNA using random hexamers (FirstStrand cDNA Synthesis kit, Invitrogen). RNA and DNA from injected embryos were obtained with a DNA/RNA ALL Prep kit (Qiagen). RNA was treated with DNase for 30 minutes as directed. cDNA from injection with EpGFPII-linked regions was made with random hexamers, with 3 μ l RNA for each 20 μ l reaction. cDNA from injection with 13-tag-linked regions was made from RNA equivalent to 3 μ l per 20 μ l, and 13-tag vector-specific primer [31] using the FirstStrand Synthesis Kit. The exception was Experiment #5, where random hexamers and the VILO cDNA Synthesis Kit (Invitrogen) were used.

Real-Time PCR

Real-Time PCR (relative quantification) was conducted using Perfecta SYBR Green Fast Mix (Quanta BioSciences) and a LightCycler 480 II (Roche). cDNA and DNA equivalent to 1.3 μ l and 1.6 μ l, respectively, per 12 μ l were used. Unless noted, all reactions were done in duplicate. (Cycling conditions are in Supplemental Table 2).

Relative levels of *SpccycD* expression with respect to the first time point was determined using the delta-delta Ct method, with ubiquitin as normalization reference [36]. Relative GFP expression from microinjected EpGFPII-linked reporters was calculated by the same method, with further normalization to the amount injected [30]. Activity levels of microinjected BAC 4013 F-18 mCherry were determined the same way, except mCherry sequence was assayed. Expression of each region-linked 13-tag reporter was found using a 13-tag unit-specific primer [31]. Activities were determined as for EpGFPII-linked regions, but were relative to that of empty reporter 1302 for each time point.

Sequence comparisons

LvcycD sequence was from: a BAC containing 17 kb upstream of exon 1; and *LvcycD* draft sequence from SpBase [37, 38]. Comparisons were made using Family Relations II [39, 40] to identify sequences of 20 bp in *SpccycD* with 90% similarity to *LvcycD*.

Fluorescence microscopy

Eggs were arrayed on 50 mm glass bottom dishes (MatTek), fertilized and injected as described above. Injected embryos were visualized with an Axiovert 200 fluorescence microscope (Zeiss).

RESULTS AND DISCUSSION

Temporal expression of *SpccycD*

The temporal profile of embryonic *SpccycD* expression was assayed by quantitative RT-PCR. As reported previously by others [16], expression commenced ~10–12 hpf (early blastula), then increased at least up to pluteus stage (72 hpf) (Supplemental Fig. 1). Interestingly, there was substantial variation between biological replicates.

We also co-assayed the temporal activities of endogenous *SpccycD* and a bacterial artificial chromosome (BAC) bearing *SpccycD* with mCherry knocked into exon 1. This BAC encompassed sequence from ~90 kb upstream of the gene to ~13 kb downstream. Both

endogenous *SpccycD* and the injected BAC exhibited similar temporal activities (Fig. 1), suggesting the information needed to regulate embryonic *SpccycD* expression is within this BAC. Our *cis*-regulatory analysis examined from ~13 kb upstream of exon 1 to ~7 kb downstream from the end of exon 5 (Fig. 2A).

Identification of *cis*-regulatory regions

We selected 22 regions spanning upstream and intronic sequence of *SpccycD* to assay for regulatory activity (Fig. 2A). The boundaries of most were chosen based on the presence of sequences of 20 bp with 90% similarity to *LvcycD* from *L. variegatus* (Fig. 2A) [37]. This criterion was based on the fact that sequence comparisons between genes in *S. purpuratus* and *L. variegatus* reliably predict *S. purpuratus* CRMs [40, 41]. Our analysis was comprehensive: all non-exonic sequence except 1 bp between the 3' end of region 10 and the 5' end of exon 5, and 2 bp between the 3' end of region 11 and the 5' end of region 21 was tested.

Candidate *cis*-regulatory regions were assayed for activity using the '13-tag' reporters developed by Nam and colleagues [31]. Representative results are in Fig. 2B and Supplemental Fig. 2. In each experiment, a region was classified as significantly active if activity at one or more time points was 2.5 times that of the mean activity of regions in the middle 40% of the distribution [31].

Several active regions were identified. Region 5, (2.4 kb) in the first half of intron 2 (Fig. 2A) showed the strongest activity, with significant activity at all tested time points from ~10–60 hpf. This activity was ~15 times greater than that of empty reporter at its peak, and at least 2 times higher than those of the next most active regions. The next most active regions were region 2 (~3.6 kb), located ~4.6 kb upstream from the beginning of exon 1; region 6 (2.7 kb), comprising the 3' half of intron 2; region 19 (4.6 kb), in intron 4; followed by region 4 (2.1 kb), which abuts exon 1; and region 17 (2.1 kb) in intron 1 (Fig. 2A). Regions 2 and 6 always showed significant activity for at least one time point when injected without region 5-linked reporter, but not always in its presence (Supplemental Fig. 2).

Temporal activity profiles of *cis*-regulatory regions

To gain further insight into the roles of each active region, temporal activity profiles were extracted from experiments in Fig. 2B and Supplemental Fig. 2 (Supplemental Fig. 3). This analysis reveals substantial inter-experimental variation in the temporal activity profiles of each region. An exception concerned region 19, as discussed below. Possible sources of this variation include biological variability, the fact that injection solutions contained different mixtures of 13-tag-linked regions, and the fact that each time point was from a separate injection plate because it was technically possible to inject a maximum of ~200 embryos per plate. To more clearly discern canonical aspects of the temporal activity patterns, the activity values across experiments were averaged (Fig. 3).

From this analysis, we found the following patterns. Region 5's activity was highest at 10–12 hpf, when *SpccycD* is initially activated. As other regions became active, region 5's activity declined somewhat, but remained significant (Fig. 2B). Region 6 likewise showed the strongest activity at ~10 hpf. During the first ~33 hours, activities of regions 5 and 6 paralleled each other, then region 6's stabilized, suggesting that region 6 contributes to maintaining *SpccycD* expression after ~33 hpf, corresponding to gastrulation and later stages.

On average, region 2's activity peaked at ~21 hpf (Fig. 3), although peak activity varied from ~12–33 hpf (Supplemental Fig. 3). Region 2's activity peak occurred after that of

regions 5 and 6. Therefore, region 2's primary role may be to activate transcription during late blastula stage.

Region 4's activity varied considerably (Supplemental Fig. 3), but on average (Fig. 3) increased to low but stable levels by ~21–33 hpf. Thus, region 4 may contribute to maintaining *SpCycD* expression.

Region 17's activity slowly increased to stability by ~21–33 hpf (Fig 3), indicating that this region may contribute to maintenance or lineage-specific activation of *SpCycD* during and after gastrulation.

Region 19's activity peaked at ~21 hpf, the mesenchyme blastula stage (Fig. 3), suggesting that this region may act as a switch that regulates *SpCycD* at the onset of gastrulation. As noted, region 19's activity showed much less variation than those of other active regions (Supplemental Fig. 3; compare Experiments #5, 2 and 3). Therefore, region 19 may be under especially strong control.

As a control, activities of region 2-linked 13-tag vectors at 12 hpf (Supplemental Fig. 4A), and 13-tag vectors linked to unique regions (Supplemental Fig. 4B) were compared. There was significantly less variation between activities of 13-tag reporters linked only to region 2 than between those linked to different regions, indicating that differences in activity among regions could mostly be attributed to region-specific differences rather than 13-tag reporter-specific differences.

Identification of candidate *cis*-regulatory modules

Since the sizes of the identified regulatory regions ranged from ~2–5 kb (Fig. 2A), additional analysis was needed to identify CRMs, which are generally only up to several hundred bp [21]. By using a combination of computational approaches to analyze each region (Fig. 4; Supplemental Figs. 5 and 6), we identified candidate CRMs within each, the activities of several of which were verified experimentally.

Region 2 contains a 0.5 kb subregion, 2-2, encompassing sequence conserved at > 90% with *LvcycD* (Fig. 4A; Supplemental Fig. 5). Experimental analysis using both 13-tag and EpGFPII-linked versions of region 2 and subregion 2-2 showed that subregion 2-2's temporal activity mirrored region 2's (Fig. 4B, panel 1; Supplemental Fig. 7). Further analysis showed that the activities of each were detected at blastula stage by fluorescence microscopy (Figs. 4A and 4C, panel 1). Together, these findings indicate that subregion 2-2 contains a CRM.

Region 4 contains two active subregions (4-1 and 4-2; Fig. 4A). Subregion 4-1 overlaps partly with conserved sequence (Fig. 4A; Supplemental Fig. 5), and bears a potential Runx site (Supplemental Fig. 5). Sequence within subregion 4-1 was previously found by chromatin immunoprecipitation to bind the Runx protein SpRunx-1, which was shown to regulate *SpCycD* [42]. Subregion 4-2 contains a 22 bp conserved sequence (Fig. 4A; Supplemental Fig. 5), and a potential Runx site [43] (Supplemental Fig. 5). When tested for activity by fluorescence microscopy, subregions 4-1 and 4-2 were both shown to be active at gastrula stage (Fig. 4C, panel 2), suggesting that both encompass CRMs.

Our analysis of the intronic regulatory regions, which contain longer stretches of sequence conservation than the upstream regions (Fig. 4A, red lines), was chiefly computational. In this analysis, a number of sequence elements of interest were identified. Among these, were potential binding sites for TCF and Runx. Wnt-TCF signaling is known to regulate cyclin D expression [44–47]; and, as noted above, the SpRunx-1 protein is known to regulate *SpCycD*.

In addition, we searched for sequences with clustered binding sites for transcription factors identified by the program Cluster-Buster, of interest because sequences where transcription factor bindings sites cluster are hypothesized to be regulatory [48–50]; and G strings, known to mark regulatory regions in other genes [51–55]. These areas are highlighted on the sequence for each region in Supplemental Fig. 5. Identities of transcription factors identified by Cluster-Buster are in Supplemental Fig. 6.

One candidate CRM in region 5 was subregion 5-1, found 6 bp upstream of a potential transcription factor cluster site to 14 bp downstream from a potential TCF binding site (Fig. 4A, Supplemental Fig. 5). However, subregion 5-1 showed only background activity (Supplemental Fig. 2, Experiments #5 and 9). This was surprising because within its boundaries, which overlapped with conserved sequence, subregion 5-1 contains 6 potential TCF and Runx sites, respectively, most of which overlap with the transcription factor cluster site. Therefore, 5-1 may be necessary but not sufficient for region 5's activity.

Within region 6, we reasoned that the 3' two-thirds of this region could contain a CRM, as most of the potential regulatory elements of interest were found in that portion (Fig. 4; Supplemental Figs 5 and 6). This subregion, 6-1, was verified to be active (Fig. 2B, panel 6; Supplemental Fig. 2, Experiments #7, 8 and 9), and its temporal activity closely resembled region 6's (Fig. 4B).

Within region 19, a sequence termed subregion 19-1, which bears few of the potential regulatory elements of interest highlighted in Supplemental Fig. 5, showed only background activity (Supplemental Fig. 2, Experiment #9), indirectly supporting the hypothesis that the highlighted sequence elements shown for this region likely mark one or more CRMs.

Conclusions

The entire *SpcycD* locus was analyzed to identify *cis*-regulatory regions and modules (CRMs) that mediate expression. We identified intronic and upstream regions that impart distinct activity patterns, and uncovered likely CRMs in several. A future aim is to determine the specific roles of each regulatory region and candidate CRM by individual deletion of each from a BAC bearing *SpcycD*. Finally, to link *SpcycD* to GRNs that control early embryogenesis, the spatial activity of each CRM should be studied.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Dr. Anthony Robertson and Alison Coluccio provided technical advice; Christine Smith provided sequencing and technical advice; Dr. Jongmin Nam of the Davidson lab generously provided 13-tag-bearing frozen cultures and technical advice; and Dr. R. Andrew Cameron of the Davidson lab provided advice on analysis of sequence found at Sphbase. This work was supported by NIH grants R01-GM070840 to J.A.C. and Maine INBRE P20-GM103423.

Abbreviations

hpf	hours post-fertilization
CRM	cis-regulatory module
GRN	gene regulatory network
5' UTR	5' untranslated region

References

1. Murray AW. Recycling the cell cycle: cyclins revisited. *Cell*. 2004; 116:221–234. [PubMed: 14744433]
2. Klein EA, Assoian RK. Transcriptional regulation of the cyclin D1 gene at a glance. *J Cell Sci*. 2008; 121:3853–3857. [PubMed: 19020303]
3. Matsushime H, Roussel MF, Ashmun RA, Sherr CJ. Colony-stimulating factor 1 regulates novel cyclins during the G1 phase of the cell cycle. *Cell*. 1991; 65:701–713. [PubMed: 1827757]
4. Resnitzky D, Reed SI. Different roles for cyclins D1 and E in regulation of the G1-to-S transition. *Mol Cell Biol*. 1995; 15:3463–3469. [PubMed: 7791752]
5. Fu M, Wang C, Li Z, Sakamaki T, Pestell RG. Minireview: Cyclin D1: normal and abnormal functions. *Endocrinology*. 2004; 145:5439–5447. [PubMed: 15331580]
6. Datar SA, Jacobs HW, de la Cruz AF, Lehner CF, Edgar BA. The *Drosophila* cyclin D-Cdk4 complex promotes cellular growth. *Embo J*. 2000; 19:4543–4554. [PubMed: 10970848]
7. Beumer TL, Roepers-Gajadien HL, Gademan IS, Kal HB, de Rooij DG. Involvement of the D-type cyclins in germ cell proliferation and differentiation in the mouse. *Biol Reprod*. 2000; 63:1893–1898. [PubMed: 11090462]
8. Adachi M, Roussel MF, Havenith K, Sherr CJ. Features of macrophage differentiation induced by p19INK4d, a specific inhibitor of cyclin D-dependent kinases. *Blood*. 1997; 90:126–137. [PubMed: 9207446]
9. Lazaro JB, Bailey PJ, Lassar AB. Cyclin D-cdk4 activity modulates the subnuclear localization and interaction of MEF2 with SRC-family coactivators during skeletal muscle differentiation. *Genes Dev*. 2002; 16:1792–1805. [PubMed: 12130539]
10. Lukasiewicz AI, Anderson DJ. Cyclin D1 promotes neurogenesis in the developing spinal cord in a cell cycle-independent manner. *Proc Natl Acad Sci U S A*. 2011; 108:11632–11637. [PubMed: 21709239]
11. Alao JP. The regulation of cyclin D1 degradation: roles in cancer development and the potential for therapeutic invention. *Mol Cancer*. 2007; 6:24. [PubMed: 17407548]
12. Jiang W, Kahn SM, Tomita N, Zhang YJ, Lu SH, Weinstein IB. Amplification and expression of the human cyclin D gene in esophageal cancer. *Cancer Res*. 1992; 52:2980–2983. [PubMed: 1533816]
13. Maeda K, Chung Y, Kang S, Ogawa M, Onoda N, Nishiguchi Y, Ikehara T, Nakata B, Okuno M, Sowa M. Cyclin D1 overexpression and prognosis in colorectal adenocarcinoma. *Oncology*. 1998; 55:145–151. [PubMed: 9499189]
14. Sherr CJ. Cancer cell cycles. *Science*. 1996; 274:1672–1677. [PubMed: 8939849]
15. Sherr CJ. The Pezcoller lecture: cancer cell cycles revisited. *Cancer Res*. 2000; 60:3689–3695. [PubMed: 10919634]
16. Moore JC, Sumerel JL, Schnackenberg BJ, Nichols JA, Wikramanayake A, Wessel GM, Marzluff WF. Cyclin D and cdk4 are required for normal development beyond the blastula stage in sea urchin embryos. *Mol Cell Biol*. 2002; 22:4863–4875. [PubMed: 12052892]
17. Fernandez-Guerra A, Aze A, Morales J, Mulner-Lorillon O, Cosson B, Cormier P, Bradham C, Adams N, Robertson AJ, Marzluff WF, Coffman JA, Genevieve AM. The genomic repertoire for cell cycle control and DNA metabolism in *S. purpuratus*. *Dev Biol*. 2006; 300:238–251. [PubMed: 17078944]
18. Kingsley PD, Angerer LM, Angerer RC. Major temporal and spatial patterns of gene expression during differentiation of the sea urchin embryo. *Dev Biol*. 1993; 155:216–234. [PubMed: 8416835]
19. Davidson EH, Rast JP, Oliveri P, Ransick A, Caletani C, Yuh CH, Minokawa T, Amore G, Hinman V, Arenas-Mena C, Otim O, Brown CT, Livi CB, Lee PY, Revilla R, Rust AG, Pan Z, Schilstra MJ, Clarke PJ, Arnone MI, Rowen L, Cameron RA, McClay DR, Hood L, Bolouri H. A genomic regulatory network for development. *Science*. 2002; 295:1669–1678. [PubMed: 11872831]
20. Endomesoderm and Ectoderm Gene Networks. Available at: <http://sugp.caltech.edu/endomes/>

21. Davidson, EH. The Regulatory Genome: Gene Regulatory Networks in Development and Evolution. Elsevier/Academic Press; Amsterdam; Boston: 2006.
22. Tanaka T, Kubota M, Shinohara K, Yasuda K, Kato JY. In vivo analysis of the cyclin D1 promoter during early embryogenesis in *Xenopus*. *Cell Struct Funct*. 2003; 28:165–177. [PubMed: 12951437]
23. Kirchhamer CV, Davidson EH. Spatial and temporal information processing in the sea urchin embryo: modular and intramodular organization of the *CyIIIa* gene cis-regulatory system. *Development*. 1996; 122:333–348. [PubMed: 8565846]
24. Makabe KW, Kirchhamer CV, Britten RJ, Davidson EH. Cis-regulatory control of the *SM50* gene, an early marker of skeletogenic lineage specification in the sea urchin embryo. *Development*. 1995; 121:1957–1970. [PubMed: 7635044]
25. Yuh CH, Davidson EH. Modular cis-regulatory organization of *Endo16*, a gut-specific gene of the sea urchin embryo. *Development*. 1996; 122:1069–1082. [PubMed: 8620834]
26. Yuh CH, Bolouri H, Davidson EH. Cis-regulatory logic in the *endo16* gene: switching from a specification to a differentiation mode of control. *Development*. 2001; 128:617–629. [PubMed: 11171388]
27. Arnone MI, Bogarad LD, Collazo A, Kirchhamer CV, Cameron RA, Rast JP, Gregorians A, Davidson EH. Green Fluorescent Protein in the sea urchin: new experimental approaches to transcriptional regulatory analysis in embryos and larvae. *Development*. 1997; 124:4649–4659. [PubMed: 9409681]
28. Minokawa T, Wikramanayake AH, Davidson EH. cis-Regulatory inputs of the *wnt8* gene in the sea urchin endomesoderm network. *Dev Biol*. 2005; 288:545–558. [PubMed: 16289024]
29. Nam J, Su YH, Lee PY, Robertson AJ, Coffman JA, Davidson EH. Cis-regulatory control of the nodal gene, initiator of the sea urchin oral ectoderm gene network. *Dev Biol*. 2007; 306:860–869. [PubMed: 17451671]
30. Revilla-i-Domingo R, Minokawa T, Davidson EH. R11: a cis-regulatory node of the sea urchin embryo gene network that controls early expression of *SpDelta* in micromeres. *Dev Biol*. 2004; 274:438–451. [PubMed: 15385170]
31. Nam J, Dong P, Tarpine R, Istrail S, Davidson EH. Functional cis-regulatory genomics for systems biology. *Proc Natl Acad Sci U S A*. 2010; 107:3930–3935. [PubMed: 20142491]
32. Foltz KR, Adams NL, Runft LL. Echinoderm eggs and embryos: procurement and culture. *Methods Cell Biol*. 2004; 74:39–74. [PubMed: 15575602]
33. Cameron RA, Oliveri P, Wyllie J, Davidson EH. cis-Regulatory activity of randomly chosen genomic fragments from the sea urchin. *Gene Expr Patterns*. 2004; 4:205–213. [PubMed: 15161101]
34. Smith J, Davidson EH. Gene regulatory network subcircuit controlling a dynamic spatial pattern of signaling in the sea urchin embryo. *Proc Natl Acad Sci U S A*. 2008; 105:20089–20094. [PubMed: 19104065]
35. Cheers MS, Etensohn CA. Rapid microinjection of fertilized eggs. *Methods Cell Biol*. 2004; 74:287–310. [PubMed: 15575612]
36. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods*. 2001; 25:402–408. [PubMed: 11846609]
37. SpBase. Available at: <http://www.spbase.org/SpBase/>
38. Cameron RA, Samanta M, Yuan A, He D, Davidson E. SpBase: the sea urchin genome database and web site. *Nucleic Acids Res*. 2009; 37:D750–754. [PubMed: 19010966]
39. Family Jewels. Available at: <http://family.caltech.edu/>
40. Brown CT, Xie Y, Davidson EH, Cameron RA. Paircomp FamilyRelationsII and Cartwheel: tools for interspecific sequence comparison. *BMC Bioinformatics*. 2005; 6:70. [PubMed: 15790396]
41. Yuh CH, Brown CT, Livi CB, Rowen L, Clarke PJ, Davidson EH. Patchy interspecific sequence similarities efficiently identify positive cis-regulatory elements in the sea urchin. *Dev Biol*. 2002; 246:148–161. [PubMed: 12027440]
42. Robertson AJ, Coluccio A, Knowlton P, Dickey-Sims C, Coffman JA. Runx expression is mitogenic and mutually linked to Wnt activity in blastula-stage sea urchin embryos. *PLoS One*. 2008; 3:e3770. [PubMed: 19020668]

43. Melnikova IN, Crute BE, Wang S, Speck NA. Sequence specificity of the core-binding factor. *J Virol.* 1993; 67:2408–2411. [PubMed: 8445737]
44. Shtutman M, Zhurinsky J, Simcha I, Albanese C, D'Amico M, Pestell R, Ben-Ze'ev A. The cyclin D1 gene is a target of the beta-catenin/LEF-1 pathway. *Proc Natl Acad Sci U S A.* 1999; 96:5522–5527. [PubMed: 10318916]
45. Rimerman RA, Gellert-Randleman A, Diehl JA. Wnt1 and MEK1 cooperate to promote cyclin D1 accumulation and cellular transformation. *J Biol Chem.* 2000; 275:14736–14742. [PubMed: 10748202]
46. Albanese C, Wu K, D'Amico M, Jarrett C, Joyce D, Hughes J, Hulit J, Sakamaki T, Fu M, Ben-Ze'ev A, Bromberg JF, Lamberti C, Verma U, Gaynor RB, Byers SW, Pestell RG. IKKalpha regulates mitogenic signaling through transcriptional induction of cyclin D1 via Tcf. *Mol Biol Cell.* 2003; 14:585–599. [PubMed: 12589056]
47. Tetsu O, McCormick F. Beta-catenin regulates expression of cyclin D1 in colon carcinoma cells. *Nature.* 1999; 398:422–426. [PubMed: 10201372]
48. Cluster Buster. Available at: <http://zlab.bu.edu/cluster-buster/>
49. Frith MC, Li MC, Weng Z. Cluster-Buster: Finding dense clusters of motifs in DNA sequences. *Nucleic Acids Res.* 2003; 31:3666–3668. [PubMed: 12824389]
50. Berman BP, Nibu Y, Pfeiffer BD, Tomancak P, Celniker SE, Levine M, Rubin GM, Eisen MB. Exploiting transcription factor binding site clustering to identify cis-regulatory modules involved in pattern formation in the *Drosophila* genome. *Proc Natl Acad Sci U S A.* 2002; 99:757–762. [PubMed: 11805330]
51. Zeller RW, Coffman JA, Harrington MG, Britten RJ, Davidson EH. SpGCF1, a sea urchin embryo DNA-binding protein, exists as five nested variants encoded by a single mRNA. *Dev Biol.* 1995; 169:713–727. [PubMed: 7781910]
52. Zeller RW, Griffith JD, Moore JG, Kirchhamer CV, Britten RJ, Davidson EH. A multimerizing transcription factor of sea urchin embryos capable of looping DNA. *Proc Natl Acad Sci U S A.* 1995; 92:2989–2993. [PubMed: 7708761]
53. Lewis CD, Clark SP, Felsenfeld G, Gould H. An erythrocyte-specific protein that binds to the poly(dG) region of the chicken beta-globin gene promoter. *Genes Dev.* 1988; 2:863–873. [PubMed: 3209071]
54. Xiang M, Lu SY, Musso M, Karsenty G, Klein WH. A G-string positive cis-regulatory element in the LpS1 promoter binds two distinct nuclear factors distributed non-uniformly in *Lytechinus pictus* embryos. *Development.* 1991; 113:1345–1355. [PubMed: 1811948]
55. Hapgood J, Patterson D. Purification of an oligo(dG).oligo(dC)-binding sea urchin nuclear protein, suGF1: a family of G-string factors involved in gene regulation during development. *Mol Cell Biol.* 1994; 14:1402–1409. [PubMed: 8289815]

- We undertook a *cis*-regulatory analysis of the *cycD* gene in the purple sea urchin.
- Both upstream and intronic regulatory sequences were uncovered.
- Together, these sequences account for expression during embryogenesis.
- Different regulatory regions mediate distinct temporal aspects of *cycD* expression.
- *cis*-regulatory modules were computationally or experimentally identified.

Fig. 1. Expression of endogenous *SpycD* and microinjected mcherry-linked BAC bearing *SpycD* plus 90 kb and 13 kb of up and downstream sequence
Relative levels of *SpycD* mRNA were measured at each indicated time point by qRT-PCR as described in the text.

Fig. 2. Identifying *cis*-regulatory sequences

A. Regions tested for CRM-containing activity. *SpycycD*, plus up 13 kb and 7 kb of upstream and downstream sequence is shown. Exons: black; potential CRM-containing regions: blue; sequences with 90% similarity to *LvcycD*: red; active regions: boxed. **B. Representative activity profiles.** Each panel is from the indicated experiment. Asterisks denote significant activity. See Supplemental Fig. 2 for additional activity profiles.

Fig. 3. Averaged temporal activity profiles

Grand means and standard deviations were calculated from the means of all experiments in Fig. 2B and Supplemental Fig 3. Small differences between time points in different experiments (for example, 45 and 47 hpf) were ignored.

Fig. 4. Identification of *cis*-regulatory modules

A. *SpycD* showing active *cis*-regulatory regions. Exons: black rectangles; active regions: blue rectangles; active and inactive subregions: blue and tan lines, respectively; conserved sequences: red; Cluster-Buster-identified sequences: gray. **B. Activities of 13-tag-linked regions 2, 2-2, 6 and 6-1.** Panel 1 shows the activities of region 2 and subregion 2-2 in co-injected embryos (one experiment). Panel 2 shows the averaged temporal activities and standard deviations of region 6 and subregion 6-1 from all presented experiments where either region was assayed. **C. Fluorescence micrographs from injection with EpGFPII-linked region 2, 2-2, 4, 4-1 or 4-2.** Brightness and contrast were adjusted equally in all images.