Gabapentin for Pain Control After Surgical Debridement of Burn Wounds

Abdul Ahad Khan¹, Muhammad Shahzad Shamim^{2*}

¹Medical College, Aga Khan University, Karachi, Pakistan

²Department of Surgery, Aga Khan University, Karachi, Pakistan

ARTICLE INFO

Article type:
Letter to Editor

Article history:
Received: 23 May 2012
Revised: 27 May 2012
Accepted: 31 May 2012

Keywords:
Gabapentin
Pain
Burns

► Please cite this paper as:

Khan AA, Shamim MS. Gabapentin for Pain Control After Surgical Debridement of Burn Wounds. *Arch Trauma Res.* 2012; **1**(2):85-6. DOI: 10.5812/atr.6489

Published by Kowsar, 2012. cc 3.0.

Dear Editor,

I read with great interest the contribution “Effect of Gabapentin on Morphine Consumption and Pain after Surgical Debridement of Burn Wounds: A Double-Blind Randomized Clinical Trial Study” by Raziet *et al.* (1). Carrying out a Randomized Controlled Trial without funding and especially in our region is an uphill task indeed. I must therefore congratulate the authors on this achievement. The trial has been well-conducted and the manuscript has been likewise prepared impressively. We would however, like to point towards a few observations.

Firstly, the authors have not mentioned how they randomized their samples. Moreover, the process of blinding was not explained in detail other than mentioning the envelope method. In this regard, it would be advised that for *randomized controlled trials* (RCT), the design and description should adhere to the Consolidated Standards of Reporting Clinical Trials’ (CONSORT) statement. Lastly, we would also like to draw the authors’ attention to a similar

study addressing the efficacy of Gabapentin in post-surgical patients conducted by Dierking *et al.* (2) published in 2004. Although the patient population was different, it also showed numerically significant reduced morphine usage in the Gabapentin group compared to placebo. Finally, we would once again congratulate the authors on the conduction and completion of this very important trial.

Authors’ Contribution

All the authors contributed in the writing and research work for this article.

Financial Disclosure

The authors declare that they have no financial interests relating to the material in the manuscript.

References

1. Rimaz S, Emir Alavi C, Sedighinejad A, Tolouie M, Kavosi S, Koochakinejad L. Effect of Gabapentin on Morphine Consumption and

* Corresponding author: Muhammad Shahzad Shamim, Department of Surgery, Aga Khan University, Karachi, Pakistan. Tel: +92-2134930051, Fax: +92-2134934294, E-mail: shahzad.shamim@aku.edu

© 2012, Kashan University of Medical Sciences. Published by Kowsar

This is an open-access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

-
- Pain after Surgical Debridement of Burn Wounds: A Double-Blind Randomized Clinical Trial Study. *Arch Trauma Res.* 2012;**1**(1):38-43.
2. Dierking G, Duedahl TH, Rasmussen ML, Fomsgaard JS, Moiniche S, Romsing J, et al. Effects of gabapentin on postoperative morphine consumption and pain after abdominal hysterectomy: a randomized, double-blind trial. *Acta Anaesthesiol Scand.* 2004;**48**(3):322-7.