Published in final edited form as:

Ann Surg. 2013 August ; 258(2): . doi:10.1097/SLA.0b013e318298b077.

Cerebrospinal Fluid A β to Tau Ratio and Postoperative Cognitive Change

Zhongcong Xie, MD, PhD^{*}, Sayre McAuliffe, BA^{*}, Celeste A. Swain, BA^{*}, Sarah A. P. Ward, BA^{*}, Catherine A. Crosby, MS^{*}, Hui Zheng, PhD[†], Janet Sherman, PhD[‡], Yuanlin Dong, MD^{*}, Yiyang Zhang, MD^{*}, Neelakantan Sunder, MB[§], Dennis Burke, MD[¶], Kevin J. Washicosky, BA^{||}, Rudolph E. Tanzi, PhD^{||}, and Edward R. Marcantonio, MD^{**}

^{*}Geriatric Anesthesia Research Unit, Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital and Harvard Medical School, Charlestown, MA

[†]MGH Biostatistics Center, Massachusetts General Hospital and Harvard Medical School, Boston, MA

[‡]Psychology Assessment Center, Massachusetts General Hospital and Harvard Medical School, Boston, MA

[§]Department of Anesthesia, Critical Care and Pain Medicine, Massachusetts General Hospital and Harvard Medical School, Boston, MA

[¶]Department of Surgery, Massachusetts General Hospital and Harvard Medical School, Boston, MA

^{||}Genetics and Aging Research Unit, MassGeneral Institute for Neurodegenerative Disease, Department of Neurology, Massachusetts General Hospital and Harvard Medical School, Charlestown, MA

^{**}Divisions of General Medicine and Primary Care and Gerontology, Department of Medicine, Beth Israel Deaconess Medical Center and Harvard Medical School, Boston, MA

Abstract

Objective—Determination of biomarker and neuropathogenesis of postoperative cognitive change (POCC) or postoperative cognitive dysfunction.

Background—POCC is one of the most common postoperative complications in elderly patients. Whether preoperative cerebrospinal fluid (CSF) β -amyloid protein (A β) to tau ratio, an Alzheimer disease biomarker, is a biomarker for risk of POCC remains unknown. We therefore set out to assess the association between preoperative CSF A β 42 or A β 40 to tau ratio and POCC.

Methods—Patients who had total hip/knee replacement were enrolled. The CSF was obtained during the administration of spinal anesthesia. Cognitive tests were performed with these participants at 1 week before and at 1 week and 3 to 6 months after the surgery. Z scores of the changes from preoperative to postoperative on several key domains of the cognitive battery were determined. We then examined the association between preoperative CSF A β 42/tau or A β 40/tau ratio and the outcome measures described earlier, adjusting for age and sex.

Results—Among the 136 participants (mean age = 71 ± 5 years; 55% men), preoperative CSF A β 42/tau ratio was associated with postoperative Hopkins Verbal Learning Test Retention [Z score 8.351; age, sex-adjusted (adj.) $P = 0.003$], and the Benton Judgment of Line= Orientation (Z score 1.242; adj. $p = 0.007$). A β 40/tau ratio was associated with Brief Visuospatial Memory Test Total Recall (Z score = 1.045; adj. $P = 0.044$).

Conclusions—Preoperative CSF A β /tau ratio is associated with postoperative changes in specific cognitive domains. The presence of the Alzheimer's disease biomarker, specifically the A β /tau ratio, may identify patients at higher risk for cognitive changes after surgery.

Keywords

A β /tau ratio; Biomarker; cognition; CSF; surgery

Postoperative cognitive change (POCC) or postoperative cognitive dysfunction (POCD)^{1,2} (reviewed in Silverstein et al³ and Terrando et al⁴) is one of the most common postoperative complications in older adults who have undergone surgery⁵ and is associated with impairments in daily functioning,⁶ risk of leaving the labor market prematurely, dependency on social transfer payments,⁷ and increased morbidity and mortality.⁷⁻⁹ However, at the present time, POCC is a clinical phenomenon and its neuropathogenesis remains largely to be determined; moreover, there have been no biomarker(s) that could identify the risk for the development of POCC. This gap in knowledge has become a barrier that prevents further studies and potential intervention of POCC.

β -Amyloid protein (A β), including A β 40 and A β 2, the key component of senile plaques in patients with Alzheimer disease (AD), and tau, the major protein component of intraneuronal neurofibrillary tangles, are the hallmarks of AD neuropathogenesis (reviewed in Querfurth and LaFerla¹⁰). Reduction of A β 40 and A β 42 levels and elevation of tau levels in cerebrospinal fluid (CSF) have been reported in AD patients^{11,12} (reviewed in Holtzman¹³). Moreover, it has been reported that the A β 42/tau ratio in CSF can better distinguish AD patients from healthy controls of the same age than A β 42 or tau alone.^{12,14} However, whether the A β 42/tau or A β 40/tau ratio in CSF is associated with POCC is unknown.

Therefore, we performed a prospective investigation to determine whether preoperative CSF A β 42/tau ratio or A β 40/tau ratio is associated with individual components, including verbal learning and visual memory, of a cognitive battery designed to evaluate POCC. We hypothesized that CSF A β 42/tau or A β 40/tau ratio will be associated with verbal learning and visual memory. The outcomes from this study may serve 2 purposes: (1) to evaluate preoperative CSF A β /tau ratio as a potential biomarker of POCC and (2) to promote more studies to determine the role of A β and tau in the neuropathogenesis of POCC.

We used the term “POCC” instead of “POCD” in the article because the studies did not assess the potential association between CSF A β 40/tau or A β 42/tau ratio and cognitive function in participants without surgery. Therefore, we were unable to determine the true incidence of POCD because the change scores could be confounded by the learning effects of the participants.

METHODS

Participants

The protocol was approved by the institutional review board of Partners Human Research Committee, Boston, MA. A total of 342 adults scheduled to have elective total hip or knee arthroplasty/replacement surgery at the Massachusetts General Hospital were asked to

participate in this study (Fig. 1). Participants were deemed eligible if they were at least 63 years old, spoke proficient English, lived within the greater Boston area, and were candidates for spinal anesthesia. Individuals who met these criteria were further screened during an initial interview that included administration of the Mini-Mental State Examination. Individuals excluded from participation were those identified as having the following: (1) existing cognitive impairment as evidenced by Mini-Mental State Examination scores below 24 (of the 30 possible points); (2) a history of neurological and psychiatric diseases including AD, stroke, and psychosis; (3) severe visual or hearing impairment; and (4) unwillingness to comply with the protocol or procedures. Consent was obtained by a study coordinator in the Pre-Admissions Clinic at Massachusetts General Hospital. A total of 136 participants were finally enrolled in the study from 2007 to 2012, and there have been no major changes in the surgery or anesthesia practice since the start of our studies in 2007. We initially targeted a sample size of 100, which would give us 80% power to detect a Pearson correlation coefficient of 0.28 or higher between the measured marker (eg, A β 42/tau ratio) and cognitive change scores with 5% type 1 error.

Baseline Assessment

Each participant who was enrolled underwent a baseline assessment, which consisted of neuropsychological testing performed 1 week before the scheduled surgery in either a hospital office setting or the participant's home. The cognitive tests included the Hopkins Verbal Learning Test Retention (HVLTRet), Hopkins Verbal Learning Total Recall (HVLTR), the Brief Visuospatial Memory Test Total Recall (BVMTR), Brief Visuospatial Memory Test Delayed Recall (BVMTRD), Brief Visuospatial Memory Test Total Learning, Brief Visuospatial Memory Test Retention, the Benton Judgment of Line Orientation (JLO), and Trail Making B (Trails B). These measures are highly sensitive to the types of cognitive impairments¹⁵⁻¹⁷ and are widely used in the field of neuropsychology.^{18,19} All neuropsychological tests were administered by the study coordinator (S.M., C.A.S., S.A.P.W., C.G., and K.W.).

Anesthesia, CSF Sample Collection, and Measurement of A β and Tau

All participants had spinal anesthesia for the scheduled surgery. One milliliter of CSF was collected by anesthesiologists at the time of spinal anesthesia administration between 7:00 AM and 10:00 AM to avoid confounding influence that resulted from the circadian changes in CSF A β levels.²⁰ A β levels (including A β 40 and A β 42) in the CSF were measured with enzyme-linked immunosorbent assay (ELISA) kits (A β 40: cat no. 292-6230; A β 42: cat no. 296-64401) (Wako, Richmond, VA) as described by Zhang et al²¹ with modification. Briefly, the collected CSF was stored in an -80-degree freezer and was thawed before measurement. Human monoclonal antibodies specific to A β 40 (BA27) or A β 42 (BC05) were coated on 96-well plates. After blocking with Block Ace, wells were incubated overnight at 4°C with test samples of CSF and then an anti-A β (α -A β -HR1) conjugated to horseradish peroxidase was added. The plates were developed with 3,3',5,5'-tetramethylbenzidine reagent, and the well absorbance was measured at 450 nm. The A β levels in the test samples were determined by comparison with the signals from unconditioned artificial CSF spiked with known quantities of A β 40 and A β 42. The levels of total tau in human CSF were determined with an ELISA kit (cat. no. KHB0041; Invitrogen, San Francisco, CA). Specifically, a monoclonal antibody specific for human tau was coated onto the wells of the provided microtiter strips. Standards of known human tau content, CSF, and control specimens were pipetted into these wells overnight. After washing, a rabbit polyclonal antibody specific for human tau was added. During the second incubation, this antibody bound to the immobilized human tau captured during the first incubation. A substrate solution was added, which was acted upon by the bound enzyme to produce a change in color. The intensity of this colored product was directly proportional to the

concentration of human tau present in the CSF. All assays were performed in triplicates, and the average from the triplicates was obtained.

Surgery

All participants had total hip or total knee replacement under spinal anesthesia by one surgeon to avoid potential confounding factors owing to varying surgery skills or different surgical practices. The participants had standardized perioperative care, including preoperative and intraoperative sedation and postoperative pain control. There were no major complications among the participants during the immediate postoperative period.

Postoperative Cognitive Testing

Each participant underwent repeat cognitive testing approximately 1 week postoperatively and then 3 to 6 months postoperatively at the participant's home or a rehabilitation facility. The same neuropsychological test battery that was administered preoperatively (see earlier) was administered by trained study coordinators. Four versions of the test battery (A, B, C, and D) were administered according to a randomized datasheet to reduce participant practice or learning effects.

Statistical Analysis

Postoperative cognitive function was represented by a change score, calculated from the raw score of the postoperative cognitive test result minus the raw score of the preoperative cognitive test result. A Z score for each of the change scores was calculated by dividing each individual change score by the total standard deviation of all of the change scores. Negative values of Z scores suggest postoperative cognitive decline except for the test of Trails B, which is a timed test where short times are indicative of better performance. Therefore, positive Z scores of Trails B suggest postoperative cognitive decline. We then applied Pearson correlation analysis to these scores and calculated Pearson correlation coefficients to analyze the relationship between preoperative CSF A β 42/tau or A β 40/tau ratio and Z score for each of the cognitive tests. We used linear regression to adjust for age and sex. *P* values less than 0.05 were considered statistically significant. The SAS (SAS Institute Inc, Cary, NC) software (version 9.2) was used for all statistical analyses.

RESULTS

Participant and Surgery Characteristics

Three hundred forty-two eligible participants were screened; among them, 215 participants provided informed consent for the study. Seventy-nine participants were subsequently excluded from the study owing to various reasons (see Fig. 1), yielding 136 participants. The demographic and clinical data of the participants are presented in Table 1. The average values of A β 40, A β 42, and tau from the participants were 9505 ± 3441 , 948 ± 641 , and 244 ± 153 pg/mL, the average A β 40/tau and A β 42/tau ratios from the participants were 50 ± 25.3 and 5 ± 4.0 , respectively. Table 2 shows the mean cognitive test scores of all participants at baseline, at 1 week, and at 3 to 6 months after the surgery.

Pearson correlation analysis showed that age was negatively associated with preoperative human CSF A β 42/tau ratio ($R = -0.199$; $P = 0.021$). The JLO was negatively associated with preoperative human CSF A β 40 ($R = -0.198$; $P = 0.021$) and A β 42 ($R = -0.183$; $P = 0.034$). Finally, Trails B was negatively associated with preoperative human CSF A β 42/tau ratio ($R = -0.309$; $P = 0.0003$). The adjusted analyses controlled for age and also for baseline cognitive performance by examining change scores in the various neurocognitive measures.

Preoperative CSF A β 42/Tau Ratio and POCC

We assessed the potential correlation between preoperative CSF A β 42/tau ratio and Z score representing changes in postoperative cognitive function. Unadjusted Pearson correlation analyses showed that the preoperative CSF A β 42/tau ratio was positively correlated with Z score of postoperative function of HVLTRet at 1 week (7.063, $P = 0.011$) and JLO at 1 week (0.915, $P = 0.024$) and 3 to 6 months (1.139, $P = 0.011$) and negatively correlated with Z score of postoperative Trails B at 1 week (-5.623 , $P = 0.019$). Linear regression analysis, after adjusting for age and sex, showed that the preoperative CSF A β 42/tau ratio was positively associated with postoperative function of HVLTRet at 1 week (8.351, $P = 0.003$), HVLTTTR at 3 to 6 months (0.833, $P = 0.046$), and JLO at 1 week (0.954, $P = 0.021$) and at 3 to 6 months (1.242, $P = 0.007$) (Table 3). After adjustment, the preoperative CSF A β 42/tau ratio was no longer negatively associated with Trails B at 1 week and was not significantly associated with any other cognitive tests (Table 3). These data suggest that the participants who had a lower preoperative CSF A β 42/tau ratio (the AD biomarker) performed worse postoperatively on HVLTRet, HVLTTTR, and JLO, measures of verbal memory and visuospatial judgment, than those with a higher preoperative CSF A β 42/tau ratio.

Preoperative CSF A β 40/Tau Ratio and POCC

Next, we assessed the potential correlation between preoperative CSF A β 40/tau ratio and Z score representing changes in postoperative cognitive function. Using unadjusted Pearson correlation analyses, we found that the preoperative CSF A β 40/tau ratio was positively correlated with Z score of postoperative function of BVMTDR (0.413, $P = 0.045$) at 3 to 6 months (Table 4). After adjusting for age and sex using linear regression, the preoperative CSF A β 40/tau ratio was still significantly correlated with Z score of postoperative function of both BVMTDR (0.418, $P = 0.046$) and BVMTTR (1.045, $P = 0.044$) at 3 to 6 months (Table 4). The preoperative CSF A β 40/tau ratio was not significantly associated with changes in other postoperative cognitive functions (Table 4). These data suggest that the participants who had a lower preoperative CSF A β 40/tau ratio had the worse performance on postoperative function of BVMTTR and BVMTDR, measures of visual memory, than those with a higher preoperative CSF A β 40/tau ratio.

DISCUSSION

We studied the association of the AD biomarker with changes in cognition after elective surgery and found that the CSF A β 42/tau ratio is associated with HVLTRet at 1 week and HVLTTTR at 3 to 6 months (Table 3). These results are consistent with the findings from the AD research that the CSF A β 42/tau ratio can be used for the diagnosis of AD and prediction of its progress and that the lower the CSF A β 42/tau ratio, the worse the cognitive function in AD patients¹² (reviewed in Holtzman¹³).

The CSF A β 40/tau ratio is not associated with any verbal functions tested in the current studies (Table 4). Instead, the CSF A β 40/tau ratio is associated with BVMTTR and BVMTDR at 3 to 6 months (after adjustment for age and sex; Table 3). These data suggest that the CSF A β 40/tau ratio could specifically represent the visual memory domain of POCC. Different from the CSF A β 42/tau ratio, the CSF A β 40/tau ratio has not been extensively reported as a biomarker of AD (reviewed in Holtzman¹³). Thus, the future studies are warranted to test a hypothesis that the CSF A β 40/tau ratio has not been recognized as a biomarker of AD because visual memory tests are not usually included in the battery of cognitive tests commonly used for diagnosing AD.^{17,18}

Interestingly, the CSF A β 40/tau and A β 42/tau ratios were associated with a different domain of POCC in our current studies. A β 40 has an identical amino acid sequence with A β 42

except for the lack of amino acids isoleucine and alanine at the C terminus. However, A β 40 and A β 42 have distinct clinical, biological, and biophysical characteristics.²² Specifically, 90% of A β is A β 40 but A β 42 is the major form of A β in senile plaques.^{23–25} A β 42 has a quicker rate of aggregation than A β 40,^{26,27} and it does so through different underlying mechanisms.^{22,28} Finally, A β 42 is more toxic to neurons than A β 40.^{29,30} Collectively, it is conceivable that A β 40 and A β 42 may have different effects on brain function, which suggest that it is also conceivable that the CSF A β 40/tau and A β 42/tau ratios may predispose patients to different domains of POCC. There are conflicting findings about the association of CSF biomarkers (eg, A β) with cognitive dysfunction.^{31–34} The findings from the current studies that the CSF A β /tau ratio was associated only with specific components of cognitive change could explain the conflicting findings, which may result from the varying cognitive tests used in previous studies.

Both the Hopkins Verbal Learning Test and the Brief Visuospatial Memory Test measure anterograde memory function and thereby the functional integrity of the medial temporal lobe system.¹⁹ Many studies have suggested that the low CSF A β level represents high brain A β levels, owing to the sequestration of A β into brain amyloid plaques^{35,36} (reviewed in Holtzman¹³ and Blennow and Zetterberg³⁷), but CSF tau levels represent brain tau levels^{35,36} (reviewed in Blennow and Zetterberg³⁷). Taken together, the findings that the CSF A β /tau ratio is associated with Hopkins Verbal Learning Test and the Brief Visuospatial Memory Test suggest that CSF and/or brain A β and tau levels may be associated with the integrity of the medial temporal lobe system. Further studies are needed to test this hypothesis by determining whether the amount of amyloid in the medial temporal lobe system is specifically associated with verbal learning and visual memory impairment (eg, by using positron emission tomographic imaging with Amyvid).

The JLO can be used to assess spatial perception and orientation, and it has been shown to be sensitive to the right hemisphere dysfunction.^{38,39} We have found that the CSF A β 42/tau ratio is associated with the postoperative cognitive function of the JLO. These findings suggest that CSF A β 42/tau ratio, the AD biomarker, is associated with increased vulnerability of certain parts of the brain and specific domains of cognitive function to the stress of surgery. It is notable that spatial perception and orientation are localized to areas of the brain that are different from those of visual memory and are reflected by differing biomarkers.

The values of human CSF A β and tau in our current studies were higher than those determined in the Alzheimer's Disease Neuroimaging Initiative studies¹² and in a recent study to assess the effects of anesthesia and surgery on human AD and inflammation biomarkers.⁴⁰ The reason for such a difference is not clear at the present time. However, there have been reports showing variable human CSF A β and tau values in different studies (reviewed in Holtzman¹³). Specifically, the values of A β and tau in our current studies were comparable with those measured in the studies by Fagan et al.⁴¹ Moreover, all CSF samples in the current study were analyzed using the same methods. Therefore, although the absolute values of CSF A β and tau may differ from those reported in other studies, the relative differences in the values between the subjects are consistent and therefore our results should be valid.

Prior studies by Tang et al.⁴⁰ and Palotas et al.⁴² have shown that anesthesia and surgery seem to modulate CSF levels of A β , tau, cytokines, and S100b; however, these studies did not attempt to determine the associations between the preoperative CSF biomarkers and postoperative cognitive function. Our current studies are different from these studies because we have focused on determining the potential association between preoperative CSF A β /tau ratio and POCC rather than the change in human CSF biomarkers after anesthesia

and surgery. Further studies are warranted in the future to determine the effects of surgery on both pre- and postoperative CSF biomarkers in the same participant.

We found that there was a significant relationship existed between POCC and A β /tau ratios in some cognitive domains. Interestingly, because POCC was generally in the positive direction, these data suggest that even within a group of cognitively normal elderly patients, A β /tau ratios may allow stratification of learning ability.

This study has several limitations. First, the CSF A β 42/tau ratio is not associated with all of the verbal functions tested. The ratio is associated only with HVLTRet and HVLTTTR but not with BVMTDR (data not shown). The CSF A β 40/tau ratio is also not associated with all of the visual functions tested. The ratio is associated only with BVMTTR and BVMTDR but not with Brief Visuospatial Memory Test Total Learning or Brief Visuospatial Memory Test Retention (data not shown). It is possible that certain aspects of memory functioning, for example, the ability to retrieve information that has effectively been stored in memory, may be more susceptible to postoperative decline in individuals with a relatively lower CSF A β /tau ratio in the current studies. Future studies should include memory measures in which semantic cues are provided to aid in the retrieval process, which may help further clarify the role of A β and tau in different aspects of cognitive dysfunction. Second, it is unknown whether the observed association between CSF A β 40/tau or A β 42/tau ratio and verbal function, visual function, spatial perception, and orientation is exclusively for postoperative cognitive function or for general cognitive function. Third, we could not draw the conclusion that the preoperative CSF A β /tau ratio is associated with POCD or postoperative cognitive improvement because the studies did not include a nonsurgical control group to quantify the learning effect. However, we were able to demonstrate that a correlation existed between preoperative CSF A β /tau ratio and change in performance in some cognitive domains. Interestingly, our data suggest that those with higher ratios had improved learning than those with lower ratios. Whether this observation translates to an ability of CSF A β /tau ratios to predict risk of postoperative cognitive decline, as suggested in a recent study of patients with normal pressure hydrocephalus,⁴³ is not yet clear. Fourth, we did not correct the *P* values for the multiple comparisons in Tables 3 and 4 because we expected that the cognitive domains tested were highly cross-correlated; therefore, the need to treat each comparison as “independent” is reduced. However, the possibility that there is no correlation among POCC domains cannot be excluded from the current studies. Finally, the average education in the participants was 17 years, which is skewed to the higher level. It is unknown whether such correlation between CSF A β /tau ratio and POCC would still exist in participants with lower education levels.

We did not use the global Z score in this study. Combining neuropsychology tests into a single test is controversial in the field of neuropsychology, and most of the POCD literature has not used this approach.⁴⁴ Creating meaningful global scores requires a sophisticated modern measurement approach that is beyond the scope of this analysis. Moreover, these data suggest that heterogeneity exists, meaning that CSF or brain levels of A β 40, A β 42, and tau may contribute only to certain, but not universal, components of POCD. Pending further studies, these results suggest that the CSF A β 42/tau and CSF A β 40/tau ratios could specifically be predictive of the verbal and visual learning domains of POCC.

CONCLUSIONS

We have found associations between preoperative CSF A β 42/tau or A β 40/tau ratio and postoperative changes in specific cognitive domains. If confirmed and extended into future studies, these associations may shed light on the currently undefined neuropathogenesis of POCC, which will, hopefully, promote more animal and human research to further

determine the neuropathogenesis and intervention of POCC, ultimately leading to safer surgery care and better postoperative outcomes for senior adults.

Acknowledgments

Author contribution is as follows: Xie, Marcantonio, Sherman, Tanzi, Sunder, and Burke: study concept and design; McAuliffe, Swain, Ward, Crosby, Washicosky, Dong, and Zhang: acquisition of data; Marcantonio, Zheng, Sherman, Tanzi, and Xie: analysis and interpretation of data; Marcantonio and Xie: drafting of the manuscript; Marcantonio, Sherman, and Xie: critical revision of the manuscript for important intellectual content; Xie: obtained funding; Sherman, Zhang, Sunder, and Burke: administrative, technical, and material support; and Xie: study supervision. Zhongcong Xie had full access to all data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

This study was supported by National Institutes of Health grants (Bethesda, MD) (K08 NS048140, R21 AG029856, R01 GM088801, R21 AG038994, and R01 AG041274); Investigator-Initiated Research Grant from Alzheimer's Association (Chicago, IL); and Cure Alzheimer's Fund (to Z.X.). Dr Marcantonio was funded in part by grants R01AG030618, P01AG031720, and a Mid-Career Investigator Award (K24AG035075) from the National Institute on Aging.

REFERENCES

1. Moller JT, Cluitmans P, Rasmussen LS, et al. Long-term postoperative cognitive dysfunction in the elderly ISPOCD1 study. ISPOCD Investigators. International Study of Post-Operative Cognitive Dysfunction. *Lancet*. 1998; 351:857–861. [PubMed: 9525362]
2. Newman MF, Kirchner JL, Phillips-Bute B, et al. Longitudinal assessment of neurocognitive function after coronary-artery bypass surgery. *N Engl J Med*. 2001; 344:395–402. [PubMed: 11172175]
3. Silverstein JH, Steinmetz J, Reichenberg A, et al. Postoperative cognitive dysfunction in patients with preoperative cognitive impairment: which domains are most vulnerable? *Anesthesiology*. 2007; 106:431–435. [PubMed: 17325500]
4. Terrando N, Brzezinski M, Degos V, et al. Perioperative cognitive decline in the aging population. *Mayo Clin Proc*. 2011; 86:885–893. [PubMed: 21878601]
5. Liu LL, Leung JM. Predicting adverse postoperative outcomes in patients aged 80 years or older. *J Am Geriatr Soc*. 2000; 48:405–412. [PubMed: 10798467]
6. Phillips-Bute B, Mathew JP, Blumenthal JA, et al. Association of neurocognitive function and quality of life 1 year after coronary artery bypass graft (CABG) surgery. *Psychosom Med*. 2006; 68:369–375. [PubMed: 16738066]
7. Steinmetz J, Christensen KB, Lund T, et al. Long-term consequences of postoperative cognitive dysfunction. *Anesthesiology*. 2009; 110:548–555. [PubMed: 19225398]
8. Monk TG, Weldon BC, Garvan CW, et al. Predictors of cognitive dysfunction after major noncardiac surgery. *Anesthesiology*. 2008; 108:18–30. [PubMed: 18156878]
9. Deiner S, Silverstein JH. Postoperative delirium and cognitive dysfunction. *Br J Anaesth*. 2009; 103(suppl 1):i41–i46. [PubMed: 20007989]
10. Querfurth HW, LaFerla FM. Alzheimer's disease. *N Engl J Med*. 2010; 362:329–344. [PubMed: 20107219]
11. Sunderland T, Linker G, Mirza N, et al. Decreased beta-amyloid1-42 and increased tau levels in cerebrospinal fluid of patients with Alzheimer disease. *JAMA*. 2003; 289:2094–2103. [PubMed: 12709467]
12. Shaw LM, Vanderstichele H, Knapik-Czajka M, et al. Cerebrospinal fluid biomarker signature in Alzheimer's disease neuroimaging initiative subjects. *Ann Neurol*. 2009; 65:403–413. [PubMed: 19296504]
13. Holtzman DM. CSF biomarkers for Alzheimer's disease: current utility and potential future use. *Neurobiol Aging*. 2011; 32(suppl 1):S4–S9. [PubMed: 22078172]
14. Welge V, Fiege O, Lewczuk P, et al. Combined CSF tau, p-tau181 and amyloid-beta 38/40/42 for diagnosing Alzheimer's disease. *J Neural Transm*. 2009; 116:203–212. [PubMed: 19142572]

15. Albert MS, Moss MB, Tanzi R, et al. Preclinical prediction of AD using neuropsychological tests. *J Int Neuropsychol Soc.* 2001; 7:631–639. [PubMed: 11459114]
16. Mickes L, Wixted JT, Fennema-Notestine C, et al. Progressive impairment on neuropsychological tasks in a longitudinal study of preclinical Alzheimer's disease. *Neuropsychology.* 2007; 21:696–705. [PubMed: 17983283]
17. Salmon DP, Thomas RG, Pay MM, et al. Alzheimer's disease can be accurately diagnosed in very mildly impaired individuals. *Neurology.* 2002; 59:1022–1028. [PubMed: 12370456]
18. Lezak, MD.; Howieson, DB.; Loring, DW. *Neuropsychological Assessment.* 4th ed.. Oxford University Press; New York: 2004.
19. Sherman, JC.; Leveroni, C.; Pollak, LN. Neuropsychological assessment. In: Stern, TA.; Rosenbaum, JF.; Fava, M.; Biederman, J.; Rauch, SL., editors. *Massachusetts General Hospital Comprehensive Clinical Psychiatry.* Mosby Elsevier; Philadelphia, PA: 2008. p. 103-119.
20. Bateman RJ, Wen G, Morris JC, Holtzman DM. Fluctuations of CSF amyloid-beta levels: implications for a diagnostic and therapeutic biomarker. *Neurology.* 2007; 68:666–669. [PubMed: 17325273]
21. Zhang B, Dong Y, Zhang G, et al. The inhalation anesthetic desflurane induces caspase activation and increases amyloid beta-protein levels under hypoxic conditions. *J Biol Chem.* 2008; 283:11866–11875. [PubMed: 18326038]
22. Bitan G, Kirkitadze MD, Lomakin A, et al. Amyloid beta-protein (Abeta) assembly: Abeta 40 and Abeta 42 oligomerize through distinct pathways. *Proc Natl Acad Sci U S A.* 2003; 100:330–335. [PubMed: 12506200]
23. Suzuki N, Cheung TT, Cai XD, et al. An increased percentage of long amyloid beta protein secreted by familial amyloid beta protein precursor (beta APP717) mutants. *Science.* 1994; 264:1336–1340. [PubMed: 8191290]
24. Iwatsubo T, Odaka A, Suzuki N, et al. Visualization of A beta 42(43) and A beta 40 in senile plaques with end-specific A beta monoclonals: evidence that an initially deposited species is A beta 42(43). *Neuron.* 1994; 13:45–53. [PubMed: 8043280]
25. Gravina SA, Ho L, Eckman CB, et al. Amyloid beta protein (A beta) in Alzheimer's disease brain. Biochemical and immunocytochemical analysis with antibodies specific for forms ending at A beta 40 or A beta 42(43). *J Biol Chem.* 1995; 270:7013–7016. [PubMed: 7706234]
26. Jarrett JT, Berger EP, Lansbury PT Jr. The carboxy terminus of the beta amyloid protein is critical for the seeding of amyloid formation: implications for the pathogenesis of Alzheimer's disease. *Biochemistry.* 1993; 32:4693–4697. [PubMed: 8490014]
27. Jarrett JT, Berger EP, Lansbury PT Jr. The C-terminus of the beta protein is critical in amyloidogenesis. *Ann N Y Acad Sci.* 1993; 695:144–148. [PubMed: 8239273]
28. Bitan G, Vollers SS, Teplow DB. Elucidation of primary structure elements controlling early amyloid beta-protein oligomerization. *J Biol Chem.* 2003; 278:34882–34889. [PubMed: 12840029]
29. Zhang Y, McLaughlin R, Goodyer C, et al. Selective cytotoxicity of intracellular amyloid beta peptide1-42 through p53 and Bax in cultured primary human neurons. *J Cell Biol.* 2002; 156:519–529. [PubMed: 11815632]
30. Mucke L, Masliah E, Yu GQ, et al. High-level neuronal expression of Abeta 1-42 in wild-type human amyloid protein precursor transgenic mice: synaptotoxicity without plaque formation. *J Neurosci.* 2000; 20:4050–4058. [PubMed: 10818140]
31. Mehta PD, Pirttila T, Patrick BA, et al. Amyloid beta protein 1-40 and 1-42 levels in matched cerebrospinal fluid and plasma from patients with Alzheimer disease. *Neurosci Lett.* 2001; 304:102–106. [PubMed: 11335065]
32. Lewczuk P, Esselmann H, Otto M, et al. Neurochemical diagnosis of Alzheimer's dementia by CSF Abeta42, Abeta42/Abeta40 ratio and total tau. *Neurobiol Aging.* 2004; 25:273–281. [PubMed: 15123331]
33. Grossman M, Farmer J, Leight S, et al. Cerebrospinal fluid profile in frontotemporal dementia and Alzheimer's disease. *Ann Neurol.* 2005; 57:721–729. [PubMed: 15852395]

34. Haense C, Buerger K, Kalbe E, et al. CSF total and phosphorylated tau protein, regional glucose metabolism and dementia severity in Alzheimer's disease. *Eur J Neurol*. 2008; 15:1155–1162. [PubMed: 18803648]
35. Tapiola T, Alafuzoff I, Herukka SK, et al. Cerebrospinal fluid (beta)-amyloid 42 and tau proteins as biomarkers of Alzheimer-type pathologic changes in the brain. *Arch Neurol*. 2009; 66:382–389. [PubMed: 19273758]
36. Seppala TT, Nerg O, Koivisto AM, et al. CSF biomarkers for Alzheimer disease correlate with cortical brain biopsy findings. *Neurology*. 2012; 78:1568–1575. [PubMed: 22517093]
37. Blennow K, Zetterberg H. Cerebrospinal fluid biomarkers for Alzheimer's disease. *J Alzheimers Dis*. 2009; 18:413–417. [PubMed: 19661632]
38. Benton AL, Varney NR, Hamsher KD. Visuospatial judgment. A clinical test. *Arch Neurol*. 1978; 35:364–367. [PubMed: 655909]
39. Calamia M, Markon K, Denburg NL, et al. Developing a short form of Benton's Judgment of Line Orientation Test: an item response theory approach. *Clin Neuropsychol*. 2011; 25:670–684. [PubMed: 21469016]
40. Tang JX, Baranov D, Hammond M, et al. Human Alzheimer and inflammation biomarkers after anesthesia and surgery. *Anesthesiology*. 2011; 115:727–732. [PubMed: 21857497]
41. Fagan AM, Roe CM, Xiong C, et al. Cerebrospinal fluid tau/beta-amyloid(42) ratio as a prediction of cognitive decline in nondemented older adults. *Arch Neurol*. 2007; 64:343–349. [PubMed: 17210801]
42. Palotas A, Reis HJ, Bogats G, et al. Coronary artery bypass surgery provokes Alzheimer's disease-like changes in the cerebrospinal fluid. *J Alzheimers Dis*. 2010; 21:1153–1164. [PubMed: 21504113]
43. Tarnaris A, Toma AK, Pullen E, et al. Cognitive, biochemical, and imaging profile of patients suffering from idiopathic normal pressure hydrocephalus. *Alzheimers Dement*. 2011; 7:501–508. [PubMed: 21757406]
44. Rudolph JL, Schreiber KA, Culley DJ, et al. Measurement of post-operative cognitive dysfunction after cardiac surgery: a systematic review. *Acta Anaesthesiol Scand*. 2010; 54:663–677. [PubMed: 20397979]

FIGURE 1. Flow diagram. The diagram shows that 342 participants were initially screened for the studies and finally 136 participants were included in the data analysis.

TABLE 1**Characteristics of the Participants (N = 136)**

Age	
Mean $\pm$ SD, yr	71 $\pm$ 5
63–69, n (%)	69 (51)
70–75, n (%)	38 (28)
76–82, n (%)	29 (21)
Male sex, n (%)	75 (55)
Race or ethnic group, n	
White	134
Black	1
Hispanic	0
Asian Indian	1
Others or unknown	0
Education, mean $\pm$ SD, yr	17 $\pm$ 4
Height, mean $\pm$ SD, cm	173 $\pm$ 10
Body weight, mean $\pm$ SD, kg	86 $\pm$ 19
ASA class, n	
I	4
II	115
III	17
Length of anesthesia, * mean $\pm$ SD, min	126 $\pm$ 24
Length of surgery, † mean $\pm$ SD, min	83 $\pm$ 18
Total hip arthroplasty/replacement, n (%)	76 (56)
Total knee arthroplasty/replacement, n (%)	60 (44)
Estimated blood loss, mean $\pm$ SD, mL	194 $\pm$ 115

ASA indicates American Society of Anesthesiologists.

* The length of anesthesia was defined from the time anesthesiologists started the spinal anesthesia in the participants to the time when the participants were sent to the postanesthesia care unit.

† The length of surgery was defined from the time of cutting the skin to the time of the closure of the skin.

TABLE 2

The Raw Scores of Cognitive Tests

	Baseline	1 wk	3–6 mo
HVLTRet	73 ± 32.7	72 ± 29.2	84 ± 26.9
HVLTTR	20 ± 4.4	21 ± 5.2	23 ± 5.0
BVMTTR	19 ± 6.2	18 ± 6.9	20 ± 6.2
BVMTDR	8 ± 2.5	7 ± 2.9	8 ± 2.7
JLO	24 ± 6.1	24 ± 6.4	24 ± 5.2
Trails B	87 ± 34.5	92 ± 44.7	80 ± 31.2

The values are average of mean ± SD from the cognitive test of HVLTRet (percentage of the words recalled), HVLTTR (total number of words recalled), BVMTTR (number of shapes recalled), BVMTDR (number of shapes recalled), JLO (number of lines recognized), and Trails B (time in seconds). The higher scores suggest better cognitive function except Trails B. The lower scores of Trails B (shorter completion time) suggest better cognitive function.

TABLE 3Correlation Between Cognitive Function and the CSF A β 42/Tau Ratio

	Unadjusted			Adjusted by Age and Sex		
	Estimate (Z Score)	Standard Error (Z Score)	P	Estimate (Z Score)	Standard Error (Z Score)	P
HVLTRet 1 wk	7.063	2.732	0.011	8.351	2.734	0.003
HVLTRet 3–6 mo	2.531	2.824	0.372	3.680	2.828	0.196
HVLTTTR 1 wk	0.474	0.398	0.236	0.412	0.408	0.314
HVLTTTR 3–6 mo	0.740	0.406	0.071	0.833	0.412	0.046
BVMTTR 1 wk	0.315	0.565	0.579	0.174	0.562	0.758
BVMTTR 3–6 mo	0.152	0.508	0.766	−0.022	0.515	0.966
BVMTDR 1 wk	0.067	0.236	0.778	0.001	0.240	0.995
BVMTDR 3–6 mo	−0.082	0.202	0.687	−0.099	0.208	0.635
JLO 1 wk	0.915	0.401	0.024	0.954	0.408	0.021
JLO 3–6 mo	1.139	0.436	0.011	1.242	0.446	0.007
Trails B 1 wk	−5.623	2.366	0.019	−4.724	2.396	0.051
Trails B 3–6 mo	1.442	2.221	0.518	1.158	2.285	0.614

The left panel of the table illustrates the results of the Pearson correlation analysis between CSF A β 42/tau ratio and cognitive function in humans. The right panel of the table shows the results of the linear regression analysis after adjustment with age and sex.

TABLE 4Correlation Between Cognitive Function and the CSF A β 40/Tau Ratio

	Unadjusted			Adjusted by Age and Sex		
	Estimate (Z Score)	Standard Error (Z Score)	P	Estimate (Z Score)	Standard Error (Z Score)	P
HVLTRet 1 wk	1.900	2.797	0.498	2.367	2.791	0.398
HVLTRet 3–6 mo	−0.950	2.918	0.745	−0.742	2.893	0.798
HVLTR 1 wk	0.259	0.400	0.519	0.267	0.404	0.509
HVLTR 3–6 mo	0.494	0.422	0.244	0.428	0.424	0.316
BVMTTR 1 wk	−0.585	0.573	0.309	−0.399	0.566	0.482
BVMTTR 3–6 mo	0.958	0.515	0.066	1.045	0.512	0.044
BVMTDR 1 wk	−0.227	0.239	0.345	−0.230	0.241	0.343
BVMTDR 3–6 mo	0.413	0.204	0.045	0.418	0.207	0.046
JLO 1 wk	−0.067	0.409	0.869	−0.146	0.411	0.723
JLO 3–6 mo	−0.289	0.465	0.536	−0.317	0.472	0.504
Trails B 1 wk	1.855	2.409	0.443	1.882	2.391	0.433
Trails B 3–6 mo	−1.712	2.291	0.457	−1.688	2.324	0.469

The left panel of the table illustrates the results of the Pearson correlation analysis between CSF A β 40/tau ratio and cognitive function in humans. The right panel of the table shows the results of the linear regression analysis after adjustment with age and sex.