Published in final edited form as:

Tetrahedron Lett. 2014 January 1; 55(1): . doi:10.1016/j.tetlet.2013.10.143.

Microwave-assisted, one-pot reaction of 7-azaindoles and aldehydes: a facile route to novel di-7-azaindolylmethanes

Md. Imam Uddin^{1,2}, Jason R. Buck^{1,2}, Michael L. Schulte², Dewei Tang², Samir A. Saleh², Yiu-Yin Cheung², Joel Harp⁷, and H. Charles Manning^{1,2,3,4,5,6,*}

¹Department of Radiology and Radiological Sciences, Vanderbilt University Medical Center, Nashville, TN 37232, USA

²Vanderbilt University Institute of Imaging Science, Vanderbilt University, Nashville, TN 37232-2310, USA

³Department of Biomedical Engineering, Vanderbilt University, Nashville, TN 37232, USA

⁴Vanderbilt Ingram Cancer Center, Vanderbilt University Medical Center, Nashville, TN 37232, USA

⁵Program in Chemical and Physical Biology, Vanderbilt University School of Medicine, Nashville, TN 37232, USA

⁶Department of Neurosurgery, Vanderbilt University Medical Center, Nashville, TN 37232, USA

⁷Department of Biochemistry and Center for Structural Biology, Vanderbilt University, Nashville, TN 37232, USA

Abstract

A novel and highly efficient synthetic method leveraging microwave-assisted organic synthesis (MAOS) to yield di-7-azaindolylmethanes (DAIMs) is reported. Under MAOS conditions, reaction of 7-azaindole with aldehydes resulted predominantly in DAIMs, as opposed to the expected 7-azaindole addition products that form at ambient temperature. Based upon studies of different indoles and azaindoles with various aromatic and aliphatic aldehydes, we herein propose a mechanism where rapid and efficient microwave heating promotes nucleophilicity of 7-azaindoles towards the corresponding alkylidene-azaindole intermediate to form the DAIM. This sequence provides a versatile approach to efficiently synthesize novel DAIMs that may be useful pharmaceuticals.

Keywords

Microwave; 7-Azaindoles; Di-7-azaindolylmethanes; One-pot; reactions; MAOS

Bisindolylmethanes, or diindolylmethanes (DIMs), are a unique class of compounds with select members exhibiting biological activity in diseased states, including chemopreventive and therapeutic properties.¹ In nature, DIMs are found in vegetables of the cruciferae family,

© 2013 Elsevier Ltd. All rights reserved.

*Corresponding author. H. Charles Manning, Tel.: +1-615-322-3793; fax: +1-615-322-0734; henry.c.manning@vanderbilt.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

particularly those of the brassica genus, such as broccoli, brussels sprouts, cabbage, and cauliflower, and are reportedly associated with reduced risk of breast, prostate, and colorectal cancer.² In cancer cells, DIMs promote apoptosis and confer protection against DNA-damage.³ For example, DIMs have been shown to inhibit estrogen-induced growth of breast cancer cell lines and human breast-tumors,⁴ with responses associated with G1 cell-cycle arrest.⁵ In leukemia cells, DIMs have been shown to inhibit extracellular signal-regulated kinase activation, which leads to apoptosis.⁶ Similarly, antitumor properties have recently been reported in pancreatic cancer using DIM-C-pPhOH (**1a**).⁷ In addition to use in oncology, DIMs have also conferred neuroprotective properties in preclinical models of Parkinson's disease.⁸ While the importance of DIMs is relatively well established in medicinal chemistry, a daughter class of this molecular scaffold is the di-7-azaindolylmethanes (DAIMs), which may possess unique biological properties (**1b**).^{9b} These studies highlight the attractiveness of these chemical classes and demonstrate the importance of developing robust, yet practical, synthetic methods. Herein, we report the development of a novel synthetic method that can be used to rapidly synthesize candidate DAIMs and DIMs using microwave irradiation.

Prior syntheses of DIMs have featured a wide variety of reagents and reaction conditions, including acid-catalyzed condensation of indoles with aldehydes under ultrasound irradiation,¹⁰ condensation of benzyl alcohols and indoles under visible light irradiation,^{11a} and metal-catalyzed reactions of indole with select aldehydes^{11b} and aldimines.^{11c} Interestingly, there are only a limited number of reported syntheses of DAIMs, these include CH₃-activation¹² and condensation of azaindole with RMgBr^{9a} or CH₂Br₂.^{9b} Unfortunately, these methods are limited by variable yields and exotic reagents. Thus, methods for rapid and efficient synthesis of DAIMs remain a synthetic challenge.

Our interest in the synthesis of azaindoles stems from our previous studies of inhibiting mutant *BRAF* in oncology. We recently reported an MAOS of Vemurafenib (PLX-4032) and PLX-4720, wherein we achieved the synthesis of these V600E-*BRAF* targeted anticancer molecules in reduced total reaction time.¹³ As part of an ongoing program to develop technology-assisted synthesis methods for novel azaindoles, we discovered DAIMs as an unexpected byproduct under MAOS conditions when the azaindole was in excess (2X) to aldehyde.

To explore this phenomenon more closely and potentially elucidate a mechanism, we examined the effects of base, solvent, and heating upon condensation of our model aldehyde and azaindole, *N*-(3–5-difluoro-4-formylphenyl)propane-1- sulfonamide (**2**) and 5-chloro-7-azaindole (**3**) (Table 1). This initial series of experiments revealed that an aqueous solvent and inorganic base were necessary for condensation (Table 1: 5 vs 1, 4, 8) and that microwave heating facilitated DAIM (**4a**) formation over the alcohol (**4b**) (Table 1: 2 vs 5).¹³ In the absence of microwave and water, the reaction gave the expected addition product (**4b**) (Table 1, entry 3). Conversely, conventional heating provided only trace amounts of **4a** as a mixture with **4b**, even after prolonged heating. Using iodine as a Lewis acid did not promote DAIM formation regardless of microwave irradiation (Table 1: 6, 7).^{15,16} In our prior work with Vemurafenib and PLX-4720, DAIM (**4a**) formation was not significant using equivalent amounts of **2** and **3** under MAOS.¹³

Prompted by these results, we examined the scope of this MAOS reaction, coupling 5-substituted-7-azaindoles with different aromatic and aliphatic aldehydes under MAOS, with K₂CO₃ as the base and MeOH:H₂O (1:1) as the solvent system (Table 2). Reaction of various 5-substituted-7-azaindoles with the sulfonamide-functionalized aldehyde **2** yielded the corresponding DAIM in good yield (Table 2: **4a**, **5**, **6**). In addition to the sulfonamide, other aliphatic and aromatic aldehydes reacted with **3** under identical reaction conditions to

give the desired DAIM product in good to excellent yield (Table 2: **5–24**). However, some reaction bias was observed regarding functional group tolerance within the aldehyde. While amido-functionalized substrates such as 4-acetamidobenzaldehyde successfully gave the corresponding DAIM (**22**), maintaining the amide functionality, the ester-functionalized aldehyde methyl 4-formylbenzoate gave compound **24** in modest yield, and only with hydrolysis of the ester functionality to the carboxylic acid (Table 2). This could suggest a possible one-pot DAIM synthesis/ester deprotection approach. Functional group tolerance for the azaindole proved intriguing as well, with the unsubstituted 7-azaindole proving recalcitrant to DAIM formation (Table 2: **21**), but rather undergoing decomposition (by TLC). This could suggest potential electronic tolerances for substituents at the 7-position of the ring, wherein electron-withdrawing groups appear preferred. Structural assignment of the new DAIMs was carried out with high-field NMR analysis, which confirmed a plane of symmetry along the central methyl carbon. These results were further supported by x-ray crystallographic analysis of compound **18** (Figure 2).

We next tested our reaction conditions towards synthesis of DIMs with select indoles and aldehydes. Substituted indoles reacted in a manner similar to their 7-azaindole counterparts, including 5-chloroindole and **2** to give the desired DIM in 62% yield (Table 3: **25**), compared to **4a** in Table 2. However, as with the 7-azaindole substrates, there seemed to be similar electronic constraints for substitutions at the 7-position of the indole ring, as highlighted by unsubstituted indole and the electron-donating hydroxyl group of the 5-hydroxyindole, both of which yielded no DIM product (Table 3: **26, 28**), but rather decomposition (by TLC). Moreover, as with the azaindoles, electron-withdrawing groups seemed to have a net positive effect overall (Table 3: **25**), though not without an apparent threshold, as with 5-nitroindole (Table 3: **27**), which gave a lower yield of 33%.

Towards a potential mechanism, we propose that the addition product (**A**) that initially forms undergoes dehydration in the presence of base and microwave irradiation to the corresponding alkylidene-azaindolene intermediate **B** (Scheme 1), an excellent Michael-acceptor for nucleophiles. Itoh and coworkers reported similar mechanistic observations for their synthesis of DIMs.¹¹ We envision the effects of microwave heating influencing the reaction by increasing nucleophilicity of the 7-azaindole towards **B** and promoting dehydration of **A** (Scheme 1).¹⁷

To test this postulated mechanism, we treated addition-product **4b** with 5-bromo-7-azaindole (**29**) and obtained the unsymmetrical DAIM-product **23** in 54% yield (Scheme 2). The formation of unsymmetrical compound **23** strongly supports the effect of microwave heating on increased nucleophilicity of 7-azaindoles since no DAIMs were observed at ambient temperature.

To verify in part the effects of microwave heating on dehydration of **A** to form intermediate **B**, we performed an ¹⁸O-labeling experiment, anticipating that rehydration in the presence of ¹⁸O-water would label intermediate **B** with ¹⁸O. Thus, treatment of alcohol **4b** in a mixture of methanol and H₂¹⁸O (1:1), in the presence of K₂CO₃, revealed that microwave heating appears to promote dehydration, followed by ¹⁸O-incorporation (>90%), as there was no ¹⁸O incorporation at room temperature (Figure 3). Mass spectrometry analysis of the samples heated under microwave and at room temperature revealed ¹⁸O-incorporation only in the microwave-heated reaction (Figure 3).

In summary, we have developed an efficient method to provide functionalized di-7-azaindolyldimethanes (DAIMs) and diindolyldimethanes (DIMs) *via* a base-promoted reaction of 7-azaindoles and indoles, respectively, with various aldehydes using microwave heating. This stepwise reaction sequence, as shown in Scheme 2, lays the framework for developing

larger libraries of asymmetrical heterodimeric azaindole and indole chemotypes. Biological application of these novel compounds is currently under investigation in our laboratory.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work has been supported by grants from the National Institutes of Health (R01 CA140628, K25 CA127349, P50 CA128323, and P30 DK058404), The Kleberg Foundation.

References and notes

- (a) Gong Y, Sohn H, Xue L, Firestone GL, Bjeldanes LF. *Cancer Res.* 2006; 66(9):4880–7. [PubMed: 16651444] (b) Riby JE, Xue L, Chatterji U, Bjeldanes EL, Firestone GL, Bjeldanes LF. *Mol Pharmacol.* 2006; 69(2):430–9. [PubMed: 16267208] (c) Tadi K, Chang Y, Ashok BT, Chen Y, Moscatello A, Schaefer SD, Schantz SP, Policastro AJ, Geliebter J, Tiwari RK. *Biochem Biophys. Res. Commun.* 2005; 337(3):1019–25. [PubMed: 16219298] (d) Chang X, Tou JC, Hong C, Kim HA, Riby JE, Firestone GL, Bjeldanes LF. *Carcinogenesis.* 2005; 26(4):771–778. [PubMed: 15661811] (e) Chang X, Firestone GL, Bjeldanes LF. *Carcinogenesis.* 2006; 27(3):541–550. [PubMed: 16199440] (f) Rahimi M, Huang KL, Tang CK. *Cancer Lett.* 2010; 295:59–68. [PubMed: 20299148] (g) Shin JA, Shim JH, Choi ES, Leem DH, Kwon KH, Lee SO, Safe S, Cho NP, Cho SD. *Eur J. Cancer Prev.* 2011; 20(5):417–425. [PubMed: 21597379] (h) Gao N, Cheng S, Budhraj A, Liu EH, Chen J, Chen D, Yang Z, Luo J, Shi X, Zhang Z. *PLoS One.* 2012; 7(2):e31783. [PubMed: 22363731] (i) Kandala PK, Wright SE, Srivastava SK. *J Pharmacol Exp. Ther.* 2012; 341:24–32. [PubMed: 22205686] (j) Singh-Gupta V, Banerjee S, Yunker CK, Rakowski JT, Joiner MC, Konski AA, Sarkar FH, Hillman GG. *Cancer Lett.* 2012; 318(1):86–92. [PubMed: 22155105]
- (a) Nachshon-Kedmi M, Yannai S, Haj A, Fares FA. *Food Chem Toxicol.* 2003; 41(6):745–752. [PubMed: 12738179] (b) Le HT, Schaldach CM, Firestone GL, Bjeldanes LF. *J. Biol. Chem.* 2003; 278(23):21136–21145. [PubMed: 12665522] (c) Wang Z, Yu BW, Rahman KM, Ahmad F, Sarkar FH. *Mol. Cancer Ther.* 2008; 7(2):341–349. [PubMed: 18281517] (d) Pappa G, Strathmann J, Löwinger M, Bartsch H, Gerhäuser C. *Carcinogenesis.* 2007; 28(7):1471–1477. [PubMed: 17331956] (e) Kim EJ, Park SY, Shin HK, Kwon DY, Surh YJ, Park JH. *J. Nutr.* 2007; 137(1):31–36. [PubMed: 17182797]
- (a) Bonnesen C, Eggleston IM, Hayes JD. *Cancer Res.* 2001; 61(16):6120–6130. [PubMed: 11507062] (b) Smith TK, Lund EK, Clarke RG, Bennett RN, Johnson IT. *J Agric Food Chem.* 2005; 53(10):3895–3901. [PubMed: 15884814]
- (a) Staub RE, Feng C, Onisko B, Bailey GS, Firestone GL, Bjeldanes LF. *Chem Res Toxicol.* 2002; 15(2):101–109. [PubMed: 11849035] (b) McDougal A, Gupta MS, Morrow D, Ramamoorthy K, Lee JE, Safe SH. *Breast Cancer Res Treat.* 2001; 66(2):147–157. [PubMed: 11437101] (c) Hong C, Firestone GL, Bjeldanes LF. *Biochem Pharmacol.* 2002; 63(6):1085–1097. [PubMed: 11931841]
- Hong C, Kim HA, Firestone GL, Bjeldanes LF. *Carcinogenesis.* 2002; 23(8):1297–1305. [PubMed: 12151347]
- (a) Rooha C, Ismael JS, Zeev E, David H, McCubrey JA, Stephen HS, Michael A, Marina K. *Cancer Research.* 2005; 65(7):2890–2898. [PubMed: 15805291] (b) Gao N, Cheng S, Budhraj A, Liu EH, Chen J, Chen D, Yang Z, Luo J, Shi X, Zhang Z. *PLoS One.* 2012; 7(2):e31783. [PubMed: 22363731]
- (a) Lee S-O, Andey T, Jin U-H, Kim K, Singh M, Safe S. *Oncogene.* 2012; 31(27):3265–3276. [PubMed: 22081070] (b) Lee S-O, Abdelrahim M, Yoon K, Chintharlapalli S, Papineni S, Kim K, Wang H, Safe SC. *Cancer Res.* 2010; 70(17):6824–6836. [PubMed: 20660371]
- Briana RM, James AM, Ryan JH, Paul JL, Stephen S, Daniel LG, Dorothy C, Ronald BT. *J. Pharmacol. Exp. Ther.* 2013; 345(1):125–138. [PubMed: 23318470]

9. (a) Giuseppe G, Mauro M, Gian PM, Sergio P, Maria OT, Silvia P, Francesco L, Francesco DA. Eur. J. Org. Chem. 2004:2411–2420.(b) Song D, Schmider H, Wang S. Org. Lett. 2002; 4(23): 4049–4052. [PubMed: 12423083]
10. Mohammad N, Leila Z, Mohammad RM. Lett. Org. Chem. 2012; 9(5):375–381. And the references cited there in.
11. (a) Nobuta T, Fujiya A, Tada N, Miura T, Itoh A. Synlett. 2012; 23(20):2975–2979.(b) Yanan Z, Qing L. Asian J. Chem. 2012; 24(11):5302–5304. And the references cited there in. (c) Chatterjee PN, Maity AK, Mohapatra SS, Roy S. Tetrahedron. 2013; 69(13):2816–2826.
12. Glen BD, Peter CJ, Stuart GL. Adv. Synth. Catal. 2003; 345:1115–1117.
13. Buck JR, Saleh S, Uddin MI, Manning HC. Tetrahedron Lett. 2012; 53:4161–4165. [PubMed: 23180892]
14. (a) Bollag G, Hirth P, Tsai J, Zhang J, Ibrahim PN, Cho H, Spevak W, Zhang C, Zhang Y, Habets G, Burton EA, Wong B, Tsang G, West BL, Powell B, Shellooe R, Marimuthu A, Nguyen H, Zhang KY, Artis DR, Schlessinger J, Su F, Higgins B, Iyer R, D'Andrea K, Koehler A, Stumm M, Lin PS, Lee RJ, Grippo J, Puzanov I, Kim KB, Ribas A, McArthur GA, Sosman JA, Chapman PB, Flaherty KT, Xu X, Nathanson KL, Nolop K. Nature. 2010; 467(7315):596–599. [PubMed: 20823850] (b) Tsai J, Lee JT, Wang W, Zhang J, Cho H, Mamo S, Bremer R, Gillette S, Kong J, Haass NK, Sproesser K, Li L, Smalley KS, Fong D, Zhu YL, Marimuthu A, Nguyen H, Lam B, Liu J, Cheung I, Rice J, Suzuki Y, Luu C, Settachatgul C, Shellooe R, Cantwell J, Kim SH, Schlessinger J, Zhang KY, West BL, Powell B, Habets G, Zhang C, Ibrahim PN, Hirth P, Artis DR, Herlyn M, Bollag G. Proc Natl Acad Sci. 2008; 105(8):3041–3046. [PubMed: 18287029]
15. Bandgar BP, Shaikh KA. Tetrahedron Lett. 2003; 44:1959–1961.
16. Sashidhara KV, Kumar M, Sonkar R, Singh BS, Khanna AK, Bhatia G. J. Med. Chem. 2012; 55(6):2769–2779. [PubMed: 22339404]
17. Uddin MI, Mimoto A, Nakano K, Ichikawa Y, Kotsuki H. Tetrahedron Letters. 2008; 49:5867–5870.

Figure 1.
Representative DIM, (**1a**) DIM-C-pPhOH, and DAIM, di-7-azaindolylmethane (**1b**).

Figure 2.
X-ray crystal structure (ORTEP drawing) of compound **18**.

Figure 3.

Mass spectrometry of **4b** at room temperature (left) and at microwave heating (right) at 130 °C, 30 min, using H₂¹⁸O in methanol (1:1) and K₂CO₃ (4 eq).

Scheme 1.

Proposed mechanism yielding DAIMs *via* nucleophilic addition of **A** to alkylidene-azaindole **B** under microwave irradiation.

Scheme 2.

Reaction of isolated addition-product intermediate **4b** with 5-bromo-7-azaindole (**29**) under microwave heating. Reaction conditions: **4b** (0.19 mmol), **29** (0.19 mmol), K_2CO_3 (1.27 mmol), MeOH:H₂O (1:1), MW heating at 130 °C for 30 min.

Table 1Evaluation of additives and reaction conditions^a

	2	3		4
Entry	Additive	Solvent	Conditions	Results ^b
1	KOH	DMSO	MW ^c	NR ^e
2	K ₂ CO ₃	MeOH H ₂ O (1:1)	MW ^c	4 : R = 3-(5-Cl-7-azaindole) (72%)
3	KOH	MeOH	RT ^d	
4	KOH	DME	MW ^c	4b : R = OH (54%)
5	K ₂ CO ₃	MeOH H ₂ O (1:1)	RT ^d	4b : R = OH (88%)
6 ^f	I ₂	MeCN	RT ^d	NR ^e
7 ^f	I ₂	MeCN	MW ^c	NR ^e
8	None	MeOH H ₂ O (1:1)	MW ^c	NR ^e

^aReaction conditions: Aldehyde (0.19 mmol), azaindole/indole (0.38 mmol), base (1.27 mmol), solvent (2.5 mL).^bIsolated yield after chromatography.^cMW heating at 130 °C, 30 min.^dRoom temperature, 48 hours.^eNR = no reaction.^fIodine (0.04 mmol).

Table 2

DAIMs synthesized using MAOS^{a,b}

19
(75%)

20
(83%)

21
(0%)

22
(47%)

23
(54%)

24
(26%)

^a Reaction conditions: Aldehyde (0.19 mmol), azaindole (0.38 mmol), K₂CO₃ (1.27 mmol), MeOH:H₂O (1:1), MW heating at 130 °C, 30 min.

^b Isolated yield after chromatography.

Table 3

DIMs synthesized using MAOS^{a,b}

25
(62%)

26
(0%)

27
(33%)

28
(0%)

^aReaction conditions: Aldehyde (0.19 mmol), indole (0.38 mmol), K₂CO₃ (1.27 mmol), MeOH:H₂O (1:1), MW heating at 130 °C, 30 min.

^bIsolated yield after chromatography.