Prevalence of nasal carriage of *S aureus* in children

Liliana Lucia Preotescu,^{1,*} Oana Streinu-Cercel²

Abstract

Asymptomatic carriage of *Staphylococcus aureus* is common, particularly in the anterior part of the nasal cavity. Apart from nasal and nasopharyngeal carriage, another frequent site for *S aureus* colonization is the skin, particularly the inguinal fold, rectum and axilla. The general prevalence of *S aureus* carriage is around 20-30% as reported by relevant studies in field literature. In children, data on nasal carriage appears to be somewhat similar to the prevalence reported in adults, ranging from 18.12 to 38.5 percent, but there are certain particularities and a wide variability between results from different countries and different studies. To determine the real prevalence of *S aureus* and MRSA strains, studies should adjust for confounding factors, as described in this article.

Keywords Nasal carriage, *Staphylococcus aureus*, MRSA, antibiotic resistance.

Asymptomatic carriage of *Staphylococcus aureus* is especially common, particularly in the anterior part of the nasal cavity, where a prevalence rate of around 20% has been reported.¹ Available literature data for *S aureus* nasal colonization in the USA showed a 32.4% prevalence of *S aureus* and 0.8% for methicillin-resistant *S aureus* (MRSA) during 2000-2001,² with similar data for 2006: 31.6% and 0.84%, respectively.³ In the UK, data from 1998 identified a 23% prevalence of *S aureus* (1.5% MRSA)⁴ while a study performed in elderly residents care homes performed during 2006-2009 reported rates of 19-22%.⁵ More recent data from a cross-sectional study in 32 206 subjects

from nine European countries (Austria, Belgium, Croatia, France, Hungary, Spain, Sweden, the Netherlands, UK) revealed a prevalence of 21.6% for *S aureus* and 0.28% for MRSA.⁶

Apart from nasal and nasopharyngeal carriage, another common site for *S aureus* colonization is the skin, particularly the inguinal fold,⁷ rectum and axilla.⁸ The bacterial strains can display an altered antibiotic sensitivity profile, particularly in elderly patients, with comorbid diseases, causes of immune suppression, previous antibiotherapy or history of prior hospital admission during the past months.⁹

In children, data on nasal carriage appears to be somewhat similar to the prevalence reported in adults, but there are certain particularities, better described in the Table below. One particular study identified the main risk factor for MRSA carriage in children as having a parent with MRSA carriage.¹⁰

Data on nasal carriage of *S aureus* is quite polymorphic and the results of epidemiological studies can vary based on a large number of intrinsic factors, such as: timeline of sample collection, season of sample collection, characteristics of subjects included in the study (data from day care centers, kindergartens, schools, versus data from stay-at-home children), age of study subjects, culture and identification technique, and so on. To determine the prevalence of *S aureus* and MRSA strains with a higher accuracy, studies should adjust for factors which might have a significant impact on study results, as described above.

Received: January 17, 2013; accepted: March 2, 2013; published online: March 5, 2013

¹MD, PhD, Assistant lecturer, Department of Infectious Diseases, Carol Davila University of Medicine and Pharmacy, Bucharest, Romania, National Institute for Infectious Diseases 'Prof. Dr. Matei Balș', Romania; ²MD, Assistant Lecturer, Department of Infectious Diseases, Carol Davila University of Medicine and Pharmacy, Bucharest, Romania; National Institute for Infectious Diseases "Prof. Dr. Matei Balș", Bucharest, Romania.

*Corresponding author: Liliana Lucia Preotescu, MD, PhD, Assistant lecturer, Department of Infectious Diseases, Carol Davila University of Medicine and Pharmacy, Bucharest, Romania, National Institute for Infectious Diseases 'Prof. Dr. Matei Balș', Romania; preolil17@yahoo.com

Article downloaded from www.germs.ro

Pre-published online on 5 March 2013

© GERMS 2013

ISSN 2248 - 2997

ISSN - L = 2248 - 2997

Country	Time of study	Number of subjects tested	Overall prevalence of <i>S aureus</i>	Prevalence of MRSA strains	Rural/urban	Type of identification
China ¹¹	2005-2007 2008-2010 Overall	2373	18.12%	4.28% 9.25% 6.28%	Urban	Culture and disk diffusion antibiogram
Gabon ¹²	February-July, 2009	500	29%	1%	Rural and semi-urban	Culture, VITEK-2 identification (Biomérieux)
Colombia ¹³	June-November 2009	104	38.5%	4.8%	Rural	Culture and disk diffusion antibiogram
Hong Kong ¹⁴	September 2009-April 2010	2211	27.6%	1.3%	Both. Day care centers	Culture and latex agglutination (Biomérieux)
Peru ¹⁵	December 2010-June 2011	452 children and adults	24.6%	0.9%	Suburban community	Culture and disk diffusion antibiogram
Russia ¹⁶	2011	63	31.7%	N/A	Urban “ecologically unfavorable district”	Random amplified polymorphic DNA
Gaza strip ¹⁰	March-July 2012	379 children-parents pair	28.4%	12%	Both	Culture, VITEK-2 identification (Biomérieux)

Table. Studies describing the prevalence of *S aureus* and MRSA strains in children.

Acknowledgement This paper is partially supported by the Sectoral Operational Programme Human Resources Development (SOP HRD), financed from the European Social Fund and by the Romanian Government under the contract number POSDRU/89/1.5/S/64109.

Conflicts of interest All authors – none to declare.

References

- Weidenmaier C, Goerke C, Wolz C. *Staphylococcus aureus* determinants for nasal colonization. Trends Microbiol 2012;20:243-50. [CrossRef] [PubMed]
- Kuehnert MJ, Kruszon-Moran D, Hill HA, et al. Prevalence of *Staphylococcus aureus* nasal colonization in the United States, 2001-2002. J Infect Dis 2006;193:172-9. [CrossRef] [PubMed]
- Graham PL 3rd, Lin SX, Larson EL. A U.S. population-based survey of *Staphylococcus aureus* colonization. Ann Intern Med 2006;144:318-25. [PubMed]
- Abudu L, Blair I, Fraise A, Cheng KK. Methicillin-resistant *Staphylococcus aureus* (MRSA): a community-based prevalence survey. Epidemiol Infect 2001;126:351-6. [CrossRef] [PubMed] [FullText]
- Horner C, Parnell P, Hall D, Kearns A, Heritage J, Wilcox M. Methicillin-resistant *Staphylococcus aureus* in elderly residents of care homes: colonization rates and molecular epidemiology. J Hosp Infect 2013;83:212-8. [CrossRef] [PubMed]
- den Heijer CD, van Bijnen EM, Paget WJ, et al. Prevalence and resistance of commensal *Staphylococcus aureus*, including methicillin-resistant *S aureus*, in nine European countries: a cross-sectional study. Lancet Infect Dis 2013;13:409-15. [CrossRef] [PubMed]
- Fritz SA, Hogan PG, Hayek G, et al. *Staphylococcus aureus* colonization in children with community-associated *Staphylococcus aureus* skin infections and their household contacts. Arch Pediatr Adolesc Med 2012;166:551-7. [CrossRef] [PubMed]
- McKinnell JA, Huang SS, Eells SJ, Cui E, Miller LG. Quantifying the impact of extranasal testing of body sites for methicillin-resistant *Staphylococcus aureus* colonization at the time of hospital or intensive care unit admission. Infect Control Hosp Epidemiol 2013;34:161-70. [CrossRef] [PubMed]
- Streinu-Cercel O. Expected sensitivity to antibiotics in bacterial infections. GERMS 2013;3:7. [CrossRef]
- Biber A, Abuelaish I, Rahav G, et al. A typical hospital-acquired methicillin-resistant *Staphylococcus aureus* clone is widespread in the community in the Gaza strip. PLoS One 2012;7:e42864. [CrossRef] [PubMed] [FullText]
- Deng JJ, Wan CM, Mu DZ, et al. [Nasal carriage of community-acquired *staphylococcus aureus* and drug sensitivity tests in healthy children in Chengdu]. Sichuan Da Xue Xue Bao Yi Xue Ban 2012;43:391-4. [PubMed]
- Ateba Ngoa U, Schaumburg F, Adegnika AA, et al. Epidemiology and population structure of *Staphylococcus aureus* in various population groups from a rural and semi urban area in Gabon, Central Africa. Acta Trop 2012;124:42-7. [CrossRef] [PubMed]

13. Rebollo-Perez J, Ordonez-Tapia C, Herazo-Herazo C, Reyes-Ramos N. Nasal carriage of Pantone Valentin leukocidin-positive methicillin-resistant *Staphylococcus aureus* in healthy preschool children. Rev Salud Publica (Bogota) 2011;13:824-32. [[CrossRef](#)]
14. Ho PL, Chiu SS, Chan MY, et al. Molecular epidemiology and nasal carriage of *Staphylococcus aureus* and methicillin-resistant *S. aureus* among young children attending day care centers and kindergartens in Hong Kong. J Infect 2012;64:500-6. [[CrossRef](#)] [[PubMed](#)]
15. Carmona E, Sandoval S, Garcia C. [The frequency and antibiotic susceptibility of *Staphylococcus aureus* from nasal swabs in an suburban marginal population in Lima, Peru]. Rev Peru Med Exp Salud Publica 2012;29:206-11. [[CrossRef](#)] [[PubMed](#)]
16. Baksheeva SS, Avetisyan LP, Chernukha M, Shaginian IA. [Genetic diversity of *Staphylococcus aureus* strains that colonize nose mucous membrane of children living in a large industrial city]. Zh Mikrobiol Epidemiol Immunobiol 2011;52:4. [[PubMed](#)]

Please cite this article as:

Preoțescu, LL, Streinu-Cercel, O. Prevalence of nasal carriage of *S aureus* in children. GERMS. 2013;3(2):49-51. doi: 10.11599/germs.2013.1035