Published in final edited form as:

Curr Gastroenterol Rep. 2009 August ; 11(4): 278–287.

Treating Pelvic Floor Disorders of Defecation: Management or Cure?

Satish S.C. Rao, MD, PhD, FRCP (LON)¹ and Jorge T. Go, MD²

¹Section of Neurogastroenterology, Division of Gastroenterology-Hepatology, Department of Internal Medicine, Iowa City, University of Iowa Carver College of Medicine, Iowa City, Iowa

²Veterans Affairs Quality Scholars (VAQS) Program and the Center for Research in the Implementation of Innovative Strategies in Practice (CRIISP) at the VA Iowa City Healthcare System, Iowa City, IA and the Division of General Internal Medicine, Department of Internal Medicine, University of Iowa Carver College of Medicine, Iowa City, Iowa

Abstract

Pelvic floor disorders that affect defecation consist of structural disorders such as rectocele and functional disorders such as dyssynergic defecation. Evaluation includes a good history and physical examination, a careful digital rectal exam, as well as physiological tests such as anorectal manometry, colonic transit study and balloon expulsion test. Defecography and dynamic magnetic resonance imaging may facilitate detection of structural defects. Their management consists of education and counseling regarding bowel function, diet, laxatives and behavioral therapies. Recently, several randomized clinical trials have shown that biofeedback therapy is effective in dyssynergic defecation. Dyssynergia may also co-exist in structural disorders such as solitary rectal ulcer syndrome or rectocele. Hence, before proceeding with surgery, neuromuscular training or biofeedback should be considered. Several surgical approaches including stapled transanal rectal resection (STARR) have been advocated, but there is lack of well controlled randomized studies and their efficacy is unproven.

INTRODUCTION

The pelvic floor is a complex muscular apparatus that is housed within the pelvic bones and serves three primary functions; defecation, micturition and sexual function. In this review, we will discuss the recent advances in the evaluation and management of disorders that affect the pelvic floor. These disorders consist of functional problems such as dyssynergic defecation or structural problems such as rectal prolapse, rectal mucosal intussusception, solitary rectal ulcer syndrome, rectocele, enterocele, or descending perineum syndrome.

PELVIC FLOOR ANATOMY AND NORMAL PHYSIOLOGY

The pelvic floor consists of superficial and deep muscle layers that interlace and envelope the rectum, bladder and uterus. The superficial muscle layers are relevant to bowel function such as the internal and external anal sphincters, perineal body and the transverse perinei muscles (1). The deep pelvic muscles, also known as levator ani, consist of the pubococcygeus, ileococcygeus and puborectalis muscles (1). It is the traction of the

Address for Correspondence: Satish S.C. Rao, MD, PhD, FRCP (LON), The University of Iowa Hospitals & Clinics, Internal Medicine, GI Division, 200 Hawkins Drive, 4612 JCP, Iowa City, IA 52242, Telephone: 319-3536602, Fax: 319-353-6399, satish-rao@uiowa.edu.

The views expressed in this article are those of the authors and do not necessarily represent the views of the Department of Veterans Affairs.

puborectalis muscle that maintains the anorectal angulation and creates a mechanical barrier for stool flow from the rectum (2). Neurological innervations, primarily from the pudendal nerve, as well as sympathetic and parasympathetic nerves, facilitate the normal function of the pelvic floor.

During normal defecation, the anal sphincters and the puborectalis muscle relax, which allows the anorectal angle to widen and the perineum to descend. Simultaneously, the voluntary effort of bearing down increases the intra-abdominal pressure, together with the contraction of the rectum and puborectalis. These complex and mixed voluntary and involuntary movements facilitates the development of a stripping wave which moves the stool from the rectum and, relaxes the pelvic floor muscles and the anus, resulting in stool evacuation.

STRUCTURAL DISORDERS OF DEFECATION

Rectal prolapse, rectal mucosal intussusception, solitary rectal ulcer syndrome (SRUS), rectocele, enterocele and descending perineum syndrome (DPS) constitute the most common structural pelvic floor disorders that affect defecation. Because these disorders predominantly affect women, a multidisciplinary team consisting of gynecologist or urogynecologist, gastroenterologist and colorectal surgeon, will be most helpful in optimizing diagnosis and management.

Rectal Prolapse

Definition, Pathogenesis, and Symptomatology—A rectal prolapse develops when all layers of the rectal wall protrude through the anus often associated with defecation. This should be differentiated from a rectal mucosal intussusception, where only the rectal mucosa protrudes within the confines of the rectum/or anal canal. The incidence of rectal prolapse has been reported to be as high as 2.5/100,000 and has a high prevalence in elderly women (3**).

Constipation with years of straining weakens the pelvic floor muscles, damaging the pudendal nerve in the process. Aside from constipation, aging and obstetric injury also contribute to pudendal neuropathy (3). The pudendal neuropathy leads to weakness of the anal sphincters. With the dilated anal sphincters offering no counter-acting forces, the rectal wall easily protrudes through the anus.

Most patients present with a rectal protrusion through the anus and/or passage of bloody stool or mucus. Fecal incontinence is seen in 20% to 100% of patients and constipation in 70% (3, 4). On inspection, the rectum is edematous and sometimes friable with an ulcerated mucosa. Rectal prolapse is associated with other defecation disorders such as dyssynergic defecation and intussusception (3). Depending on the degree of protrusion, either up to the anal verge and/or inability to push back the prolapse, it is graded into four types: grade 1= up to anal verge; grade 2=prolapse outside the anus but reduces spontaneously; grade 3=prolapses outside the anus but can be manually reduced; and grade 4=prolapse cannot be reduced manually.

Diagnosis: The diagnosis is made by a careful perineal examination while asking the patient to strain as if to defecate. If no abnormality is detected, it can be useful to have the patient strain in the lavatory. Although defecography will demonstrate the prolapse through the anus, this test is not essential if perineal examination has already exposed the abnormality. However, defecography may also reveal an obtuse anorectal angle and other anatomical defects such as rectocele, sigmoidocele, or enterocele (3). These findings suggest weakness of the pelvic floor and may help the surgeon to better plan a surgical intervention. MR

(magnetic resonance) defecography may provide additional details, especially of adjacent pelvic floor structures and co-existing problems (5).

Management: Among the structural anorectal disorders, rectal prolapse has the clearest indication for surgery. There are currently two accepted approaches: trans-abdominal and perineal. The trans-abdominal approach involves mobilization of the rectum, with or without anterior resection of the rectum followed by fixation of the rectum to the promontory (rectopexy) (3). Since patients with rectal prolapse usually have associated enteroceles and/or genital prolapse, the trans-abdominal approach offers the advantage of obliterating the pouch of Douglas (6).

However, since patients with rectal prolapse are elderly and poor surgical candidates, perineal approaches have been considered in these patients. Among the perineal techniques, the Delorme's procedure or perineal proctosigmoidectomy (Altemeier) has been commonly recommended (3). The Delorme's procedure consists of stripping the mucosa of the prolapsed rectum, plicating the remaining muscular layer and anastomosing the remaining mucosa together (3). In contrast, proctosigmoidectomy involves resecting the prolapsed rectum from below, leaving at least 1 cm of rectum close to the dentate line and then performing a colo-anal anastomosis (3). Lately, advancement in laparoscopy has allowed wider use of the abdominal approach in the elderly population. A recent study found that open and laparoscopic rectopexy have comparable outcomes in terms of recurrence, morbidity and length of stay (7).

Rectal Mucosal Intussusception

Definition, Pathogenesis, and Symptomatology—Rectal mucosal intussusception is also known as “occult rectal prolapse” or “internal procidentia.” Here, the rectal mucosa prolapses but the full thickness of the rectal wall does not protrude through the anus. Rectal intussusception itself is not pathologic since 50–60% of healthy subjects may demonstrate this phenomenon on defecography (3, 8). In symptomatic patients, the rectal intussusception is larger when compared to asymptomatic counterparts (8). Furthermore, rectal intussusception rarely leads to a full-thickness rectal prolapse and is considered by many as a consequence of untreated dyssynergic defecation (3).

Most patients with rectal intussusception complain of a feeling of incomplete evacuation. Patients usually have other structural abnormalities such as rectocele, SRUS, pudendal neuropathy or sphincter damage that may cause more symptoms such as soiling, bloody discharge, and/or fecal incontinence (3, 9).

Diagnosis: Rectal examination detects only a third of patients with rectal intussusception, making defecography the standard test for establishing this diagnosis (3). However, rectal mucosal folds can sometimes be misread as rectal intussusception (3). Recently, dynamic three-dimensional CT, three-dimensional dynamic anorectal ultrasonography and dynamic MRI have shown promise as alternative imaging options (10–12).

Management: The role of surgery in rectal intussusception is controversial at best.

Patients are usually started on fiber-enriched diet and laxatives, followed by biofeedback therapy to correct the underlying dyssynergia. Several studies showed favorable results with this approach (3). Some studies suggest avoiding surgery altogether because of the following: (i) rectopexy was found to correct incontinence but made constipation symptoms worse; (ii) only a few patients improve after surgery; (iii) some patients have poor prognosis after surgery, especially those with preoperative diarrhea, fecal incontinence, and descending perineum syndrome and proximal intussusception (13). Surgery should only be

performed on patients with large intussusception with refractory symptoms and in whom all conservative and behavioral approaches have failed over several years.

Recently, a panel of experts recommended stapled transanal rectal resection (STARR) as a novel approach to restoring normal anatomy in patients with large symptomatic rectal intussusception and rectocele after failing conservative therapy (14**). In principle, the rectovaginal septum is strengthened and the redundant tissues from the anterior and posterior rectal walls are resected (14). Complications from STARR include: bleeding, hematoma, urinary retention, severe pain, dehiscence, infection, fecal incontinence, recto-vaginal fistula, necrotizing fasciitis, peritonitis, and fecal urgency (14).

Rectocele

Definition, Pathogenesis, and Symptomatology—Rectocele is a protrusion of the rectal wall, usually anterior and towards the vagina (Figure 1). However, posterior rectoceles may also occur. Using a subset the Pelvic Floor Distress Inventory questionnaire, the Pelvic Floor Disorders Network (PFDN) found the prevalence of pelvic floor disorders in the US to be 23.7% (15**).

Rectocele is believed to be caused by weakness of the pelvic floor and rectovaginal septum. Obstetric injury from multiple vaginal deliveries is a common cause (3, 16). Rectoceles are also seen in postmenopausal women, especially if they are nulliparous (3, 16). Childbirth and/or excessive and inappropriate straining weaken the vaginal septum, pelvic floor muscles, and damage the pudendal nerve. This process lowers the pelvic floor and makes the vaginal outlet bigger. Since the vagina cannot close on straining, this creates a pressure gradient between the rectum (high pelvic pressure) and the vagina (low atmospheric pressure) (3). The high abdominal pressure generated during chronic straining pushes the rectal wall towards the vaginal, producing a rectocele. Hysterectomy, dyssynergic defecation and intussusception are often associated with rectoceles (3) and whether they are a cause or an effect is unclear.

Aside from pelvic floor weakness and a rectocele, patients usually have non-specific symptoms. Patients can present with pelvic pain, feeling of prolapse, dragging, constant pressure, backache, constipation, fecal soiling, urologic complaints and dyspareunia (3). Vaginal splinting that consists of digital support can sometimes help with either initiating or completing defecation.

Diagnosis: Diagnosis is apparent on inspection and digital examination. Anterior bulging of the rectal wall is more evident on straining. Digital rectal palpation may also confirm the diagnosis. In the last decade, the Pelvic Organ Prolapse Quantification (POP-Q) has been used to grade pelvic organ prolapse (17). It takes 6 different measurements of the female genital area, grading genital prolapse into 4 standard stages.

Defecography can detect rectocele, document its size and identify concomitant retention (3). However, its usefulness in clinical work-up is limited. Eighty percent of asymptomatic controls may show small rectoceles (3, 8). Hence, clinical correlation is of paramount importance and one should search for another etiology before labeling dyssynergia as due to rectocele. Only rectoceles larger than 3 cm or with significant contrast retention is considered clinically significant (3). MRI can also provide good visualization of the rectocele and pelvic floor movements (3). There are no physiologic findings that are specific for a rectocele (3). However, dyssynergic pattern of defecation on anorectal manometry was seen with increasing frequency in patients with rectocele (60%) compared to those without (24%) (18).

Management: The first-line treatments are fiber-rich diet, bulk-forming agents and laxatives to ensure easy defecation.

For both rectocele and rectal mucosal intussusception, surgical treatment carries frequent recurrences (3). Recent studies have shown the importance of detecting “occult diseases” like psychological disorders (anxiety and depression), functional disorders (dyssynergic defecation, rectal hyposensation, pudendal neuropathy) and organic disorders (urogynecological disorders, solitary rectal ulcer) (19**). In these patients, conservative therapy in the form of psychological counseling and behavioral modification, including posture and education of the defecation process and biofeedback therapy are recommended before considering surgery (19).

However, surgery is considered appropriate in patients with a large rectocele (>4cm) that causes a bulge in the vagina or a feeling of prolapse and after a failure of medical therapy. The success rate of surgery is reportedly 75%, with the vaginal approach associated with a lower recurrence rate than transanal approach (3). However, complications such as constipation and sexual dysfunction are very common (3). In patients with dyssynergic defecation associated with rectocele, the transanal approach is ideal (3). However, the transanal approach may compromise anal sphincter pressures (3). The STARR procedure, described in the section on rectal intussusception, has also been performed with some success in patients with rectoceles (14). Although reducing a rectocele size may improve rectal emptying, it may not help with overall symptoms.

Solitary rectal ulcer syndrome (SRUS)

Definition, Pathogenesis, and Symptomatology—The annual incidence of SRUS is 1 to 3.6 per 100,000; 80% of patients are less than 50 years of age (3, 20). The gender distribution is either equal or slightly higher in females (20). SRUS is often associated with rectal mucosal intussusception as revealed by defecography (3). Mucosal ulceration is believed to be due to forceful straining against an immobile or a non-relaxing pelvic floor together with trauma from digital manipulations as well as ischemic necrosis of the prolapsing rectal mucosa. Spontaneous resolution of SRUS during pregnancy suggests that hormones may play a role in its pathogenesis (21).

Patients usually present with rectal bleeding, rectal pain, mucus discharge, straining and tenesmus and a feeling of incomplete evacuation. About 55% of patients present with constipation, 20%–40% with diarrhea, and 25% are asymptomatic (3). A quarter of the patients are misdiagnosed and treated as inflammatory bowel disease (3). Excessive rectal bleeding has been previously reported (3). Obsessive-compulsive disorder has been described in patients with SRUS (3).

Diagnosis: Sigmoidoscopy with rectal biopsy is the gold standard for diagnosis. On endoscopic inspection, the ulcer in SRUS is usually a small, shallow lesion with a white slough or hyperemic mucosa on the anterior wall of the rectum (3). The lesions can be multiple (30%), ulcerated (57%), polypoid (25%) and with patches of hyperemic mucosa (18%) (3). Histologically, the mucosa appears elongated with distorted glands at the base, with an edematous fibroblast-rich lamina propria and thickened inner circular muscular layer (22). When the glands migrate down to the submucosa, bleeding may occur. Pathognomonic features in SRUS include: decussation of the two muscularis layers, nodular induration of the inner layer, and grouping of outer longitudinal layers into bundles (22). Biopsy is needed to differentiate SRUS from ulcers that are caused by non-steroidal anti-inflammatory agents, malignancy and sometimes rectal endometriosis (3). Defecography may show concomitant morphological abnormalities like intussusception in 45% to 80% of subjects (3). However, it

is not essential for establishing a diagnosis. Barium enema may show nodularity of the rectal mucosa, thickening of the rectal folds, strictures, polypoid lesions or ulcerations (3).

Management: Behavioral advice remains the mainstay of treatment. The most important step is to advise the patient not to strain and stop digital rectal maneuvers. These recommendations may improve symptoms in 67% of patients (3, 23). A high-fiber diet showed a variable response rate of 19% to 70%, suggesting that although diet helps, it is often not enough (3, 24). Local treatment with topical steroids and sulphasalazine has been generally ineffective (3). Although there is limited data, sucralfate enemas and topical human fibrin sealant have been tried (3).

To date, biofeedback therapy appears to be the most effective therapy; although RCTs are lacking. Seventy-five percent (12/16) of patients with SRUS had symptomatic improvement after biofeedback therapy and 31% (5/16) had ulcer resolution on sigmoidoscopy (25). Mucosal flow improved in patients who felt subjectively better after biofeedback. Another study prospectively followed 11 patients with refractory SRUS treated with biofeedback (26**). Eighty-two percent (9/11) had dyssynergic defecation. After biofeedback therapy, straining effort and stool frequency improved. Five patients discontinued digital maneuvers and bleeding ceased in 56%. Ulcer healing was reported in 10 patients: 4 had complete healing, 2 had >50% healing and 4 had <50% healing.

Constipation is a known complication of surgery and since constipation is the underlying mechanism for this problem, surgery should only be performed in highly selected cases (3). The surgical options include; rectopexy, anterior rectopexy and Delorme's procedure (3).

Enterocoele

Definition, Pathogenesis, and Symptomatology—In enterocoele, a peritoneum-lined sac (usually the small bowel) herniates into the space between the vagina and rectum and produce symptoms of obstructed defecation. The epidemiology of enterocoele is currently unknown (3). However, there is a high correlation between women who have had hysterectomy and enterocoeles (27). Usually, patients with enterocoeles have concomitant anorectal morphological abnormalities like rectocoele, excessive perineal descent and rectal mucosal intussusception (3, 27).

Patients usually present with either pelvic pain or heaviness (on standing), difficult defecation and feeling of incomplete evacuation. Upon lying down, the pain and heaviness disappears as the bowel repositions itself into the abdominal cavity.

Diagnosis: Enterocoele is best diagnosed with defecography. Since enterocoele usually involves herniated small bowel, it is crucial that the small bowel is opacified with oral contrast (3). Other imaging techniques like dynamic three-dimensional CT, anorectal ultrasound and dynamic MRI may facilitate its diagnosis (10–12).

Management: Improving defecation mechanics and biofeedback therapy are the initial treatment of choice for patients with enterocoeles (3). However, some patients might benefit from surgery, especially those having persistent pelvic pain and constant urge to defecate. When surgery is indicated, the goal is to obliterate the pouch of Douglas (3). Abdominal approach (either open or laparoscopic) and transvaginal approach have been tried in the past (3).

However, post-surgical outcomes remain poor. In a study of 25 patients with enterocoele, 5 had surgery; only 1 had complete resolution and 4 showed partial relief of symptoms (27).

Similarly, despite adequate surgical correction of the defect, symptoms of obstructed defecation persisted in 75% of the patients being followed up to 85 months (28).

Descending perineum syndrome (DPS)—Descending perineum syndrome (Figure 1) is also known as “excessive perineal descent.” It is characterized by ballooning of the perineum several centimeters below the bony outlet of the pelvis during a straining effort (29). Clinically, patients presents with painful defecation, impaired defecation, excessive straining, a sense of incomplete, or fecal incontinence (29, 30). Hysterectomy is a known predisposing factor in the development of DPS (29). DPS is indicative of pelvic floor weakness and usually associated with other anorectal disorders such as rectal prolapse, rectocele, SRUS or enterocele (30).

According to Parks’ and colleagues criteria, DPS can be diagnosed on physical examination (29). The patient is asked to lie on a left lateral position and the anal canal is examined for rapid descent of more than 3 cm during a straining effort (29). However, since this method is not physiologic (given the patient is on a lateral decubitus position), DPS is preferably diagnosed using defecography (31). Diagnostic criteria for DPS on defecography include: (i) perineal descent is more than 4 cm at rest; or (ii) during a maximal push effort, the perineal descent is more than 3 cm from the value measured at rest (31). When available, dynamic MR imaging is also a good diagnostic option (5, 31).

There is no surgical option for isolated DPS. Treatment consists mainly of correcting excessive straining, use of an artificial device (a polycarbonate plate placed under the toilet seat to support the perineum during defecation) and biofeedback therapy (31). Although controlled trials are lacking, success with biofeedback therapy is around 50% (31).

DYSSYNERGIC DEFECATION

Definition, Pathogenesis, and Symptomatology

Dyssynergic defecation has been described in the literature with a plethora of other terms such as anismus, pelvic floor dyssynergia, obstructive defecation, paradoxical puborectalis contraction, pelvic outlet obstruction, and spastic pelvic floor syndrome (32**). However, the term dyssynergic defecation is now recommended by most experts (33**). In tertiary care setting, the prevalence of dyssynergic defecation among patients with chronic constipation is about 40–50% (34).

It was suggested that paradoxical anal contraction or involuntary anal spasm during defecation was the main cause of this problem. However, simple myectomy or chemo-paralysis of the anal sphincter with Botulinum toxin lead to a minimal improvement only in patients with dyssynergia, suggesting that other mechanisms may play a role (32). Later studies have found that a failure of recto-anal coordination, manifesting as either impaired rectal contraction, paradoxical anal contraction, or inadequate anal relaxation were primarily responsible for this condition (35).

In a large series of well characterized patients with dyssynergic defecation, the most common symptoms were excessive straining (84%), feeling of incomplete evacuation (76%), abdominal bloating (74%), passage of hard stools (65%) and less than 3 bowel movements per week (62%) (34). Although these patients would not readily admit to their doctors, digital maneuvers such as disimpaction or vaginal splinting to relieve symptoms were frequently encountered (~50%) (32).

Diagnosis of Dyssynergic Defecation

Medical History and Physical Examination—Diagnosis of dyssynergic defecation starts with a good history, physical examination and digital rectal exam. The rectal exam should include a careful inspection of the perianal skin for excoriation, fissures, skin tags or hemorrhoids. Perianal sensation and anocutaneous reflex can be examined by stroking the perianal skin with a cotton bud. When asked to bear down on digital exam, patients with dyssynergia may demonstrate paradoxical contraction of the external sphincter, puborectalis, ineffective perineal descent, or impaired push effort or a combination.

Laboratory Tests, Imaging and Physiologic Tests—Underlying metabolic and pathologic disorders should be excluded first by using routine laboratory blood tests, colonoscopy and/or barium enema. Physiologic tests are essential for the diagnosis of dyssynergic defecation, especially anorectal manometry (ARM) and balloon expulsion test (32). ARM provides information on rectal and anal pressures at rest and during simulated defecation (32). It also tests for rectal compliance, rectal sensation and recto-anal reflexes (32). For example, it can exclude Hirschsprungs disease. However, one should exercise caution when diagnosing dyssynergic defecation based on ARM alone. An ARM performed with the patient lying in the left lateral position can sometimes show a dyssynergic pattern, but when the patient is seated on a commode and particularly when provided with a sensation of stooling, most normal subjects exhibit a normal pattern of defecation (36–38). Similarly, many fail to expel the artificial stool in the lying position but can expel the balloon easily on a sitting position (36–38). Thus, dyssynergia is best diagnosed by asking the subject to evacuate in a sitting position after evoking a sensation of stooling (36–38). Four patterns of dyssynergia have been described with ARM (32). In addition, the defecation index (DI) -calculated from ARM-generated pressures - can serve as a simple and useful quantitative measure of recto-anal coordination during defecation (32, 37).

Rome III Criteria for Diagnosis of Dyssynergic Defecation—Dyssynergic defecation is diagnosed when a patient fulfills Rome III criteria for functional constipation and exhibits dyssynergic pattern of defecation on ARM or electromyogram (EMG) and demonstrates one other quantifiable measure of abnormal defecation such as an abnormal balloon expulsion test or a prolonged delay in colonic transit or incomplete evacuation during defecography (33**).

The balloon expulsion test provides information on the patient's ability to expel an artificial stool. A normal individual can pass a 50-ml water-filled balloon placed in the rectum within 1 minute. Although this test has 80–90% specificity for dyssynergia (32, 39), its sensitivity is only 50% (32).

Since slow transit constipation may co-exist with dyssynergic defecation, a colonic transit study may prove to be useful. A colonic transit is considered abnormal if more than five markers (20% markers retained) are present on plain abdominal film taken 5 days (120 hours) after ingestion of a sitzmarker® capsule containing 24 radio-opaque markers.

Other Imaging Modalities for Dyssynergic Defecation—Several imaging modalities may identify dyssynergic defecation. Defecography is performed by placing barium paste into the patient's rectum. A video fluoroscopic imaging system records the anatomic and functional changes as the patient tries to evacuate the barium. Patients with dyssynergic defecation shows poor activation of the levator muscles, prolonged retention of the contrast material, inability to expel the barium or absence of a stripping wave in the rectum (32). Defecography is only considered an adjunct to clinical and manometric assessment of anorectal function. Endoanal and dynamic pelvic MRI (MRI Defecography) is the only

imaging study that can evaluate pelvic floor anatomy in dynamic motion (32). Recently, 3-dimensional dynamic anorectal ultrasonography (echodefecography) has been tested against standard defecography and it showed good agreement for detecting rectocele and dyssynergia (12).

Management of Dyssynergic Defecation

Standard Treatment and other Medical Options—Standard treatment consists of avoiding constipating medications (i.e. narcotics), increasing fiber and fluid intake, exercise activity and timed toilet training. A fiber intake of 20 grams per day is recommended. Currently, there is insufficient data to recommend synthetic fiber supplements. Only psyllium, a natural fiber supplement, was given a grade B recommendation by the American College of Gastroenterology (ACG) task force (40) and a systematic review (41). Medications that promote bowel movement like stool softeners (sodium and calcium docusate compounds), stimulant laxatives (senna, bisacodyl, castor oil), osmotic laxatives (salts of magnesium, phosphate and sulfate, lactulose, sorbitol, mannitol, polyethylene glycol) can be useful (41). A recent study reported relief of constipation in 52% of patients treated with polyethylene glycol compared to 11% treated with placebo (42). Lubiprostone, a chloride channel-2 activator, at 24 ug twice daily was more effective than placebo in decreasing straining, improving stool frequency, and relieving symptoms of chronic constipation (43, 44), but whether these medications help dyssynergic defecation is not known.

Biofeedback Therapy—Biofeedback therapy is proven to be effective in patients with dyssynergic defecation (45**). The main purpose of biofeedback is to restore the normal pattern of defecation using an instrument-based learning process that is based on “operant conditioning” techniques (46, 47). There are two primary goals: (i) correct the underlying dyssynergia involving the abdominal, rectal and anal muscles; and (ii) improve rectal sensory perception in patients with impaired rectal sensation.

Biofeedback consists of improving abdominal push effort, manometric guided pelvic floor relaxation, followed by simulated defecation training. This procedure involves inserting a manometric probe into the patient’s rectum. This would capture rectal and anal pressure readings on a monitor, providing the patient instant feedback regarding their performance and helps them to learn quickly. Patients are then taught to coordinate their abdominal push effort showing a rise in intra-rectal pressure and synchronized relaxation of the anal sphincter depicted by decreased anal pressure on the monitor. About 10–15 maneuvers are usually attempted in a single session. Next, a balloon in the rectum is distended with 60cc of air to provide the patient a sensation of rectal fullness or the desire to defecate. Patients are asked to attempt defecation while observing the pressure changes in the monitor. These maneuvers are then repeated 5–10 times. Lastly, an artificial stool (50-ml water-filled balloon) is placed in the rectum. The patient is asked to sit on a commode and attempt defecation. The patient is taught to relax the pelvic floor muscles and to correct their posture and breathing techniques. Gradually, the patient learns to coordinate the defecation maneuver and be able to successfully expel the balloon.

Since about half of patients with dyssynergic defecation have impaired rectal sensation, rectal sensory training is also beneficial. The goal of this training is to improve sensory perception to improve awareness for stooling (47). This is accomplished by repeated inflations and deflations of the rectal balloon, establishing newer thresholds for rectal perceptions in the process.

Depending on their needs, the number of biofeedback sessions and the length of each session should be customized for each patient. Typically, each biofeedback training session

usually takes an hour. Patients are asked to visit the clinic every 2 weeks with an average of 4–6 training sessions. Aside from manometry, biofeedback therapy has been tried using different instruments. EMG biofeedback, balloon defecation training, and home training devices have been described (32).

There are four randomized clinical trials evaluating the use of biofeedback in patients with dyssynergic defecation (Table 1) (45, 48–50**). All of these studies showed that biofeedback is superior to laxatives, sham feedback, standard therapy, placebo, and diazepam. Figure 2 shows the manometry tracings of a patient with dyssynergic defecation before and after biofeedback therapy. It is notable that the rectal and anal pressures normalized after biofeedback.

Despite its proven effectiveness, biofeedback is only offered in a few motility centers in the US. To treat dyssynergic patients in the community, a home-based biofeedback program maybe important. Although there are studies that have demonstrated the feasibility of home training, there is no controlled trial that has compared home training with the standard office biofeedback (32).

CONCLUSIONS

Pelvic floor disorders that include dyssynergic defecation and anatomical abnormalities of the anorectum are very common, particularly in women. The role of biofeedback is increasingly recognized not only in patients with dyssynergic defecation but also in others with rectal mucosal intussusception, rectocele, SRUS and enterocele. This underscores the importance of performing physiologic tests such as anorectal manometry and balloon expulsion test to properly diagnose dyssynergic defecation. Correcting the underlying pathophysiological dysfunction offers patients better control of symptoms when compared to performing surgical correction of the anatomic defects. Further research is required to define the phenotype of these patients, examine predisposing factors and examine the correlation of symptoms with diagnostic tests and longitudinal follow-up. Most importantly, randomized controlled trials are urgently needed to test current and evolving therapies for these disorders.

Acknowledgments

Grant Support: Dr. Satish Rao is supported by NIH grant RO1 DK 57100-05

References and Recommended Reading

Papers of particular interest, published recently, have been highlighted as:

*Of importance

**Of major importance

1. Raizada V, Mittal RK. Pelvic floor anatomy and applied physiology. *Gastroenterol Clin North Am.* 2008 Sep.37(3):493, 509, vii. [PubMed: 18793993]
2. Azpiroz F, Fernandez-Fraga X, Merletti R, et al. The puborectalis muscle. *Neurogastroenterol Motil.* 2005 Jun; 17(Suppl 1):68–72. [PubMed: 15836457]
- 3**. Felt-Bersma RJ, Stella MT, Cuesta MA. Rectal prolapse, rectal intussusception, rectocele, solitary rectal ulcer syndrome, and enterocele. *Gastroenterol Clin North Am.* 2008 Sep.37(3):645, 68, ix. This review article talks about structural disorders of defecation in great detail and surgical techniques used in their treatment. [PubMed: 18794001]

4. Madoff RD, Mellgren A. One hundred years of rectal prolapse surgery. *Dis Colon Rectum*. 1999 Apr; 42(4):441–50. [PubMed: 10215042]
5. Bharucha AE. Update of tests of colon and rectal structure and function. *J Clin Gastroenterol*. 2006 Feb; 40(2):96–103. [PubMed: 16394868]
6. D'Hoore A, Penninckx F. Laparoscopic ventral recto(colpo)pexy for rectal prolapse: Surgical technique and outcome for 109 patients. *Surg Endosc*. 2006 Dec; 20(12):1919–23. [PubMed: 17031741]
7. Slawik S, Soulsby R, Carter H, et al. Laparoscopic ventral rectopexy, posterior colporrhaphy and vaginal sacrocolpopexy for the treatment of recto-genital prolapse and mechanical outlet obstruction. *Colorectal Dis*. 2008 Feb; 10(2):138–43. [PubMed: 17498206]
8. Shorvon PJ, McHugh S, Diamant NE, et al. Defecography in normal volunteers: Results and implications. *Gut*. 1989 Dec; 30(12):1737–49. [PubMed: 2612988]
9. Dvorkin LS, Knowles CH, Scott SM, et al. Rectal intussusception: Characterization of symptomatology. *Dis Colon Rectum*. 2005 Apr; 48(4):824–31. [PubMed: 15785903]
10. Okamoto N, Maeda K, Kato R, et al. Dynamic pelvic three-dimensional computed tomography for investigation of pelvic abnormalities in patients with rectocele and rectal prolapse. *J Gastroenterol*. 2006 Aug; 41(8):802–6. [PubMed: 16988771]
11. Bruscianno L, Limongelli P, Pescatori M, et al. Ultrasonographic patterns in patients with obstructed defaecation. *Int J Colorectal Dis*. 2007 Aug; 22(8):969–77. [PubMed: 17216218]
12. Murad-Regadas SM, Regadas FS, Rodrigues LV, et al. A novel three-dimensional dynamic anorectal ultrasonography technique (echodefecography) to assess obstructed defecation, a comparison with defecography. *Surg Endosc*. 2008 Apr; 22(4):974–9. [PubMed: 17705074]
13. Sieleznoff I, Malouf A, Cesari J, et al. Selection criteria for internal rectal prolapse repair by delorme's transrectal excision. *Dis Colon Rectum*. 1999 Mar; 42(3):367–73. [PubMed: 10223758]
- 14*. Corman ML, Carriero A, Hager T, et al. Consensus conference on the stapled transanal rectal resection (STARR) for disordered defaecation. *Colorectal Dis*. 2006 Feb; 8(2):98–101. This paper encourages the use of a novel procedure called STARR, to reverse the anatomic disorders of the pelvic floor. [PubMed: 16412068]
- 15*. Nygaard I, Barber MD, Burgio KL, et al. Prevalence of symptomatic pelvic floor disorders in US women. *JAMA*. 2008 Sep 17; 300(11):1311–6. This study highlights the prevalence of pelvic floor disorders in the US using subjects who participated in the nationwide NHANES study (National Health and Nutrition Examination Survey). [PubMed: 18799443]
16. Beevors MA, Lubowski DZ, King DW, et al. Pudendal nerve function in women with symptomatic utero-vaginal prolapse. *Int J Colorectal Dis*. 1991 Feb; 6(1):24–8. [PubMed: 2033349]
17. Bump RC, Mattiasson A, Bo K, et al. The standardization of terminology of female pelvic organ prolapse and pelvic floor dysfunction. *Am J Obstet Gynecol*. 1996 Jul; 175(1):10–7. [PubMed: 8694033]
18. Mellgren A, Lopez A, Schultz I, et al. Rectocele is associated with paradoxical anal sphincter reaction. *Int J Colorectal Dis*. 1998; 13(1):13–6. [PubMed: 9548094]
- 19*. Pescatori M, Spyrou M, Pulvirenti d'Urso A. A prospective evaluation of occult disorders in obstructed defecation using the 'iceberg diagram'. *Colorectal Dis*. 2007 Jun; 9(5):452–6. This paper discusses the role of "occult disorders" in patients with structural anorectal defects and its effects on surgical outcomes. There is increasing awareness that these psychological, functional and organic disorders should be treated first before considering any surgery. [PubMed: 17504343]
20. Haray PN, Morris-Stiff GJ, Foster ME. Solitary rectal ulcer syndrome--an underdiagnosed condition. *Int J Colorectal Dis*. 1997; 12(5):313–5. [PubMed: 9401849]
21. Sood SK, Garner JP, Amin SN. Spontaneous resolution of solitary rectal ulcer syndrome during pregnancy: Report of a case. *Dis Colon Rectum*. 2008 Jul; 51(7):1149–52. [PubMed: 18392597]
22. Kang YS, Kamm MA, Engel AF, et al. Pathology of the rectal wall in solitary rectal ulcer syndrome and complete rectal prolapse. *Gut*. 1996 Apr; 38(4):587–90. [PubMed: 8707093]
23. Ho YH, Ho JM, Parry BR, et al. Solitary rectal ulcer syndrome: The clinical entity and anorectal physiological findings in singapore. *Aust N Z J Surg*. 1995 Feb; 65(2):93–7. [PubMed: 7857237]

24. Tjandra JJ, Fazio VW, Church JM, et al. Clinical conundrum of solitary rectal ulcer. *Dis Colon Rectum*. 1992 Mar; 35(3):227–34. [PubMed: 1740066]
25. Jarrett ME, Emmanuel AV, Vaizey CJ, et al. Behavioural therapy (biofeedback) for solitary rectal ulcer syndrome improves symptoms and mucosal blood flow. *Gut*. 2004 Mar; 53(3):368–70. [PubMed: 14960517]
- 26**. Rao SS, Ozturk R, De Ocampo S, et al. Pathophysiology and role of biofeedback therapy in solitary rectal ulcer syndrome. *Am J Gastroenterol*. 2006 Mar; 101(3):613–8. Although an uncontrolled study, this article emphasizes the role of biofeedback in the management of SRUS. [PubMed: 16464224]
27. Takahashi T, Yamana T, Sahara R, et al. Enterocoele: What is the clinical implication? *Dis Colon Rectum*. 2006 Oct; 49(10 Suppl):S75–81. [PubMed: 17106819]
28. Oom DM, van Dijk VR, Gosselink MP, et al. Enterocoele repair by abdominal obliteration of the pelvic inlet: Long-term outcome on obstructed defaecation and symptoms of pelvic discomfort. *Colorectal Dis*. 2007 Nov; 9(9):845–50. [PubMed: 17608820]
29. Pucciani F, Boni D, Perna F, et al. Descending perineum syndrome: Are abdominal hysterectomy and bowel habits linked? *Dis Colon Rectum*. 2005 Nov; 48(11):2094–9. [PubMed: 16228840]
30. Harewood GC, Coulie B, Camilleri M, et al. Descending perineum syndrome: Audit of clinical and laboratory features and outcome of pelvic floor retraining. *Am J Gastroenterol*. 1999 Jan; 94(1):126–30. [PubMed: 9934742]
31. Wexner, SD.; Stollman, N. *Diseases of the Colon*. CRC Press; 2006. Perineal descent syndrome; p. 127
- 32**. Rao SS. Dyssynergic defecation and biofeedback therapy. *Gastroenterol Clin North Am*. 2008 Sep; 37(3):569–86. This is a comprehensive review article that talks about the pathophysiology, symptomatology, diagnosis and treatment options for patients with dyssynergic defecation. [PubMed: 18793997]
- 33**. Bharucha AE, Wald A, Enck P, et al. Functional anorectal disorders. *Gastroenterology*. 2006 Apr; 130(5):1510–8. This report offers a consensus and diagnostic criteria for functional anorectal disorders such as fecal incontinence, anorectal pain, and disorders of defecation. [PubMed: 16678564]
34. Rao SS, Tuteja AK, Vellema T, et al. Dyssynergic defecation: Demographics, symptoms, stool patterns, and quality of life. *J Clin Gastroenterol*. 2004 Sep; 38(8):680–5. [PubMed: 15319652]
35. Rao SS, Welcher KD, Leistikow JS. Obstructive defecation: A failure of rectoanal coordination. *Am J Gastroenterol*. 1998 Jul; 93(7):1042–50. [PubMed: 9672327]
36. Rao SS, Patel RS. How useful are manometric tests of anorectal function in the management of defecation disorders? *Am J Gastroenterol*. 1997 Mar; 92(3):469–75. [PubMed: 9068472]
37. Rao SS, Hatfield R, Soffer E, et al. Manometric tests of anorectal function in healthy adults. *Am J Gastroenterol*. 1999 Mar; 94(3):773–83. [PubMed: 10086665]
38. Rao SS, Azpiroz F, Diamant N, et al. Minimum standards of anorectal manometry. *Neurogastroenterol Motil*. 2002 Oct; 14(5):553–9. [PubMed: 12358684]
39. Minguez M, Herreros B, Sanchiz V, et al. Predictive value of the balloon expulsion test for excluding the diagnosis of pelvic floor dyssynergia in constipation. *Gastroenterology*. 2004 Jan; 126(1):57–62. [PubMed: 14699488]
40. Brandt LJ, Prather CM, Quigley EM, et al. Systematic review on the management of chronic constipation in north america. *Am J Gastroenterol*. 2005; 100(Suppl 1):S5–S21. [PubMed: 16008641]
41. Ramkumar D, Rao SS. Efficacy and safety of traditional medical therapies for chronic constipation: Systematic review. *Am J Gastroenterol*. 2005 Apr; 100(4):936–71. [PubMed: 15784043]
42. Dipalma JA, Cleveland MV, McGowan J, et al. A randomized, multicenter, placebo-controlled trial of polyethylene glycol laxative for chronic treatment of chronic constipation. *Am J Gastroenterol*. 2007 Jul; 102(7):1436–41. [PubMed: 17403074]
43. Johanson JF, Morton D, Geenen J, et al. Multicenter, 4-week, double-blind, randomized, placebo-controlled trial of lubiprostone, a locally-acting type-2 chloride channel activator, in patients with chronic constipation. *Am J Gastroenterol*. 2008 Jan; 103(1):170–7. [PubMed: 17916109]

44. Tuteja AK, Rao SS. Lubiprostone for constipation and irritable bowel syndrome with constipation. *Expert Rev Gastroenterol Hepatol*. 2008 Dec; 2(6):727–33. [PubMed: 19090733]
- 45**. Rao SS, Seaton K, Miller M, et al. Randomized controlled trial of biofeedback, sham feedback, and standard therapy for dyssynergic defecation. *Clin Gastroenterol Hepatol*. 2007 Mar; 5(3): 331–8. This is one of 4 randomized controlled trials that proved the efficacy of biofeedback in the treatment of dyssynergic defecation. [PubMed: 17368232]
46. Rao SS. The technical aspects of biofeedback therapy for defecation disorders. *Gastroenterologist*. 1998 Jun; 6(2):96–103. [PubMed: 9660527]
47. Rao SS, Welcher KD, Pelsang RE. Effects of biofeedback therapy on anorectal function in obstructive defecation. *Dig Dis Sci*. 1997 Nov; 42(11):2197–205. [PubMed: 9398795]
- 48**. Chiarioni G, Whitehead WE, Pezza V, et al. Biofeedback is superior to laxatives for normal transit constipation due to pelvic floor dyssynergia. *Gastroenterology*. 2006 Mar; 130(3):657–64. [PubMed: 16530506]
- 49**. Chiarioni G, Salandini L, Whitehead WE. Biofeedback benefits only patients with outlet dysfunction, not patients with isolated slow transit constipation. *Gastroenterology*. 2005 Jul; 129(1):86–97. [PubMed: 16012938]
- 50**. Heymen S, Scarlett Y, Jones K, et al. Randomized, controlled trial shows biofeedback to be superior to alternative treatments for patients with pelvic floor dyssynergia-type constipation. *Dis Colon Rectum*. 2007 Apr; 50(4):428–41. References 48–50 were randomized controlled trials that supported the use of biofeedback therapy in dyssynergic defecation. [PubMed: 17294322]

Figure 1. Images on defecography at rest (A) and during straining (B); Figure A reveals an anorectal angle of 123 degrees and the anus at rest. Figure B shows a perineal descent of 5.2 cm from the pubococcygeal line (white arrow), showing descending perineum syndrome (DPS) together with an anterior rectocele (black arrow).

Figure 2.
Rectal and anal pressure changes during anorectal manometry in a subject with dyssynergic defecation before (A) and after (B) biofeedback therapy

Table 1

Summary of the randomized controlled trials of biofeedback therapy for dyssynergic defecation

	Chiarioni et al (48)	Rao et al (45)	Chiarioni et al (49)	Heymen et al (50)
Trial Design	Biofeedback versus polyethylene glycol, 14.6 g	Biofeedback versus standard versus sham biofeedback	Biofeedback for slow transit versus dyssynergia	Biofeedback versus diazepam, 5 mg, versus placebo
Subjects and randomization	104 women 54 biofeedback 55 polyethylene glycol	77 (69 women) 1:1:1 distribution	52 (49 women) 34 dyssynergia 12 slow transit 6 mixed	84 (71 women) 30 biofeedback 30 diazepam 24 placebo
Duration and number of biofeedback sessions	3 months and 1 year, 5 weekly, 30-minute training sessions performed by physician investigator	3 months, biweekly, 1 hour, maximum of six sessions over 3 months, performed by biofeedback nurse therapist	Five weekly 30- minute training sessions, performed by physician investigator	Six biweekly, 1- hour sessions
Primary outcomes	Global improvement of symptoms Worse=0 No improvement=1 Mild=2 Fair=3 Major improvement=4	<ol style="list-style-type: none"> 1 Presence of dyssynergia 2 Balloon expulsion time 3 Number of complete spontaneous bowel movements (CSBM) 4 Global satisfaction 	Symptom improvement None=1 Mild=2 Fair=3 Major=4	Global symptom relief
Dyssynergia corrected or symptoms improved	79.6% reported major improvement at 6 and 12 months, 81.5% reported major improvement at 24 months	Dyssynergia corrected at 3 months in 79% with biofeedback versus 4% sham and 6% in standard group; CSBM=biofeedback group versus sham or standard, $p<0.05$	71% with dyssynergia and 8% with slow transit alone reported fair improvement in symptoms	70% improved with biofeedback compared with 38% with placebo and 30% with diazepam
Conclusions	Biofeedback was superior to laxatives	Biofeedback was superior to sham feedback and standard therapy	Biofeedback benefits dyssynergia and not slow transit constipation	Biofeedback is superior to placebo and diazepam