Published in final edited form as:

Org Lett. 2013 June 21; 15(12): 3134–3137. doi:10.1021/ol401337p.

Synthesis of Quaternary α -Methyl α -Amino Acids by Asymmetric Alkylation of Pseudoephedrine Alaninamide Pivaldimine

Cedric L. Hugelshofer[†], Kevin T. Mellem, and Andrew G. Myers^{*}

Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA 02138

Abstract

The utility of pseudoephedrine as a chiral auxiliary for the alkylative construction of quaternary α -methyl α -amino acids is demonstrated. The method is notable for the high diastereoselectivities of the alkylation reactions, for its versatility with respect to electrophilic substrate partners, and for its mild hydrolysis conditions, which provide α -amino acids without salt contaminants. Alternatively, α -amino esters can be obtained by direct alcoholysis.

(1*S*,2*S*)-Pseudoephedrine (*R*)-alaninamide pivaldimine (**1**) or its enantiomer serve as substrates in a new method for the alkylative construction of quaternary α -methyl α -amino acids. These substrates can be prepared in high yield by coupling of the appropriate stereoisomers of pseudoephedrine¹ and *N*-Boc alanine by the mixed anhydride method (pivaloyl chloride)² followed by *N*-Boc deprotection (HCl) and *tert*-butylimine formation (see Supporting Information). Two methods were developed to form the *N*-*tert*-butyl imine derivatives cleanly and in quantitative yield, which was essential to achieve high yields in the subsequent alkylation reactions. The first method involved adding pivaldehyde (2.0 equiv) to a stirring suspension of pseudoephedrine alaninamide (1 equiv) and activated 4Å MS in a mixed solvent of benzene and dichloromethane at 23 °C. Evaporation of the solvents after 50 min afforded a white solid, which was held under vacuum (<1 Torr) at 35 °C overnight to remove excess pivaldehyde. The product (99% yield, est. 95% purity by ¹H and ¹³C NMR) was used without further purification. A second successful protocol involved initial synthesis of pivaldehyde *N*-propyl imine as a reagent for transimination, a more facile and rapid process than imine formation from the corresponding aldehyde.³ A mixture of pivaldehyde *N*-propyl imine (5.0 equiv) and pseudoephedrine alaninamide (1 equiv) was stirred in dry benzene at 23 °C under moderate vacuum (200 mmHg) for 30 min, during which time gas was observed to evolve from the reaction mixture (presumably *N*-propylamine). Concentration afforded a white solid, which was held under vacuum (<1 Torr) at 35 °C to remove all traces of the transimination reagent. The product, obtained in 99% yield (est. 95% purity by ¹H and ¹³C NMR), was used without further purification in subsequent alkylation reactions. These methods were also effective for the preparation of (1*S*,2*S*)-pseudoephedrine (*S*)-alaninamide pivaldimine and its enantiomer, which proved

^{*}myers@chemistry.harvard.edu.

[†]Current address: Department of Chemistry, Ludwig-Maximilians-Universität München, Butenandtstrasse 5-13, 81377 München, Germany.

Supporting Information Available Full experimental procedures, characterization data, and ¹H and ¹³C NMR spectra for all synthesized compounds. This material is available free of charge via the Internet at <http://pubs.acs.org>.

to be mismatched substrates in pivaldimine alkylations (vide infra). In contrast to these successful methods for imine formation, attempts to form the *N*-*tert*-butyl imine by combining the alaninamide substrate with pivaldehyde and magnesium sulfate, or with pivaldehyde, magnesium perchlorate, and trimethylorthorformate (catalyst and stoichiometric dehydrating agent, respectively) were not satisfactory with respect to conversion and product purity, and efforts to purify samples of the imine only led to increased contamination with its hydrolysis product.

Enolization-alkylation of substrate **1** was optimally achieved by the following protocol. A solution of (1*S*,2*S*)-pseudoephedrine (*R*)-alaninamide pivaldimine (1 equiv) in dry tetrahydrofuran (THF) was transferred to a flask containing flame-dried lithium chloride (6.0 equiv), and the resulting slurry was cooled to -78°C . A solution of lithium diisopropyl amide (LDA) in THF (2.2 equiv) was then added slowly down the side of the flask by cannula or syringe so as to allow the solution of base to cool before reaching the substrate solution. After completed addition and further stirring at -78°C for 5 minutes, the reaction flask was transferred to an ice bath for 10 minutes before cooling to -50°C . An electrophile (2.5 equiv) was then added to the cold reaction solution, and the ensuing alkylation reaction was monitored by TLC (reaction times typically ranged from 1.5–3.5 h). Upon completed reaction, a solution of 1 *N* hydrochloric acid was added to the reaction mixture to induce hydrolysis of the *tert*-butyl imine function within the alkylated product, which typically occurred in less than 3 h at 23°C . Table 1 summarizes results from alkylation reactions using six different electrophiles. In all cases, diastereoselectivities equaled or exceeded 19:1, and the products, isolated in 83–95% yield by flash-column chromatography, were solids. We established that the benzylation product of entry 1 had the configuration depicted by comparison with a sample of known configuration, prepared by an independent route (see Supporting Information). The diastereoisomer that is formed arises from replacement of the α -CH bond by α -C-benzyl with retention of configuration. This alkylation product and two others whose stereochemistry was established unambiguously (shown in equation 2 of Scheme 1 and in Scheme 2 below) were found to form a homochiral series. The products of entries 2–6 of Table 1 were presumed to have formed analogously.

Table 2 summarizes results from three parallel alkylation reactions using the diastereomeric substrate (1*S*,2*S*)-pseudoephedrine (*S*)-alaninamide (**2**), otherwise conducted as described in the paragraph above. Surprisingly, in all three cases the major product was the same as that formed using substrate **1**, although the stereoselectivities and yields were lower, making it clear that substrate **2** is mismatched.⁴

These findings can be rationalized by arguments that extend from our earlier studies of the enolization of α,α -dialkyl pseudoephedrine and pseudoephedrine amide enolates, summarized in Figure 1.⁵ Briefly, both matched and mismatched substrates are proposed to form the same *E*-enolate intermediate (with the enoxy and α -imino groups in *trans* disposition), which then undergoes alkylation predominantly or exclusively in the usual sense.⁶ Enolization of the mismatched substrate is believed to be less *E*-selective, however, because *E*-enolization requires approach of the base along a trajectory impeded by the auxiliary. Interestingly, if we are correct in this proposal, then formation of the *Z*-enolate from the mismatched substrate must remain a higher energy pathway in spite of the fact that it would arise from deprotonation along a more favorable trajectory. We speculate that an important factor may be a developing repulsive electronic interaction between the enolate oxygen atom and the α -imino lone pair in the transition state for formation of the *Z*-enolate.

As depicted in Scheme 1, it proved possible to assemble cyclic α -amino acid derivatives containing an α -quaternary center in a single operation using biselectrophiles such as 3-bromopropyl trifluoromethane-sulfonate (equation 1), (*R*)-3-chloro-2-methylpropyl

trifluoromethane sulfonate (equation 2) and α,α' -dibromo-*o*-xylene (equation 3). Due to their chromatographic instability (believed to be a consequence of facile $N \rightarrow O$ acyl transfer), products from the latter two alkylations were directly subjected to transacylation with lithium benzyloxide, a useful transformation we discuss in greater detail below.

As a concluding alkylation result, in Scheme 2 below we summarize a successful α -allylation of the matched substrate **1**, which required development of an alternative workup method (using hydroxylamine in lieu of acid to cleave the *tert*-butyl imine function of the alkylated product). Interestingly, hydrolysis of the imine function of the allylated product under the usual conditions (1 *N* HCl) led to a significant by-product (Scheme 3, aminal **7**, accompanied by an unidentified minor diastereomeric aminal by-product in a 7:1 ratio, respectively). Crystallization afforded a single crystal of pure **7** suitable for X-ray analysis (see Supporting Information). As depicted in Scheme 3, by-product **7** presumably arises from an aza-Cope rearrangement followed by cyclization.⁷

An exceptional and highly useful feature of the present study was the finding that α -quaternary α -amino amides of pseudoephedrine undergo hydrolysis to afford α -amino acids simply upon refluxing in aqueous dioxane (salt-free conditions, Table 3), whereas treatment with lithium alkoxides affords α -amino esters (Table 4, and Scheme 1 above). In the former case, the pseudoephedrine auxiliary can be easily recovered in high yield by a simple extractive isolation procedure, whereas in the latter it can be isolated chromatographically.

Prior auxiliary-based methods for α -alkylation of alanine derivatives have generally achieved stereochemical control of both the enolate geometry and the nascent quaternary carbon center by incorporating the alanine substrate within a rigid heterocyclic framework, and liberation of the α -amino acid typically requires harsh conditions, in some cases resulting in destruction of the auxiliary.⁸ The present work differs in these respects. Advances in asymmetric phase-transfer catalysis have also achieved highly enantioselective alkylations of alanine derivatives.⁹ Determination of the most appropriate methodology for a given specific application will be context-dependent, but we believe that the present work offers a potentially useful new option for the stereodefined construction of α -methyl α -amino acids.^{10,11,12}

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge the NSF (CHE-1152205) and NIH (CA-047148) for financial support of this research. We also wish to express our sincere appreciation to Dr. Shao-Liang Zheng of Harvard University for X-ray crystallographic analyses.

References

1. Morales MR, Mellem KT, Myers AG. *Angew. Chem. Int. Ed.* 2012; 51:4568–4571.
2. Vaughan JR Jr, Osato RL. *J. Am. Chem. Soc.* 1951; 73:5553–5555.
- 3 (a). Smith GEP Jr, Bergstrom FW. *J. Am. Chem. Soc.* 1934; 56:2095–2098.(b) Cordes EH, Jencks WP. *J. Am. Chem. Soc.* 1961; 84:826–831.(c) Leach BE, Leussing DL. *J. Am. Chem. Soc.* 1971; 93:3377–3384.
4. Ko SY, Lee AWM, Masamune S, Reed LA III, Sharpless KB, Walker FJ. *Tetrahedron.* 1990; 46:245–264.

5. Kummer DA, Chain WJ, Morales MR, Quiroga O, Myers AG. *J. Am. Chem. Soc.* 2008; 130:13231–13233. [PubMed: 18788739]
6. The *E*-enolate obtained from the matched diastereomer could be trapped in the form of a cyclic siloxane whose stereochemistry was determined by NOE experiments. See Supporting Information for details.
7. This interesting transformation suggests a possible new approach to transfer aminoallylation. See, for example: Sugiura M, Mori C, Kobayashi SJ. *Am. Chem. Soc.* 2006; 128:11038–11039.
8. For selected references describing chiral auxiliary-based methods for the asymmetric synthesis of quaternary α -methyl α -amino acids, see: Schöllkopf U, Hausberg HH, Hoppe I, Segal M, Reiter U. *Angew. Chem. Int. Ed.* 1978; 17:117–119. Seebach D, Aebi JD, Naef R, Weber T. *Helv. Chim. Acta.* 1985; 68:144–154. Williams RM, Im J. *J. Am. Chem. Soc.* 1991; 113:9276–9286. Berkowitz DB, Smith MK. *J. Org. Chem.* 1995; 60:1233–1238. Alonso F, Davies SG, Elend AS, Haggitt JL. *J. Chem. Soc., Perkin Trans.* 1998; 1:257–264. Chinchilla R, Galindo N, Nájera C. *Synthesis.* 1999:704–717. Lu T, Lin CJ. *Org. Chem.* 2011; 76:1621–1633.
9. For leading references, see: Kano T, Sakamoto R, Mii H, Wang Y, Maruoka K. *Tetrahedron.* 2010; 66:4900–4904. Jew S, Jeong B, Lee J, Yoo M, Lee Y, Park B, Kim MG, Park H. *J. Org. Chem.* 2003; 68:4514–4516. [PubMed: 12762758] Ooi T, Takeuchi M, Kameda M, Maruoka K. *J. Am. Chem. Soc.* 2000; 122:5228–5229.
10. For recent reviews of the asymmetric synthesis of quaternary α -amino acids, see: Cativiela C, Díaz-de-Villegas M. *Tetrahedron: Asymmetry.* 2007; 18:569–623. Cativiela C, Ordoñez M. *Tetrahedron: Asymmetry.* 2009; 20:1–63. [PubMed: 20300486]
11. For selected references describing other methods for the asymmetric synthesis of quaternary α -methyl α -amino acids, see: Trost BM, Dogra K. *J. Am. Chem. Soc.* 2002; 124:7256–7257. [PubMed: 12071719] Vachal P, Jacobsen EN. *J. Am. Chem. Soc.* 2002; 124:10012–10014. [PubMed: 12188665] Masumoto S, Usuda H, Suzuki M, Kanai M, Shibasaki M. *J. Am. Chem. Soc.* 2003; 125:5634–5635. [PubMed: 12733893] Smith ND, Wohlrab AM, Goodman M. *Org. Lett.* 2005; 7:255–258. [PubMed: 15646971]
12. Efforts to extend the present method to alkylate higher α -alkyl pivaldimines (replacing α -methyl with α -benzyl or α -ethyl) were not successful.

Figure 1.

Proposed rationale for the stereochemical outcome of alkylations of matched and mismatched diastereomers of pseudoephedrine alaninamide pivaldimine.

Scheme 1. Synthesis of cyclic amino acid derivatives.^a^a Diastereomeric ratios were determined by ¹H NMR analysis of the crude alkylation reaction mixtures.

Scheme 2. Optimized synthesis of α-allyl quaternary amide **6**.

Scheme 3.
Aza-Cope rearrangement of an α -allylation product.

Table 1Alkylation of (1*S*,2*S*)-pseudoephedrine (*R*)-alaninamide – matched case.

entry ^a	RX	crude dr ^b	isolated yield (%)
1 ^c	BnBr	19:1	95
2	BrCH ₂ CO ₂ <i>t</i> Bu	19:1	95
3 ^d	EtI	19:1	91
4	<i>m</i> -CH ₃ OBnBr	19:1	89
5	<i>i</i> BuOTf	19:1	84
6	CyCH ₂ OTf	19:1	83

^a 2.2 equivalents of base and 2.5 equivalents of electrophile were used in all cases.^b Diastereomeric ratios were determined by ¹H NMR analysis of the crude reaction mixtures.^c Conducted on a 2.1-g scale.^d The alkylation reaction was conducted at 0 °C.

Table 2Alkylation of (1*S*,2*S*)-pseudoephedrine (*S*)-alaninamide – mismatched case.

entry ^a	RX	crude dr ^b	isolated yield (%)
1	BnBr	8:1	69
2	BrCH ₂ CO ₂ tBu	19:1	60
3 ^c	EtI	13:1	72

^a2.2 equivalents of base and 2.5 equivalents of electrophile were used in all cases.^bDiastereomeric ratios were determined by ¹H NMR analysis of the crude reaction mixtures.^cThe alkylation reaction was conducted at 0 °C.

Table 3

Hydrolysis of pseudoephedrine amides to form quaternary α -methyl α -amino acids.

entry	R	isolated yield, α -amino acid (%)	isolated yield, auxiliary (%)
1	Bn	99 ^a	99
2	Allyl	97 ^a	94
3	CyCH ₂ -	99	98

^a>98% ee, as determined by chiral HPLC analysis.

Table 4

Synthesis of quaternary α -methyl α -amino esters.

entry	ROH	base	isolated yield, α -amino ester (%)	isolated yield, auxiliary (%)
1	CH ₃ O H	NaOCH 3	88	91
2	BnOH	nBuLi	84	83