Published in final edited form as:

J Rheumatol. 2013 May ; 40(5): . doi:10.3899/jrheum.121075.

Trends in Serious Infections in Rheumatoid Arthritis

Orla Ni Mhuircheartaigh, MB BCH¹ [Resident (Internal Medicine)], Eric L. Matteson, MD² [Professor of Medicine], Abigail B. Green, BA³, and Cynthia S. Crowson, MS⁴
Division of Rheumatology Mayo Clinic College of Medicine 200 1st St Rochester, MN, 55905, USA

¹ Department of Internal Medicine, Mayo Clinic College of Medicine, Rochester, MN, USA

² Division of Rheumatology, Mayo Clinic College of Medicine, Rochester, MN, USA

³ Department of Health Sciences Research, Mayo Clinic College of Medicine, Rochester, MN, USA

⁴ Department of Health Science Research and Division of Rheumatology, Mayo Clinic College of Medicine, Rochester, MN, USA

Abstract

Objective—To examine trends in the rates of serious infections among patients diagnosed with rheumatoid arthritis (RA) in 1995-2007 compared to rates previously reported from the same geographical area diagnosed in 1955-1994.

Methods—A population-based inception cohort of patients with RA in 1995-2007 was assembled and followed through their complete medical records until death, migration, or December 31, 2008. All serious infections (requiring hospitalization or IV antibiotics) were recorded. Person-year (py) methods were used to compare rates of infection.

Results—Among 464 patients with incident RA in 1995-2007, 54 had 1 serious infection (178 total). These were compared to 609 patients with incident RA in 1955-1994 (290 experienced 1 serious infection; 740 total). The rate of serious infections declined from 9.6 per 100py in the 1955-1994 cohort to 6.6 per 100py in the 1995-2007 cohort. However, serious gastrointestinal infection rates increased from 0.5 per 100py in the 1955-1994 cohort to 1.25 per 100py in the 1995-2007 cohort. Among patients with a history of serious infection, the rate of subsequent infection increased from 16.5 per 100py in 1955-1994 to 37.4 per 100py in 1995-2007. There was an increase in the rate of serious infections in patients who received biologic agents, but this did not reach significance.

Conclusions—Aside from gastrointestinal infections, the rate of serious infections in patients with RA has declined in recent years. However, the rate of subsequent infections was higher in recent years than previously reported.

Keywords

Rheumatoid arthritis; infection; biologic agents

Introduction

Patients with rheumatoid arthritis [RA] have increased susceptibility for infections (1). The reasons for the increased infection risk in patients with RA are multifactorial. Probable causes include the underlying immunologic disturbance associated with the disease process, immunosuppressive therapy used for treatment of RA, and other co-existing risk factors for infection which may be more common in patients with RA (2). The infection risk concern has been heightened in recent years as clinicians have struggled to evaluate the possible impact of the introduction of biologic agents for the treatment of many rheumatological conditions (3-6). In addition, the approach to the management of RA has evolved toward more aggressive therapy in order to prevent disease progression and complications. It is unclear whether these trends have had an impact on infection risk.

This has led to recent publications, by us and others, examining the risk factors for infection in patients with RA and providing scoring systems to evaluate the risk of infection in these patients (7-9). However, recent trends in the rates and types of infection among patients with RA have not been evaluated. The objective of this study was to compare the rates and types of serious infection among patients diagnosed with RA in 1995-2007 to rates previously reported among patients with RA from the same geographical area who were diagnosed in 1955-1994.

Materials and Methods

Study population

We performed a retrospective longitudinal cohort study comparing infection rates in Olmsted County, MN, residents age 18 years old with incident RA in 1995-2007 with our previous cohort of Rochester, MN residents with incident RA in 1955-1994(1). Patients in both cohorts fulfilled 1987 American College of Rheumatology (ACR) criteria for RA (10). Identification of patients with incident RA has been described previously (11). All study subjects were followed up through their entire (inpatient and outpatient) medical record, until death, migration from the county, or the date of study end (December 31, 1999 for the 1955-1994 cohort and December 31, 2008 for the 1995-2007 cohort).

These RA cases were identified using the data resources of the Rochester Epidemiology Project (REP), a diagnostic indexing and medical records linkage system that affords access to medical records from all sources of care for community residents (12). This study was approved by the institutional review boards of the Mayo Clinic and the Olmsted Medical Center.

Data collection

Data on all episodes of serious infection requiring hospital admission or IV antibiotics occurring after the RA incidence date were collected according to a prespecified and pretested detailed protocol. Data for the 1995-2007 cohort were collected by an abstractor who collected data for the 1955-1994 cohort.

The operational definitions for each infection type were as follows: bacteremia/septicemia, isolation of a pathogenic microorganism from one or more blood cultures, with fever ($>38^{\circ}\text{C}$); septic arthritis, positive microbiologic culture from joint aspirate fluid in the presence of suggestive clinical features; urinary tract infection, including pyelonephritis and urosepsis, isolation of $>100,000$ colony-forming units/ml of urine in the presence of suggestive clinical features; pneumonia, presence of new infiltrates, consolidation, or effusion seen by chest radiography and suggestive clinical features; osteomyelitis, clinical suspicion with confirmation by definite radiologic findings or positive bone culture. Lower

respiratory tract infections, skin and soft tissue infections, and acute gastrointestinal infections could be included on the basis of a physician's diagnosis and relevant clinical findings alone, but microbiologic culture results were recorded if available. Skin and soft tissue infections included cellulitis, abscesses, wound infections, herpes zoster, and diabetic foot infections. Gastrointestinal infections included gastroenteritis, diverticulitis, infective colitis, *Clostridium difficile*, and vancomycin-resistant enterococci. Opportunistic infections included cytomegalovirus, cryptococcus, mycobacterium tuberculosis, nontuberculosis mycobacterium, vancomycin-resistant enterococci, histoplasmosis, blastomycosis, coccidioidomycosis, cryptococcosis, endemic mycosis, nocardiosis/actinomycosis, listeriosis, toxoplasmosis, pneumocystis, legionellosis, salmonellosis, aspergillosis, candidemia, progressive multifocal leukoencephalopathy and optic neuritis. Intra-abdominal infections could be included on the basis of clinical findings alone, and comprised acute cholecystitis, ascending cholangitis, suppurative appendicitis, and peritonitis. The category "other infections" included episodes of otitis media and sinusitis that required hospitalization, eye infections, male and female genital tract infections, and acute hepatitis. Data on urinary tract infections (other than those classified as urosepsis/acute pyelonephritis), was not recorded. Patients who fulfilled criteria for more than 1 infection simultaneously were classified in both categories, except in the case of septicemia, which was classified in a single category referred to as septicemia with a notation of the accompanying infectious condition (e.g., pneumonia with septicemia, urinary tract infection with septicemia).

Information on potential confounding factors for infection (rheumatoid factor positivity, diabetes mellitus, leukopenia, smoking status, alcoholism, chronic lung disease, cancer and extra-articular manifestations of RA (ExRA)), along with dates of onset, was ascertained. Leukopenia was defined as white blood cell counts < 4000/ml on 2 or more occasions. Manifestations of severe ExRA included pericarditis, pleuritis, Felty's syndrome, glomerulonephritis, vasculitis, peripheral neuropathy, scleritis, and episcleritis (13, 14). Data were also collected on start and stop dates of medication used at any point during follow up, including commonly used disease-modifying antirheumatic drugs (DMARDs) such as methotrexate, hydroxychloroquine, other DMARDs (gold, sulfasalazine, azathioprine, cyclophosphamide, cyclosporine, D-penicillamine or leflunomide), biologic agents and corticosteroids.

Data on vital status was also collected. Case fatality was defined as a death within 30 days of a serious infection.

Data analysis

Baseline characteristics of the study population were summarized using descriptive statistics. Cumulative incidence adjusted for the competing risk of death was used to compare the occurrence of characteristics between cohorts appropriately accounting for differences in length of follow-up (15). Cumulative incidence rates were compared using the methods of Gray (16). Incidence rates for infections were calculated by dividing the total number of events by the number of person-years of follow-up. Rate ratios (RR) were obtained by dividing infection incidence rates in RA patients in the 1995-2007 cohort by those in patients with RA in the 1955-1994 cohort, and 95% confidence intervals (CI) for these RR were calculated. Due to the differential length of follow-up in the 2 time periods, cumulative incidence rates were also computed for infections to provide comparison of infection rates at the same length of follow-up in the two time periods.

Comparisons of infection rates between time periods were also performed after adjusting for infection risk factors. Adjustment for risk factors was performed by first calculating the infection risk score we previously developed at the RA diagnosis and at the beginning of

each subsequent year of follow-up for each patient (7). The Anderson-Gill adaptation of the Cox model allowing inclusion of multiple events in the same patient was used to compare the rate of development of serious infections between the time periods after adjusting for the risk score (17). The risk score adjustment was performed using a time-dependent covariate to represent the risk score, which changed at the beginning of each yearly interval throughout follow-up.

The rate of infections during biologic treatment was calculated as the number of infections that occurred between the start and stop dates of biologic treatment divided by the length of follow-up from the start to the stop of biologic treatment. Infection rates in patients with biologic exposure included the infections and follow-up after the stop of biologic treatment in patients who were exposed to biologics.

Results

The cohort of patients with incident RA in 1995-2007 was comprised of 464 patients. These patients were compared with 609 patients with incident RA in 1955-1994. The mean age at RA incidence for the early cohort was 58.0 years (73% female) and 56.0 years (69% female) for the later cohort (Table 1). The mean follow-up time was 12.7 years for the early cohort and 5.9 years for the later cohort, corresponding to 7,730 total person-years (py) and 2,715 py, respectively. Rates of smoking and comorbidities were similar in the two time periods, except for diabetes mellitus, which occurred more frequently among patients in the 1995-2007 cohort compared to the 1955-1994 cohort. The risk score for serious infections was significantly higher among patients with incident RA in 1995-2007 compared to those with incident RA in 1955-1994 ($p=0.015$).

More patients in the 1995-2007 cohort received the non-biological DMARDs methotrexate and hydroxychloroquine, biologic agents, and more patients were exposed to corticosteroids compared with patients in the 1955-1994 cohort. However, patients in the 1995-2007 cohort were less likely to receive other non-biologic DMARDs than patients in the 1955-1994 cohort.

In the 1995-2007 cohort, 54 patients had 1 serious infection (178 total infections) and in the 1955-1994 cohort, 290 patients experienced 1 serious infection (740 total infections). The rate of all serious infections in patients in the 1995-2007 cohort was 6.6 per 100 py, which was less than the rate of 9.6 per 100 py that was seen in patients in the 1955-1994 cohort (RR: 0.69, 95% CI: 0.58, 0.80; Table 2). The rate of first infection was also lower in the 1995-2007 cohort (2.0 per 100 py) compared to the 1955-1984 cohort (3.8 per 100 py). Similar differences in rates were noted when comparing the cumulative incidence of first serious infection at 10 years after RA (15.6%; 95% confidence interval [CI]: 11.2 - 20.0% for 1995-2007 vs. 34.5%; 95% CI: 30.4 - 38.6% for 1955-1994). Among patients with a serious infection, the risk scores for serious infections for the beginning of the year of follow-up when the first serious infection occurred were somewhat higher among patients in the 1995-2007 cohort (median: 8.4%; 25th percentile, 75th percentile: 3.8%, 36.0%) compared to the 1955-1994 cohort (median: 6.5%; 25th percentile, 75th percentile: 3.0%, 23.8%; $p=0.059$). Adjustment for the serious infection risk score had no effect on the difference in infection rates between the cohorts.

Bacteremia/septicemia, pneumonias and skin/ soft tissue infections were the most common types of infections in both time periods. Bacteremia/septicemia rates decreased from 0.78 infections per 100 py in the 1955-1994 cohort to 0.52 infections per 100 py in the 1995-2007 cohort (RR: 0.68, 95% CI: 0.36, 1.16). The rate of serious pneumonia infections decreased from 3.10 to 1.99 infections per 100 py (RR: 0.65, 95% CI: 0.47, 0.85). Skin/ soft tissue

infections decreased from 2.37 to 1.51 infections per 100 py (RR: 0.64, 95% CI: 0.45, 0.89). Septic arthritis, osteomyelitis, lower respiratory infections, urosepsis/pyelonephritis, intra-abdominal infections and other infections all followed a similar decline in infection rates between the two time periods.

In contrast, the rates of serious gastrointestinal infections increased between the two groups from an infection rate of 0.49 to 1.25 infections per 100 py, respectively (RR: 2.55, 95% CI: 1.60, 4.04). The rates of opportunistic infections increased significantly (RR: 9.35, 95% CI: 3.22, 41.4), as did the rates of clostridium difficile infections (RR: 12.34; 95% CI 2.84, 176.5). The only opportunistic infections that occurred in our cohort were clostridium difficile and vancomycin-resistant enterococci.

The overall rate of all subsequent infections in patients who developed at least one previous serious infection requiring hospitalization was increased from 26.3 to 65.1 infections per 100 py, (RR: 2.48, 95% CI: 2.02, 3.01). Similarly, the rate of second serious infection among patients with a first serious infection was increased from 16.5 per 100 py in 1955-1994 to 37.4 per 100 py in 1995-2007. Note that the rate of all subsequent infections (i.e., not including the initial infection) declined when estimated among all RA patients, not just those with an initial infection (5.8 vs. 4.6 infections per 100 py; RR: 0.79; 95% CI: 0.64, 0.95).

In the 1995-2007 cohort, 96 patients received biologic agents at some point; approximately 95% were TNF inhibitors, so analyses of subtypes of biologic agents were not performed. The rate of infections during treatment with biologic agents was 8.2 infections per 100 py (95% CI: 4.9, 12.8 per 100 py) compared with 6.4 infections per 100 py (95% CI: 5.4, 7.5 per 100 py) for those not on biologic agents (RR: 1.31; 95% CI 0.78-2.01). Similarly, the rate of infections for patients who were ever exposed to biologic agents (during or after use of biologic agents) was 7.9 infections per 100 py (95% CI: 5.3, 11.1 per 100 py) compared with 6.4 infections per 100 py (95% CI: 5.4, 7.5 per 100 py) for those who had never used (or prior to use of) biologic agents (RR: 1.24; 95% CI 0.82-1.79).

Case fatalities occurred in 90 (12%) of the 740 infections in the 1955-1994 cohort and 7 (4%) of the 178 infections in the 1995-2007 cohort, indicating a substantial decrease in case fatality rates ($p < 0.001$).

Discussion

The overall rate of serious infections in our population has declined among patients diagnosed with RA in more recent years, while the rate of subsequent serious infections in those patients with a history of at least one serious infection has increased. The principal sites of infections remained consistent between the two cohorts; however, there were significantly increased rates of gastrointestinal infections and opportunistic infections, predominately clostridium difficile in the 1995-2007 cohort. We noted a similar burden of comorbidities related to infection risk in the 2 cohorts, except for an increase in diabetes mellitus among patients in the 1995-2007 cohort compared to the 1955-1994 cohort. More patients in the 1995-2007 cohort were exposed to anti-rheumatic medications including certain DMARDS (such as methotrexate and hydroxychloroquine), biologic agents and corticosteroids. A possible increased rate of serious infections was observed in patients who received biologic agents; however, due to the limited number of patients on biologic agents in this study, a definitive conclusion about the use of biologic agents and the rate of serious infections could not be made.

Patients with RA are known to be at increased susceptibility for infections (1, 18). Active inflammatory disease may confer a higher risk of infection (19). Alterations in the cellular immune system, including alterations in the RA related T-cell functions likely also

contribute to the infection risk (18, 20, 21). Other risk factors for infection include advancing age, leukopenia and comorbidities (chronic lung disease, alcoholism, dementia/Alzheimer's disease and diabetes mellitus) (2, 7). Treatment options for RA can also place patients at increased risk for serious infections. These include corticosteroids, DMARDs and biologic agents (2, 22-24). As well, severity indices of RA such as increased sedimentation rate, ExRA and rheumatoid factor positivity are all predictive of the development of serious infections in these patients (2).

In the past decade, more aggressive use of conventional DMARDs as well as the introduction of biologic agents have contributed to better control of RA activity. The decrease in the number of serious infections in patients with RA may potentially be due to decreased inflammation. While we did not directly assess disease activity, we noted that more of the patients who were diagnosed recently had received corticosteroids (90% vs 46%), methotrexate, hydroxychloroquine and biologic agents compared to the patients diagnosed earlier. This more aggressive treatment is likely associated with a lessened inflammatory burden, but additional associated potential safety risks of such agents.

It is unclear why the rate in gastrointestinal infections and particularly clostridium difficile infection has risen in more recently diagnosed patients. The rate of clostridium difficile infection in the general community has increased in recent years, likely due to a combination of factors (25). Therefore, the increased risk of this infection in patients with RA may simply reflect the overall increase in this infection in the general population. There is some evidence of an increased risk of clostridium difficile in patients on biologic agents (26, 27); however, further studies are needed to clarify this.

Although the rates of subsequent serious infections in patients who have had a previous serious infection requiring hospitalization was increased, we speculate this was likely a result of overall declines in hospitalization rates (28). For instance, some of the infections that required hospitalization in the 1960s would not require hospitalization in the 1990s as these infections can now be treated outside the hospital due to the development of more potent antibiotics and more accessible diagnostic tests. Therefore, patients in the later cohort who require hospitalization for an initial infection are fewer but generally of poorer health than those who have been managed in the community, and are hence more likely to develop a subsequent serious infection requiring hospitalization. For this reason, although the rate of subsequent infection appears to be increasing in the subset of patients who had an initial infection, it has actually declined overall. However, the decrease in the overall rate of subsequent infections was smaller than the decrease in the rate of initial infections, which resulted in an apparent increase in the rate of subsequent infections when estimated using the subset of patients who experienced an initial serious infection.

Strengths of this study include the longitudinal population-based study design with extensive follow-up and the use of complete in-patient and out-patient medical records, providing complete ascertainment of study outcomes for all study subjects. In addition, the data for both time periods was collected using the same criteria, as well as the same abstractors ensuring comparable data between the two cohorts of patients.

As with all studies, our results need to be interpreted in light of potential limitations. Some ethnic groups are underrepresented in the catchment area of Olmsted County, Minnesota, but results are generally reflective of the US population (5). The impact of disease activity on infection risk was not assessed, as disease activity scores (e.g., DAS28) were not available. In addition, the follow-up time in the 1995-2007 cohort was substantially shorter than in the 1955-1994 cohort, which might influence the rates of infection. However, comparisons of cumulative incidence at a specific RA duration yielded similar results. In

addition, disease duration was not associated with infection rates in our cohort, so the differences in length of follow-up between cohorts were unlikely to influence our findings. Finally, the rates of serious infections over time are influenced by secular trends in the threshold for hospitalization, the use of diagnostic tools, the prevalence of antibiotic-resistant strains, the use of influenza and pneumococcal vaccines and other factors, which could not be accounted for in this observational study.

In conclusion, there has been a decline in the rates of serious infections in patients with RA diagnosed in recent years, despite some increases in comorbidities which could predispose to infection. This decline is most likely due to declining hospitalization rates in the general population over this time period, but better control of the inflammation associated with RA is likely to also play a role. The rate of gastrointestinal infections such as *Clostridium difficile* has risen in recent years, probably related to the increase in this infection in the general population. Further studies to assess the relationship between the severity of inflammation and its impact on serious infections in RA patients would be beneficial.

Acknowledgments

Funding:

This study was funded by a grant from Genentech, Inc. and made possible by grants from the National Institutes of Health (NIAMS R01 AR46849) and the Rochester Epidemiology Project (R01 AG034676 from the National Institute on Aging).

References

1. Doran M, Crowson C, Pond G, O'Fallon W, Gabriel S. Frequency of infection in patients with rheumatoid arthritis compared with controls: A population-based study. *Arthritis & Rheumatism*. 2002; 46(9):2287–93. [PubMed: 12355475]
2. Doran M, Crowson C, Pond G, O'Fallon W, Gabriel S. Predictors of infection in rheumatoid arthritis. *Arthritis & Rheumatism*. 2002; 46(9):2294–300. [PubMed: 12355476]
3. Baghai M, Osmon DR, Wolk DM, Wold LE, Haidukewych GJ, Matteson EL. Fatal sepsis in a patient with rheumatoid arthritis treated with etanercept. *Mayo Clinic Proceedings*. 2001; 76(6): 653–6. [PubMed: 11393506]
4. Phillips K, Husni ME, Karlson EW, Coblyn JS. Experience with etanercept in an academic medical center: are infection rates increased? *Arthritis Rheum*. 2002; 47(1):17–21. [PubMed: 11932873]
5. Wallis RS. Biologics and infections: lessons from tumor necrosis factor blocking agents. *Infect Dis Clin North Am*. 2011; 25(4):895–910. [PubMed: 22054762]
6. Hernandez-Cruz B, Garcia-Arias M, Ariza Ariza R, Martin Mola E. [Rituximab in rheumatoid arthritis: a systematic review of efficacy and safety]. *Reumatol Clin*. 2011; 7(5):314–22. [PubMed: 21925447]
7. Crowson CS, Hoganson DD, Fitz-Gibbon PD, Matteson EL. Development and validation of a risk score for serious infection in patients with rheumatoid arthritis. *Arthritis Rheum*. 2012; 64(9):2847–55. [PubMed: 22576809]
8. Curtis JR, Xie F, Chen L, Muntner P, Grijalva CG, Spettell C, et al. Use of a disease risk score to compare serious infections associated with anti-tumor necrosis factor therapy among high- versus lower-risk rheumatoid arthritis patients. *Arthritis Care Res (Hoboken)*. 2012; 64(10):1480–9. [PubMed: 22833479]
9. Strangfeld A, Eveslage M, Schneider M, Bergerhausen HJ, Klopsch T, Zink A, et al. Treatment benefit or survival of the fittest: what drives the time-dependent decrease in serious infection rates under TNF inhibition and what does this imply for the individual patient? *Ann Rheum Dis*. 2011; 70(11):1914–20. [PubMed: 21791449]
10. Arnett FC, Edworthy S, Bloch DA, McShane DJ, Fries JF. Subcommittee ARAC. The 1987 revised ARA criteria for rheumatoid arthritis (RA). *Arthritis & Rheumatism*. 1987; 30:S17.

11. Myasoedova E, Crowson CS, Kremers HM, Therneau TM, Gabriel SE. Is the Incidence of Rheumatoid Arthritis Rising? Results from Olmsted County, Minnesota, 1955-2007. *Arthritis Rheum.* 2010; 62(6):1576–82. PMID: PMC2929692. [PubMed: 20191579]
12. Melton LJ 3rd. History of the Rochester Epidemiology Project. *Mayo Clin Proc.* 1996; 71(3):266–74. [PubMed: 8594285]
13. Turesson C, Jacobsson L, Bergstrom U. Extra-articular rheumatoid arthritis: prevalence and mortality. *Rheumatology.* 1999; 38(7):668–74. [PubMed: 10461483]
14. Myasoedova E, Crowson CS, Turesson C, Gabriel SE, Matteson EL. Incidence of extraarticular rheumatoid arthritis in Olmsted County, Minnesota, in 1995-2007 versus 1985-1994: a population-based study. *J Rheumatol.* 2011; 38(6):983–9. [PubMed: 21459933]
15. Gooley TA, Leisenring W, Crowley J, Storer BE. Estimation of failure probabilities in the presence of competing risks: new representations of old estimators. *Statistics in Medicine.* 1999; 18(6):695–706. [PubMed: 10204198]
16. Gray R. A class of K-sample tests for comparing the cumulative incidence of a competing risk. *Annals of Statistics.* 1988; 16:1141–54.
17. Gill R. Understanding Cox's regression model. *Experientia Suppl.* 1982; 41:187–99. [PubMed: 6754425]
18. Mikuls TR. Co-morbidity in rheumatoid arthritis. *Best Pract Res Clin Rheumatol.* 2003; 17(5): 729–52. [PubMed: 12915155]
19. Baum J. Infection in rheumatoid arthritis. *Arthritis Rheum.* 1971; 14(1):135–7. [PubMed: 5542366]
20. Wagner UG, Koetz K, Weyand CM, Goronzy JJ. Perturbation of the T cell repertoire in rheumatoid arthritis. *Proc Natl Acad Sci U S A.* 1998; 95(24):14447–52. [PubMed: 9826720]
21. Koetz K, Bryl E, Spickschen K, O'Fallon WM, Goronzy JJ, Weyand CM. T cell homeostasis in patients with rheumatoid arthritis. *Proc Natl Acad Sci U S A.* 2000; 97(16):9203–8. [PubMed: 10922071]
22. Lacaille D, Guh DP, Abrahamowicz M, Anis AH, Esdaile JM. Use of nonbiologic disease-modifying antirheumatic drugs and risk of infection in patients with rheumatoid arthritis. *Arthritis Rheum.* 2008; 59(8):1074–81. [PubMed: 18668604]
23. Bongartz T, Sutton AJ, Sweeting MJ, Buchan I, Matteson EL, Montori V. Anti-TNF antibody therapy in rheumatoid arthritis and the risk of serious infections and malignancies: systematic review and meta-analysis of rare harmful effects in randomized controlled trials. *JAMA.* 2006; 295(19):2275–85. [PubMed: 16705109]
24. Furst DE. The risk of infections with biologic therapies for rheumatoid arthritis. *Semin Arthritis Rheum.* 2010; 39(5):327–46. [PubMed: 19117595]
25. Khanna S, Pardi DS, Aronson SL, Kammer PP, Orenstein R, St Sauver JL, et al. The epidemiology of community-acquired *Clostridium difficile* infection: a population-based study. *Am J Gastroenterol.* 2012; 107(1):89–95. [PubMed: 22108454]
26. Laret V, Sion C, Bataille C, Hustinx R, Reginster M. [Clinical case of the month: clostridium difficile colitis]. *Rev Med Liege.* 2006; 61(11):750–2. [PubMed: 17191741]
27. Schneeweiss S, Korzenik J, Solomon DH, Canning C, Lee J, Bressler B. Infliximab and other immunomodulating drugs in patients with inflammatory bowel disease and the risk of serious bacterial infections. *Aliment Pharmacol Ther.* 2009; 30(3):253–64. [PubMed: 19438424]
28. HCUP Reports. Healthcare Cost and Utilization Project (HCUP). Agency for Healthcare Research and Quality. Rockville, MD: Dec. 2012 www.hcup-us.ahrq.gov/reports.jsp

Table 1

Characteristics of 609 incident patients with rheumatoid arthritis (RA) diagnosed in 1955-1994 and 464 incident patients with RA diagnosed in 1995-2007 *

Variable	RA	RA
	1955 - 1994 (n=609)	1995 - 2007 (n=464)
Age, mean $\pm$ SD years	58.0 $\pm$ 15.1	55.6 $\pm$ 15.5
Female	445 (73)	320 (69)
Length of follow-up, mean $\pm$ SD years	12.7 $\pm$ 9.4	5.9 $\pm$ 3.5
Rheumatoid factor positivity	392 (65)	306 (66)
Infection risk score at RA incidence	2.6 $\pm$ 2.1	3.3 $\pm$ 3.0
Ever Smoker	317 (55)	235 (51)
Diabetes mellitus		
At RA incidence	25 (4)	48 (10)
Ever during follow-up	63 (9)	77 (21)
Chronic lung disease		
At RA incidence	67 (11)	74 (16)
Ever during follow-up	113 (20)	99 (24)
Alcoholism		
At RA incidence	13(2)	37 (8)
Ever during follow-up	42 (6)	41 (9)
Ischemic heart disease		
At RA incidence	20 (3)	25 (5)
Ever during follow-up	73 (13)	46 (14)
Cancer		
At RA incidence	24 (4)	28 (6)
Ever during follow-up	72 (13)	63 (20)
Leukopenia - ever during follow-up	102 (15)	56 (18)
Extraarticular RA * - ever during follow-up	78 (8)	22 (6)
Medication use - ever during follow-up		
Methotrexate	133 (18)	299 (73)
Hydroxychloroquine	221 (37)	297 (68)
Other non-biological DMARDs	215 (36)	104 (30)
Biologic agents	3 (0.2)	96 (29)
Corticosteroid (PO or IV)	312 (46)	376 (90)

DMARD = Disease modifying anti-rheumatic drug; n= number; SD = standard deviation

* Except where indicated otherwise, values are the number (%) of patients. Percentages for "ever during follow-up" are estimates of cumulative incidence at 10 years of follow-up.

* includes pericarditis, pleuritis, Felty's syndrome, glomerulonephritis, vasculitis, peripheral neuropathy, scleritis, and episcleritis

Table 2

All infections requiring hospitalization in 609 patients diagnosed with rheumatoid arthritis (RA) in 1955-1994 compared with 464 patients diagnosed with RA in 1995-2007

Infection Type	Patients, n		Infections, n		Incidence/100 person-years (all events/person-years)		Rate ratio [*]	95% confidence interval
	1955-1994	1995-2007	1955-1994	1995-2007	1955-1994	1995-2007		
Total	290	54	740	178	9.57	6.56	0.69	0.58, 0.80
Bacteremia/septicemia	53	10	60	14	0.78	0.52	0.68	0.36, 1.16
Septic arthritis	20	3	27	3	0.35	0.11	0.36	0.08, 0.92
Osteomyelitis	11	0	13	0	0.17	0.00	0.11	0.00, 0.59
Pneumonia	155	28	240	54	3.10	1.99	0.65	0.47, 0.85
Lower respiratory tract	57	10	89	10	1.15	0.37	0.33	0.16, 0.59
Urosepsis/pyelonephritis	27	4	35	5	0.45	0.18	0.44	0.15, 0.96
Skin/soft tissue	109	21	183	41	2.37	1.51	0.64	0.45, 0.89
Gastrointestinal infections [†]	26	13	38	34	0.49	1.25	2.55	1.60, 4.04
Intra-abdominal	25	5	26	5	0.34	0.18	0.59	0.20, 1.33
Other	24	8	29	12	0.38	0.44	1.21	0.59, 2.26
Opportunistic Infections [‡]	3	6	3	11	0.04	0.41	9.35	3.22, 41.4
Clostridium difficile	1	3	1	6	0.01	0.22	12.34	2.84, 176.5

^{**}Any urinary tract infection that required hospitalization was classified as urosepsis/pyelonephritis.

CI = confidence interval; NA = not applicable.

^{*} Obtained by dividing infection incidence rates in 1995-2007 RA patients by those in 1955-1994 RA patients.

[†] includes gastroenteritis, diverticulitis, infective colitis, clostridium difficile, and vancomycin-resistant enterococci

[‡] includes clostridium difficile and vancomycin-resistant enterococci