Published in final edited form as:

J Comp Neurol. 2013 March 1; 521(4): . doi:10.1002/cne.23204.

Global Expression Profiling of Globose Basal Cells and Neurogenic Progression Within the Olfactory Epithelium

Richard C. Krolewski^{1,2}, Adam Packard^{1,2}, and James E. Schwob^{1,*}

¹Department of Anatomy and Cellular Biology, Tufts University School of Medicine, Boston, Massachusetts 02111

²Program in Cellular, Molecular, and Developmental Biology, Sackler School of Graduate Biomedical Sciences, Tufts University School of Medicine, Boston, Massachusetts 02111

Abstract

Ongoing, lifelong neurogenesis maintains the neuronal population of the olfactory epithelium in the face of piecemeal neuronal turnover and restores it following wholesale loss. The molecular phenotypes corresponding to different stages along the progression from multipotent globose basal cell (GBC) progenitor to differentiated olfactory sensory neuron are poorly characterized. We used the transgenic expression of enhanced green fluorescent protein (eGFP) and cell surface markers to FACS-isolate $\Delta Sox2-eGFP(+)$ GBCs, *Neurog1-eGFP(+)* GBCs and immature neurons, and $\Delta OMP-eGFP(+)$ mature neurons from normal adult mice. In addition, the latter two populations were also collected 3 weeks after olfactory bulb ablation, a lesion that results in persistently elevated neurogenesis. Global profiling of mRNA from the populations indicates that all stages of neurogenesis share a cohort of >2,100 genes that are upregulated compared to sustentacular cells. A further cohort of >1,200 genes are specifically upregulated in GBCs as compared to sustentacular cells and differentiated neurons. The increased rate of neurogenesis caused by olfactory bulbectomy had little effect on the transcriptional profile of the *Neurog1-eGFP(+)* population. In contrast, the abbreviated lifespan of $\Delta OMP-eGFP(+)$ neurons born in the absence of the bulb correlated with substantial differences in gene expression as compared to the mature neurons of the normal epithelium. Detailed examination of the specific genes upregulated in the different progenitor populations revealed that the chromatin modifying complex proteins LSD1 and coREST were expressed sequentially in upstream $\Delta Sox2-eGFP(+)$ GBCs and *Neurog1-eGFP(+)* GBCs/immature neurons. The expression patterns of these proteins are dynamically regulated after activation of the epithelium by methyl bromide lesion.

Keywords

microarray; neurogenesis; regeneration; Neurog1; Sox2; LSD1; coREST

© 2012 Wiley Periodicals, Inc.

*CORRESPONDENCE TO: James E. Schwob, Department of Anatomy and Cellular Biology, Tufts University School of Medicine, 136 Harrison Ave., Boston, MA 02111. jim.schwob@tufts.edu.

Additional Supporting Information may be found in the online version of this article.

CONFLICT OF INTEREST

The authors declare that no conflict of interest exists.

ROLE OF AUTHORS

All authors had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis. Study concept and design: R.C.K., A.P., J.E.S.; Acquisition of data: R.C.K., A.P.; Analysis and interpretation of data: R.C.K., J.E.S.; Drafting of the article: R.C.K., J.E.S.; Statistical analysis: R.C.K.; Obtained funding: R.C.K., J.E.S.

The olfactory epithelium (OE) maintains a population of several million olfactory sensory neurons, each of which expresses only one of the greater than 1,000 olfactory receptor (OR) genes (Buck and Axel, 1991). Conversely, each odorant receptor is expressed by neurons within a spatially limited epithelial territory, or zone, the number of which may reach into the hundreds (Iwema et al., 2004; Miyamichi et al., 2005). Adult stem and progenitor cells are responsible for the lifelong replacement of dead or damaged neurons due to normal turnover, environmental insults, or various forms of experimental injury (Schwob, 2002). Two populations of cells within the OE: globose and horizontal basal cells (GBCs and HBCs, respectively), are neurocompetent and multipotent depending on conditions. These cells maintain the tissue throughout the lifetime of the organism and are activated robustly after various forms of injury to repopulate both the neuronal and nonneuronal cell types of the OE (Graziadei and Graziadei, 1979; Caggiano et al., 1994; Schwob et al., 1995; Chen et al., 2004; Leung et al., 2007). During both development and regeneration, cells of the epithelium transition from highly proliferative multipotent cells to neurons that choose the one odorant receptor among the set that is appropriate for, and available to, neurons within that region of the OE (Schwob, 2002; McClintock, 2010).

The progenitor cells apparently express a defined sequence of transcription factors during neurogenesis that seem to mark distinct stages en route to producing the differentiated cell types of the OE. This progression has been identified by immunohistochemical analysis (which directly assesses whether or not there is overlap between specific transcription factors), genetic epistasis in mutant mice, and the timing of expression after, and in response to, various types of lesion (Cau et al., 1997, 2002; Manglapus et al., 2004; Leung et al., 2007; Iwai et al., 2008; Guo et al., 2010; Packard et al., 2011). Analysis in embryonic and adult rodent epithelium indicates that Sox2 expression marks GBCs that are “upstream” stem and progenitor cells, including ones that are multipotent (Kawauchi et al., 2005; Beites et al., 2005; Guo et al., 2010; Tucker et al., 2010). Sox2 marks the primitive placodal cells during invagination of the olfactory pit, and it is expressed by all GBCs immediately after activating lesions of the adult epithelium, at a stage before the appearance of markers of differentiated or committed cells (Kawauchi et al., 2005; Guo et al., 2010). The basic helix-loop-helix transcription factors Ascl1 (also known as Mash1), Neurog1, and NeuroD1 are expressed in that order downstream of the multipotent Sox2-only GBCs. This cascade progresses as GBCs become first neuron-committed and then neuron-producing mitotic progenitors and precursors during development and during the regeneration of the OE (Cau et al., 1997, 2002; Murray et al., 2003; Manglapus et al., 2004; Packard et al., 2011). That sequence was evident when assessing the consequences of targeted gene mutation (e.g., Ascl1 knockout and Neurog1 knockout) in the embryo and by the timing of gene expression during the reconstitution of the methyl bromide (MeBr)-injured OE (Cau et al., 1997, 2002; Murray et al., 2003; Manglapus et al., 2004; Beites et al., 2005; Guo et al., 2010). While Neurog1 and NeuroD1 are coexpressed in many GBCs in the normal or lesioned-recovering adult OE, Ascl1 is wholly nonoverlapping with NeuroD1 (Packard et al., 2011). The hierarchical relationships among this small set of transcription factors provides a convenient framework for approximating the different functional capacities of these cells. However, we lack a comprehensive understanding of the molecular phenotype of various stages in the neurogenic progression, which is required for understanding how those differences come about.

Moreover, damage to the neuronal population or the OE as a whole activates stem and progenitor cells that respond by expanding the population of actively dividing GBCs and, in the case of HBCs after epithelial lesion, by acquiring multipotency. While the response to injury may be mediated in part by the activity of these same transcription factors, it is not known whether more downstream neuronal precursors behave differently or acquire

different characteristics as a consequence of a radically different developmental history (Manglapus et al., 2004; Leung et al., 2007; Guo et al., 2010).

Beyond the canonical expression of the aforementioned transcription factors, very little is known about the epigenetic influences responsible for the vastly different transcriptional profile of upstream progenitors as compared to differentiating neurons. For example, the choice of odorant receptor by differentiating neurons looks to be subject to feedback and epigenetic control. The first sign of OR expression, as monitored by the accumulation of a marker protein encoded by a bicistronic OR-IRES-marker construct, is seen after the neuron is “born” (has completed its terminal mitosis) and after some further lag relative to the expression of general neuronal markers (Iwema and Schwob, 1993).

To that end, the current experiments analyzed the global expression profile of defined stages in the progression of GBCs as they apparently commit, proliferate, and differentiate into neurons with or without lesion-induced activation. Various cell populations can be isolated for microarray profiling due to the availability of transgenic reporter mice that mark defined stages in the hierarchy of GBCs as well as stages of neuronal differentiation; in these mice enhanced green fluorescent protein (eGFP) expression is linked with that of transcription factors and proteins that have traditionally been used to define these stages. McClintock, Getchell and their laboratories have already demonstrated the usefulness of this approach for mature olfactory sensory neurons using $\Delta OMP-eGFP$ mice (Shetty et al., 2005; Getchell et al., 2006; Sammeta et al., 2007). For this study, we extended that analysis to more upstream progenitors and focused specifically on the neurogenic progression in the OE. Using a combination of transgenic reporter mice and fluorescence-activated cell sorting (FACS), we isolated populations enriched for upstream multipotent progenitor GBCs (defined by Sox2 expression), immediate neuronal precursor GBCs (defined by Neurog1 expression), and olfactory sensory neurons (defined by olfactory marker protein [OMP] expression). Olfactory bulbectomy of transgenic animals was used to test the hypothesis that lesion-induced activation drastically changes the expression profile of Neurog1 and OMP-expressing cells. Additionally, we report for the first time expression profiles for cell populations enriched for two types of GBCs, Sox2(+) GBCs and Neurog1(+) GBCs. These data shed new light on the pathways involved with and genes important for the progression from upstream multipotent progenitor to differentiated olfactory sensory neuron.

MATERIALS AND METHODS

Animals

Wildtype F1 males were bred in-house from parental strains (129S1/SvImJ $\times$ C57BL/6J) acquired from the Jackson Laboratory (Bar Harbor, ME). A number of gene-targeted transgenic mouse lines were also used. *Neurog1-eGFP* BAC transgenic mice were generated by the GENSAT project (Gong et al., 2003) and maintained as heterozygotes by successive matings to FVB/NJ mice or 129S1/ SvImJ (Jackson Laboratory); BAC RP23-457E22 (Gensat BX561) was modified by the insertion of eGFP immediately downstream of the initiation codon of the *Neurogenin 1* (*Neurog1*) gene via recombineering, then purified BAC DNA was injected into the pronuclei of fertilized oocytes (Gong et al., 2003). Multiple transgenic lines were evaluated and the one supplied by the GENSAT project matched the reported expression pattern of *Neurog1* expression.

$\Delta OMP-eGFP$ mice were generously provided by Dr. Peter Mombaerts (Potter et al., 2001) and used as heterozygotes generated by outcrosses of homozygous males to CD-1 females; in this case, the entire open reading frame for OMP was removed during the recombination/insertion of GFP from the initiation methionine codon onward. The use of heterozygotes is intended to eliminate concerns regarding the distortions that occur in signal transduction and

olfactory function as a consequence of the total absence of OMP. While there are few published data on OMP heterozygotes (Youngentob and Margolis, 1999; Youngentob et al., 2001, 2003; Reisert et al., 2007; Kwon et al., 2009), those that are in the literature suggest that heterozygosity has no physiological consequence, as the slope and recovery kinetics of EOGs recorded in heterozygotes are indistinguishable from the wildtype control (Ivic et al., 2000). Moreover, haploinsufficiency is a rare consequence of gene deletion (Wilkie, 1994). Furthermore, immunostaining for OMP in heterozygotes is as robust as in wildtype animals (for example, Fig. 1). Finally, as shown by the results below, the gene expression profile for the eGFP-expressing mature olfactory neurons (from the heterozygote mice) shows substantial overlap with the profile of normal olfactory mucosa, which is dominated by mature olfactory neurons that are wild-type for the OMP gene (see Fig. 3). Thus, olfactory sensory neurons (OSNs) from heterozygous animals have been used as the normal control for comparing gene expression between them and homozygous knockout animals in other publications (Sammata et al., 2007).

$\Delta Sox2-eGFP$ mice were generously provided by Dr. Mahendra Rao (Ellis et al., 2004) and were maintained as an inbred colony. Heterozygous offspring of outcrosses to FVB/NJ females were used; in this mouse the entire open reading frame and 1 kb of the 3' untranslated region was replaced with an eGFP-loxP-neomycin resistance-loxP cassette. No phenotype has been observed by us in the OE of the $\Delta Sox2-eGFP(+/-)$ mice or even $\Delta Sox2(+/-)/\Delta Pax6(+/-)$ double heterozygotes despite close examination (Packard and Schwob, data not shown).

The age and gender of the mice of these strains used in the microarray analysis are listed as part of the GEO record (<http://www.ncbi.nlm.nih.gov/geo/>) as required by NIH guidelines and are available as record GSE37581. All of them were within the age range of 8–12 weeks and there were no systematic differences in the percentage of male or female mice between the groups.

All animals were housed in a heat- and humidity-controlled, AALAC-accredited vivarium operating under a 12:12-hour light/dark cycle. All protocols for the use of vertebrate animals were approved by the Committee for the Humane Use of Animals at Tufts University School of Medicine, where the animals were housed and experiments were conducted. All mice were maintained on a 12-hour light/dark cycle with ad libitum access to food and water.

Olfactory bulbectomy and methyl bromide lesion

The right olfactory bulb was removed in mice of the *Neurog1-eGFP* and $\Delta OMP-eGFP$ transgenic lines as previously described (Schwob et al., 1992). In brief, 8–12-week-old mice were anesthetized by intraperitoneal injection of 0.6 ml/kg of an induction cocktail (43 mg/ml ketamine, 9 mg/ml xylazine, 1.5 mg/ml acepromazine), supplemented as needed by 0.5 ml/kg of a maintenance cocktail (95 mg/ml ketamine, 1.9 mg/ml acepromazine). The bulb was exposed by removal of the overlying bone. The dura was lanced with a sterile 27G needle and the bulb was removed by aspiration. Hemostasis was achieved by filling the ablation cavity with Oxycel and the animals were killed 3 weeks after the surgery as described in the next section. Only septum and turbinates ipsilateral to the bulbectomized side were isolated for analysis; tissue from the unoperated side was discarded. Unoperated, age-matched animals were used as the respective controls for each transgenic line subjected to bulbectomy. In addition, 129 × B6 F1 male mice were used for the immunohistochemical analysis of coREST complex expression.

Other 12-week-old 129 × B6 F1 male mice were exposed to methyl bromide (MeBr) gas at a concentration of 180 ppm for 8 hours as previously described (Chen et al., 2004), and used at the time points listed in the results for purposes of coREST complex expression.

Cell dissociation and FACS

Detailed FACS protocols have been reported from our laboratory (Chen et al., 2004). Briefly, for all of the target populations except for the GBC_{MPPs}, normal and bullectomized adult animals were injected with an overdose of the induction anesthesia cocktail and then perfused intracardially with low-Ca²⁺ Ringer's solution (140 mM NaCl, 5 mM KCl, 10 mM HEPES, 1 mM EDTA, 10 mM glucose, and 1 mM sodium pyruvate, pH 7.2) when deeply anesthetized. The nasal mucosa was separated into the septum and individual turbinate scrolls, which were then incubated with 0.05% trypsin/EDTA (Gibco, Grand Island, NY) in low-Ca²⁺ Ringer's solution for 15 minutes at 37°C, followed by a cocktail of dissociation enzymes (collagenase/hyaluronidase/trypsin inhibitor/papain; 1 mg/ml, 1.5 mg/ml, 0.1 mg/ml, 15 µl/ml, respectively; Worthington Biochemical, Freehold, NJ, and Sigma, St. Louis, MO) in Ringer's solution (140 mM NaCl, 5 mM KCl, 10 mM HEPES, 1 mM CaCl₂, 1 mM MgCl₂, 10 mM glucose, and 1 mM sodium pyruvate, pH 7.2) for 30 minutes at 37°C with occasional trituration. Dissociated cells were treated with DNase I (Worthington) and subsequently filtered through 120 µm and 35 µm nylon mesh before staining and FACS. The foregoing protocol was modified for purposes of isolating GBC_{MPPs} by substituting a purely mechanical dissociation (without proteolytic digestion, but including DNase I) in order to take advantage of their simpler, roughly spherical shape and their lack of attachment to the basal lamina, making them more likely to be released intact from the tissue by comparison with Sustentacular (Sus) cells and HBCs, and to minimize the proteolytic digestion of cell surface antigens used for the sort. It was anticipated that purely mechanical dissociation would compromise yield, but would likely fracture those other cell types and eliminate their contribution to the transcriptional profile.

For certain populations, the dissociated cells (either enzymatic or mechanical) were stained in suspension with anti-CD54 antibody or anti-E-cadherin antibody for 30 minutes at 4°C, washed by centrifugation, and incubated with the appropriate allophycocyanin (APC)-conjugated secondary antibody before undergoing FACS.

FACS was performed on a MoFlo cell sorter (Cytomation, Fort Collins, CO) running Summit software (Dako, Carpinteria, CA). Dissociated cells from OE of wildtype and transgenic mice were prepared as described above and resuspended in staining solution containing Hank's buffered salt solution (Ca²⁺/Mg²⁺-free HBSS; Invitrogen, Aukland, NJ) with 25 mM HEPES and 1 mM EDTA. An Innova 90 argon plasma laser (Coherent, Santa Clara, CA) was used to excite the cells at 488 nm and the samples were gated in FL1 (for GFP-positivity) and FL6 (for APC-positivity). Additional gating on forward- and sidescatter and on exclusion of propidium iodide (PI) was used to limit the sort to live cells (Jang et al., 2007). Cells were collected in HBSS/HEPES/EDTA solution on ice. After collection, cells were spun for 10 minutes at 200g to pellet the cells.

RNA purification, quality control, labeling, and bead array processing

Cell pellets were resuspended in Buffer ZR (Zymo Research, Orange, CA, Quick RNA Microprep Cat. No. R1050) and the manufacturer's instructions were followed exactly. RNA was eluted with water and DNase treated for 30 minutes at 37°C. RNA was then purified using the DNA-free RNA kit from Zymo Research (Cat. No. R1013). Samples were run on an Agilent 2100 Bioanalyzer (Palo Alto, CA) to determine quality and select intact, nondegraded samples. Three biological replicates (2–7 animals per replicate) were sent to the Keck Microarray Facility (Yale University) for RNA amplification, cRNA production,

labeling, and hybridization to Illumina MouseWG-6 v2.0 bead arrays. Illumina Genome Studio software was used to export both raw data and background corrected, quantile-normalized data.

Microarray data: quality control and analysis

Quality control and processing of microarray data was performed using the BioConductor packages *lumi* and *limma* for R (Smyth, 2004; Gentleman et al., 2004; Smyth et al., 2005; Du et al., 2008). Fluorescence intensity values were exported to Microsoft Excel in which data were formatted for subsequent analysis with GenePattern hierarchical clustering modules (Reich et al., 2006). The *limma* package was used to generate pairwise comparisons of the biological comparisons of interest. Gene lists for each comparison were filtered based on an adjusted $P < 0.05$. When smaller gene lists were required for analysis or study of specific candidates, an additional criterion of a fold-change greater than 2 was used.

Filtered lists described above were then input into the DAVID (Database for Annotation, Visualization and Integrated Discovery; <http://david.niaid.nih.gov>) functional annotation clustering tool (e.g., Fig. 4, Table 4). DAVID default settings (which include numerous gene ontology classifications and functional domain classifications for all annotated genes) were used for this analysis, with the additional inclusion of Biocarta and Panther pathway databases for the clustering analysis. Upregulated gene lists were input into DAVID separately for each population analyzed such that the resulting enrichment terms correspond to the first condition listed in any given comparison. The DAVID analysis tool treats all inputted genes equally—magnitude of fold change or expression value from the Illumina array is not taken into account for this portion of the analysis. Enrichment of a particular annotation cluster indicates that this annotation is overrepresented, or enriched in the gene list for that given population compared to the background level expected based on the frequency of occurrence of genes with the same annotation in the mouse genome. Resulting enrichment clusters were considered for further analysis if the enrichment score was >1.3 (as recommended by the database designers) (Dennis et al., 2003; Huang et al., 2009).

The expression data obtained for each type of sort has been submitted to GEO (<http://www.ncbi.nlm.nih.gov/geo/>) as required by NIH guidelines and is available as record GSE37581.

Tissue processing and immunohistochemistry

Animals were deeply anesthetized with the induction cocktail and transcardially flushed with phosphate-buffered saline (PBS) pH 7.2, then perfused with either 4% paraformaldehyde (Fisher Scientific, Suwanee, GA) in 0.05 M sodium phosphate buffer, pH 7.2, or Zamboni's Fixative. The OE was dissected and the tissue blocks were postfixed under vacuum for 2 hours. Tissues were then rinsed with PBS, equilibrated with 30% sucrose in PBS, and then frozen in OCT compound (Miles, Elkhart, IN). The olfactory mucosa was sectioned on a cryostat (Leica) in the coronal plane, 8- μ m sections were collected onto "Plus" slides (Fisher Scientific), and stored at -20°C for future applications.

Standard laboratory protocols were used to detect the expression pattern of individual proteins in normal and transgenic OE. Adequate labeling with a number of the antibodies requires a set of treatments on the sections prior to incubation with the antibodies. Briefly, frozen sections were rinsed in PBS for 5 minutes to remove the OCT and then boiled in 0.01 M citric acid buffer (pH 6.0) for 10 minutes using a commercial food steamer. After cooling, sections were rinsed with PBS briefly before incubating with block (10% serum + 5% nonfat dry milk + 4% bovine serum albumin [BSA] + 0.1% Triton X-100) for 15 minutes at room temperature. In all cases the sections were incubated with primary antibodies overnight at

4°C. Bound primary antibodies were visualized by the corresponding fluorescently conjugated secondary antibodies.

Antibody specificity

Please see Table 1 for a list of all antibodies used. Information on the antibodies is derived from the manufacturers' description and our own data.

The chicken anti-GFP and rabbit anti-GFP antibodies stain various transgenic reporter mice bearing the eGFP transgene, they do not stain wildtype littermates.

The rat anti-NCAM antibody recognizes a triplet of glycoproteins at the neural cell surface that is identified as neural BSP2, which is a synonym for NCAM (Abcam, Cambridge, MA; Datasheet ab19782).

The goat anti-OMP antibody obtained from Wako (Osaka, Japan) recapitulates the canonical staining pattern for OMP described for the goat anti-OMP originally distributed by Frank Margolis (Keller and Margolis, 1975; Baker et al., 1989; Verhaagen et al., 1989, 1990; Schwob et al., 1992) and has been extensively characterized previously in this journal (Rodriguez-Gil and Greer, 2008).

The anti-Neurog1 antibody detects a single band of ~19 kDa on western blots of mouse brain tissue extract (Santa Cruz Biotechnology, Santa Cruz, CA; Datasheet sc-19231), and the post-MeBr lesion timing, number, and pattern of labeled cells fit with ISH studies (Manglapus et al., 2004).

Tuj1 is a mouse monoclonal antibody that is highly reactive with neuron-specific Class III β -tubulin (β III) and does not label β -tubulin found in glial cells.

The mouse anti-Ki67 antibody reacts with 345/395 kDa doublet on western blots corresponding to Ki-67 Ag; B56 binding to cells is blocked by clone MIB 1, the canonical α -Ki67 antibody (BD Biosciences, San Jose, CA; Datasheet 556003).

The mouse anti-CK14 monoclonal antibody VP-C410 labels basal cells in skin, selectively, with a cytoplasmic distribution (Vector Laboratories, Burlingame, CA; Datasheet VP-C410). The staining pattern with VP-C410 is completely equivalent to that observed with another anti-CK14 monoclonal, LL-002, which labels a single spot in 2D IEF-SDS-PAGE western blot with a molecular weight and isoelectric point consistent with CK14 (Holbrook et al., 1995). The choice of VP-C410 was predicated on the retention of antigenicity despite formaldehyde fixation.

The rabbit anti-CK14 antibody gives an identical staining as above mouse α -CK14 and stains A431 cells and skin or squamous cell carcinoma (Lab Vision, Fremont, CA; Datasheet).

The rat anti-E-cadherin antibody blocks both the aggregation of mouse embryonal carcinoma cells and the compaction of preimplantation embryos. The antibody disrupts confluent monolayers of MDCK epithelial cells (Sigma Datasheet).

The anti-CD-54 antibody shows inhibition in the HSB2 adhesion assay by blocking adhesion of target cells to ICAM-1 (R&D Systems, Minneapolis, MN; product datasheet AF583) and a single band at 90 kDa molecular weight is detected on western blot of HUVEC cell lysate stimulated with tumor necrosis factor- α (TNF α ; Sawa et al., 2007).

The rabbit anti-LSD1 (lysine-specific demethylase 1) antibody detects a single band at 105 kDa on western blot of P815 cell lysate (AbCam; Datasheet ab62582).

The rabbit anti-coREST (corepressor of REST[RE1 silencing transcription factor 1]) antibody detects a single band of 66 kDa on western blot of A431 lysate (Millipore, Bedford, MA; Datasheet 07–455).

The mouse anti-Mash1 antibody detects a single band at 34 kDa on western blot of fetal rat brain (BD Biosciences, San Jose, CA; Datasheet 556604).

The goat anti-Sox2 antibody detects a single band of 34 kDa in western blot of mouse and human embryonic stem cell lysates (clone Y-17; Datasheet sc-17320), and antibody staining on embryonic tissue from Mash1 knockout OE replicates the published Sox2 mRNA pattern by ISH (Kawauchi et al., 2005). Conditional deletion of Sox2 extinguishes immunolabeling in mouse cortex by the antibody (Favaro et al., 2009).

The following secondary antibodies were used: Alexa-Fluor488-donkey antigoat IgG, AlexaFluor488-donkey antimouse IgG, AlexaFluor488-donkey antirabbit IgG, AlexaFluor594-donkey antigoat IgG, AlexaFluor594-donkey antimouse IgG (all from Invitrogen/Molecular Probes, Eugene, OR) each used at a dilution of 1:250. Additionally, AlexaFluor488-conjugated streptavidin and Alexa-Fluor594-conjugated streptavidin (Invitrogen/Molecular Probes) were used at dilutions of 1:250 for conventional immunofluorescence. The remaining secondary antibodies were used at a dilution of 1:50 and were purchased from Jackson ImmunoResearch: biotin-conjugated donkey antigoat IgG, biotin-conjugated donkey antimouse IgG, biotin-conjugated donkey antirabbit IgG, FITC-conjugated donkey antigoat IgG, FITC-conjugated donkey antirabbit IgG, AMCA-conjugated donkey antimouse IgG, Texas Red-conjugated donkey antigoat IgG.

RESULTS

Characterizing the populations isolated by FACS

For purposes of expression profiling each of the following three cell populations were isolated from the normal OE via FACS by utilizing a combination of cell surface markers and expression of endogenous eGFP in various transgenic mouse lines: 1) putatively multipotent progenitor GBCs (GBC_{MPP}); 2) immediate neuronal precursor GBCs (GBC_{INP}) and immature neurons; and 3) mature neurons. Sus cells and normal olfactory mucosa were also isolated and used as needed as control populations. In addition, we also sorted GBC_{INP}/immature neurons as well as mature neurons from epithelium activated by removal of one olfactory bulb (bulbectomy or OBX) 3 weeks prior to harvest. The global transcriptional profile for each population was compared to control and to one another in biologically meaningful ways. As a consequence, we identified a set of genes that differentiates GBCs from other cell types, as well as the different stages within the neurogenic progression. A subset of the pathways and genes identified by transcriptional profiling is highlighted and examined in more detail to validate and expand upon genes that inform the regulation of this neurogenic process. The following describes the characteristics of the gene expression patterns in the epithelium of the different transgenic mouse strains and validates our identification of the three stages.

Differentiated OSNs were isolated on the basis of eGFP expression from the $\Delta OMP-eGFP$ mouse line, which has been used for this purpose elsewhere (Potter et al., 2001; Sammeta et al., 2007). The onset of OMP expression is conventionally used as a marker of full neuronal maturity since it seems to correlate with the elaboration of cilia and is reciprocal with markers of neuronal immaturity such as GAP43 (Verhaagen et al., 1989, 1990; Schwob et

al., 1992). We used heterozygous mice to avoid the confounding functional consequence of completely eliminating functional OMP protein (Youngentob and Margolis, 1999; Youngentob et al., 2001, 2003; Reisert et al., 2007; Kwon et al., 2009). In these animals the vast majority, if not all, of the eGFP(+) cells coexpress endogenous OMP, which indicates that expression of eGFP is confined to mature neurons and is not expressed in advance of maturation (Fig. 1). As expected, eGFP expression is also reciprocal with markers of immature neurons, such as high levels of NCAM expression, as is the OMP itself. Animals that were analyzed 3 weeks after bulbectomy show the stratification of immature neurons (marked by intense expression of NCAM) and maturing OSNs (marked by OMP and GFP). Post-hoc immunocytochemical analysis of eGFP(+) cells isolated by FACS indicates that this is a highly enriched population, as only 3% are contaminating Sus cells (marked by staining for CK18) (data not shown).

Immature postmitotic neurons and the population of GBC_{INPs} were isolated on the basis of eGFP expression from the *Neurog1-eGFP* BAC transgenic line (Gong et al., 2003). *Neurog1* is known to be downstream of *Ascl1* by analysis of gene mutation and epistatic relationships and on the basis of timing during embryonic development or recovery after epithelial lesion in the adult; it likely marks the GBCs whose daughter cell(s) become postmitotic neurons (Cau et al., 2002; Manglapus et al., 2004). In the *Neurog1-eGFP* mice, a discrete row of eGFP(+) cells lines the basal region of the OE. Because GFP has the potential to persist beyond the lifetime of *Neurog1* protein, we compared the population of GFP-labeled cells to the expression of both *Neurog1* itself and markers of immature olfactory neurons. When compared to endogenous *Neurog1* protein, we find that eGFP is expressed in the majority of *Neurog1*(+) cells (Fig. 1). However, eGFP is not limited to proliferative GBCs nor the *Neurog1*(+) population, as it is also expressed in many intensely NCAM(+) immature neurons, indicating that *Neurog1*-driven eGFP does perdure relative to the *Neurog1* protein (Fig. 1). eGFP is undetectable beyond the immature neuronal stage, i.e., OMP(+) cells are not labeled with anti-GFP (Fig. 1). The populations of eGFP(+) and of OMP(+) cells remain discrete and reciprocal in the *Neurog1-eGFP* mice even when neurogenesis is accelerated by ablation of the olfactory bulb. Thus, little or no overlap is seen between the GFP expression and OMP labeling of basal and apical cell populations, respectively, even though the OE of the lesioned mice is thinner, proliferation is accelerated, the population of immature neurons is expanded, and mature neurons are fewer (Fig. 1). In fact, accelerated neurogenesis resulting from bulbectomy highlights the differences between stages of differentiation in olfactory neurogenesis, as shown by stratification of NCAM(+) postmitotic neurons as a distinct layer apical to some TuJ1(+) progenitors that express β III tubulin (neuron-specific tubulin) (data not shown and Packard et al., 2011). eGFP(+) cells isolated by FACS from *Neurog1-eGFP* mice are also highly enriched, as only 5% of the cells are contaminating Sus cells seen by staining for CK18. In addition to the aforementioned populations from the normal OE, we also used the same FACS strategy to isolate and compare eGFP(+) GBC_{INPs}/immature neurons (from *Neurog1-eGFP* mice) with mature OMP(+) neurons (from Δ OMP-*eGFP* mice) following unilateral bulbectomy. Thus, the impact of accelerated neurogenesis was also addressed.

The population of GBC_{MPPs} was enriched by FACS on the basis of eGFP expression in the Δ Sox2-*eGFP* mice and lack of cell surface antigen labeling. The Δ Sox2-*eGFP* transgenic line expresses eGFP under control of the endogenous Sox2 locus via ES cell knockin and faithfully reproduces the expression pattern of Sox2 protein (Ellis et al., 2004). However, eGFP expression in the Δ Sox2-*eGFP* animals is complex, as is the distribution of the Sox2 protein. GBCs, HBCs, and Sus cells all express Sox2 protein, albeit at different levels (Guo et al., 2010). Thus, the most heavily eGFP-labeled cells are Sus cells at the apex of the epithelium, which are also most heavily labeled for Sox2 protein. HBCs and GBCs are less strongly labeled either by anti-Sox2 antibodies or by eGFP expression. Confocal microscopy

highlights these three distinct populations of eGFP(+) cells (Fig. 2). $\Delta Sox2$ -eGFP(+) GBCs are those basal cells that are CK14(-) and Ki67(+).

Thus, the three distinct cohorts of cells that express eGFP—Sus cells, GBCs, and HBCs—can be identified and enriched by FACS following staining with E-cadherin (which strongly labels Sus cells, but not GBCs; data not shown) and CD54 (which strongly labels HBCs, but not GBCs; Carter et al., 2004; Guo et al., 2010) by taking eGFP intensity and marker labeling into account as well as dissociation protocol and scatter-gate criteria (Fig. 2; Table 2). Post-FACS analysis of the three cohorts of Sox2-eGFP(+) cells demonstrates the expected enrichment: >95% of cells in the strongly positive eGFP population (designated +++) are CK18(+), >95% of cells in the moderately positive eGFP (designated +)/CD54(+) population are CK14(+) (unpubl. data). Thus, the Sus cell and HBC sorts are highly purified when the sorting strategy targets their corresponding surface antigens (Table 2). In contrast, the other eGFP(+) population—namely, GBCs—were sorted following purely mechanical dissociation as the E-cadherin(-)/CD54(-)/eGFP(+) cells, i.e., by excluding eGFP(+++) cells and all those that stain with surface markers that characterize Sus cells and HBCs (Fig. 2; Table 2). Because the sorting strategy targets unstained cells, the E-cadherin(-)/CD54(-)/eGFP(+) gating is not completely effective at eliminating all Sus cell contaminants from that fraction, most likely because E-cadherin is clipped from the surface of some Sus cells during the tissue dissociation, and ~30% of the sorted cells are CK18(+), indicating contamination by Sus cells or fragments. However, both the fall-off in eGFP intensity in these CK18(+) cells and the scatter-gate settings, which exclude cells of complex shape, imply that any Sus cells among the sorted population were damaged during dissociation and suggest that other readily soluble constituents, such as mRNA, were at risk for leaking out. Indeed, expression analysis demonstrates little contribution of Sus cells to the transcriptional profile of the E-cadherin(-)/CD54(-)/eGFP(+) population (see below).

Taken together, the cells isolated by FACS from these transgenic animals under normal and bulbectomized conditions represent a five-step neurogenic progression: 1) upstream $\Delta Sox2$ -eGFP(+) GBC_{MPP}; 2) *Neurog1*-eGFP(+) GBC_{NP} and immature neurons from the epithelium of control, unoperated transgenic animals (Neurog1-Ctrl); 3) putatively activated *Neurog1*-eGFP(+) GBC_{NP} and immature neurons from the epithelium of bulbectomized mice (Neurog1-OBX); 4) newly mature ΔOMP -eGFP(+) neurons from the epithelium of bulbectomized mice (OMP-OBX); and 5) the pool of ΔOMP -eGFP(+), fully mature neurons from control, unoperated transgenic mice (OMP-Ctrl) (Fig. 1, 2; Table 2). RNA samples from three additional populations of cells were available as controls and comparisons: Normal olfactory mucosa, $\Delta Sox2$ -eGFP (+++) Sus cells, and mouse reference RNA (Stratagene, La Jolla, CA; Cat. No. 740100). Sus cells were also isolated as a highly pure population (>95% CK18-positive on post-hoc analysis) using their optical complexity, size, and high level of GFP expression in $\Delta Sox2$ -eGFP mice.

Global analysis of isolated populations

As a first step towards understanding the global transcriptional differences among the stages in the neurogenic progression, the general quality of the dataset and relationships between the samples were evaluated, lists of significantly regulated targets were generated for a variety of pairwise comparisons, and the significantly regulated genes were evaluated by functional annotation clustering. (Note on terminology: due to partial redundancy intentionally built into the Illumina Bead array technology, the same gene may be represented on multiple beads each targeting a different region of the specific mRNA. In what follows, “target” refers to an individual bead sequence corresponding to a portion of the annotated gene, while “gene” refers to an actual unique gene. As a consequence, the number of targets overrepresents the number of actual unique genes by a small percentage.)

Sample relations hierarchical clustering was applied to >7,000 of the most highly regulated targets across the datasets to cluster all the samples into a dendrogram. Most of the replicates clustered together as expected (Fig. 3). In contrast to that general rule, the samples from the control, unoperated *Neurog1-eGFP* mice and the bulbectomized *Neurog1-eGFP* mice were intermingled (Neurog1-Ctrl and Neurog1-OBX, respectively; Fig. 3). The intermingling suggested that the differences in gene expression between treatments were insignificant for these cell populations. In contrast, the samples from the control, unoperated Δ OMP-*eGFP* mice (OMP-Ctrl) and the bulbectomized Δ OMP-*eGFP* mice (OMP-OBX) clustered according to treatment condition, indicating that the between-treatment differences were more significant than the among-replicate differences. Other results included a close clustering of all the Δ OMP-*eGFP* enriched conditions with the normal olfactory mucosa (Normal OM), which is a likely consequence of the high percentage of cells in the normal olfactory mucosa that are OMP(+) olfactory sensory neurons. In addition, the GBC-containing samples (Neurog1-Ctrl, Neurog1-OBX, and Sox2-GBC) all clustered more closely together with each other than with any of the other cell types profiled, including the Sus cells, which are distant (Fig. 3). Thus, contamination with Sus cell mRNA was, indeed, not a significant factor in the overall transcriptional profile of the Sox2-GBC population, since the Sox2-GBC population would have clustered more closely with the Sus population than with any of the other samples had contamination contributed to the expression profile.

In order to determine which genes and pathways are upregulated or downregulated in specific populations, a series of pairwise comparisons between different cell populations were generated by linear modeling of the microarray data (Smyth, 2004). Three biological replicates were generated for each cell population (as described in Materials and Methods) and used to calculate fold changes with corresponding adjusted *P*-values (to correct for multiple hypothesis testing). The numbers of targets that differ significantly in the various pairwise comparisons are listed in Table 3 and parallels the distance observed on the dendrogram (Fig. 3). The pairwise comparisons also offer insight into a number of important questions related to the gene expression changes that occur during the transitions from multipotent progenitor to differentiated neurons.

As expected from the dendrogram analysis, the pairwise comparisons demonstrate substantial differences in the transcriptional profile between cell types and between stages of neurogenesis (Table 3); for example, some 6,976 targets differ significantly between the transcriptional profiles of Neurog1-Ctrl vs. OMP-Ctrl and 10,640 for Sox2-GBC vs. Sus cells. Fewer differentially expressed targets were observed when comparing the sequential stages of neurogenesis corresponding to Sox2-GBC vs. Neurog1-Ctrl (3,143 differences). Perhaps the most unanticipated results emerge from the pairwise comparisons of the same types of cells from the epithelium of bulbectomized vs. control mice. In the case of the pairwise comparison of *Neurog1-eGFP*(+) cells between Neurog1-OBX vs. Neurog1-Ctrl mice, the expression of only an exceptionally small number of genes differs significantly between the two conditions (63, as compared to >1,500 for each of the other pairwise comparisons) (Table 3). These data indicate that *Neurog1-eGFP*(+) cells do not alter their gene expression profile despite the acceleration of neurogenesis caused by bulb ablation. In contrast, gene expression in the OMP-Ctrl dataset is substantially unlike that of the OMP-OBX cells, such that recently maturing OMP(+) neurons of the bulbectomized mice differ from the stable population of control mice by >1,500 targets (Table 3). The large number of differences most likely reflects the abbreviated lifespan and the corresponding transience of the OMP population born in the absence of the bulb as compared to control (Schwob et al., 1992).

Transitions from one stage to the next in the progression from GBC to mature OSN are known to be identifiable on the basis of immunohistochemical markers, and a comparison of

the level of expression of some of these is useful. For example, one of the best known and most used is the upregulation of the somatic expression of GAP-43 in association with the expression of neurotubulin and NCAM, which is then followed by the downregulation of GAP-43 somatic staining roughly contemporaneous with the onset of immunodetectable OMP labeling (Verhaagen et al., 1989, 1990; Meiri et al., 1991; Schwob et al., 1992, 1995). The levels of GAP-43 gene expression are also regulated in a dynamic manner and correspond to the changes in the levels of GAP-43 immunolabeling in the somata. Thus, the levels of GAP-43 mRNA are low in the Sox2-GBCs, rise 1.4-fold in the Neurog1-Ctrl dataset before reaching a peak in the Neurog1-OBX population that is 1.6-fold that of the Sox2 GBCs. GAP-43 levels then decline about 1.4-fold to the OMP-OBX group and reach a nadir in the OMP-Ctrl cells, wherein GAP-43 mRNA is roughly 8-fold less than the level in the Neurog1-Ctrl cells. The dip in mRNA levels from Neurog1-OBX to OMP-OBX might be less sharp than might be expected from the immunostaining pattern. However, several factors influence the steepness of the decline as the neurons make the transition from immaturity to maturity. First, neurons that are double-labeled for GAP-43 and OMP reach significant numbers on the bulbectomized side of the epithelium, because the termination of axon growth and final neuronal maturation are not directly coupled (Monti Graziadei, 1983; Chuah and Farbman, 1983; Schwob et al., 1992). Second, the disappearance of GAP-43 from the cell body is presumed to signal the shutting off of the GAP-43 gene, but there are no data to prove that is so. Any persistence of GAP-43 expression during a transition period of finite length during that transition to final full differentiation would blunt the slope of the initial decline.

Significantly enriched annotation clusters in the neurogenic progression

Gene expression analysis by annotation clustering provided an open-ended approach to characterizing the different types of genes that were differentially expressed during the several transitions from multipotent GBC (Sox2-GBC) to mature OSN (OMP-Ctrl). Thus, the comparisons used for the annotation clustering of the regulated genes include comparisons to normal mucosa, comparisons to Sus cells, and comparisons across the several stages within the neuronal lineage. Since the normal mucosa is dominated by mature OSNs, the Sus cell population is perhaps the best control, because it is highly enriched, substantially less mitotically active, and biologically distinct from the neuronal lineage from an early stage. Normal OM is the least manipulated population, least subject to experimental artifact, and is an important validator for the gene clusters that emerge.

The identifiers of significantly regulated genes were input into the DAVID Bioinformatics database for functional annotation clustering (<http://david.abcc.ncifcrf.gov/home.jsp>) (Dennis et al., 2003; Huang et al., 2009). This tool compares the gene ontology classifications across all of the significantly regulated genes and returns annotation clusters of grouped genes that share ontology assignments. An enrichment score and a *P*-value for a particular cluster are calculated by comparing the index cluster to the chance of generating that cluster by random choice among the list of genes in the mouse. Higher enrichment scores indicate a more biologically relevant association for a given list of genes. An enrichment score of >1.3 is generally recommended as a cutoff indicating that the annotation cluster is biologically significant (Huang et al., 2009). The most highly enriched annotation clusters comparing cell types at different stages in the neurogenic progression are listed in Table 4.

Our accumulated understanding of the biology of neurogenesis in the OE suggests that GBCs should be proliferating and expressing genes characteristic of neuronal progenitors, while immature neurons should be expressing genes associated with neurogenesis and neuronal differentiation, and mature neurons should be expressing numerous olfactory

receptor genes and genes within the olfactory signal transduction cascade, but not genes related to progenitor cell biology or mitotic activity. In keeping with prediction, annotation clusters related to *mitosis*, *cell division*, *neuron development*, *neurogenesis* are highly enriched in the several GBC populations (Sox2-GBC, Neurog1-Ctrl, Neurog1-OBX) when compared to normal mucosa, Sus cells, or more differentiated cell types. The comparison of the Sox2-GBC and the Neurog1-Ctrl populations with normal mucosa produces a highly similar set of enriched terms for these two separate comparisons. For example, *neuron development/ nervous system development and organelle biogenesis/ cellular component biogenesis* show up in both of these GBC-containing populations (Table 4). When Neurog1-Ctrl or Sox2-GBCs are compared to Sus cells, *protein kinase inhibitor* is one of the most highly enriched terms and two of the genes in this annotation cluster are p21 (CKDN1A) and p57 (CDKN1C), two cyclin-dependent kinase inhibitors (CDKIs) known to play a role in inhibiting neuronal cell migration, inducing cell cycle exit, and acting as an antiapoptotic factors for neurons (Harms et al., 2007; Itoh et al., 2007). The finding that CDKIs are enriched in these populations alongside terms relating to mitosis and cell division appears contradictory. However, the enrichments for mitotic progression and quiescence likely reflect complexity and heterogeneity within the sorted populations. For example, some among the population of Sox2-expressing GBCs also express p27, another CDKI (Guo et al., 2010), and the Neurog1-Ctrl population included differentiating neurons, which also express several CDKIs (Legrier et al., 2001; Guo et al., 2010). More refined analysis and additional subdivision of the datasets will be needed to parcel out these effects.

The finding that many of the genes enriched in Neurog1-Ctrl and Sox2-GBC populations are identical when both are compared to Sus cells provides further evidence that these two populations are highly similar and require additional data mining and analysis and/or a more finegrained sorting strategy to resolve differences between the two datasets. When these two populations are compared to one another directly, terms such as *blood vessel morphogenesis* are enriched, and within the set of genes that contribute to this annotation cluster is FGFR2, a receptor for FGF ligands, both of which are expressed in the OE and likely involved in the proliferation-inducing effects of FGFs on GBCs (DeHamer et al., 1994; Hsu et al., 2001; Barraud et al., 2007). *Lung development* is another term associated with Sox2-GBCs. In this cluster are transcription factors such as Gata6, Foxa1, Id1, and growth factor-related genes BmprII and Pdgf-alpha. Gata6 has been shown to be involved in responding to retinoic acid signaling, and Id1 has been shown to affect the stability of tissue-specific basic helix-loop-helix (bHLH) factors, making these two proteins candidates for studies of whether or not they play analogous roles in the OE (Vinals et al., 2004; Capo-Chichi et al., 2005). The importance of TFG β /BMP-signaling is already firmly established in the OE by work from the Calof laboratory with the use of transgenic mice and in vitro models (Shou et al., 1999, 2000; Wu et al., 2003).

Differences in the annotation terms from other comparisons highlight additional biologically interesting differences between these populations. The Neurog1-Ctrl vs. OMP-Ctrl comparison provides a direct comparison between an immature neuron-enriched population (Neurog1-Ctrl) and a mature neuronal population (OMP-Ctrl). *Ribosome* was the most highly enriched term in the comparison, suggesting that the population of GBC_{INPs} and immature neurons is committing significant cellular resources to protein synthesis. *Mitosis* was also highly enriched in the Neurog1-Ctrl population, as would be expected for the comparison with an entirely postmitotic mature neuron population. *Chromatin assembly or disassembly* was also enriched in Neurog1 populations vs. OMP-Ctrl. The complex regulation of expression of OR genes and other key proteins related to axon targeting in the OE makes it likely that epigenetic processes are critical during the transitions from progenitor cell to mature neuron. Thus, it has been suggested that epigenetic mechanisms are thought to be involved in the global repression of all but the single receptor gene to ensure

the singularity of OR expression (McClintock, 2010). Some of the upregulated genes (or targets) known to be involved in epigenetic regulation of gene expression were investigated further and will be described in detail below.

Identifying the genes expressed at defined stages of the neurogenic progression

That the Sox2-GBC and Neurog1-Ctrl datasets show extensive similarity prompted us to refine the lists of upregulated genes with reference to their overlap with or exclusion from the OMP-Ctrl dataset. The goals of this analysis were to: 1) identify genes that are limited to certain cell populations as a way to identify potential unique markers for that population, and 2) identify pathways that typify distinct stages of the neurogenic progression. The Sox2-GBC, Neurog1-Ctrl, and OMP-Ctrl datasets were each compared to Sus cells. Genes that were significantly upregulated from those comparisons were used to generate a 3-way Venn diagram in order to define the unique vs. shared genes for each population (Fig. 4). The area of each region of the Venn diagram is roughly proportional to the number of genes there (Fig. 4). The area colored in red represents the 817 genes specific to Sox2-GBCs, the area in green represents the 979 genes specific to Neurog1-Ctrl, and the area in light blue represents the 1,686 genes specific to OMP-Ctrl dataset. A large number of the genes identified by comparison with Sus cells overlap all three populations (purple: 2,168), in line with the clustering observed on the dendrogram (Fig. 3). As might be predicted, only a very few genes are found in the OMP-Ctrl and Sox2-GBC datasets, but not in the Neurog1-Ctrl population (dark blue: 168; Fig. 4).

To address the functional significance of the differentially regulated genes among the three populations, the lists were input into DAVID as described above to identify the functional annotation clusters within each comparison (Fig. 4). A number of the DAVID terms match the biology expected for each of the various cell types. For example, GBC-specific genes (orange: 1,256) would include those genes found in the Sox2 and Neurog1 datasets and excluded from the OMP one; this set is enriched for terms such as *mRNA processing* and *transcription regulation*. Moreover, genes expressed in a more downstream population and found in both Neurog1 and OMP gene lists (yellow: 1,128) are significantly enriched for the functional annotation clusters expected of maturing olfactory neurons: *olfactory receptor* and *integral to membrane*. All three populations (purple: 2,168) contained clusters that included *olfactory transduction*, *neuron development*, and *cilium morphogenesis*. Interestingly, it is worth noting that neither the genes that were exclusive to the Sox2-GBC set, nor the ones exclusive to the Neurog1-Ctrl sets had particularly high enrichment scores for functional annotation clusters (all <3). Nonetheless, the clusters that were significantly upregulated in Sox2-GBC included: *negative regulation of cell migration*, *blood vessel morphogenesis*, *insulin-like growth factor binding*; and for Neurog1-Ctrl included *ribosome*, *nuclear lumen*, and *RNA biosynthetic process*. The list of exclusive OMP-Ctrl genes was highly enriched for *olfactory receptor activity* and *intrinsic to membrane*—two annotation terms that reflect the large number of odorant receptor genes in this list.

Expression of chromatin-modifying genes

The enrichment of chromatin modifying genes within the GBC populations prompted us to examine this functional annotation cluster in detail to nominate specific genes for expression analysis that may be important functionally in olfactory neurogenesis. Examination of the gene lists highlighted expected and unexpected results. For example, six members of the coREST-corepressor complex—*LSD1*, *coREST(RCor1)*, *HDAC2*, *Zmym2*, *Ehmt2*, *Cdy12*—out of a total of 18 identified components were differentially regulated across the datasets. The expression pattern of HDAC2 has been described previously by immunohistochemistry in OE (MacDonald et al., 2005) and appears in a pattern that is consistent with the microarray data reported here. The average expression value for these genes plotted for each condition

clearly shows that these components are expressed at a high level in more primitive progenitor cell types than in maturing OSNs (Fig. 5). The coREST complex is known to participate in the repression of neuronal differentiation genes in nonneuronal and neuronal progenitor cell types (Andres et al., 1999; Shi et al., 2005). One of the components, LSD1, has been shown to affect the global methylation patterns, cell-specific gene expression, and the proliferative status of neuronal progenitor cells as a result of its demethylase activity (Wang et al., 2007, 2009a; Sun et al., 2010). The remainder have not been analyzed in detail either in vitro or in vivo. However, in the neurogenic progression dataset the components of the coREST complex are not precisely coregulated. For example, LSD1 was higher in Sox2 in this setting, while coREST was higher in Neurog1.

As a consequence, we examined the expression of LSD1 and coREST more closely in normal, unoperated OE, in the context of heightened neurogenesis in the OE of bulbectomized mice, and during the regeneration of the OE as a whole following MeBr lesion. We used two strategies: colabeling with a standard panel of cell-type-specific antibodies and coexpression with GFP in transgenic marked mice.

LSD1 and coREST expression in OE dominated by neuronal turnover

Expression of LSD1 was examined by immunohistochemistry in the epithelium of unoperated animals (the control condition) and in the epithelium on the unoperated side and on the ablated side of unilaterally bulbectomized mice. LSD1(+) cells are concentrated in the basal region of the normal OE but are CK14(–), and the vast majority are also NCAM(–) (Fig. 6). On occasion, an LSD1(+)/NCAM(+) cell will be apical to the rest (Fig. 6). We compared anti-LSD1 staining to eGFP labeling in both the *Neurog1-eGFP* and in the $\Delta OMP-eGFP$ mouse strains (Fig. 6). Based on comparison with eGFP expression in these lines, most of the LSD1(+) cells are upstream within the neurogenic progression. With regard to *Neurog1-eGFP* mice, some, more deeply placed LSD1(+) basal cells do not express detectable levels of eGFP (Fig. 6). With regard to $\Delta OMP-eGFP$ mice, none of the LSD1(+) cells—either basal or the rare apical ones—contain detectable eGFP (Fig. 6).

The expression of coREST protein was different. coREST labels a population of cells about 1–2 cell layers above the basal lamina, with the strongest staining in more basally situated cells and decreasing intensity in more apically located cells (Fig. 7). In normal transgenic *Neurog1-eGFP* mice, coREST heavily labels the nuclei of a set of cells near the base of the epithelium, which are also eGFP(+); none of the coREST(+) cells are deep to the band of eGFP(+) basal cells and immature neurons (Fig. 7). Indeed, in some of the basally situated eGFP(+) cells of the *Neurog1-eGFP* mice coREST is undetectable by immunostaining. In addition to the densely labeled basal group, some more lightly labeled, coREST(+) cells are found apical to the eGFP(+) cells (Fig. 7). The expression pattern of coREST closely mimics that of HDAC2 (MacDonald et al., 2005; and data not shown). In response to bulbectomy, the number of coREST(+) cells dramatically expands in parallel with the expansion of the eGFP(+) population in the *Neurog1-eGFP* mice. A complementary pattern is observed with respect to eGFP labeling in the $\Delta OMP-eGFP$ mice. After bulbectomy, most of the heavily labeled coREST(+) cells are eGFP(–) in the $\Delta OMP-eGFP$ mice, which would be expected from the reciprocal distribution of eGFP in the *Neurog1-eGFP* vs. $\Delta OMP-eGFP$ mice. However, a low level of coREST is detectable (by comparison with the Sus cells) in many, perhaps most of the eGFP(+) cells in the $\Delta OMP-eGFP$ mice (Fig. 7).

The foregoing analysis of LSD1 and coREST suggest that LSD1 anticipates but overlaps with the expression of coREST. Thus, most coREST-expressing cells are immature neurons, some of which are also labeled by LSD1. However, the more basally situated LSD1-expressing cells are likely to be upstream GBCs. Indeed, by comparing with Sox2 and Ascl1, markers that identify subsets of upstream GBCs (Cau et al., 2002; Manglapus et al.,

2004; Guo et al., 2010), the overwhelming majority of LSD1(+) basal cells were found to be Sox2(+) or Ascl1(+), indicating that LSD1 is expressed in many GBCs upstream of the INP stage of the neurogenic progression (Fig. 8).

LSD1 and coREST expression in the OE recovering after MeBr lesion

The coexpression of LSD1 in GBCs suggests that this demethylase may play a role during normal epithelial maintenance. In contrast to the normal or bullectomized setting, direct injury to the OE has a more complex effect on the stem and progenitor populations, as all of the different cell types need to be regenerated during the reconstitution of the epithelium (Schwob et al., 1995; Schwob, 2002). For example, Sox2 becomes universal after MeBr lesion, and the sequential expression of *Ascl1*, *Neurog1*, and *NeuroD1* heralds the appearance of more differentiated neuronal progenitors and eventually neurons. As a consequence, time after lesion can be used to highlight particular stages in the neurogenic progression for purposes of comparison with the expression of LSD1. Given extensive coexpression of Ascl1 and LSD1 in unlesioned epithelium, we predicted that LSD1 expression should closely mirror that of Ascl1 after epithelial injury caused by the inhalation of MeBr gas. However, the temporal pattern of LSD1 expression was substantially more complicated than that of Ascl1 (Fig. 9). The day after lesion, LSD1 expression was largely undetectable. In tissue harvested 2–4 days postlesion, LSD1 stains the nuclei of HBCs, a cell population in which LSD1 is not found in the OE of unlesioned or bullectomized mice. In the case of the post-MeBr OE, double labeling with LSD1 and CK14, a marker of HBCs clearly shows LSD1(+) nuclei surrounded by cytoplasmic CK14 staining in cells directly opposed to the basal lamina as well as CK14(+) cells that have accumulated suprabasally (Fig. 9B – D). This staining pattern shifts dramatically back toward the normal pattern at later timepoints following lesion. At 5 days postlesion, LSD1 first begins to emerge in CK14(–) cells but has not completely disappeared from CK14(+) cells at this timepoint. At 7 days the balance has shifted further: there are more CK14 (–)/LSD1(+) cells. By 14 days postlesion, the expression of LSD1 is found almost exclusively in GBCs, reminiscent of the pattern from normal epithelium, although in a larger number of cells.

In the MeBr-lesioned OE, coREST-expressing cells are few and superficially located during the earliest stages in recovery (Fig. 10). At 4 days after lesion, numerous cells superficial to the monolayer of HBCs are weakly positive and a smaller percentage express coREST strongly that are also CK14(–) (Fig. 10D). At 5 days after lesion the coREST(+) cells are more heavily labeled and stand out more sharply by comparison with the coREST(–) ones; at this time, coREST labels primarily suprabasal and apical cells (Fig. 10E). While a rare coREST(+) cell will also be CK14(+), they are far less common than CK14(–)/coREST(+) cells. At 7 days after lesion, the overwhelming majority of coREST(+) cells remain CK14(–) and have begun to stratify in the middle of the epithelium (not shown). By 14 days after lesion the pattern of coREST labeling is similar to that of the normal epithelium: a band of coREST(+) cells is positioned roughly 1–2 cell layers above the basal lamina, and the intensity of labeling decreases in cells that are found at more apical positions in the epithelium (Fig. 10F).

DISCUSSION

We have isolated different stages in the differentiation of neurons from the GBC progenitors and subsequently analyzed and compared the transcriptomes for these various populations to that of Sus cells, to the olfactory mucosa as a whole, and among themselves. Four major findings emerge from the work described above. First, all stages in the neurogenic progression, from GBC_{MPP} to differentiated neuron, share a substantial portion of their transcriptomes and express greater than 2,100 genes in common. Second, the cells within the

GBC_{MPP} sort and the ones within the GBC_{INP}/immature neuron sort share greater than 1,200 genes that are unique to these progenitor cells. Third, the transcriptional profiles of *Neurog1-eGFP*-expressing cells are essentially identical whether isolated from the normal epithelium or isolated from the epithelium during the enhancement of neurogenesis by which the OE responds to bullectomy. Fourth, components of the transcriptional corepressor and chromatin modifying coREST complex, which are expressed in the GBC populations, are dynamically regulated after bullectomy and MeBr lesion. LSD1 is expressed in upstream GBCs, while coREST appears to be expressed in more downstream GBCs and immature neurons.

The findings reported above and conclusions derived from the resulting expression data depend primarily on the ability 1) to identify and purify reproducibly the populations of interest; 2) to establish meaningful and interpretable comparisons of those datasets; and 3) validate the genes identified as regulated in these various comparisons. With regard to purification, three different transgenic mouse strains that we and others have shown faithfully express the reporter in the appropriate cell types were used (Potter et al., 2001; Sammeta et al., 2007; Agustsson et al., 2009; Guo et al., 2010). As with all transgenic mice in which the endogenous gene locus includes a separate cistron encoding eGFP rather than fusion of the reporter protein to the targeted protein, perdurance (a.k.a. lingering expression) of eGFP into more downstream cell types becomes an issue (Aubert et al., 2003). While perdurance is potentially a problem when studying each population in isolation, we have minimized that concern by studying a progression of cell types as they transition successively through the various stages we obtained (Manglapus et al., 2004; Packard, 2010). Thus, we used analysis of overlapping and nonoverlapping genes that are regulated at each step of the progression as a means to identifying interesting targets for further validation and analysis. With regard to issues of interpretation, the conceptual framework established by previous genetic, functional, and expression studies mentioned above (including the characterization described here) has assigned unique and varied biological roles for the various cells populations examined in this study. As a result, the genes upregulated in these comparisons can be interpreted according to the proposed functional capacity of the various cells. For example, previous studies have placed Sox2-expressing GBCs upstream of *Neurog1*-expressing GBCs with OMP(+) neurons as the terminal stage in the differentiation of the postmitotic progeny of this lineage (Guo et al., 2010; Packard et al., 2011). As a consequence, meaningful hypotheses can be developed that are based on the genes identified in the different types of progenitors (by comparison with each other) and in the progenitors as compared to differentiated neurons. With regard to validation, published work in the field, our global analysis of the datasets, and the expression patterns of genes not previously examined in the OE all illustrate the usefulness of the microarray expression data. In contrast to previously described global expression data for OSNs (Sammeta et al., 2007), we performed all isolations and comparisons using three separate preparations of multiple animals per replicate (the number of animals per replicate was determined by the yield of RNA and varied from 2 to greater than 4) in an effort to highlight average population differences between given cell types that will be less susceptible to differences among animals or subtle differences in methodology. Beyond that, we also confirmed expected upregulation of numerous genes in several populations based on the literature (i.e., hundreds of ORs in the OMP populations, cell-type specific transcription factors in the progenitor populations, and axon guidance, growth, and sorting factors in immature neurons).

The finding that a large cohort of genes are common to the neurogenic lineage when all the different stages are compared to Sus cells suggests that these cell types may be using common pathways in slightly different ways to effect functional differences or that posttranscriptional regulation plays a significant role in functionally differentiating these

cells from one another. This finding actually parallels data described for a comprehensive analysis of the transcriptomes of various stages in the development and differentiation of the spermatocyte lineage (Johnston et al., 2008). This study described between 7,812 and 9,888 genes expressed at each of 10 distinct stages. The total number of unique genes expressed in the testes as a whole was 11,699—implying a significant amount of overlap in the cohort of genes expressed in the spermatocyte lineage. Our data, although not involving as many distinct stages, reveals a similar trend of lineage-specific overlap of upregulated genes. The shared genes for the neurogenic lineage also reinforce the data that describe Neurog1(+) GBCs and Ascl1(+) GBCs (the latter of which are included in the Sox2-GBC) as increasingly fate-committed neuronal progenitors (Murray et al., 2003; Guo et al., 2010). One would expect that an upstream fate choice would result in a large cohort of genes that are exclusively expressed in more downstream committed precursors for that lineage and distinguishes these cells from the nonneuronal lineage.

Within the neurogenic lineage, we also found that populations enriched for two classes of GBCs—Sox2-GBCs and Neurog1-Ctrl—share more than 1,400 significantly upregulated genes. Many cells among the GBC population share the properties of a highly activated, highly mitotic population (Caggiano et al., 1994; Huard and Schwob, 1995; Guo et al., 2010). Nonetheless, analysis in the embryo and adult has suggested that GBCs that express Sox2 vs. Neurog1 (the subsets isolated here) represent two functionally distinct populations (Cau et al., 1997, 2002; Manglapus et al., 2004; Kawauchi et al., 2005; Guo et al., 2010). Based on transcription factor expression, retroviral lineage tracing, retroviral gene transduction, and transplant analysis, among the Sox2(+) GBCs are ones that are actively multipotent, which may or may not be true of the subset that are also Ascl1(+) (Huard et al., 1998; Goldstein et al., 1998; Chen et al., 2004; Manglapus et al., 2004; Guo et al., 2010; Packard et al., 2011; Schwob lab, unpubl. data). Moreover, Sox2 expression by GBCs is largely unaffected by the absence of Ascl1 and thus not epistatic to it (Krolewski, Packard and Schwob, unpublished data). In contrast, the Neurog1(+) GBCs are a distinct group, epistatic to Ascl1, and later to emerge during embryogenesis or tissue regeneration than Sox2-expressing GBCs and thus likely to be downstream of them (Cau et al., 1997, 2002; Manglapus et al., 2004; Beites et al., 2005; Guo et al., 2010; Packard et al., 2011).

The current microarray data suggest that the apparent diversity of function does not manifest as dramatic transcriptional differences, and that other genes may show more graded differences in transcription or alternate methods of regulation as compared with the sharper differences seen in the foregoing stage-specific transcription factors. Indeed, the extensive transcriptional overlap could be explained in part by related findings comparing embryonic stem cells with tissue-specific progenitor and differentiated cell types (Golan-Mashiach et al., 2005). In this case, ES cells and adult stem cells express a general set of genes representative of the different cell types, which these progenitors are capable of differentiating into, and then gradually decrease the expression of “stemness” genes as they differentiate. An analogous kind of regulation in the OE could explain the large number of commonly upregulated genes found throughout the neurogenic progression, while the uniquely expressed genes for the upstream GBCs could be those that are being turned off as these cells differentiate.

The use of bullectomy to activate and expand the population of Neurog1-eGFP(+) cells also highlights a characteristic that would be expected of neuronal-committed, fate-restricted proliferating progenitors. The extremely small number of genes that differ between cells that express this gene (i.e., cells of this type) in normal, unoperated OE (Neurog1-Ctrl) vs. the epithelium following bullectomy (Neurog1-OBX) indicates that their transcriptional profiles are essentially identical. Thus, the Neurog1(+) cells are acting as an “effector” population that is minimally responsive to the lesion environment per se, but rather increases

neuronal numbers in the same manner as during normal maintenance of the OE. In contrast, a much greater degree of transcriptional diversity was observed between OMP-GFP(+) cells harvested 3 weeks after bullectomy (OMP-OBX) as compared to those from the epithelium of an animal with intact bulbs, i.e., the normal epithelium (OMP-Ctrl). Our data complement and parallel previous findings from other laboratories (Shetty et al., 2005; Getchell et al., 2006). That the OMP(+) neurons in the bullectomized setting have more recently transitioned to maturity and have a more abbreviated lifespan by comparison with the OMP(+) neurons in the normal epithelium is key to the interpretation of that heterogeneity (Carr and Farbman, 1992; Schwob et al., 1992). Two explanations emerge from the differences in the history of the OMP(+) neurons in these two settings. Either the differences in transcriptional profile indicate that OMP expression encompasses different stages of differentiation, i.e., young OMP(+) neurons are functionally different from old OMP(+) neurons. Some evidence can be adduced for that notion (Schwartz et al., 1992; Ukhanov et al., 2010). Alternatively, the difference in transcriptome in the OMP(+) neurons with bullectomy may reflect their failure to contact their olfactory bulb target and acquire trophic support, and are preparing to die on account of that failure. Additional analysis will be needed to ascertain the relative contribution of these two possibilities.

How do the transcriptional differences between the different stages in the neurogenic progression come about? Obviously, differential expression of major transcription factors, such as Sox2, Ascl1, Neurog1, and NeuroD1, will have major consequences for the pattern of gene expression at each of these stages. However, epigenetic regulators and chromatin modifications are likely responsible for an additional layer of control by modulating the accessibility of genes involved in maintaining a progenitor phenotype or initiating a neuronal differentiation program in concert with the canonical neurogenic transcription factors. We show in this work that two known regulators of chromatin accessibility—histone and DNA methylation—are expressed in GBCs and immature neurons and are dynamically regulated in response to methyl bromide lesion (Wang et al., 2007, 2009a; Cunliffe, 2008). Although coREST and LSD1 are commonly found in the same complex and generally contribute to gene repression by coordinated demethylation of histones (Qureshi et al., 2010), the expression data shown here indicate that their expression need not be tightly linked; LSD1 is prominent in stem and progenitor cells, while coREST is primarily found in immature neurons. Another component of the complex, HDAC2, and its distribution have previously been described in adult OE (MacDonald et al., 2005). The published data along with the findings reported here for coREST suggest that coREST and HDAC2 are expressed in the OE in a primarily LSD1-independent complex. Other studies of pituitary organogenesis and regulation global demethylation have suggested functions for LSD1 independent of coREST-containing complexes (Wang et al., 2007, 2009a). Likewise, the finding that LSD1 expression is dynamically expressed in activated HBCs early after MeBr lesion and then restricted to GBCs at later timepoints after lesion (and in normal OE) places the demethylase at crucial regulatory points for the proliferation and possibly cell fate choice of HBCs and GBCs. It will be of obvious importance to ascertain whether the coREST complex plays a functional role in the chromatin-mediated silencing that apparently contributes to the regulation of odorant receptor genes (Magklara et al., 2011).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors thank Nikolai Schnittke and Dr. Woochan Jang for assistance with FACS and RNA preps, Allen Parmelee and Stephen Kwok for technical assistance with FACS, and Po Kwok Tse for excellent technical assistance. The authors also thank the Schwob and Gill lab members for helpful discussions about the data.

Grant sponsor: National Institutes of Health (NIH)R21 DC008568, R01 DC002167 (to J.E.S.), F30 DC010276 (to R.C.K.) to the TUSM/T-NEMC Center for Neuroscience Research (P30 NS047243).

LITERATURE CITED

- Agustsson T, Nilsson M, Henriksson G, Arnelo U, Juto JE, Lundell L. Treatment of postoperative esophagorespiratory fistulas with dual self-expanding metal stents. *World J Surg.* 2009; 33:1224–1228. [PubMed: 19363689]
- Andres ME, Burger C, Peral-Rubio MJ, Battaglioli E, Anderson ME, Grimes J, Dallman J, Ballas N, Mandel G. CoR-EST: a functional corepressor required for regulation of neural-specific gene expression. *Proc Natl Acad Sci U S A.* 1999; 96:9873–9878. [PubMed: 10449787]
- Aubert J, Stavridis MP, Tweedie S, O'Reilly M, Vierlinger K, Li M, Ghazal P, Pratt T, Mason JO, Roy D, Smith A. Screening for mammalian neural genes via fluorescence-activated cell sorter purification of neural precursors from Sox1-gfp knock-in mice. *Proc Natl Acad Sci U S A.* 2003; 100(Suppl 1):11836–11841. [PubMed: 12923295]
- Baker H, Grillo M, Margolis FL. Biochemical and immunocytochemical characterization of olfactory marker protein in the rodent central nervous system. *J Comp Neurol.* 1989; 285:246–261. [PubMed: 2760264]
- Barraud P, He X, Zhao C, Ibanez C, Raha-Chowdhury R, Caldwell MA, Franklin RJ. Contrasting effects of basic fibroblast growth factor and epidermal growth factor on mouse neonatal olfactory mucosa cells. *Eur J Neurosci.* 2007; 26:3345–3357. [PubMed: 18088275]
- Beites CL, Kawauchi S, Crocker CE, Calof AL. Identification and molecular regulation of neural stem cells in the olfactory epithelium. *Exp Cell Res.* 2005; 306:309–316. [PubMed: 15925585]
- Buck L, Axel R. A novel multigene family may encode odorant receptors: a molecular basis for odor recognition. *Cell.* 1991; 65:175–187. [PubMed: 1840504]
- Caggiano M, Kauer JS, Hunter DD. Globose basal cells are neuronal progenitors in the olfactory epithelium: a lineage analysis using a replication-incompetent retrovirus. *Neuron.* 1994; 13:339–352. [PubMed: 8060615]
- Capo-Chichi CD, Rula ME, Smedberg JL, Vanderveer L, Parmacek MS, Morrissey EE, Godwin AK, Xu XX. Perception of differentiation cues by GATA factors in primitive endoderm lineage determination of mouse embryonic stem cells. *Dev Biol.* 2005; 286:574–586. [PubMed: 16162334]
- Carr VM, Farbman AI. Ablation of the olfactory bulb upregulates the rate of neurogenesis and induces precocious cell death in olfactory epithelium. *Exp Neurol.* 1992; 115:55–59. [PubMed: 1728573]
- Carter LA, MacDonald JL, Roskams AJ. Olfactory horizontal basal cells demonstrate a conserved multipotent progenitor phenotype. *J Neurosci.* 2004; 24:5670–5683. [PubMed: 15215289]
- Cau E, Gradwohl G, Fode C, Guillemot F. Mash1 activates a cascade of bHLH regulators in olfactory neuron progenitors. *Development.* 1997; 124:1611–1621. [PubMed: 9108377]
- Cau E, Casarosa S, Guillemot F. Mash1 and Ngn1 control distinct steps of determination and differentiation in the olfactory sensory neuron lineage. *Development.* 2002; 129:1871–1880. [PubMed: 11934853]
- Chen X, Fang H, Schwob JE. Multipotency of purified, transplanted globose basal cells in olfactory epithelium. *J Comp Neurol.* 2004; 469:457–474. [PubMed: 14755529]
- Chuah MI, Farbman AI. Olfactory bulb increases marker protein in olfactory receptor cells. *J Neurosci.* 1983; 3:2197–2205. [PubMed: 6355401]
- Cunliffe VT. Eloquent silence: developmental functions of Class I histone deacetylases. *Curr Opin Genet Dev.* 2008; 18:404–410. [PubMed: 18929655]
- DeHamer MK, Guevara JL, Hannon K, Olwin BB, Calof AL. Genesis of olfactory receptor neurons in vitro: regulation of progenitor cell divisions by fibroblast growth factors. *Neuron.* 1994; 13:1083–1097. [PubMed: 7946347]

- Dennis G Jr, Sherman BT, Hosack DA, Yang J, Gao W, Lane HC, Lempicki RA. DAVID: database for annotation, visualization, and integrated discovery. *Genome Biol.* 2003; 4:P3. [PubMed: 12734009]
- Du P, Kibbe WA, Lin SM. lumi: a pipeline for processing Illumina microarray. *Bioinformatics.* 2008; 24:1547–1548. [PubMed: 18467348]
- Ellis P, Fagan BM, Magness ST, Hutton S, Taranova O, Hayashi S, McMahon A, Rao M, Pevny L. SOX2, a persistent marker for multipotential neural stem cells derived from embryonic stem cells, the embryo or the adult. *Dev Neurosci.* 2004; 26:148–165. [PubMed: 15711057]
- Gentleman RC, Carey VJ, Bates DM, Bolstad B, Dettling M, Dudoit S, Ellis B, Gautier L, Ge Y, Gentry J, Hornik K, Hothorn T, Huber W, Iacus S, Irizarry R, Leisch F, Li C, Maechler M, Rossini AJ, Sawitzki G, Smith C, Smyth G, Tierney L, Yang JY, Zhang J. Bioconductor: open software development for computational biology and bioinformatics. *Genome Biol.* 2004; 5:R80. [PubMed: 15461798]
- Getchell ML, Li H, Vaishnav RA, Borders AS, Witta J, Subhedar N, de Villiers W, Stromberg AJ, Getchell TV. Temporal gene expression profiles of target-ablated olfactory epithelium in mice with disrupted expression of scavenger receptor A: impact on macrophages. *Physiol Genomics.* 2006; 27:245–263. [PubMed: 16882882]
- Golan-Mashiach M, Dazard JE, Gerecht-Nir S, Amariglio N, Fisher T, Jacob-Hirsch J, Bielorai B, Osenberg S, Barad O, Getz G, Toren A, Rechavi G, Itskovitz-Eldor J, Domany E, Givol D. Design principle of gene expression used by human stem cells: implication for pluripotency. *FASEB J.* 2005; 19:147–149. [PubMed: 15498892]
- Gong S, Zheng C, Doughty ML, Losos K, Didkovsky N, Schambra UB, Nowak NJ, Joyner A, Leblanc G, Hatten ME, Heintz N. A gene expression atlas of the central nervous system based on bacterial artificial chromosomes. *Nature.* 2003; 425:917–925. [PubMed: 14586460]
- Graziadei GA, Graziadei PP. Neurogenesis and neuron regeneration in the olfactory system of mammals. II. Degeneration and reconstitution of the olfactory sensory neurons after axotomy. *J Neurocytol.* 1979; 8:197–213. [PubMed: 469573]
- Guo Z, Packard A, Krolewski RC, Manglapus GL, Harris M, Schwob JE. Expression of Pax6 and Sox2 in adult olfactory epithelium. *J Comp Neurol.* 2010; 518:4395–4418. [PubMed: 20852734]
- Harms C, Albrecht K, Harms U, Seidel K, Hauck L, Baldinger T, Hubner D, Kronenberg G, An J, Ruscher K, Meisel A, Dirnagl U, von Harsdorf R, Endres M, Hortnagl H. Phosphatidylinositol 3-Akt-kinase-dependent phosphorylation of p21(Waf1/Cip1) as a novel mechanism of neuroprotection by glucocorticoids. *J Neurosci.* 2007; 27:4562–4571. [PubMed: 17460069]
- Holbrook EH, Szumowski KE, Schwob JE. An immunochemical, ultrastructural, and developmental characterization of the horizontal basal cells of rat olfactory epithelium. *J Comp Neurol.* 1995; 363:129–146. [PubMed: 8682932]
- Hsu P, Yu F, Feron F, Pickles JO, Sneesby K, Mackay-Sim A. Basic fibroblast growth factor and fibroblast growth factor receptors in adult olfactory epithelium. *Brain Res.* 2001; 896:188–197. [PubMed: 11277992]
- Huang, da W.; Sherman, BT.; Lempicki, RA. Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc.* 2009; 4:44–57. [PubMed: 19131956]
- Huard JM, Schwob JE. Cell cycle of globose basal cells in rat olfactory epithelium. *Dev Dyn.* 1995; 203:17–26. [PubMed: 7647371]
- Itoh Y, Masuyama N, Nakayama K, Nakayama KI, Gotoh Y. The cyclin-dependent kinase inhibitors p57 and p27 regulate neuronal migration in the developing mouse neocortex. *J Biol Chem.* 2007; 282:390–396. [PubMed: 17092932]
- Iwai N, Zhou Z, Roop DR, Behringer RR. Horizontal basal cells are multipotent progenitors in normal and injured adult olfactory epithelium. *Stem Cells.* 2008; 26:1298–1306. [PubMed: 18308944]
- Iwema CL, Fang H, Kurtz DB, Youngentob SL, Schwob JE. Odorant receptor expression patterns are restored in lesion-recovered rat olfactory epithelium. *J Neurosci.* 2004; 24:356–369. [PubMed: 14724234]
- Jang W, Kim KP, Schwob JE. Nonintegrin laminin receptor precursor protein is expressed on olfactory stem and progenitor cells. *J Comp Neurol.* 2007; 502:367–381. [PubMed: 17366606]

- Johnston DS, Wright WW, Dicandoloro P, Wilson E, Kopf GS, Jelinsky SA. Stage-specific gene expression is a fundamental characteristic of rat spermatogenic cells and Sertoli cells. *Proc Natl Acad Sci U S A*. 2008; 105:8315–8320. [PubMed: 18544648]
- Kawauchi S, Shou J, Santos R, Hebert JM, McConnell SK, Mason I, Calof AL. Fgf8 expression defines a morphogenetic center required for olfactory neurogenesis and nasal cavity development in the mouse. *Development*. 2005; 132:5211–5223. [PubMed: 16267092]
- Keller; Margolis, Keller A.; Margolis, FL. Immunological studies of the rat olfactory marker protein. *J Neurochem*. 1975; 24:1101–1106. [PubMed: 805214]
- Kwon HJ, Koo JH, Zufall F, Leinders-Zufall T, Margolis FL. Ca extrusion by NCX is compromised in olfactory sensory neurons of OMP mice. *PLoS One*. 2009; 4:e4260. [PubMed: 19165324]
- Lee MG, Wynder C, Cooch N, Shiekhatter R. An essential role for CoREST in nucleosomal histone 3 lysine 4 demethylation. *Nature*. 2005; 437:432–435. [PubMed: 16079794]
- Legrier ME, Ducray A, Propper A, Chao M, Kastner A. Cell cycle regulation during mouse olfactory neurogenesis. *Cell Growth Differ*. 2001; 12:591–601. [PubMed: 11751454]
- Leung CT, Coulombe PA, Reed RR. Contribution of olfactory neural stem cells to tissue maintenance and regeneration. *Nat Neurosci*. 2007; 10:720–726. [PubMed: 17468753]
- MacDonald JL, Gin CS, Roskams AJ. Stage-specific induction of DNA methyltransferases in olfactory receptor neuron development. *Dev Biol*. 2005; 288:461–473. [PubMed: 16288735]
- Magklara A, Yen A, Colquitt BM, Clowney EJ, Allen W, Markenscoff-Papadimitriou E, Evans ZA, Kheradpour P, Mountoufaris G, Carey C, Barnea G, Kellis M, Lomvardas S. An epigenetic signature for monoallelic olfactory receptor expression. *Cell*. 2011; 145:555–570. [PubMed: 21529909]
- Manglapus GL, Youngentob SL, Schwob JE. Expression patterns of basic helix-loop-helix transcription factors define subsets of olfactory progenitor cells. *J Comp Neurol*. 2004; 479:216–233. [PubMed: 15452857]
- McClintock TS. Achieving singularity in mammalian odorant receptor gene choice. *Chem Senses*. 2010; 35:447–457. [PubMed: 20460312]
- Meiri KF, Bickerstaff LE, Schwob JE. Monoclonal antibodies show that kinase C phosphorylation of GAP-43 during axonogenesis is both spatially and temporally restricted in vivo. *J Cell Biol*. 1991; 112:991–1005. [PubMed: 1705561]
- Metzger E, Wissmann M, Yin N, Muller JM, Schneider R, Peters AH, Gunther T, Buettner R, Schule R. LSD1 demethylates repressive histone marks to promote androgen-receptor-dependent transcription. *Nature*. 2005; 437:436–439. [PubMed: 16079795]
- Miyamichi K, Serizawa S, Kimura HM, Sakano H. Continuous and overlapping expression domains of odorant receptor genes in the olfactory epithelium determine the dorsal/ventral positioning of glomeruli in the olfactory bulb. *J Neurosci*. 2005; 25:3586–3592. [PubMed: 15814789]
- Monti Graziadei GA. Experimental studies on the olfactory marker protein. III. The olfactory marker protein in the olfactory neuroepithelium lacking connections with the forebrain. *Brain Res*. 1983; 262:303–308. [PubMed: 6839159]
- Murray RC, Navi D, Fesenko J, Lander AD, Calof AL. Widespread defects in the primary olfactory pathway caused by loss of Mash1 function. *J Neurosci*. 2003; 23:1769–1780. [PubMed: 12629181]
- Packard, A. PhD dissertation. Boston, MA: Tufts University; 2010. Regulation of neurogenesis in the adult olfactory epithelium: analysis of Pax6, Sox2, NeuroD1 and stem cell activation.
- Potter SM, Zheng C, Koos DS, Feinstein P, Fraser SE, Mombaerts P. Structure and emergence of specific olfactory glomeruli in the mouse. *J Neurosci*. 2001; 21:9713–9723. [PubMed: 11739580]
- Qureshi IA, Gokhan S, Mehler MF. REST and CoREST are transcriptional and epigenetic regulators of seminal neural fate decisions. *Cell Cycle*. 2010; 9:4477–4486. [PubMed: 21088488]
- Reich M, Liefeld T, Gould J, Lerner J, Tamayo P, Mesirov JP. GenePattern 2.0. *Nat Genet*. 2006; 38:500–501. [PubMed: 16642009]
- Reisert J, Yau KW, Margolis FL. Olfactory marker protein modulates the cAMP kinetics of the odour-induced response in cilia of mouse olfactory receptor neurons. *J Physiol* 585. 2007; (Pt 3):731–740.

- Rodriguez-Gil DJ, Greer CA. Wnt/Frizzled family members mediate olfactory sensory neuron axon extension. *J Comp Neurol*. 2008; 511:301–317. [PubMed: 18803244]
- Saleque S, Kim J, Rooke HM, Orkin SH. Epigenetic regulation of hematopoietic differentiation by Gfi-1 and Gfi-1b is mediated by the cofactors CoREST and LSD1. *Mol Cell*. 2007; 27:562–572. [PubMed: 17707228]
- Sammata N, Yu TT, Bose SC, McClintock TS. Mouse olfactory sensory neurons express 10,000 genes. *J Comp Neurol*. 2007; 502:1138–1156. [PubMed: 17444493]
- Schwartz GA, Deutsch G, Gattley DM, Crandall JE. Glycoconjugates are stage- and position-specific cell surface molecules in the developing olfactory system. 2: Unique carbohydrate antigens are topographic markers for selective projection patterns of olfactory axons. *J Neurobiol*. 1992; 23:130–142. [PubMed: 1527523]
- Schwob JE. Neural regeneration and the peripheral olfactory system. *Anat Rec*. 2002; 269:33–49. [PubMed: 11891623]
- Schwob JE, Szumowski KE, Stasky AA. Olfactory sensory neurons are trophically dependent on the olfactory bulb for their prolonged survival. *J Neurosci*. 1992; 12:3896–3919. [PubMed: 1403089]
- Schwob JE, Youngentob SL, Mezza RC. Reconstitution of the rat olfactory epithelium after methyl bromide-induced lesion. *J Comp Neurol*. 1995; 359:15–37. [PubMed: 8557844]
- Shetty RS, Bose SC, Nickell MD, McIntyre JC, Hardin DH, Harris AM, McClintock TS. Transcriptional changes during neuronal death and replacement in the olfactory epithelium. *Mol Cell Neurosci*. 2005; 30:583–600. [PubMed: 16456926]
- Shi YJ, Matson C, Lan F, Iwase S, Baba T, Shi Y. Regulation of LSD1 histone demethylase activity by its associated factors. *Mol Cell*. 2005; 19:857–864. [PubMed: 16140033]
- Shou J, Rim PC, Calof AL. BMPs inhibit neurogenesis by a mechanism involving degradation of a transcription factor. *Nat Neurosci*. 1999; 2:339–345. [PubMed: 10204540]
- Shou J, Murray RC, Rim PC, Calof AL. Opposing effects of bone morphogenetic proteins on neuron production and survival in the olfactory receptor neuron lineage. *Development*. 2000; 127:5403–5413. [PubMed: 11076761]
- Smyth GK. Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mol Biol*. 2004; 3 Article3.
- Smyth GK, Michaud J, Scott HS. Use of within-array replicate spots for assessing differential expression in microarray experiments. *Bioinformatics*. 2005; 21:2067–2075. [PubMed: 15657102]
- Sun G, Alzayady K, Stewart R, Ye P, Yang S, Li W, Shi Y. Histone demethylase LSD1 regulates neural stem cell proliferation. *Mol Cell Biol*. 2010; 30:1997–2005. [PubMed: 20123967]
- Tucker ES, Lehtinen MK, Maynard T, Zirlinger M, Dulac C, Rawson N, Pevny L, Lamantia AS. Proliferative and transcriptional identity of distinct classes of neural precursors in the mammalian olfactory epithelium. *Development*. 2010; 137:2471–2481. [PubMed: 20573694]
- Ukhanov K, Corey EA, Brunert D, Klasen K, Ache BW. Inhibitory odorant signaling in mammalian olfactory receptor neurons. *J Neurophysiol*. 2010; 103:1114–1122. [PubMed: 20032232]
- Verhaagen J, Oestreicher AB, Gispén WH, Margolis FL. The expression of the growth associated protein B50/GAP-43 in the olfactory system of neonatal and adult rats. *J Neurosci*. 1989; 9:683–691. [PubMed: 2918383]
- Verhaagen J, Oestreicher AB, Grillo M, Khew-Goodall Y-S, Gispén WH, Margolis FL. Neuroplasticity in the olfactory system: differential effects of central and peripheral lesions on the primary olfactory pathway on the expression of B-50/GAP43 and the olfactory marker protein. *J Neurosci Res*. 1990; 26:31–44. [PubMed: 2141653]
- Vinals F, Reiriz J, Ambrosio S, Bartrons R, Rosa JL, Ventura F. BMP-2 decreases Mash1 stability by increasing Id1 expression. *EMBO J*. 2004; 23:3527–3537. [PubMed: 15318167]
- Wang J, Scully K, Zhu X, Cai L, Zhang J, Prefontaine GG, Kronen A, Ohgi KA, Zhu P, Garcia-Bassets I, Liu F, Taylor H, Lozach J, Jayes FL, Korach KS, Glass CK, Fu XD, Rosenfeld MG. Opposing LSD1 complexes function in developmental gene activation and repression programmes. *Nature*. 2007; 446:882–887. [PubMed: 17392792]
- Wang J, Hevi S, Kurash JK, Lei H, Gay F, Bajko J, Su H, Sun W, Chang H, Xu G, Gaudet F, Li E, Chen T. The lysine demethylase LSD1 (KDM1) is required for maintenance of global DNA methylation. *Nat Genet*. 2009a; 41:125–129. [PubMed: 19098913]

- Wang Y, Zhang H, Chen Y, Sun Y, Yang F, Yu W, Liang J, Sun L, Yang X, Shi L, Li R, Li Y, Zhang Y, Li Q, Yi X, Shang Y. LSD1 is a subunit of the NuRD complex and targets the metastasis programs in breast cancer. *Cell*. 2009b; 138:660–672. [PubMed: 19703393]
- Wilkie AO. The molecular basis of genetic dominance. *J Med Genet*. 1994; 31:89–98. [PubMed: 8182727]
- Wu HH, Ivkovic S, Murray RC, Jaramillo S, Lyons KM, Johnson JE, Calof AL. Autoregulation of neurogenesis by GDF11. *Neuron*. 2003; 37:197–207. [PubMed: 12546816]
- Youngentob SL, Margolis FL. OMP gene deletion causes an elevation in behavioral threshold sensitivity. *Neuroreport*. 1999; 10:15–9. [PubMed: 10094125]
- Youngentob SL, Margolis FL, Youngentob LM. OMP gene deletion results in an alteration in odorant quality perception. *Behav Neurosci*. 2001; 115:626–631. [PubMed: 11439451]
- Youngentob SL, Kent PF, Margolis FL. OMP gene deletion results in an alteration in odorant-induced mucosal activity patterns. *J Neurophysiol*. 2003; 90:3864–3873. [PubMed: 12917392]

Figure 1.

Tissue expression and FACS profiles in the neurogenic progression. Tissues harvested from normal *Neurog1-eGFP* (A,B,E,F,I,J) and Δ *OMP-eGFP* (C,D,G,H,K,L) mice euthanized 3 weeks post-bulbectomy were stained for various antigens to illustrate the different stages from which RNA was collected for microarray analysis and the resulting FACS profiles. (A–D) The expression of the eGFP transgene relative to the targeted locus is shown; standard fluorescence microscopy of coronal sections. A,B: eGFP(+) cells in normal *Neurog1-eGFP* mice encompass the pool of immediate neuronal precursors among the GBCs as well as immature neurons. A: Tissue sections from normal adult *Neurog1-eGFP* mice stained for Neurog1 and eGFP demonstrate that eGFP is expressed in basal cells and immature neurons. The asterisks indicate examples of Neurog1(+)/eGFP(+) cells. 78% of Neurog1(+) cells are also eGFP(+) in unoperated, normal adult OE. The arrow illustrates an example of a Neurog1(+)/eGFP(–) cell and the double arrow indicates a pair of them. While a minority, cells of this type can be found with regularity, presumably due to the fact that GFP has not yet accumulated to detectable levels in them. Many of the eGFP(+) cells are immature neurons, as shown by their lack of Neurog1 expression, more apical position in the epithelium, and the dendrite extending from the cell body toward the surface. B: Sections from transgenic animals killed 3 weeks after unilateral bulbectomy show a dramatic increase in eGFP expression and an expansion apically of the band of Neurog1(+) cells, due to persistently elevated neurogenesis. In the bulbectomized setting, 70% of Neurog1(+) cells are eGFP(+). C,D: eGFP(+) cells in the Δ *OMP-eGFP* mice encompass the pool of mature neurons in normal epithelium (C) and a population of maturing neurons in bulbectomized epithelium (D). C: Colabeling of OMP and GFP show that the vast majority of OMP(+) mature neurons are eGFP(+). D: After bulbectomy, the transgene maintains fidelity to OMP(+) neurons after bulbectomy. E–H: The expression of the eGFP transgene relative to NCAM, which marks all neurons, and immature neurons most intensely; confocal microscopy of coronal sections. E: In normal *Neurog1-eGFP* animals, eGFP is detectable in

some immature neurons, as shown by costaining with NCAM. F: In *Neurog1-eGFP* mice 3 weeks after bulbectomy, the population of eGFP(+) neurons is expanded, in keeping with the persistent acceleration of neurogenesis. G: In normal $\Delta OMP-eGFP$ mice, there is minimal overlap of eGFP and the intensely NCAM(+) immature neurons. H: In bulbectomized $\Delta OMP-eGFP$ mice the population of eGFP(+) neurons is smaller and remains largely but not completely complementary to the intensely NCAM(+) immature neurons. Due to the abbreviated lifespan of neurons born after bulbectomy, the eGFP(+) cells in these animals, represent a smaller population of newly maturing neurons rather than a fully mature population. I–L: FL1 (eGFP) vs. FL4 (no stain) FACS profiles of dissociated cells from the corresponding conditions. The percentage of eGFP(+) cells in each condition is listed in or near the outlined region. The percentages are predicted by and reflect the prevalence of eGFP(+) cells observed in the tissue for both transgenic strains and both conditions (normal vs. bulbectomy). Arrowheads indicate basal lamina. A magenta-green version of the photomicrographs is available online as Supporting Figure 1. Scale bars = 25 mm in D (applies to A–D); 25 μ m in H (applies to E–H).

Figure 2.

Isolation of eGFP(+) GBCs from $\Delta Sox2-eGFP$ mice for expression profiling. **A–C:** $\Delta Sox2-eGFP$ mice express GFP in GBCs, HBCs, and Sus cells. Coronal sections of OE were stained to demonstrate the cell types expressing GFP. **A:** eGFP is expressed in two distinct populations of basal cells, HBCs as shown by colabeling with CK14, and GBCs based on exclusion from CK14 (arrows). **B:** GBCs make up the vast majority of proliferating basal cells, and eGFP colocalizes with Ki67, a marker of proliferation, in these cells (arrows). **C:** GFP is expressed in GBCs but not the immature neurons downstream as shown by exclusion from NCAM(+) cells. **D–F:** Strategy for enriching eGFP(+) GBCs from the $\Delta Sox2-eGFP$

mice by FACS. $\Delta Sox2-eGFP$ animals were dissociated according to a nonstandard protocol to enrich for GBCs and separate them from HBCs and Sus cells. In this case, the olfactory epithelium was subjected to mechanical disruption without benefit of proteolytic digestion, and the released cells were stained with anti-CD54 (to label HBCs and remove them from the sort) and anti-E-cadherin (to label Sus cells and remove them from the sort). The use of mechanical disruption to isolate a population enriched for GBCs exploits the complex architecture and the junctional attachments of Sus cells and HBCs (as compared to the round, simple GBCs), as well as preserving the cell surface antigens CD54 and E-cadherin for sorting purposes. D: Scatter-gating was used to separate larger and more optically complex cells—characteristic of Sus cells and HBCs—from small, optically simple cells—a property typical of GBCs (the area of the profile outlined in white); cells falling within the indicated gate were then subject to sorting on the basis of GFP and cell surface labeling with the two surface markers in E,F. E,F: FACS profiles of dissociated olfactory epithelial cells from $\Delta Sox2-eGFP$ mice stained concurrently for the surface markers CD54 (which labels HBCs) and E-cadherin (which labels Sus cells), detected with two separate allophycocyanin (APC)-labeled secondary antibodies in the FL6 channel. E: The populations enriched in Sus cells (areas outlined in orange) and in HBCs (outlined in blue) are shifted well away from the unstained populations, as confirmed by post-hoc analysis (not shown). F: Following scatter-gating as indicated in D, the population of low-intensity eGFP(+) cells that do not stain for CD54 and E-cadherin was isolated and found, on post-hoc analysis, to be largely free of intact Sus cells or HBCs (see text). Note that with the mechanical dissociation protocol a large percentage of the Sus cells show up with decreased eGFP intensity (orange box) relative to the staining in vivo or the FACS profile after enzymatic digestion, presumably because these cells have leaked some soluble GFP as a result of the dissociation. A magenta-green version of the photomicrographs is available online as Supporting Figure 2. Scale bar = 25 μ m in C (applies to A–C).

Sample relations based on 7482 genes with $sd/mean > 0.1$ **Figure 3.**

Dendrogram of sample relations. A global comparison of gene expression profiles across the individual samples highlights the degree to which the replicates in each type of cell-sort cluster together and the similarity between that group and all of the others. The samples enriched for different types of GBCs clustered together (blue and purple), while the samples enriched for olfactory sensory neurons clustered together (red and black). The control cell type from the olfactory epithelium that is not part of the neurogenic progression, sustentacular cells (SUS), did not cluster closely with any of the samples from the neurogenic progression. For the two kinds of Neurog1 samples (NgnCtrl, which derive from unoperated Neurog1-eGFP mice, and NgnOBX, which derive from epithelium ipsilateral to the lesion in unilaterally bulbectomized Neurog1-eGFP mice), the variation between replicates within a group was as great as the variation between control and bulb-ablated groups. The replicates of all other cell types segregate together in relative isolation. Signal intensity values from each bead array replicate were normalized and background corrected in Illumina Genome Studio. Sample relations based on >7,000 of the most highly regulated genes were determined using the lumi package for R (Du et al., 2008). mREF designates mouse reference RNA (Stratagene), SUS designates sustentacular cells, NgnCtrl designates Neurog1-Ctrl (unoperated *Neurog1-eGFP* mice), NgnOBX designates Neurog1-OBX

(*Neurogl-eGFP* mice harvested 3 weeks post-unilateral bulbectomy), OMP designates OMP-Ctrl (unoperated heterozygote $\Delta OMP-eGFP$ mice), OMPGFP_OBX designates OMP-OBX (heterozygous $\Delta OMP-eGFP$ mice harvested 3 weeks post-bulbectomy), Normal_OM, normal olfactory mucosa.

Figure 4.

Overlapping and distinct upregulated genes in the neurogenic progression. Sox2-GBC, Neurog1-Ctrl, and OMP-Ctrl expression data were individually compared to the gene profile for Sus cells. The lists of significantly upregulated genes for each pairwise comparison were used to generate a Venn diagram that depicts the exact number of genes unique to, or shared among, the various conditions. The area for each region is roughly proportional to the number of genes in that area to give a representation of the overlap between different datasets. The red area represents genes uniquely upregulated in Sox2-GBCs (see Materials and Methods and the Results sections for a description of the FACS protocol used to

generate this set of cells) but not upregulated in Neurog1-Ctrl (from unoperated *Neurog1-eGFP* mice) or OMP-Ctrl (from unoperated heterozygote $\Delta OMP-eGFP$ mice). Green represents those unique to Neurog1-Ctrl, light blue represents those unique to OMP-Ctrl. Overlapping gene sets are depicted as areas of different colors: orange designates the number of genes that are upregulated in both Sox2-GBC and Neurog1-Ctrl, yellow represents those upregulated in both Neurog1-Ctrl and OMP-Ctrl, dark blue represents those upregulated in both Sox2-GBC and OMP-Ctrl. The central purple area represents those genes upregulated in all three conditions.

Figure 5.

coREST complex components are regulated as differentiation proceeds. A list of proteins known to interact with or participate in the coREST corepressor complex was compiled from the literature (Lee et al., 2005; Metzger et al., 2005; Shi et al., 2005; Saleque et al., 2007; Wang et al., 2007, 2009a,b; Cunliffe, 2008). The set of genes that was significantly upregulated at one or more stages within the neurogenic progression (as compared to Sus cells) was interrogated for inclusion of putative coREST components. Six of the coREST-associated genes were significantly upregulated in Sox2-GBC, Neurog1-Ctrl, or Neurog1-OBX samples. The normalized expression values for samples in the neurogenic progression were grouped by gene and plotted as a bar graph, which clearly show that the coREST complex components are upregulated in progenitor cells and decrease in expression as differentiation proceeds.

Figure 6.

LSD1 is expressed in GBCs and excluded from mature OSNs. Tissue sections from normal adult olfactory epithelium were examined by triple-label immunohistochemistry. **A:** LSD1 expression is detected in the nuclei of basal cells (arrows) and occasional apical cells (arrows with asterisks). **B:** Staining for CK14, a marker of HBCs. **C:** Staining for NCAM, which is detectable only on neurons and not GBCs (Packard et al., 2011). **D:** Merged image for all three markers demonstrates that LSD1(+) cells near the basal lamina are NCAM(−) and CK14(−), indicating that these are GBCs. **E:** The overlap of GFP and LSD1 staining in the *Neurog1-eGFP* mice is consistent with the expression profile. However, several of the

LSD1(+) cells are not colabeled by detectable expression of eGFP and sit below the band of GBCs and neurons that stain with anti-GFP (arrows). **F**: Likewise, the complete absence of eGFP(+)/LSD1(+) cells in the $\Delta OMP-eGFP$ mice also fits with the gene expression data, and substantiates the fall-off in expression as the sensory neurons mature. Arrowheads mark the basal lamina. Scale bars = 25 μm in D (applies to A–C); 25 μm in F (applies to E).

Figure 7.

coREST is expressed in more mature cells than LSD1. coREST protein is expressed in eGFP(+) cells in the *Neurog1-eGFP* mice but extends beyond that stage into the early phase of OMP expression. **A,B:** coREST labeled nuclei in unlesioned, control epithelium (Control) are confined to the deeper part of the neuronal zone of the OE. Many coREST(+) cells in control *Neurog1-eGFP* mice are also eGFP(+). However, some coREST(+) neurons, which are a bit more lightly labeled, are found apical to the layer of eGFP(+) cells, suggesting that expression extends beyond the time that LSD1 expression is no longer detectable (cf., Fig. 6E,F). **C,D:** The population of coREST(+) cells is dramatically expanded along with

eGFP(+) population in the bulsectomized *Neurog1-eGFP* mice. However, a substantial population of coREST(+)/eGFP(−) neurons are found above the belt formed by the Neurog1-eGFP(+) cells; although more lightly labeled, these are evident by comparison with the lack of nuclear staining in the Sus cells in the most apical layer of the epithelium. **E,F:** In complementary fashion, coREST(+) neurons are concentrated below the eGFP(+) neurons in the bulsectomized *ΔOMP-eGFP* mice. However, at least some of the eGFP(+) neurons show low but detectable (relative to the absence of staining in the apically located Sus cells) levels of coREST (arrows), which serves as a further indication that coREST expression remains detectable in cells that are further along in the neurogenic progression than is the case for LSD1. Arrowheads mark the basal lamina. Scale bar = 25 μm in F (applies to all).

Figure 8.

LSD1 is coexpressed with Ascl1 and Sox2 in GBCs. Tissue sections from normal animals were stained for LSD1 and Ascl1 (A–C) or LSD1 and Sox2 (D–F). A–C: The vast majority of Ascl1(+) (a.k.a. Mash1) cells are also LSD1(+) as seen in the merged image (B) of LSD1 and Ascl1 staining. Arrows indicate double-labeled cells. However, some of the LSD1(+) cells lack detectable Ascl1 (arrows with asterisks). D–F: Many LSD1(+) GBCs are also Sox2(+) as seen in the merged image (E). Arrows indicate colocalization. Arrowheads mark the basal lamina. A magenta-green version of the photomicrographs is available online as Supporting Figure 8. Scale bar = 25 μ m in F (applies to all).

Figure 9.

LSD1 is first expressed in HBCs followed by GBCs during recovery after MeBr lesion. Tissue sections from animals sacrificed at the indicated times after MeBr lesion were stained for LSD1 to determine the dynamics of the expression in relationship to HBCs, identified by CK14 staining. All panels are counterstained with Hoechst 33258 and insets show areas at higher magnification without the nuclear stain. **A:** Very few LSD1(+) cells are present at 1 day after lesion. **B-D:** At 2, 3, and 4 days after lesion, LSD1 is expressed in many CK14(+) cells. Asterisks indicate CK14(-)/LSD1(-) cells. White triangles indicate examples of CK14(+)/LSD1(+) cells. **E,F:** At later timepoints after lesion, CK14(-)/LSD1(+) GBCs reappear (arrows) and the majority of CK14(+) cells are LSD1(-) (brackets and insets). Bent arrow marks an apically situated LSD1(+) cell. Outlined arrowheads indicate the basal lamina. Scale bars = 25 μ m in F (applies to all); 10 μ m in inset of F (applies to all insets).

Figure 10.

coREST is strongly expressed at later timepoints after MeBr lesion. Tissue sections from animals sacrificed at increasing times after MeBr lesion were stained for coREST to determine the dynamics of the expression in relation to HBCs, identified by CK14 staining. All panels are costained with Hoechst 33258. **A–C:** coREST(+) cells are very rare at 1–3 days after MeBr lesion and, where expressed, the intensity of staining is weak. **D:** At 4 days after lesion, more superficially located CK14(+) cells are weakly labeled by anti-coREST antibody, and strongly coREST(+) on occasion. Their position is reminiscent of the CK14 cells that have downregulated p63 expression and detached from the basal lamina as they activate to multipotency after MeBr lesion (Packard et al., 2011). **E,F:** At 5 and 14 days after lesion, the vast majority of strongly labeled coREST(+) cells are CK14(–). Arrowheads indicate basal lamina. Scale bar = 25 μ m in F (applies to all).

TABLE 1

Antibodies Used in the Study

Primary antibody	Source and catalog number	Immunogen and preparation
Chicken α -GFP	AbCam ab13970	Recombinant full length protein. This anti-GFP antibody recognizes the enhanced form of GFP, and all of the fluorescent proteins made by this algae, including yellow FP.
Rabbit α -GFP	AbCam ab6556	Highly purified recombinant full length GFP made in <i>Escherichia coli</i>
Rat α -NCAM	Abcam ab19782 (H28–123)	Glycoprotein fraction from neonatal mouse brain
Goat α -OMP	Wako 544–10001	Multiple immunizations of rodent olfactory marker protein
Mouse α -Ngn1	Santa Cruz, sc-19231	20mer peptide sequence of mouse Ngn1 (ARLQPL-ASTSGLSVPARSAK)
Mouse α -TuJ1	Covance MMS-435P	Microtubules derived from rat brain.
Mouse α -Ki67	BD-Biosciences, 556003 (clone B56)	22 aa Ki-67 repeat motif (APKEKAQPLEDLASFQELSQ)
Mouse α -CK14	Vector #VP-C410	Synthetic peptide corresponding to C-terminus region of human keratin 14 (GKVVSTHEQVLRITKN) conjugated to thyroglobulin.
Rabbit α -CK14	LabVision #RB-9020-P1	A synthetic peptide of 15 amino acid residues from the C-terminus of human keratin 14.
Rat α -E-cadherin	Sigma U 3254	Rat embryonal carcinoma line PCC4 AZA R1 was used as immunogen. The antibody is a monoclonal IgG1 provided in rat ascites fluid from the DECMA-1 hybridoma line. Monoclonal Anti-E-Cadherin was selected against the mouse E-Cadherin.
Goat α -CD54	R&D systems, AF583	Recombinant extracellular domain from rat, affinity-purified (Accession Q00238, Gln 28 - Thr 493)
Rabbit α -LSD1 (a.k.a. KDM1)	Abcam 62582	Synthetic 16 amino acid peptide (QAPPEEENESEPEEPS) from near the centre of human KDM1 / LSD1.
Rabbit α -coREST	Millipore Cat #07–455	GST fusion of human coREST aa 109–293
Mouse α -Mash1 (a.k.a. Ascl1)	BD-Biosciences, 556604 (clone 24B72D11.1; Lo et al, 1991)	Recombinant-full length rat MASH1/ASCL1 protein, affinity chromatography-purified
Goat α -Sox2	Santa Cruz, sc-17320	Amino acids 277–293 of human Sox2 affinity-purified. YLPGEVPEPAAPSLH

TABLE 2

Sorting Strategy and Cell Types Isolated for Illumina Bead Array Profiling

Population of interest	Transgenic line/condition	Sorting strategy	Purity/enrichment
Neurogenic Progression Datasets			
GBCMPPs	Sox2-eGFP heterozygotes (hets), normal olfactory mucosa	eGFP (+), E-cad (-)/>CD54 (-), PI (-), FSC/SSC low	<1% CK14 contamination, significant CK18 (+) contamination
GBC _{INP} /immature neurons	Neurog1-eGFP hets, normal olfactory mucosa	eGFP (+), PI (-), FSC/SSC low	5% CK18 (+) contamination
Activated GBC _{INP}	Neurog1-eGFP hets, 3wks post OBX	eGFP (+), PI (-), FSC/SSC low	–
Newly differentiating mature neurons	Δ OMP-eGFP hets, 3wks post OBX	eGFP (+), PI (-), FSC/SSC low	–
Mature neurons	Δ OMP-eGFP hets, normal olfactory mucosa	eGFP (+), PI (-), FSC/SSC low	3% CK18 (+) contamination
Differentiated Cell Datasets			
Normal olfactory mucosa	Control mice, olfactory mucosa	PI (-)	N/A
Sus cells	Sox2-eGFP hets, normal olfactory mucosa	eGFP (+++), PI (-), CD54 (-)	> 95% CK18 (+)

Cell types isolated for expression profiling with Illumina Bead Chips (Mouse Whole Genome-6) are listed. The table lists the source tissue, the FACS sorting protocol used for isolation, and the purity of the preparation as determined by post-hoc analysis of cytospun cells from the sorting conditions.

TABLE 3

Number of Bead Sets Significantly Regulated in Pairwise Comparisons

Comparison			Upregulated	Downregulated
Sox2-GBC	vs.	Normal	3133	2749
Neurog1-Ctrl	vs.	Normal	3607	2932
OMP-Ctrl	vs.	Normal	1384	1582
Neurog1-Ctrl	vs.	OMP-Ctrl	3628	3348
Neurog1-Ctrl	vs.	Neurog1-OBX	38	25
Neurog1-OBX	vs.	OMP-OBX	2394	2342
OMP-OBX	vs.	OMP-Ctrl	743	844
Sox2-GBC	vs.	Neurog1-Ctrl	1285	1858
Sox2-GBC	vs.	Sus	5299	5341

Pairwise comparisons between each set of biologically meaningful replicates were made using the limma package for R (Smyth, 2004). The number of gene sequences that were significantly regulated with an adjusted *P*-value of < 0.05 are listed for each comparison. “Upregulated” refers to the number of probe sequences with a significantly higher expression level in the first listed sample in each row, while “Downregulated” refers to the number of probe sequences with a significantly higher expression level in the second listed sample in each row. The numbers slightly overrepresent the number of actual genes for each comparison because some genes have more than one sequence represented within the bead array. The abbreviations used for the purposes of discussion and comparison between various cell populations are as follows (name in *italics* refers to the population of interest described in the first column of Table 2): “Sox2-GBC” = *GBCMPP*, “Normal” = *normal olfactory mucosa*, “Neurog1-Ctrl” = *GBCINP/Immature Neurons*, “OMP-Ctrl” = *Mature Neurons*, “Neurog1-OBX” = *Activated GBCINP*, “OMP-OBX” = *Newly Differentiating Mature Neurons*, “Sus” = *Sus Cells*. In general, “-Ctrl” and “-OBX” are used to distinguish animals from the same transgenic mouse line that were subjected to no surgical intervention (“-Ctrl”) and those that were subjected to bulbectomy (“-OBX”).

TABLE 4

Functional Annotation Clusters Upregulated in the Neurogenic Progression

Sox2GBC vs Normal OM		
	Functional Annotation Cluster	Enrichment Score
Cluster 1:	Negative regulation of cytoskeleton organization	3.25
Cluster 2:	Neuron development	2.74
Cluster 3:	Regulation of cellular component biogenesis	2.70
Cluster 4:	Cell migration	2.56
Cluster 5:	Protein amino acid phosphorylation	2.50

Sox2GBC vs Sus		
	Functional Annotation Cluster	Enrichment Score
Cluster 1:	Cytoskeleton	7.66
Cluster 2:	Synaptic vesicle	7.43
Cluster 3:	Neuron development	6.60
Cluster 4:	Protein kinase inhibitor	4.73
Cluster 5:	Cytoplasmic vesicle	3.96

Neurogl vs Normal OMus		
	Functional Annotation Cluster	Enrichment Score
Cluster 1:	Nervous system development	5.00
Cluster 2:	Organelle organization and biogenesis	4.92
Cluster 3:	Neurogenesis	4.43
Cluster 4:	Neuron Recognition	3.38
Cluster 5:	Cytoskeletal Binding protein	2.65
Cluster 6:	Establishment and/or maintenance of chromatin architecture	2.51

Neurogl vs sus		
	Functional Annotation Cluster	Enrichment Score
Cluster 1:	Olfactory transduction	9.10
Cluster 2:	Synaptic vesicle	6.81
Cluster 3:	Neuron development	6.79
Cluster 4:	Protein kinase inhibitor	4.36
Cluster 5:	Synaptic transmission	4.32

Sox2GBC vs Neurogl		
	Functional Annotation Cluster	Enrichment Score
Cluster 1:	Glutathione transferase activity	5.57
Cluster 2:	Lung development	4.18
Cluster 3:	Membrane fraction	3.99
Cluster 4:	Aldehyde dehydrogenase (NAD)	2.97
Cluster 5:	Glutathione metabolic process	2.97
Cluster 6:	Cytochrome P450	2.72
Cluster 7:	Blood vessel morphogenesis	2.63

Neurogl vs OMP		
	Functional Annotation Cluster	Enrichment Score
Cluster 1:	Ribosome	27.87
Cluster 2:	M phase	6.15
Cluster 3:	Small ribonucleoprotein	5.76
Cluster 4:	Chromatin assembly or disassembly	5.67
Cluster 5:	Cellular macromolecular complex subunit organization	4.99

NeurogOBX vs OMPOBX		
	Functional Annotation Cluster	Enrichment Score
Cluster 1:	Ribosome	22.80
Cluster 2:	M phase	11.76
Cluster 3:	Nuclear lumen	11.73
Cluster 4:	Cell division	11.43
Cluster 5:	Kinetochore	9.30
Cluster 6:	Nucleotide binding	6.98
Cluster 7:	RNA splicing	5.00
Cluster 8:	Steroid biosynthesis	4.23
Cluster 9:	Macromolecular complex assembly	4.02
Cluster 10:	Methylation	4.01
Cluster 11:	Transcription regulation	3.87

For each listed pairing, upregulated gene sequences from the first sample listed were input into the DAVID functional annotation clustering tool at high stringency (<http://david.abcc.ncifcrf.gov/home.jsp>) (Dennis et al., 2003; Huang et al., 2009). A representative term for each cluster was chosen for the top functional annotation clusters and is listed along with the corresponding enrichment scored for the biologically meaningful comparisons.