Published in final edited form as:

Chem Rev. 2013 September 11; 113(9): 6949–6985. doi:10.1021/cr400121w.

Properties and Reactivity of Gaseous Distonic Radical Ions with Aryl Radical Sites

Peggy E. Williams, Bartłomiej J. Jankiewicz, Linan Yang, and Hilikka I. Kenttämäa*

Department of Chemistry, Purdue University, West Lafayette, Indiana 47906

1. Introduction

The reactivity of carbon-centered distonic radical ions has been of interest for decades. The existence of distonic radical ions was first postulated by Gross and McLafferty in the early 1970's.^{1,2} In 1978, Bouma, MacLeod and Radom reported experimental results that supported theoretical predictions of the existence of a stable ring-opened ethylene oxide distonic radical cation.³ Ions with spatially separated charge and radical sites were coined as “distonic ions” by Yates, Bouma, and Radom⁴ in 1984. They later refined this definition⁵ to correspond to radical ions generated by ionization of a zwitterion, ylide, or diradical. Eberlin and co-workers later introduced the term “distonoid”, meaning distonic like, to encompass any radical ion that displays distonic “character” (i.e., ions with a high degree of discrete (non-mandatory) charge-spin separation) and is over-looked as a result of the strict distonic ion definition.⁶ However, the “distonoid” classification is not commonly used by the scientific community. Currently, the term “distonic” is widely accepted and used to denote ions with formally separated charge and radical sites even if they do not fall into the formal definition.⁷ According to the conventional valence bond description, the charge and radical sites are on adjacent atoms in α -distonic ions while they are separated by one and two atoms in β - and γ -distonic ions, respectively. A vast amount of experimental and theoretical studies were dedicated to distonic ions from the 1980's to 1990's, which have been previously reviewed separately by Hammerum and Kenttämäa.^{8,9,10}

The focus of this review is on gaseous ions with one or more aryl radical sites, a subgroup of distonic radical cations. The interest in these distonic ions was initially sparked by the limited knowledge on the reactivity of neutral phenyl radicals and their diradical counterparts in spite of the vast amount of research dedicated to these reactive intermediates.^{11–70} Many such mono- and diradicals have been investigated as they are thought to play a vital role in numerous fields, including combustion,^{11–13} polymerization,^{14–16} atmospheric chemistry,^{17–19} interstellar chemistry,²⁰ organic synthesis⁸ and the biological activity of certain drugs.^{21–33} In the 1990's, the formation of such aromatic diradicals in naturally occurring anti-tumor antibiotics was associated with their DNA-cleaving ability.^{21–33} The two radical sites are thought to abstract a hydrogen atom from each strand of double stranded DNA, thus causing irreversible DNA cleavage. Since then, theoretical and experimental research on aryl mono- and diradicals has boomed. An area of special interest has been the mechanistic understanding of hydrogen atom abstraction by these radicals from small organic and biological molecules both in solution^{34–45} and in the gas phase.^{46–70} However, the ability to predict the rates of such seemingly simple reactions has proven challenging due to a poor understanding of the nature of the transition states for these reactions. Further, the examination of the chemical

*hilikka@purdue.edu.

properties of neutral radicals is a challenge due to the difficulty to cleanly generate them both in solution and in the gas phase.

In order to address the above difficulties, studies were carried out in the early 1990's on distonic radical cations' ion-molecule reactions inside mass spectrometers as ions can be easily manipulated in this environment.^{46–70} Distonic radical cations that have a phenyl radical site spatially separated from a chemically inert charge site were found to almost exclusively undergo radical reactions at the radical site(s) in the gas phase^{46–70} and lately also in solution.⁴⁵ Hence, examination of these distonic ions will provide information on the properties of phenyl mono- and diradicals.

A special benefit of using mass spectrometry to study above species is that the desired charged radical can be isolated before examining its reactivity. Hence, the precursors to any products formed in these gas-phase experiments are known, which is not always true for solution experiments wherein highly reactive molecules cannot be isolated. The chemical properties of many aryl mono-, di- and triradicals have been successfully examined in mass spectrometers by using this 'distonic ion approach'.^{49,50,52} The results obtained in these studies provide valuable information on the relative reactivities of mono- and polyradicals, which would otherwise not be available. This paper reviews the current knowledge of the properties and reactivity of distonic radical ions with aryl radical sites and the mechanisms of these reactions. Distonic phenyl radical ions generated within peptides are not included due to space limitations and also since these radicals are usually generated as precursors to less reactive nonaromatic peptide radicals that are the true interest of the researchers. However, this is an important and exciting new field of distonic ion research that should be reviewed separately.

2. Experimental Methods

For decades, Fourier transform ion cyclotron resonance (FT-ICR) mass spectrometry, developed by Comisarow and Marshall in 1974,⁷¹ has been the technique of choice for the study of gas-phase ion-molecule reactions. These mass spectrometers enable isolation of reactant ions and the variation of reaction time in order to determine their ion-molecule reaction products and reaction rates. Until recently, nearly all ion-molecule reactions of distonic ions were studied in an FT-ICR.

FT-ICR mass spectrometers are ion traps that allow storage of ions for long time periods due to the ions' cyclotron motion perpendicular to the static magnetic field (B) in the instrument and due to a potential energy well between two cell plates in the direction parallel to the magnetic field. Ion generation can be achieved through external and internal electron ionization⁷² (EI), chemical ionization^{73,74} (CI), electrospray ionization^{75–78} (ESI), atmospheric pressure chemical ionization^{79,80} (APCI), matrix-assisted laser desorption ionization^{81–84} (MALDI) and other ionization techniques. Broadband frequency-sweep^{71,85,86} (chirp) and stored-waveform inverse Fourier transform⁸⁷ (SWIFT) excitations are the two most commonly used excitation methods for ion manipulation, such as ion isolation by ejection of unwanted ions from the trap and ion detection by forcing ions to move coherently and on larger cyclotron orbits. Low-energy excitation techniques, such as sustained off-resonance collision-activated dissociation^{88,89} (SORI-CAD), can be employed to gently kinetically excite isolated ions that upon collisions with an inert gas form desired new ions or structurally informative fragment ions. MSⁿ experiments involving multiple consecutive ion isolation and collision-activated dissociation (CAD) or ion-molecule reaction events are easily carried out in FT-ICRs simply by changing the excitation pulse sequence via computer software,⁹⁰ allowing generation of distonic ions via multiple-reaction

processes, conclusive determination of their structures via CAD and/or ion-molecule reactions, and detailed study of their ion-molecule reactions and their products.

All events in a single-cell FT-ICR take place in the same space, which increases the probability for undesirable ion-molecule reactions between, for example, the reaction product ions and the reactant ions' neutral precursor. However, the use of an external ion source^{91,92–94} or a differentially pumped dual cell⁹¹ allows the generation of the ions and examination of their reactions in separate locations, thus preventing unwanted ion-molecule reactions. Numerous distonic ions have been examined in dual cell FT-ICR instruments by transferring the ions of interest into the adjacent clean cell for reactivity studies.

Mass spectrometers having three⁹⁵ or five quadrupoles^{6,96–101} also have been used to examine distonic ions, but these instruments are limited to MS/MS and MS/MS/MS experiments, respectively. However, three-dimensional quadrupole ion traps (QIT) can perform MSⁿ experiments, and these instruments have been used to explore CAD and ion-molecule reactions of distonic ions.⁹⁵ Similar to FT-ICR instruments, ions in quadrupole ion traps are manipulated and studied using a consecutive series of events. In general, ions can be generated in the trap or introduced from an external ion source, and isolated inside the trap by the application of rf pulses to eject unwanted ions. The ion of interest is then allowed to react with reagents introduced into the trap via pulsed valves, laser desorption, or by mixing with the helium buffer gas. The last method allows for direct measurement of reaction rate constants. After a preset reaction time, all ions are ejected out of the trap to an external electron multiplier for detection. Instead of ejection, product ions may be isolated and subjected to MSⁿ experiments. Gronert has previously reviewed studies of ion-molecule reactions in quadrupole ion traps.^{102,103} Most recently, modifications made to commercial linear quadrupole ion trap mass spectrometers (LQIT) have facilitated the study of gas-phase ion-molecule reactions in these instruments.^{104,105} The most important modification involved adding a mixing system to introduce neutral reagents into the trap via a syringe drive.

3. Ion Structure Determination

The late Bob Squires once wrote: “Any synthetic method without structural verification or some means for chemically characterizing the product is of dubious worth.”¹⁰⁶ Confirmation that a desired distonic radical ion was truly generated is indeed the most important component of studying distonic radical ions. The structures of gaseous distonic ions can be probed using a variety of indirect methods, including (1) determining their heats of formation and comparing to calculated values, (2) measuring CAD mass spectra, (3) studying ion-molecule reactions, and (4) by performing theoretical calculations.^{107,108} Measured heats of formation (ΔH_f) data are sometimes used to probe ion structures. The ions of interest and reference ions may have similar structures if their measured ΔH_f values are identical, and they definitely have different structures if ΔH_f values are different.^{107,109} ΔH_f for aromatic carbon-centered radical anions can be experimentally determined by measurement of the threshold energy for the formation of the radical anion from an activated precursor.¹⁰⁷ These measurements are critically dependent upon the precursor from which the ion is formed.¹¹⁰ Activation methods employed in these measurements include EI, CAD, and photodissociation (PD). As with any experimental method, there are limitations, including the fact that the structures of the neutral products of the dissociation are usually not known but assumed.¹⁰⁹ Both positive and negative ion approaches to these measurements are extremely sensitive to small amounts of isomeric impurities (in some cases as little as 1–2%) in the precursor ions.^{110,111} Further, the accuracy of these measurements is affected by the internal and/or kinetic energy distribution of the precursor ion.^{107,109}

CAD has become a standard technique used in the structural determination of distonic ions.^{112,113} During a CAD event, an isolated ion is translationally excited and subjected to collisions with a neutral inert gas, typically helium, nitrogen, air or argon. Collisions with the gas dampen the ions' motion and convert some of their kinetic energy into internal energy. Multiple collisions may be needed for the ions to gain enough internal energy to overcome fragmentation barriers.^{107,109,114} These experiments typically produce many fragment ions, which makes CAD a powerful tool for ion structure determination.⁶ For example, Eberlin and Gozzo⁹⁶ used a pentaquadrupole (MS³ experiments) to prove that transfer of ionized methylene (CH₂⁺) from ring-opened oxirane radical cation to ethylpyridines occurs at the nitrogen atom in the pyridine ring. Further, since CAD of the isomeric N-methylene-n-ethylpyridinium radical cations, where n = 2, 3, or 4, yields structurally characteristic fragment ions, the isomers can be differentiated. For example, the N-methylene-2-ethylpyridinium radical cation predominantly fragments to lose a hydrogen atom and ethylene whereas the N-methylene-4-ethylpyridinium radical cation fragments to lose a methyl radical (see Figure 1).

Another example is provided by a 1993 study on reactions of the pyridine radical cation with ethene carried out by Gross and co-workers to demonstrate the utility of high-pressure ion trapping in FT-ICR mass spectrometry.¹¹⁵ Under high-pressure conditions, the electrophilic pyridine radical cation was found to add to ethene to produce a β-distonic ion (Scheme 1). CAD of the ion (Figure 2) resulted in a facile hydrogen atom loss as well as loss of C₂H₄ to regenerate the pyridine radical cation, a fragmentation reaction typical for β-distonic radical cations.¹¹⁶ The β-distonic ion was distinguished from its α-distonic isomer by preparing the α-isomer by reaction of methylcyclopropane radical cation with pyridine and measuring its CAD mass spectrum (Figure 3). The CAD mass spectra of the α- and β-distonic ions show differing abundances of fragment ions of m/z 79 and m/z 80 (Figures 2 and 3). The three isomeric ethylpyridine radical cations (Figure 4) are also readily distinguished from each other and from the isomeric α- and β-distonic ions based on CAD. However, one important caveat of CAD is that some distonic ions may have internal energies nearing the barrier to fragmentation prior to activation, which may cause them to isomerize if the barrier to isomerization is lower than the fragmentation barrier. Isomeric ions also may rearrange after activation and prior to fragmentation.

Because of the limitations of CAD in differentiation of isomeric ions, reactions between distonic radical cations and different neutral molecules have been studied extensively in order to find reactions that can be used to unambiguously identify distonic radical cations. These reactions have been previously reviewed.^{97,102,103,117–119} Conventional radical cations predominantly undergo electron or proton transfer reactions. Unlike their conventional counterparts, distonic radical contain spatially separated charge and radical sites, and electron transfer reactions are typically not observed due to the low recombination energies of their charged sites. Since radical-like reactivity is often the main characteristic distinguishing distonic radical cations from their conventional isomers, reagents that readily donate atoms or groups to neutral radicals have been used for the identification of distonic radical cations and for the examination of their reactivity, including dimethyl disulfide,⁴⁶ dimethyl diselenide,⁵¹ *tert*-butyl isocyanide,⁵⁴ allyl bromide,⁶ allyl iodide,⁵⁵ cyclohexane,⁶⁷ decaline,¹²⁰ tetrahydrofuran,⁵⁹ and dioxygen.¹¹⁵ For example, examination of the reactivity of the β-distonic ion formed upon the reaction of the pyridine radical cation with ethene as discussed above, as well as the three isomeric ethylpyridine radical cations, toward dioxygen revealed that only the β-distonic ion reacts with dioxygen to produce a stable oxygen adduct (Figure 5).¹¹⁵ The lack of reactivity of the ethylpyridine radical cations and the α-distonic ion was suggested to be due to the delocalization of the radical site into the aromatic ring in these species. Eberlin and co-workers later generated and differentiated the three conventional and a total of eight distonic ethylpyridine radical cation isomers by using

multiple stage pentaquadrupole MS² and MS³ experiments.¹²¹ Examination of the reactivity of the isomeric ethylpyridine radical cations toward dioxygen confirmed the above finding¹¹⁵ that the β -distonic *N*-ethylenepyridinium cation reacts extensively while the α -distonic ion is completely unreactive. Interestingly, the α -distonic *meta*- and *para*-methyl substituted *N*-methylenepyridinium ions were found to readily react with dioxygen while the *ortho*-analog was unreactive. Ab initio calculations found the spin densities for the α -distonic ions to be largely localized on the *N*-methylene radical site.¹²¹ Hence, high spin densities on the radical sites of distonic ions do not necessarily lead to radical coupling reaction with oxygen.

Lastly, but definitely not least, a plethora of structural properties of distonic ions, including bond angles, bond lengths, and charge and odd spin distributions, have been probed using quantum chemical calculations. Further, thermochemical properties, such as electron affinities, acidities, and enthalpies of formation, as well as singlet-triplet splittings, transition state (TS) energies, heats of reaction, and many other useful values have been predicted computationally.^{67–70}

4. Properties and Reactivity

4.1 Charged Aryl Monoradicals

Distonic cations' radical reactions have been of interest for a long time.^{8–10,46–70} The reactivity of many α -distonic ions was first examined⁶ in the gas phase but the location of the charged site adjacent to the carbon radical center makes it difficult to determine the extent to which the charge site (for example, via inductive effects) influences the reactivity of the radical site. Therefore, aliphatic distonic ions with more than one atom separating the charge and radical sites were explored.^{9,10} However, the flexibility of the aliphatic chain of these distonic ions did not prohibit the neutral reagent from simultaneously interacting with the charged and radical sites.^{9,10} Hence, many recent studies have focused on the reactivity of rigid aromatic carbon-centered σ -type radicals with a spatially separated chemically inert charge site. Their reactions with a variety of neutral reagents have been explored in the gas phase by FT-ICR mass spectrometry.^{49,50,52} The results of these studies are summarized in this section.

4.1.1 Barriers for Radical Reactions—The significance of avoided curve crossings on transition states of chemical reactions is well known^{122–124} and has been reviewed by Donahue¹²⁵ in 2003. The research discussed herein requires a brief discussion on this topic. Since the work by Heitler and London,¹²⁶ it has become widely accepted to treat chemical reactions as avoided curve crossings. Three theories based on avoided curve-crossings have been developed to understand the factors that cause the barrier heights of radical-molecule reactions to vary from one reaction to another: the Marcus theory, the covalent (Heitler-London) curve crossing model, and the ionic curve crossing model (Figure 6).

Marcus theory originates from the work by Evans and Polanyi¹²⁷ who were the first to propose that barrier heights and reaction enthalpies are directly proportional to one another. Marcus expanded upon their work and developed the most widely used form of this theory to explain electron transfer reactions in weakly coupled systems¹²⁸ and later atom transfer reactions in strongly coupled systems.¹²⁹ The Marcus theory assumes that the barrier to reaction, E_{AB} , is dependent upon a single extrinsic parameter, the free energy release (ΔE_0), such that $E_{AB} = E(1 + \Delta E_0/E)^2/4$, where E is the average of the intrinsic barriers for the two corresponding identity reactions, $E = (E_{AA} + E_{BB})/2$.¹³⁰ The presence of intrinsic barriers renders the Marcus theory difficult to test, however, the general principle proposed by Evans and Polanyi¹²⁷ is easily tested as the changes in barrier heights should be directly proportional to changes in reaction enthalpy so long as the central atom does not change.

The covalent curve crossing model developed by Pross and Shaik^{131,132} was derived from the valence-bond formalism of London,¹²⁶ and has been modified and used to describe a multitude of chemical reactions,^{131–136} including hydrogen atom abstraction. Recognizing the importance of excited electronic states in the formation of the reaction barrier, this theory treats the evolution of the excited states along the reaction coordinate and predicts barrier dependence on the singlet-triplet splitting of the breaking and forming covalent bonds.

The third theory, the ionic curve crossing model, is most relevant to the work reviewed herein. This theory emerged from the covalent curve crossing model discussed above. After discovering that the transition state energy for reactions of polar radicals is directly proportional to the ionic energy of the separated reactants, Anderson and coworkers modified^{130,137–139} the covalent curve crossing model by defining the barrier height of a polar radical-molecule reaction as an avoided crossing of the ground state and a hypothetical ionic excited state. The ionic energy of the separated reactants was initially expressed as the difference between the vertical ionization energy of the cleaving bond and the vertical electron affinity of the radical site of the reactant radical. The vertical ionization energy (IE_v) of the cleaving bond is the energy required to remove an electron from the bond to form a vibrationally excited state with the same geometry as the precursor molecule. The vertical ionization energy of the cleaving bond is often approximated by the molecular ionization energy of the reactant.⁶⁷ The vertical electron affinity (EA_v) of the radical site of the reactant radical is the energy released when an electron is added to the radical site without geometry change. The ionic curve crossing model predicts that the energy of the transition state increases as the vertical IE of the breaking bond increases and as the vertical EA of the radical site decreases. This model has been used in a number of experimental studies to explain relative rates of radical reactions, including several reviewed here.^{50,52,56,58,62,67}

4.1.2 Charged Phenyl Radicals—Among the variety of charged aryl monoradicals whose ion-molecule reactions have been studied, distonic anilinium radical cations were the first class to be investigated in detail.^{47,48} This class of distonic ions was the first to enable the bimolecular reactivity of aryl radicals to be examined in the gas phase. CAD of 2-, 3-, and 4-iodoanilinium cations in a dual-cell FT-ICR mass spectrometer was used to generate^{47,48} the 2-, 3-, and 4-dehydroanilinium cations (**1** – **3**; for 4-dehydroanilinium cation, see Scheme 2), which were found to be stable towards isomerization by using energy-resolved CAD experiments (Figure 7).⁴⁸ Examination of their ion-molecule reactions revealed that they all abstract a thiomethyl group from dimethyl disulfide, which is indicative of a distonic ion structure.^{46,140} Radicals **1** – **3** were also found to abstract an iodine atom from 2-iodopropane, add to cyclohexene, and readily donate a proton to pyridine (for 4-dehydroanilinium cation, see Figure 8). On the other hand, ionized aniline was found to be unreactive toward dimethyl disulfide, 2-iodopropane, and pyridine, which provided conclusive evidence that the ions generated upon CAD of protonated 2-, 3- and 4-iodoanilines are structurally different from ionized aniline.⁴⁷ The reactivity observed for the distonic ions **1** – **3** is consistent with the reactivity of neutral phenyl radicals toward dimethyl disulfide^{35,141,142} and alkyl iodides.^{143–145}

In the following years, several (4-dehydrophenyl)phosphonium and -sulfonium cations (**4** – **12**, Chart 1) were studied in order to assess how different inert charge sites located remote from the radical site, as well as substitution, influence the reactivity of charged phenyl radicals.^{49,50} In general, these radicals were generated in a dual cell FT-ICR by first subjecting the appropriate halobenzene to electron ionization and allowing the resulting radical cation to react with a nucleophile to generate the charged site. The substituted halobenzene cations were then subjected to SORI-CAD using argon target gas to induce

homolytic cleavage of the weak carbon-iodine bond in order to form the desired charged phenyl radical (Scheme 3). The structures of the charged phenyl radicals were verified by ion-molecule reactions with dimethyl disulfide and dimethyl diselenide as well as CAD experiments. The type of charge site appeared not to influence reactivity. Radicals **4** - **8** and **11** (Chart 1) were found to undergo characteristic thiomethyl and iodine atom abstraction reactions with dimethyl disulfide and allyl iodide, respectively. Radicals **4** and **7** reacted more efficiently with dimethyl disulfide and allyl iodide than radicals **5** and **6**, suggesting that electron-withdrawing substituents can enhance the reactivity of charged phenyl radicals, in agreement with the ionic curve crossing model proposed by Anderson and co-workers.^{130,137–139} The *meta*-isomer of **5** (Chart 1) was found to react faster with dimethyl disulfide and allyl iodide than the *para*-isomer (**5**), suggesting that the closer proximity of an electron-withdrawing substituent to a radical site enhances radical reactivity. Again, this finding is in agreement with the ionic curve crossing model. Further, radical **11** reacts more efficiently with these two reagents than radical **5**, suggesting that additional electron-withdrawing substituents on the same ring as the radical site enhance radical reactivity. These findings are consistent with the reactivity of neutral phenyl radicals toward the same reagents in solution.^{35,141–146} In agreement with the ionic curve crossing model, electron-withdrawing substituents have been found to enhance the reactivity of neutral phenyl radicals in solution. For example, thiomethyl abstraction has been observed to occur three times faster for *para*-nitrodehydrobenzene than dehydrobenzene in solution.¹⁴⁶

Inspired by above observations, radicals **4**, **7**, **9**, **10**, and **12** (Chart 1) were allowed to react with 1,4-cyclohexadiene, phenol, thiophenol, benzeneselenol, and tetrahydrofuran in the gas phase.⁵⁰ Slow hydrogen atom abstraction was the major reaction observed, except for the reactions with phenol. Phenol reacts with radicals **9** and **12** by very slow addition as well as fast phenyl radical abstraction followed by slow adduct formation. Fluorine substitution was found to drastically enhance the rate of all hydrogen atom abstraction reactions. The greatest rate increase was observed in reactions of the fluorinated monoradicals with tetrahydrofuran, which was studied as a simple model of the sugar moiety in DNA. These findings suggested that the sugar moiety in DNA might be sensitive to substituents in the attacking radical, indicating that incorporation of fluorine atoms in anticancer drugs that cleave DNA through radical mechanisms could lead to greater anti-tumor activity. The exothermicity of hydrogen atom abstraction by charged phenyl radicals was explored by calculating the homolytic phenyl C-H bond dissociation energies of the dimethylphenylsulfonium ion **7** and those of methane and benzene (using isodesmic reactions and the ROMP2/6-31G(p)//ROHF/3-21G(p)+ZPE level of theory), and using these energies to estimate the enthalpy changes for hydrogen atom abstraction by **7**. The results suggest that the hydrogen atom abstraction reactions are highly exothermic, despite of their low rates,⁵⁰ indicating that a high barrier exists on the reaction coordinate. To explore whether fluorine substitution affects the exothermicity of hydrogen atom abstraction reactions by phenyl radicals, the same process was used to calculate the exothermicity of hydrogen atom abstraction by (3-fluorophenyl)sulfonium ion from benzene and methane. The introduction of a fluorine atom was found to have a significant effect on the exothermicity of these hydrogen atom abstraction reactions, in agreement with the above observation of rate enhancement with fluorine substitution.

A wide variety of distonic *N*-phenyl-pyridinium radical cations were also studied in order to examine how the polarity of the transition state may affect the reactivity of these radicals.^{47–67} In general, these charged phenyl radicals were formed in the gas phase in FT-ICR mass spectrometers by allowing radical cations of halobenzenes react with a variety of pyridine derivatives, leading to the formation of *N*-phenylpyridinium cations, which upon SORI-CAD generate the desired charged phenyl radicals (Schemes 4a and 4b). Fluorinated pyridine was used to generate a number of these charged phenyl radicals in order to be able

to distinguish between a brominated radical cation precursor and the resulting substitution product since bromine and pyridine have the same molecular weight. Examination of the reactivity of these radicals suggested that the fluorine substitution does not affect the reactivity (Scheme 4b).⁵⁵ All the above radicals were allowed to react with a variety of neutral reagents, *i.e.*, allyl iodide, dimethyl disulfide, *tert*-butyl isocyanide, tributyltin hydride, benzeneselenol, tetrahydrofuran, thiophenol, 4-fluorothiophenol and pentafluorothiophenol. The major reactions observed for allyl iodide were iodine atom and allyl abstractions, while thiomethyl abstraction occurred from dimethyldisulfide and cyano and hydrogen cyanide abstractions from *tert*-butyl isocyanide. Hydrogen atom abstraction was the major reaction observed for tributyl tinhydride, benzeneselenol, tetrahydrofuran, thiophenol, 4-fluorothiophenol, and pentafluorothiophenol, but addition reactions were also found to occur with some of these reagents. The polar character of the reacting system was reported to be the key factor controlling the reactivity of the phenyl radicals in these exothermic hydrogen atom abstraction reactions.^{55,56,58} The calculated vertical electron affinity (EA_v) of the radical site and vertical ionization energy (IE_v) of the substrate were found to be useful measures of the electron deficiency of the radical and the electron-donating character of the substrate, respectively. The reactivities of the charged phenyl radicals were found to correlate with their EA_v s for any given substrate. In addition, the reactivity of a given charged phenyl radical toward different substrates correlated with the IE_v s of the substrates. While substitution plays a major role in controlling the reactivity of charged phenyl radicals through polar effects, it was reported to exert only a minor influence on the enthalpy change of exothermic hydrogen atom abstraction reactions, even when charged rather than neutral substituents are involved.^{55,56,58}

Reactions of charged radicals with a methyl group adjacent to the radical site were found not to follow the expected pseudo-first order kinetics due to isomerization of the σ -radical to a methylene π -radical via a hydrogen atom shift (Scheme 5). However, this deviation from pseudo-first order kinetics was not observed when the methyl group was separated from the radical site by a carbon atom.⁵⁵

Both fluorinated and unsubstituted pyridinium substituents serve as electron withdrawing groups, making radicals more electrophilic. In general, electron- withdrawing substituents were found to enhance the reactivity of positively charged phenyl radicals most when located in a position *ortho*- with respect to the radical site, followed by *meta*- and then *para*-positions.⁵⁵

The 3-dehydropyridinium cation (**17**, Chart 3) with the positive charge in the same ring as the radical site, as opposed to the radicals discussed above, was found to react faster than the other charged phenyl radicals discussed thus far. This was attributed to the great calculated EA_v (6.53 eV at the B3LYP/6-31+G(d)+ZPVE level of theory) of this radical compared to those of the other phenyl radicals (5.11 – 5.82 eV at the B3LYP/6-31+G(d)+ZPVE level of theory).⁵⁸ These findings further emphasize the importance of the ionic surface in controlling the barrier height in radical abstraction reactions, in agreement with the work by Donahue et al.¹³⁰

Atom abstraction reactions are not the only reactions observed for charged and neutral phenyl radicals. Addition/elimination reactions have been observed to compete with hydrogen atom abstraction in solution.¹⁴⁷ Also these reactions have been found to be facilitated by electron-withdrawing substituents on the phenyl radical. Indeed, phenyl radicals with electron-withdrawing substituents were reported to add to C=C bonds much faster than they undergo hydrogen atom abstraction in solution. In 2001, five charged phenyl radicals (**13** - **16**, Chart 2) were allowed to react with thiophenol, benzaldehyde, toluene, aniline, phenol, and benzene in the gas phase in order to explore how substituents affect

competitive addition and hydrogen atom abstraction reactions in the gas phase.⁵⁷ In some cases, formation of stable adducts was observed. This was rather surprising as adducts are not frequently observed at low pressure in the gas phase. Under these conditions, adducts typically undergo fragmentation to form products or dissociate back to reactants before they can be stabilized via photon emission. Electron-withdrawing substituents were found to enhance both addition and substitution reactions but the addition reactions were more sensitive to substituent effects. The rate enhancements caused by substituents were rationalized by polarization of the transition state for both hydrogen atom abstraction and addition.⁵⁷ Both the hydrogen atom abstraction reactions and addition reactions were calculated to be quite exothermic. The degree to which hydrogen atom abstraction competes with addition was found to be directly proportional to the homolytic bond dissociation energy of the breaking R-H bond of the hydrogen atom donor.⁵⁷

4.1.2 Dehydropyridinium, Dehydroquinolinium, and Dehydroisoquinolinium

Cations—In the 2000's, following the observation of the enhanced reactivity of the 3-dehydropyridinium cation, numerous related dehydropyridinium, -quinolinium and -isoquinolinium cations (**17** – **23**, Chart 3) with varied distances between the formally charged heteroatom and the radical site were examined.^{67,68,148} In general, the distonic ions were generated in a dual-cell FT-ICR by subjecting appropriate protonated precursors to sustained off-resonance collision-activated dissociation (SORI-CAD) to cleave either a weak C-I or C-NO₂ bond. Hydrogen atom abstraction reactions were the sole reactions observed when distonic ions **17** - **23** were allowed to react with tetrahydrofuran and 2-methyltetrahydrofuran, while cyano radical and hydrogen cyanide abstraction reactions were observed with *tert*-butyl isocyanide.¹⁴⁸ The trends observed for the reactivities of these radicals were found not to arise from differences in reaction enthalpy, size of the radical, or the position of the radical site in the aromatic ring system. Analogous to the results discussed above, the reactivity of these radicals directly correlates with the calculated vertical electron affinities (EA_vs) of their radical sites. Hence, polarization of the transition state was found to be the major reactivity controlling parameter for also these distonic ions, providing further support to the validity of the ionic curve crossing model of Anderson and co-workers.^{130,137–139} The distance between the formally positively charged heteroatom and the radical site as well as the presence of substituents were found to affect the magnitude of the EA_v of the radical site.¹⁴⁸

Until 2008, only a small number of charged aryl radicals with a rather narrow range of EA_vs had been examined, all suggesting an important relationship between the EA_v of the radical site of the radical and the efficiency by which the radical abstracts a hydrogen atom. Hence, a systematic gas-phase study was conducted by examining the efficiency of hydrogen atom abstraction from 18 hydrogen atom donors, whose vertical molecular ionization energies (IE_vs) ranged from 8.8 up to 10.64 eV, by 29 aryl radicals whose calculated (UB3LYP/6-31+G(d)//UB3LYP/6-31+G(d) level of theory) EA_v values ranged from 3.31 up to 6.69 eV.⁶⁷ The radicals' reactivity toward cyclohexane and isopropanol was found to depend solely on the magnitude of the EA_v of the radical site (Figure 9) with no obvious dependence on the size of the aryl radical or calculated reaction enthalpy of the hydrogen atom abstraction reaction. Further, the natural logarithm of the reaction efficiencies for hydrogen atom abstraction from 15 different hydrogen atom donors by radical **14** (Figure 10) and from 14 different hydrogen atom donors by radical **15** was found to correlate with the molecular IE_vs of the hydrogen atom donors. Finally, the natural logarithm of the reaction efficiency of hydrogen atom abstraction by all the 29 aryl radicals from 18 different hydrogen atom donors was found to directly correlate with the difference between the IE_v of the hydrogen atom donor and the EA_v of the radical site of the radicals (Figure 11). However, no correlation was found between the efficiencies of hydrogen atom abstraction and the homolytic bond dissociation energies of the hydrogen atom donors. These findings

demonstrate that the molecular IE_v of the hydrogen atom donor can be used as a reactivity predictor instead of the IE_v of the breaking bond of the hydrogen atom donor. Further, it can be concluded that the reactivity of aryl monoradicals can be “tuned” by structural changes that alter either the EA_v of the radical site or the molecular IE_v of the hydrogen atom donor, in agreement with the ionic curve crossing theory.^{130,137–139}

In 2009, a thorough computational and experimental investigation was conducted on the reactivity of gaseous 2-, 3-, and 4-dehydropyridinium cations (**17**, **24** and **25**) toward a variety of organic reagents.⁶⁸ The structures of **17**, **24** and **25** were verified to be distonic based on predominant thiomethyl abstraction from dimethyl disulfide, and they were differentiated from each other based on the observed large differences in their hydrogen atom abstraction efficiencies from ethanol and methanol in an FT-ICR. As expected, the three radicals abstract an iodine atom from allyl iodide and hydrogen atom from cyclohexane, methanol, ethanol, tetrahydrofuran and 1-pentanol. The relative rates of hydrogen atom abstraction from methanol, ethanol, and tetrahydrofuran by **17**, **24** and **25** were found to correlate with the calculated activation enthalpies (G3MP2B3 and MPW1K levels of theory), $IE_v - EA_v$, and EA_v (B3LYP/aug-cc-pVTZ//B3LYP/cc-pVTZ level of theory) of the radicals, with a general reactivity trend of **17**, **24** and **25**. On the other hand, the calculated reaction enthalpies for hydrogen atom abstraction from methanol and ethanol did not reflect the observed differences in the reaction rates. Despite similar homolytic C-H bond dissociation energies¹⁴⁹ of the α -C-H bond in methanol and ethanol, charged radicals were found to react more efficiently with ethanol than methanol, likely due to the lower (by about ~ 0.32 eV) IE_v of ethanol. Based on the calculated (G3MP2B3) transition state structures for the reactions of **17**, **24** and **25** with methanol, the distance between the radical site and the hydrogen atom being abstracted is much longer than the distance between the hydrogen atom being abstracted and the α -carbon of methanol, indicating a more reactant-like, or “early”, transition state. Hence, the barriers for these reactions should correlate with calculated $IE_v - EA_v$, as was observed. These findings are consistent with the previous studies on the reactivity of charged phenyl radicals discussed above.

Unexpectedly, reactions of the charged radical **25** (with the radical site in the *para*-position with respect to the formal charge site) with tetrahydrofuran, *tert*-butyl isocyanide, toluene, and phenol resulted in unusual products. For example, while radicals **17** and **24** were found to exclusively undergo hydrogen atom abstraction from tetrahydrofuran and CN abstraction from *tert*-butyl isocyanide, radical **25** underwent several additional reactions, which were identified as abstraction of CH_2 , C_2H_3 , CHO, and C_2H_3O from tetrahydrofuran (confirmed by elemental composition measurements and examination of reactions of d₈-tetrahydrofuran), and HCN, secondary HCN and secondary C_4H_8 abstraction from *tert*-butyl isocyanide. These unusual reactions were rationalized by taking into consideration the ionized carbene-type resonance structure of **25** (Scheme 6). The reactions with tetrahydrofuran were suggested to occur via an initial nucleophilic addition of tetrahydrofuran to the dehydrocarbon atom (the most electron deficient carbon) of **25** rather than a radical attack by **25** on tetrahydrofuran. The proposed mechanisms for the CH_2 , C_2H_3 , CHO, and C_2H_3O abstractions are shown in Scheme 7. The resonance structure (Scheme 6) proposed to be responsible for the unusual products of **25** also exists for **24**, but a comparison of the calculated (G3MP2B3) barrier heights for addition of tetrahydrofuran to **24** versus hydrogen atom abstraction from tetrahydrofuran by **24** (-11.9 vs. -17.4 kcal/mol) suggest that hydrogen atom abstraction is the kinetically favored reaction. The calculated (G3MP2B3) transition state structures for hydrogen atom abstraction by **17**, **24** and **25** indicate that a stabilizing hydrogen-bonding interaction between the oxygen atom of tetrahydrofuran and the N-H proton of **24** biases **24** toward hydrogen atom abstraction. The ionized carbene-type resonance structure of **25** was also presumed to be partly responsible for its unusual reactivity toward *tert*-butyl isocyanide, toluene, and phenol. The nature of

charge/spin delocalization in the transition states was proposed to be the reactivity controlling parameter dictating whether CN or HCN abstraction was favored. Analysis of Mulliken spin density calculations (B3LYP/cc-pVDZ//B3LYP/6-31G(d)) for the intermediate resulting from the addition of *tert*-butyl isocyanide to **17**, **24** and **25** (Figure 12) indicated that the α -spin density for **25** is predominantly distributed over the heavy atoms, favoring HCN abstraction (Scheme 8), while the α -spin densities for **17** and **24** predominantly reside on the benzylic carbon atom. The same logic was employed to rationalize the unusual reactivity of **25** toward toluene and phenol, leading to the proposed mechanisms shown in Scheme 9 for these reactions. These results indicate that charge delocalization into the π -electron system containing the radical site in these distonic ions can facilitate nucleophilic addition to the aromatic ring and that hydrogen-bonding interactions in the transition state can greatly facilitate their radical reactions.

Blanksby and co-workers¹⁵⁰ have examined the reactivity of the *N*-methylated derivative of radical **24** toward 2-butyne in the gas phase in order to gain insights into the mechanism, products, and kinetics by using a modified Thermo Scientific linear quadrupole ion trap that was coupled to FT-ICR (LQIT/FT-ICR). The *N*-methyl-4-iodopyridinium precursor cation was dissolved in methanol, introduced into the ESI source of the instrument, mass selected and subjected to low-energy CAD with helium collision gas, which resulted in the homolytic cleavage of the weak C-I bond to generate the *N*-methyl-4-dehydropyridinium cation. The cation was isolated and allowed to react with 2-butyne from 0.1 up to 1 s. 2-Butyne was introduced via the helium mixing system of the modified instrument. The flow rates of helium and the reagent were used to control the concentration of the reagent in the ion trap. The *N*-methyl-4-dehydropyridinium cation readily reacted with 2-butyne to predominantly form adduct – H as well as some adduct – CH₃ product ions. Transition state calculations (B3LYP/6-31G(2df,p) level of theory) aided in the elucidation of the mechanisms proposed for the reactions (Scheme 10). Addition of the radical cation to 2-butyne was calculated to occur without a barrier. This finding is in agreement with the earlier studies discussed above suggesting that addition reactions of charged radicals can be facile.

4.1.3 Substituted Dehydropyridinium Cations—Following the observation that stabilizing hydrogen-bonding interaction in the transition state affects the calculated barrier height and reactivity of the 2-dehydropyridinium cation,⁶⁸ the gas-phase reactivity of a series of substituted 2- and 3-dehydropyridinium cations (**17**, **24**, **26** - **31**; Chart 3) toward cyclohexane, methanol, and tetrahydrofuran was investigated in order to probe the generality of hydrogen bonding as a reactivity controlling parameter for charged aryl radicals.⁶⁹ Hydrogen atom abstraction was the sole reaction observed. A correlation was found between the reaction efficiencies of the radicals with cyclohexane and their calculated EA_vs (Figure 13), consistent with the studies discussed above. On the other hand, no correlation was found between the calculated EA_vs of the radicals and the reaction efficiencies measured for methanol and tetrahydrofuran that are capable for hydrogen bonding, as opposed to cyclohexane. However, a correlation was found between the measured reaction efficiencies and the calculated activation enthalpies (G3MP2B3 and MPW1K) for hydrogen atom abstraction from the methyl group of methanol and the α -carbon of tetrahydrofuran (Figure 14). Further, examination of the calculated (MPW1K) transition state structures suggested that hydrogen bonding interactions exist in the transition states with methanol (Figure 15) and tetrahydrofuran for radicals **24**, **26** – **28** and **31**. Radicals **17**, **29** and **30** cannot have hydrogen-bonding interactions in their transition states due to the lack of a hydrogen atom donor near the radical site. The stabilizing hydrogen-bonding interactions observed for **24**, **26** – **28** and **31** lower the barrier to hydrogen atom abstraction, which helps rationalize the lack of correlation between the reactivity of the radicals toward methanol and tetrahydrofuran and their calculated EA_vs. These hydrogen-bonding interactions involve

either the NH proton or the amino/hydroxyl hydrogens of the radicals. The efficiency of the reaction of radical **31** with methanol is nearly five times greater than that of radical **30** (no hydrogen bonding in TS) due to hydrogen bonding in the transition state. The length of the hydrogen bond in the transition state was found to reflect the extent to which hydrogen bonding stabilizes the transition state. Shorter hydrogen bonds indicate stronger interactions between the radical cation and the neutral molecule, thus causing a greater lowering in the barrier to hydrogen atom abstraction and resulting in enhanced reactivity. This study confirmed that hydrogen bonding in transition states can control radical reactivity.

In 2012, a study of the reactivity of a variety of substituted 4-dehydropyridinium cations toward tetrahydrofuran was conducted in order to understand better the nature of the non-radical reactions occasionally observed for 4-dehydropyridines upon interaction with tetrahydrofuran.⁷⁰ The reactions of several of the radicals with tetrahydrofuran yielded CH₂, C₂H₃, CHO, and C₂H₃O abstraction products presumed to occur via non-radical mechanisms and rationalized by the ionized carbene-type resonance structure of the 4-dehydropyridinium cation.⁶⁸ However, the radicals with a calculated EA lower than that of the 4-dehydropyridinium cation were found to solely undergo hydrogen atom abstraction. As the EA_v of the radical site increased, the rate of the dominant non-radical reaction (CH₂ abstraction) was found to increase and eventually even dominate over hydrogen atom abstraction. When a substituent capable of hydrogen bonding was adjacent to the radical site, hydrogen atom abstractions alone were observed. The results of this study implicated two key factors that strongly influence the occurrence and rate of non-radical reactions for positively charged pyridyl radicals: (1) polar effects, reflected by the magnitude of the calculated (G3MP2B3) EA_v of the radical site and (2) hydrogen bonding in the transition state.

4.2 Charged Aryl Diradicals

Arynes, a class of reactive intermediates with two formally unpaired electrons, have been extensively studied both theoretically and experimentally for well over 100 years.^{151–153} Some arynes possess diradical character. Diradicals, as defined by Salem in 1972,¹⁵⁴ are molecules possessing two formally unpaired electrons occupying two degenerate or nearly degenerate orbitals (Chart 4). Based on the IUPAC convention,¹⁵⁵ diradicals are molecular species having two unpaired electrons, in which at least two different electronic states with different multiplicities, electron-paired (singlet state) or electron-unpaired (triplet state), can be identified.

ortho-Benzyne **32** was first proposed to be a reactive intermediate in the decomposition of lithiated halobenzenes in the early 1940's.¹⁵⁶ Several decades of work by Wittig,^{156,157,158} Roberts^{159–161} and Huisgen^{162–164} was necessary to obtain solid evidence for its existence. Since 1963, the ionization energy of *ortho*-benzyne **32** has been measured in a mass spectrometer¹⁶⁵ and it has been isolated in low-temperature matrices^{166,167} and fully characterized by IR, UV and mass spectrometry.^{168–173} In the 1990's, a revival of interest in aryne chemistry was catalyzed by the discovery that the DNA cleaving ability of some naturally occurring compounds is a result of formation of biologically active benzyne warheads.³² At that time, little was known about the chemical and physical properties of *meta*- and *para*-benzynes (**33** and **34**, respectively).^{21–29,30–33} Since then, these reactive intermediates have been extensively studied.^{174–181} Recent studies on the electronic structures and reactivity of charged pyridine- and quinoline-based arynes in the gas phase are discussed below.

4.2.1 *ortho*-Benzynes—Despite the wealth of information on the structure and reactivity of neutral *ortho*-benzynes, knowledge of neutral and charged pyridine and (iso)quinoline

based *ortho*-benzynes (**35** – **39**; Chart 4) remains relatively limited. In 2000, Debbert and Cramer¹⁵³ systematically compared the structure and energetics of isomeric benzyne, pyridynes, and pyridinium cations (Chart 4). Calculated singlet-triplet (S-T) splittings, at various levels of theory, were reported for **32** and **35** – **38** and are compared in Table 1 to the experimentally determined¹⁸² S-T splitting for **32**. These results predict a 15.5 kcal/mol difference in the S-T splittings for the *ortho*-benzyne **32** and the pyridyne **35**, and a 7.7 kcal/mol difference for **32** and another pyridyne **36** with the C=C moiety further away from the nitrogen atom. While the effect of a heteroatom on the S-T splittings of *ortho*-pyridynes is substantial, it is interesting to note that the calculated S-T splittings of both the neutral and charged 3,4-didehydropyridynes (**36** and **38**, respectively) were predicted to be nearly the same as for **32**, indicating that this heteroatomic effect only occurs over a short distance. The large magnitude of these calculated S-T energy splittings indicate strong coupling of the radical sites. Since the magnitude of the S-T splitting has been proposed to affect the radical reactivity of singlet diradicals, which has been attributed to the partial uncoupling of their radical electrons in the transition state of radical reactions, these *ortho*-arynes should react more like alkynes than radicals.¹⁸³ The optimized heavy-atom bond lengths for the singlet and triplet states of **32** and **35** – **38** as calculated at the BPW91/cc-pVDZ and CAS(8,8)/cc-pVDZ levels of theory are shown in Figure 16. The addition of a nitrogen and protonated nitrogen into **32** was predicted to change all of the bond lengths. The dehydrocarbon atom separation for the singlet states of **35** and **36** was predicted to be larger at the BPW91 level of theory than the CAS(8,8) level of theory. Interestingly, the predicted dehydrocarbon atom separations of **37** and **38** (with a protonated nitrogen) are nearly identical to that of **32**.

The 1,2-didehydropyridinium cation **39** has been the subject of numerous experimental and theoretical investigations beginning as early as 1961.^{98–101,174,184–191} For example, mass spectrometric studies and DFT calculations by Turek¹⁹¹ and Gozzo⁹⁹ found that the 2-pyridyl cation, a resonance form of **39**, is more stable than the isomeric 3- and 4-pyridyl cations. These findings suggest that the 2-pyridyl cation is significantly stabilized by the sp² nitrogen orbital overlap with the empty sp² orbital on the adjacent carbon. Polar cycloaddition reactions have been reported for the 2-pyridyl cation (**39**) in a mass spectrometer by Eberlin et al.^{98,100,101} In solution, the reaction of **39** with furan has been reported by Bunnett to result in the formation of 2-(2-pyridyl)furan **40**.^{189,192} The mechanism of this reaction remains unclear as at least two possible pathways could lead to this product (Scheme 11).^{189,193} On the other hand, *ortho*-benzyne (**32**) has been found to undergo Diels-Alder reactions with furan in solution.¹⁵¹ Reactions of 2,3- and 3,4-didehydropyridines with furan, followed by a lithium amalgam reduction, were found to afford quinoline and isoquinoline, respectively.¹⁹⁴ In contrast, the reactivity of the charged diradicals **37** and **38** has remained largely unexplored until recently.

Beginning in 2008, the gas-phase reactions of **37** – **39** (Chart 4) with furan, tetrahydrofuran, cyclohexane, allyl iodide, dimethyl disulfide, and *tert*-butyl isocyanide were investigated in the Kenttämäa laboratories.^{195,196} Furan was chosen as one of the reagents since the reactions of *ortho*-benzynes with furan have been studied earlier in solution, as discussed above. The remaining five reagents have become a “gold standard” for fundamental reactivity studies in the Kenttämäa laboratories as their examination provides useful information on the properties of mono- and diradicals. Diradicals **37** and **38** were generated in a dual cell FT-ICR via self-chemical ionization of respective 2,3- and 3,4-pyridinecarboxylic anhydride precursors, transferring the ionized precursors into the adjacent clean cell, and subjecting them to SORI-CAD by using an argon collision gas. This process resulted in the loss of CO and CO₂ from the ionized precursors to afford the desired diradicals. The 2-pyridyl cation **39** was similarly generated via HI loss from 2-iodopyridinium precursor cation upon SORI-CAD.

As discussed above, phenyl radicals in general tend to abstract a hydrogen atom from tetrahydrofuran and cyclohexane, an iodine atom from allyl iodide, thiomethyl from dimethyl disulfide, and CN and HCN from *tert*-butyl isocyanide via radical mechanisms. In contrast, mostly different, non-radical reactions were observed for **37** – **39** upon interaction with these reagents, such as H₂O and CH₂O abstraction from tetrahydrofuran, CH₂=C=CH₂ abstraction from allyl iodide, and CH₂S and CH₃SH abstraction from dimethyl disulfide. These non-radical reactions were proposed to occur via addition-elimination mechanisms (for examples on analogous benzyne, see Scheme 12), in agreement with previously reported solution studies.^{151,152} In general, **37** and **39** were found to react with similar efficiencies, despite the large differences in their EA_vs, and much faster than **38** except for *tert*-butyl isocyanide. It is not clear at this time why their reactivity does not reflect their calculated (UBLYP/aug-cc-pVDZ//UBLYP/cc-pVDZ level) EA_vs. A slow iodine atom abstraction was observed for **37** from allyl iodide. This reaction likely occurs via nucleophilic addition of the iodine atom of allyl iodide to the benzyne followed by a homolytic cleavage of a C-I bond, and hence is not an indication of radical reactivity. These sort of non-radical reactions have been observed previously for *meta*-benzyne analogs, as discussed below. Reports on *ortho*-arynes undergoing radical reactions are rare¹⁹⁷ since radical reactions are hindered by the large S-T splittings of these diradicals.

Further, the gas-phase reactivity of a series of quinoline and isoquinoline based *ortho*-benzyne (**41** – **45**; Chart 4) toward the above mentioned reagents was investigated.^{198,199} In agreement with the results discussed above, addition, addition/elimination and small molecule abstraction reactions were observed. For example, H₂O and CH₂O abstractions from tetrahydrofuran were observed instead of the hydrogen atom abstraction typically observed for phenyl radicals. The addition/elimination pathways were presumed to occur via an initial nucleophilic addition of the reagent to the electrophilic benzyne moiety (Scheme 12). Diradicals **44** and **45** have much larger calculated S-T splittings than **41** – **43**. However, they were observed to react more efficiently than the other diradicals, likely due to their large EA_vs. This finding supports the hypothesis that the benzyne studied undergo non-radical reactions, which are not affected by the magnitude of the S-T splittings of the benzyne. Indeed, the reaction efficiencies of **41** – **45** correlate with their EA_vs. Radicals **42** and **43** have similar calculated EA_vs (and S-T splittings). Radicals **44** and **45** also have similar calculated EA_vs (and S-T splittings) but these values are much greater than those of **42** and **43**.¹⁷⁹ Hence, it is not surprising that similar reaction efficiencies, products and product branching ratios were observed for the reactions of **42** and **43** with all reagents, and also for the reactions of **44** and **45** with all reagents. Proposed mechanisms for the reactions of the diradicals with tetrahydrofuran are shown in Scheme 12. In general, the reactivity studies described here have shown that the reactivity of *ortho*-benzyne derivatives is predominantly influenced by the magnitude of the electron affinity of the radical sites.

4.2.1 meta-Benzyne—The *meta*-benzyne, **33**, is arguably the most fundamentally interesting member of the aryne family, especially in terms of reactivity. However, experimental and theoretical studies on **33** did not begin until in the early 1960s.^{151–153} The first attempt to generate **33** in the gas phase was reported by Fisher and Lossing in 1963.¹⁶⁵ However, pyrolysis of 1,3-diiodobenzene resulted in thermal decomposition of the precursor instead of formation of **33**, and mass spectral analysis of the products suggested that an enediyne, rather than **33**, was formed. Berry and co-workers²⁰⁰ were the first to successfully generate **33** in 1965 by flash vacuum pyrolysis of benzenediazonium-3-carboxylate. The pyrolysis products were monitored using time-resolved mass spectrometry and flash-absorption optical spectroscopy. The lifetime of **33** was found to be 400 – 600 μs. The fragmentation patterns and lifetime of **33** were found to differ substantially from those measured for the *ortho*- and *para*-isomers, indicating that the *meta*-benzyne had indeed been

formed.²⁰¹ Since then, Sander and co-workers have reported numerous studies on the characterization of **33** and its halo- and hydroxy-substituted analogues in low temperature matrices^{178,202–213} by IR and UV-Vis spectroscopy as well as by quantum chemical calculations.

Despite nearly a half century of theoretical and experimental investigations,^{178,179} **33** has not received the same high degree of attention as its *ortho*- and *para*-benzyne siblings, rendering this isomer to be the least well understood in the benzyne family. Through-space spin interactions via back orbital overlap in *meta*-benzynes were indicated by the early computational studies of Hoffmann et al.¹⁸³ Later computational studies by Noell,²¹⁴ Thiel²¹⁵ and Dewar²¹⁶ predicted that **33** has an “open” structure with significant diradical character and is relatively stable. Significant progress in experimental studies of **33** was not made until the 1990s. Measurement of the heat of formation,^{217–219} S-T splitting,¹⁸² and IR spectrum^{202,204} of **33** led to a renewed interest in *meta*-benzynes.^{203,220–236}

The most controversial issue of *meta*-benzyne research has been its structure, and specifically, whether it has a bicyclic²⁰¹ or diradicaloid^{237,238} structure (Chart 5). In 1978, Washburn reported evidence for the existence of **33** and **33a** (Chart 5) in solution reactions through trapping and labeling experiments.²⁰¹ More recent studies probing the possibility of a bicyclic structure have been reported by Hess,²³⁹ Kraka²⁴⁰ and Winkler.²⁰³ Hess predicted the bicyclic **36a** to be more stable than the diradicaloid **33b**, whereas Kraka and Winkler predicted that **33** has a diradicaloid rather than a bicyclic structure. Most recently, however, vibrational frequencies from negative ion photoelectron spectroscopy¹⁸² and infrared absorption^{90a} spectra of matrix-isolated **33** and its analogues,^{210,211} as well as high-level calculations,²⁰³ have conclusively shown that **33** is best described by the monocyclic, σ -allylic (diradicaloid) structure **33b**.

The absolute heat of formation^{111,217} (122 ± 3.1 kcal/mol), electron affinity¹⁸² (0.852 ± 0.011 eV) and S-T splitting¹⁸² (-21.0 ± 0.3 kcal/mol) of **33** were experimentally measured by Squires and co-workers. **33** is $\sim 9 - 15$ kcal/mol less stable than its *ortho*-isomer and $\sim 13 - 21$ kcal/mol more stable than its *para*-isomer.¹⁵³ The large S-T splitting indicates that the through-space bonding interaction in *meta*-benzyne is substantial and results in a strong preference for a singlet ground state. Experimental and computational work by Chen and co-workers shows that even the 4 kcal/mol stabilizing interaction between the singlet-coupled diradical electrons in *para*-benzyne (**34**) and its analogues severely hinders radical reactivity because part of this stabilizing interaction is lost in the transition state.^{241,242} Hence, the much larger S-T splitting of **33** was expected to completely block radical reactions.

Related nitrogen-containing heteroaromatic *meta*-benzyne analogs (**46** – **48** and **50** – **52**; Chart 5) have been much less studied in comparison to their parent **33**. In 2011, Wenthold and co-workers²⁴³ experimentally determined the absolute heat of formation (130 ± 3 kcal/mol) of 2,4-didehydropyridine (**46**) by using energy-resolved CAD of deprotonated 2-chloropyridine. The structures and energetics of *meta*-benzyne (**33**), isomeric pyridynes **46** and **47** (Chart 5), isomeric pyridinium cations **50** – **52** (Chart 5), and their hydroxy-substituted analogues **53** – **55** (Chart 5), were computationally examined by Debbert, Johnson and Cramer.^{153,244} S-T splittings calculated at several levels of theory were reported for **33**, **46** – **48**, **50** – **52** and the hydroxy-substituted derivatives of **46** – **48**. These values are shown in Table 2, with the experimentally determined¹⁸² S-T splitting of **33**. Incorporation of a nitrogen atom into the aromatic ring of **33** results in a smaller S-T splitting for **47** and **48** but a larger S-T splitting for **46**. The substantially decreased S-T splitting of **48** is likely due to the destabilizing four-electron interactions between the nitrogen lone pair and the adjacent radical sites, which destabilize the singlet state and cause the singlet and triplet states to be nearly degenerate. Protonation of **46** results in a slightly

smaller S-T splitting (**50**), while protonation of **47** results in a somewhat larger S-T splitting (**51**) and protonation of **48** in a substantially larger S-T splitting (**52**). The protonated nitrogen atom in **52** removes the destabilizing interactions that occur in the singlet state of **48**. Addition of a hydroxyl substituent to the aromatic ring of **33** results in a substantial increase in the S-T splitting for **54** and **55**, but the S-T splitting of **53** is nearly identical to that of **33**. These findings demonstrate that the S-T splittings of *meta*-benzynes are very sensitive to structural changes, and suggests that their reactivity also may be sensitive to such changes.

Substitution was also found to affect the geometry of *meta*-benzynes, especially the distance between the dehydrocarbon atoms in the singlet states.^{153,244} Substitution of **33** with a hydroxyl group results in a slight decrease in the calculated dehydrocarbon atom distance (**53** and **55**), while no change was observed for **54** (Figure 17). The C-O bond in **53** and **55** was calculated to be shorter than that of phenol (1.381 Å), while no real effect was found on the C-O bond length in **54**.²⁴⁴ The shortening of the C-O bond in **53** was proposed to arise from stabilizing π -interactions between the oxygen lone pair and the cyclopropenium cation moiety in the zwitter ionic resonance structure of the bicyclic structure (**33c**, Chart 5; Scheme 13), while the C-O bond shortening in **55** arises from stabilizing inductive interactions between the electron-withdrawing oxygen atom and the allyl anion moiety. Incorporation of a neutral or protonated nitrogen atom into the aromatic ring affects all of the bond lengths and bond angles of **33**.¹⁵³ The dehydrocarbon atom separation provides insights into the strength of the interactions between the radical sites, making this distance of the greatest interest. DFT and CAS(8,8) theories predict significantly shorter dehydrocarbon atom separations for **47** and **48** (Chart 5) and a longer dehydrocarbon atom separation for **46** than for **33** (Figure 18). Protonation of the nitrogen atom in **46** and **47** was predicted to result in a smaller dehydrocarbon atom separation (**50** and **51**, respectively) and a larger dehydrocarbon atom separation for **48** (**52**).¹⁵³ The increased dehydrocarbon atom separation in **52** arises from the loss of the destabilizing interactions between the nitrogen lone pair and the adjacent radical sites in the singlet state of **48**. All these findings show that *meta*-benzynes have a remarkably flexible structure that can be greatly influenced by relatively small structural changes.

Throughout the mid to late 1990's, Wenthold and Squires synthesized *meta*-benzyne anion (**49**) and examined its reactivity (Table 3) in the gas phase in a flowing afterglow triple quadrupole apparatus.^{245,246} **49** was generated by subjecting the *meta*-bis(trimethylsilyl)benzene to fluoride-induced desilylation to afford the *meta*-(trimethylsilyl)phenyl anion, which upon subsequent reaction with F₂ yields the desired *meta*-benzyne anion.²⁴⁵ **49** was easily distinguished from the lower energy *ortho*-isomer as sequential abstraction of two sulfur atoms from CS₂ was observed for **49** but only electron transfer was observed for the *ortho*-isomer. **49** was also found to react with nitric oxide by sequential addition of two NO molecules, while no reaction was observed for the *ortho*-isomer.²⁴⁵ The basicity of the *meta*-benzyne anion was evident in its reactions with acids. Further, **49** was found to abstract an oxygen atom from N₂O and O₂ to produce dehydrophenoxide anions. Overall, the negative ion chemistry of the *meta*-benzyne anion **49** appears to dominate its reactivity toward the reagents examined. For example, while proton abstraction and nucleophilic reactions were observed for **49**, radical reactions, such as hydrogen or halogen atom abstraction, did not occur.

A *meta*-benzyne anion analogue, the 3,5-didehydrobenzoate anion (**56**; Chart 6), was examined in the Kenttämä laboratories in 2003.²⁴⁷ **56** was generated in an FT-ICR by electron ionization of triallyl-1,3,5-benzenetricarboxylate followed by CAD. The S-T splitting of **56**, calculated at the CASPT2 and CCSD(T) levels of theory, was predicted to be

–20.2 and –20.6 kcal/mol, respectively (using the UB3LYP geometry). These values are nearly identical to the measured S-T splitting of the *meta*-benzyne (**36**) (-21.0 ± 0.3 kcal/mol). Hence, theory predicts that carboxylate substitution at the *meta*-position of *meta*-benzyne has no effect on the relative energies of the singlet and triplet states. Comparison of the reactivity of **56** with that of the 3-dehydrobenzoate anion toward several electrophilic reagents revealed net radical abstractions at rates similar to those of the analogous monoradical.²⁴⁷ Surprisingly, **56** was found to abstract a bromine atom from carbon tetrabromide and bromotrichloromethane, and an iodine atom from iodine, in spite of its large S-T splitting. However, **56** did not react with hydrogen atom donors by hydrogen atom abstraction. When **56** and the 3-dehydrobenzoate anion were allowed to react with nucleophilic reagents, no reactivity was observed. The negatively charged substituent of **56** appears to bias its reactivity toward electrophilic reagents. Interestingly, **56** was found to react with the strongly electrophilic boron trifluoride to form a new *meta*-benzyne, (3,5-didehydrophenyl)trifluoroborane, by a charge-site substitution mechanism. This reaction provides an easy access to negatively charged *meta*-benzynes with different charged moieties.

The gas-phase reactivity of many positively charged *meta*-benzyne analogs has been examined in the Kenttämä laboratories since the late 1990's. In 1997, a general approach to synthesize and characterize positively charged phenyl diradicals in the gas phase was reported and utilized to examine the properties of a distonic *meta*-benzyne cation, the N-(3,5-didehydrophenyl)-3-fluoropyridinium cation (**57**; Chart 6).²⁴⁸ Similar to the methods discussed above for charged phenyl radical generation,^{49,52} **57** was generated in a dual-cell FT-ICR by ionizing 1,3,5-tribromobenzene by electron ionization and allowing the molecular ion to react with neutral 3-fluoropyridine to afford the N-(3,5-dibromophenyl)-3-fluoropyridinium cation. After transfer into the adjacent clean cell, the N-(3,5-dibromophenyl)-3-fluoropyridinium cation was subjected to SORI-CAD with an argon target or photodissociation (at 266 nm using a Nd-YAG laser), inducing homolytic cleavage of both carbon-bromine bonds to yield the *meta*-benzyne analogue **57**.²⁴⁸ *meta*-Benzyne **57** was found to react slowly with dimethyl diselenide by adduct formation, whereas charged phenyl radicals typically react with this reagent by SeCH₃ abstraction.^{49,50,52} Reaction of **57** with *tert*-butyl isocyanide occurs via both HCN abstraction and abstraction of two CN groups.²⁴⁸ The *ortho*-isomer of **57** was generated using an analogous procedure and its reactivity toward several reagents was compared to the reactivity of **57**.²⁴⁹ The *ortho*-isomer was found to abstract HSCH₃ from dimethyl disulfide, whereas **57** is unreactive toward this reagent. Addition followed by elimination of formaldehyde was the primary reaction observed when the *ortho*-isomer was allowed to react with 2,3-dimethoxy-1,3-butadiene; however, addition followed by elimination of methanol was observed for **57**. Despite the expectation that **57** might undergo both radical and non-radical reactions like the 3,5-didehydrobenzoate anion (**56**), no obvious radical reactions were reported. Overall, *meta*-benzyne **57** was found to be much less reactive than both its *ortho*-isomer and a related monoradical.

In order to better understand the electrophilic nature of *meta*-benzynes, the gas-phase reactivity of a series of positively charged *meta*-benzyne analogues was investigated (Chart 6) in an FT-ICR.²⁵⁰ The 3,5-didehydrobenzoyl cation (**58**) was found to undergo rapid, non-radical substitution reactions with a number of nucleophiles (Scheme 14), such as tetrahydrofuran, dimethyl disulfide, trimethylphosphine, triethylamine, and pyridine, to afford new *meta*-benzyne analogues, with rates that correlate with the proton affinity of the nucleophile as well as that of the leaving group.²⁵⁰ The fastest substitutions were found to occur when the nucleophile was more basic than the leaving group. The new *meta*-benzyne analogues generated from these reactions can also undergo further substitution reactivity at

the charge site. Interestingly, the charge site substitution was only observed in the presence of the *meta*-benzyne moiety, which is inconsistent with a mechanism involving ipso-substitution at the charge site. Hence, an addition/elimination mechanism was proposed to account for the necessity of the *meta*-benzyne moiety (Scheme 15). The existence of the addition intermediate was demonstrated by using a *meta*-benzyne with a *tert*-butyl substituent (**63**; Chart 6) since this benzyne formed a stable adduct that could be detected and whose reactions were examined. As expected, this intermediate showed nucleophilic reactivity due to the highly reactive phenide moiety. Although this substitution reactivity is an important component of the reactivity of charged *meta*-benzyne analogues, not all analogues of *meta*-benzynes undergo these reactions. Even though nucleophiles can add to **33**, it does not have a leaving group to eliminate. Charged *meta*-benzynes, such as **51** and **66** (Chart 6), which also do not have a leaving group, cannot undergo these addition/elimination reactions either. The electrophilic addition/elimination mechanism proposed for these reactions has little similarity to the concerted radical reactions considered by Chen et al.²⁴² Hence, the electrophilic reactivity represents a new and unexplored aspect of *meta*-benzynes' reactivity. It should be noted here that nucleophilic addition reactions of related *ortho*-benzynes are common, as discussed above, and that if the *ortho*-benzyne analog contains a leaving group, these reactions can be used to generate novel *meta*-benzyne analogs. For example, the N-(3,4-didehydrophenyl)-3-fluoropyridinium cation forms a new *meta*-benzyne analog (**64**; Chart 6) upon reaction with triethylamine.

In 2003, the gas-phase reactivity of the two isomeric chloro-substituted distonic *meta*-benzyne cations **62a** and **62b** (Chart 6) was examined and compared to that of **57** (Chart 6) in order to explore the effects of substitution on *meta*-benzynes' reactivity (the nonfluorinated analogs, *e.g.*, **59** - **61**, were also studied).²⁴⁷ The S-T splittings for various chloro-substituted benzynes were calculated at varying levels of theory and are shown in Table 4. In general, all three benzynes examined react via electrophilic pathways. However, the chloro-substituent was found to increase the electrophilicity of the *meta*-benzyne moiety as the chloro-substituted *meta*-benzyne analogues were found to react significantly faster (as much as two orders of magnitude in some cases) than the unsubstituted *meta*-benzyne **57**. For example, reactions of **57**, **62a** and **62b** with pyridine result in a charge-site substitution product, with the reaction occurring an order of magnitude faster for **62a** and **62b** than for **57**. **62a** and **62b** also abstract a thiomethyl group from dimethyl disulfide an order of magnitude faster than **57**. Interestingly, **62a** and **62b** react at nearly identical rates despite having a predicted ~4 kcal/mol difference in their S-T splitting, indicating that the reactions of these *meta*-benzynes occur via ionic rather than radical mechanisms. This mechanism avoids the costly uncoupling of the diradical electrons of these singlet diradicals in the transition states of radical reactions.

Following the above research, seven *meta*-benzyne derivatives (**51**, **65** and **68** – **71**, Chart 7) were examined both computationally and experimentally in order to explore the relationship between their structures and reactivity.²⁵¹ (U)BLYP/cc-pVDZ calculations for the parent *meta*-benzyne (**33**) indicated that the (singlet state) potential energy surface for the dehydrocarbon atom separation is very flat. The corresponding (U)BLYP/cc-pVDZ surfaces for **51**, **68** and **69** are also very flat²⁵¹ (Figure 19) even though the dehydrocarbon atom separations for **51** and **69** are predicted to be substantially smaller than either for the *meta*-benzyne (**33**) or **68**. Similar geometry differences for singlet *meta*-benzynes have previously been rationalized^{153,244,252} by taking into account the stabilizing or destabilizing effects of heteroatoms and substituents on their zwitterionic resonance structures (Scheme 16). Computational results are summarized in Table 5. In general, the NH⁺ group at C-R₃ in **51** stabilizes the allyl anion portion of the molecule and was predicted to shorten the dehydrocarbon atom separation (DAS) of **51**, as shown in Figure 19. Hydroxyl substitution

at R₂ in **51** destabilizes the allyl anion part of the molecule due to unfavorable π,π -interactions with the oxygen atom's lone pair electrons and was predicted to result in an increased DAS for **68**, while substitution at the same position with a less powerful π -donating substituent, chlorine atom, was predicted not to change DAS for **69** (Figure 19). Hydroxyl substitution at C-8 in **65** to yield **70** was predicted to have a similar effect as for radical **51**; however, this substitution so drastically flattens the potential energy surface that there is only ~0.4 kcal/mol difference in energy between structures for which the dehydrocarbon atom separation is 1.6 and 2.0 Å (Figure 20). Based on these results, *meta*-benzynes with large DAS, such as **68**, **70** and **71**, may be expected to undergo radical reactions, such as hydrogen atom abstraction from tetrahydrofuran, much faster than the other *meta*-benzynes. This was indeed observed. While radical reactions occur much more slowly for **51** than the corresponding monoradical, e.g., hydrogen and iodine atom abstractions from tetrahydrofuran and allyl iodide, respectively, occur with efficiencies of 0.7% and 10% for **51** but 38% and 76% for the monoradical, respectively,²⁵⁰ the hydroxy-substituted diradical **68** abstracts a hydrogen atom from tetrahydrofuran much faster than **51** (efficiency = 30%), at an efficiency that parallels that of the corresponding monoradical. Iodine atom abstraction from allyl iodide also occurs very quickly (efficiency = 32%). These results unambiguously demonstrate that appropriate substituents can be used to “tune” the reactivity of *meta*-benzynes from mildly carbocationic to radical-like.

In 2012, the Kenttämä laboratories reported on the gas-phase reactivity of an interesting distonic diradical cation, the 4,5-didehydroisoquinolinium cation **72** (Chart 7), with one carbon atom between the radical sites like in *meta*-benzynes but the radical sites in close spatial proximity.²⁵³ **72** (S-T splitting = -0.6 kcal/mol) is a positively charged analog of 1,8-didehydronaphthalene, which has shown interesting chemical properties^{254–257} attributed to its nearly degenerate singlet and triplet states (S-T splitting = -0.9 kcal/mol).²⁵⁸ Examination of the gas-phase reactions of **72** suggested that they mostly occur via radical mechanisms, as opposed to the *meta*-benzyne analogs discussed above. Calculation of the enthalpy changes for hydrogen atom abstraction reactions of **72** with tetrahydrofuran (UMPW1K/6-31+G(d,p)//UMPW1K/6-31+G(d,p) level of theory) predicted that hydrogen atom abstraction by the C4 radical site is favored by ~2 kcal/mol over abstraction by the C5 radical site. This finding was confirmed by experimental examination of reactions of **72** with cyclohexane. The product of the first hydrogen atom abstraction was isolated and allowed to react further with cyclohexane in order to determine the efficiency of the second hydrogen atom abstraction (efficiency = 2%). This efficiency is comparable to the efficiency of hydrogen atom abstraction by the 5-dehydroisoquinolinium cation (efficiency = 3%) and substantially different from that of the 4-dehydroisoquinolinium cation (efficiency = 18%). Hence, the C4 radical site was experimentally proven to abstract the first hydrogen atom from cyclohexane, followed by the C5 radical site.²⁵³ Analogous to the reactions with tetrahydrofuran and cyclohexane, the reactions of **72** with dimethyl disulfide and allyl iodide are likely to be initiated by atom or group abstraction by the C4 radical site.

4.2.1 *para*-Benzynes—The first studies on *para*-benzynes were reported in the early 1960's.^{151–153} Berry and co-workers attempted to generate the *para*-benzyne (**34**, Chart 8) in the gas phase by flash vacuum pyrolysis of benzenediazonium-4-carboxylate, and monitored the products by time-resolved mass spectrometry and flash-absorption optical spectroscopy.^{200,259} While the authors claimed that the formation of a *para*-benzyne had been observed, its long lifetime of more than 2 minutes makes these results speculative^{68b,112} since the lifetimes of *ortho*- and *meta*-benzynes in the gas phase have been reported to be on the order of milli- and microseconds, respectively.²⁰⁰ In 1972, Jones and Bergman reported that gas-phase pyrolysis of an enediyne yielded a mixture of isomeric enediynes, which could only be explained by the formation of the symmetrical, cyclic *para*-

benzyne (**34**) as an intermediate.²⁶⁰ Since then, the process of cycloaromatization of enediyne has been known as the Bergman cyclization reaction.

By the 1990s, several theoretical^{214,215,261–263} and experimental studies^{237,264,265} regarding *para*-benzyne and its derivatives had appeared. However, the great interest in *para*-benzyne analogues over the last 20 years is related to their identification as the reactive intermediates of the very potent naturally occurring enediyne anti-tumor antibiotics.^{32,179,266} When intercalated into double-stranded DNA, the enediyne “warhead” of these anti-tumor antibiotics undergoes Bergman cyclization to form *para*-benzyne analogs. Each radical site of the diradical is thought to abstract a hydrogen atom from the sugar-phosphate backbone of each stand of the DNA double helix, resulting in irreversible DNA scission and eventual apoptosis of the cell.^{32,179,266} The absolute heat of formation^{111,217} (127 ± 3 kcal/mol), electron affinity¹⁸² (1.265 ± 0.008 eV) and S-T splitting¹⁸² (-3.9 ± 0.4 kcal/mol) of **34** (Chart 8) were measured by Squires and co-workers. While the S-T splitting of **34** is relatively small in comparison to the *ortho*- and *meta*-isomers, the experimental and computational work of Chen and coworkers showed that even this weak interaction between the diradical electron pair in **34** and its analogues severely impedes radical reactivity.^{241,242} Related heteroaromatic *para*-benzynes (**73** and **74**; Chart 8) and their analogues have been much less studied than **34**. The structures and energetics of **34** and the pyridynes **73** and **74** have been compared by Debbert and Cramer.¹⁵³ Calculated S-T splittings, at various levels of theory, were reported for **34**, **73** and **74** (Table 6) and compared to the experimentally determined¹⁸² S-T splitting for **34**. Incorporation of a nitrogen atom into the aromatic ring of **34** was predicted to increase the S-T splitting, which is associated with a larger diradical stabilization energy of the singlet state than the triplet state caused by the energetically favorable mixing of a zwitterionic mesomer into the overall electronic structure of 2,5-didehydropyridine and -pyridinium cation.¹⁵³

In 1994, Wenthold and Squires successfully synthesized the *para*-benzyne anion and examined its reactivity (Table 7) in the gas phase.^{245,246} The *para*-benzyne anion was generated in a flowing afterglow triple quadrupole mass spectrometer^{267,268} by using the same method employed to generate the *meta*-benzyne anion discussed above.²⁴⁵ The *para*-benzyne anion was readily distinguished from the lower-energy *ortho*-isomer as sequential abstraction of two sulfur atoms from CS₂ was observed for the *para*-benzyne anion, while only electron transfer was observed for the *ortho*-isomer. The *para*-benzyne anion was found to react with nitric oxide by sequential addition of two NO molecules, while no reaction was observed for the *ortho*-isomer.²⁴⁵ The dominance of the anion site of the *para*-benzyne anion in these reactions is consistent with its strongly basic and nucleophilic character. Reactions of the *para*-benzyne anion with a series of alkyl halides were also investigated in order to determine whether halogen atom and/or alkyl group abstractions, such as those reported by Kenttämä and co-workers for negatively charged *meta*-benzyne analogs,^{47–49,52} would take place. However, no halogen atom or alkyl group abstractions were observed. The *para*-benzyne anion was also allowed to react with several molecules with relatively weak C-H bonds, such as methanol, dimethyl ether, toluene, and cyclohexa-1,3-diene, to determine whether hydrogen atom abstraction would take place. However, this was not observed. Instead, proton transfer from methanol, cyclohexa-1,3-diene, and toluene to the anion was observed. Overall, the negative ion chemistry of the *para*-benzyne anion appears to dominate its reactivity toward the reagents examined, just as was observed for its -isomer, *meta*.^{245,246}

Most recently, the formation and reactivity of three gaseous charged *para*-benzyne analogues (**74** – **76**; Chart 8) were examined.²⁶⁹ Diradicals **74** – **76** were generated in an FT-ICR by using two consecutive SORI-CAD events to homolytically cleave C-I and/or C-NO₂ bonds in N-deutero-1,4-diiodoisoquinolinium, 5-iodo-8-nitroisoquinolinium, and 2,5-

diiodopyridinium cations, respectively. When diradicals **74** - **76** were allowed to react with cyclohexane, tetrahydrofuran, allyl iodide, dimethyl disulfide, and *tert*-butyl isocyanide, no radical reactivity was observed. Hence, the diradicals did not behave like *para*-benzyne analogues expected to display reactivity similar to that of related monoradicals.

Experimental and computational results indicate that the three *para*-benzyne analogs were never generated due to ring-opening of their monoradical precursors upon the second SORI-CAD event.

4.3 Reactions of Aryl Mono- and Diradicals with Biological Molecules

Radical attack on various biological molecules has been shown to play an important role in a number of biological processes, such as oxidative damage of proteins,²⁷⁰ carcinogenesis, and chemotherapy via drug-mediated cleavage of DNA.^{30–33} Among the various radicals involved in DNA degradation, hydroxyl radicals have been the most thoroughly studied. As they are abundant in human cells and highly reactive, they pose a great hazardous potential.^{271–273} Hydroxyl and other electrophilic radicals are known to undergo addition at position C5 of the C5–C6 double bond of pyrimidine bases and at positions C4, C5 and C8 in purine bases.^{274–276} As mentioned above, some antitumor drugs containing the enediyne warhead generate *para*-benzyne analogs that are believed to abstract a hydrogen atom from the sugar moiety of each strand of double-stranded DNA to irreversibly cleave DNA.^{30–33,277,278} This has generated a lot of interest in the properties of such diradicals. Unfortunately, experimental studies on their reactivity are limited by the difficulty to cleanly generate them as well as their short lifetimes in solution.

Experimental studies on the reactivity of (nucleophilic) phenyl radicals toward DNA components in solution have revealed that these radicals abstract hydrogen atoms from several sites in monosaccharides but display slightly different selectivity than the electrophilic hydroxyl radicals.^{29,275} Further, both hydroxyl radicals and phenyl radicals were found to predominantly add to nucleobases, and, more specifically, preferentially to the C5–C6 double bond of cytosine, uracil, thymine, and their derivatives. The nucleophilic phenyl radicals studied prefer addition at the C-6 carbon while the electrophilic hydroxyl radicals prefer addition at the C-5 carbon.^{272,273,275} In order to further explore the fundamental factors controlling the reactivity of phenyl radicals toward biomolecules, the distonic ion approach has been employed.^{61,63,279–284} Positively charged phenyl radicals were generated in an FT-ICR mass spectrometer by using the methods described above. Hydrogen atom abstraction was the only reaction observed for 2-deoxy-D-ribose,⁶¹ in qualitative agreement with the previously reported solution results discussed above.^{30,285} The reactivity of the positively charged phenyl radicals was found to correlate with the electron affinities of the radical sites, as expected.^{130,137–139} Hence, hydrogen atom abstraction from 2-deoxy-D-ribose in DNA can be facilitated by increasing the electrophilicity of the radical. Increasing the electrophilicity of the radicals examined also led to an increase in reactivity toward the nucleobases uracil, 1-methyl uracil, adenine, thymine and 1-methylthymine.⁶¹ Addition was reported to be the predominant reaction pathway for adenine and uracil, analogous to the reactivity of the hydroxyl radical in solution,³⁰ whereas hydrogen atom abstraction predominates for thymine. These results confirmed that phenyl radicals can damage DNA directly via hydrogen atom abstraction from the sugar moiety as well as indirectly via attack at the nucleobase, as proposed earlier.²⁷⁵

During 2005 – 2011, the reactivity of a large number positively charged phenyl radicals (e.g., radicals **14**, **15**, and **17**; Chart 2 and 3) toward amino acids was investigated using the distonic ion approach in an FT-ICR mass spectrometer.^{63,64,283} Again, the reactivity of the charged phenyl radicals was found to correlate with their calculated EA_vs. Examination of

the reactions of the charged phenyl radicals with glycine, alanine, valine, proline, cysteine and methionine revealed unprecedented abstraction of various functional groups in addition to the expected hydrogen atom abstraction.⁶³ These abstractions include NH₂ abstraction for all amino acids except proline, SH abstraction from cysteine, and SCH₃ abstraction from methionine, all of which were presumed to occur via a nucleophilic addition-elimination pathway rather than a radical mechanism. The rate of these functional group abstractions was found to increase with increasing electrophilicity of the radical sites in the radicals. The ratio of hydrogen atom to amino group abstraction was reported to increase as the size of the amino acid's side chain increases, implying that a certain amount of hydrogen atoms are abstracted from the alkyl side chain of the amino acids instead of just from the α -carbon (the thermochemically favored site). Similar reactivity trends were observed for phenylalanine, tyrosine, and tryptophan.^{63,64} However, the phenyl radicals examined were found to predominantly undergo addition to the aromatic ring of these aromatic amino acids followed by fragmentation. Radical addition to the aromatic ring of tyrosine followed by C α – C β bond cleavage, yielding either a side chain abstraction product or loss of OH (if addition occurs at the *para*-position of tyrosine), had not been reported previously (Schemes 17 and 18). Numerous positively charged phenyl radicals were later allowed to react with several partially isotope labeled amino acids.²⁸¹ Results obtained for partially deuterium-labeled leucine and proline indicated that only a fraction of the hydrogen atom abstractions, if any, occurred at the α -carbon of the amino acid. Instead, the majority of hydrogen atoms are abstracted from the alkyl side chain. Labeling of the lysine side chain with ¹⁵N revealed that positively charged phenyl radicals abstract NH₂ from both the N-terminus and the side chain of lysine as both NH₂ and ¹⁵NH₂ abstractions were observed. The mechanism of NH₂ abstraction was examined computationally. The results indicate that nucleophilic addition of the amino group to the radical carbon is kinetically favored, implying that the radical carbon in positively charged phenyl radicals is the most likely nucleophilic addition site for amino acids.

In 2006, the reactions of several aryl diradicals (e.g., radical **65** and its N-methylated derivative as well as the isomeric 4,6-didehydroisoquinolinium cation and its N-methylated derivative) with monosaccharides were examined.²⁸² Hydrogen atom abstraction was the major reaction observed for aryl diradicals with small S-T splittings, while those with large S-T splittings predominantly formed addition products via non-radical mechanisms. The relative reactivities of diradicals having similar S-T splittings were reported to be controlled by their calculated EA_{v,s}. It was concluded that a large EA_v can counterbalance the radical-reactivity hindering effect of the singlet coupling of the diradical electrons. The results of these studies revealed that S-T splitting and EA_v, two of the four reactivity-controlling parameters found by that time to affect the reactivity of aryl diradicals toward simple organic substrates,^{247,251} also affect the reactivity of diradicals toward monosaccharides. These results also imply that some aryl diradicals are not capable of damaging DNA by hydrogen atom abstraction.

Most biological molecules, such as peptides, nucleosides and oligonucleotides, are nonvolatile, thermally labile compounds that readily decompose upon heating before evaporation occurs. Hence, their introduction as neutral molecules into the high vacuum environment of a mass spectrometer is difficult. In order to overcome this obstacle and effectively evaporate neutral, intact biomolecules into the gas phase, laser-induced acoustic desorption (LIAD) was employed.^{286,287} LIAD was reviewed in 2012.²⁸⁸ LIAD enabled the expansion of the study of the reactivity of aryl mono- and diradicals to peptides and oligonucleotides, which facilitated the acquisition of more relevant information about protein and DNA degradation by aryl radicals. In 2009, two charged phenyl radicals, the *N*-phenyl-3-dehydropyridinium cation (the *N*-phenyl derivative of **17**; Chart 3) and the *N*-(3-dehydrophenyl) pyridinium cation (**14**, Chart 2), were allowed to react with several

dipeptides desorbed by LIAD into an FT-ICR mass spectrometer.²⁸⁰ The results of these experiments demonstrated that the N-terminal amino acid of dipeptides is more susceptible to radical attack than the C-terminal amino acid. Hydrogen atom abstraction was the predominant reaction for all dipeptides examined. The ability of a phenyl radical to damage dipeptides was found to increase with increasing (calculated) EA_v of the radical site, as expected. In most cases, radical reactivity toward a given amino acid residue appeared to be influenced by both the location of the amino acid in the peptide as well as the other amino acids present, implying that gas-phase radical reactivity studies of free amino acids are not good indicators for reactivity toward the same amino acids in dipeptides.

A study on the reactivity of the N-methyl-6,8-didehydroquinolinium cation (Scheme 19) toward amino acids and dipeptides was reported in 2010. LIAD was used to desorb the neutral dipeptides into the FT-ICR mass spectrometer.²⁸⁴ Reactions of the diradical with free amino acids resulted in the abstraction of two hydrogen atoms, H_2O abstraction, addition followed by elimination of CO_2 , addition followed by elimination of $HCOOH$ and formation of a stable adduct. When allowed to react with dipeptides, these same pathways and one additional pathway, addition followed by C-terminal (R_C) amino acid elimination (addition – $CO - NH=CHR_C$) (Scheme 19), were observed. Most of the reactions of this diradical are thought to be initiated by hydrogen atom abstraction, radical addition to an aromatic ring, or nucleophilic attack by an amino group of a dipeptide at the diradical, analogous to reactions of related monoradicals. Abstraction of two hydrogen atoms by the diradical from peptides (Scheme 20), as well as addition of the diradical to an aromatic ring if present in the C-terminal peptide, leading to N-terminal (R_N) amino acid elimination via loss of CO and $NH=CHR_N$ (Scheme 21), likely occur via radical mechanisms. It should be noted that addition to aromatic rings of amino acids and peptides by the monoradicals was found to lead to side-chain abstraction.²⁸⁴ Carbon 6 (C-6) of the diradical appears to be its most reactive site regardless of the reaction mechanism. After reaction of this radical site, the unquenched radical site at C-8 reacts to afford products not observed for analogous monoradicals. The N-methyl-6,8-didehydroquinolinium cation was reported to react similarly with aliphatic dipeptides and those aromatic dipeptides not containing an aromatic amino acid in the C-terminus.

In order to improve the understanding of interactions of aryl diradicals with DNA, dinucleoside phosphates were successfully desorbed into the gas phase in an FT-ICR by using LIAD, and the reactivity of aryl mono- and diradicals toward these net-neutral substrates were studied.^{280,283} The structure of the aryl diradical was demonstrated to have a major effect on its reactivity toward dinucleoside phosphates. As true for the reactivity of *meta*-benzyne analogs toward simple organic substrates,^{247,251,282} the reactivity of the *meta*-benzyne analogs studied (5,7- (**65**) and 6,8-didehydroquinolinium cation (**71**; Chart 7)) also appeared to be governed by not only the S-T splitting and EA_v , but also the dehydrocarbon atom separation. The structure of the dinucleoside phosphate was also found to influence its reactions with aryl (bi)radicals. Thymine was the least susceptible to radical attack among the nucleobases studied, while cytosine was the only base to undergo skeletal fragmentation (loss of $HNCO$) upon diradical attack. This fragmentation was not observed when cytosine was allowed to react with monoradicals. The same skeletal fragmentation has been reported for CAD of an ionic cytosine adduct of an enediyne.²⁸⁹ These findings indicate that the purine bases, adenine and guanine, are most susceptible to (bi)radical attack, while thymidine is the least susceptible.

4.4 Charged Aryl Triradicals

Aromatic carbon-centered σ,σ,σ -triradicals are reactive intermediates with three formally unpaired electrons (Chart 9). According to Salem and Rowland, they can be defined as

molecules with three electrons distributed in three (nearly) degenerate orbitals.^{290,291} The complex electronic states of triradicals make them extremely interesting to study experimentally and theoretically. However, the first theoretical and experimental studies on such triradicals were reported only 14 and 10 years ago, respectively.^{292,293} Therefore, it is not surprising that in a review on arynes published in 2003, only a short note was devoted to trihydrobenzenes.¹⁷⁹ Interestingly, much more was known about tetrahydrobenzenes at that time.¹⁷⁹

The first computational studies carried out at the CCSD/DZP and BLYP/6-31G* levels of theory predicted that 1,2,4-trihydrobenzene (**78**; Chart 9) and 1,3,5-trihydrobenzene (**79**; Chart 9) are higher in energy than 1,2,3-trihydrobenzene (**77**, Chart 9) by 4.0 and 11.8 kcal/mol, respectively.²⁹² Based on the calculated energy of an isodesmic reaction of **77** and benzene, the delocalization of the three electrons over three σ -type orbitals lowers the energy of the system by 3.9 kcal/mol.²⁹² The first experimental studies on trihydrobenzenes were carried out by Wenthold and co-workers who measured the gas-phase heat of formation ($\Delta H_{f,298}$) of **78** to be 179.1 ± 4.6 kcal/mol.²⁹³ Based on these calculations, **78** has a 2A_1 ground state, with a 2B_2 state higher in energy by only 1 kcal/mol. Analysis of the homolytic bond dissociation energies of benzene indicated stabilizing interactions between the two electrons in *meta*-positions in **78**.²⁹³ Therefore, triradical **78** is best described as a phenyl radical weakly interacting with a *meta*-benzyne moiety.

In the following years, several computational studies on the trihydrobenzenes **77**, **78** and **79**, discussed separately or all together, were reported by Krylov et al.,^{291, 294–298} Sander et al.,^{297,299,300} Wenthold,^{302,303} Nguyen et al.,³⁰⁴ and, more recently, by Fishtik³⁰⁵ and Chattopadhyay et al.^{306–308} In agreement with Wenthold's predictions,²⁹³ the electronic ground state of 1,3,5-trihydrobenzene (**79**) was assigned to be 2A_1 and a closely lying 2B_2 excited state was calculated to be higher in energy by 1 kcal/mol by using various computational methods.^{291,294,295,304} Sander and co-workers recently demonstrated that the ground state of 1,3,5-trihydrobenzene is not affected by substitution of the three hydrogen atoms with three fluorine atoms.³⁰⁰ The ground state of the least studied isomer, the 1,2,4-trihydrobenzene (**78**), was calculated by Krylov and co-workers to be $^2A'$.²⁹⁴ The electronic ground state of 1,2,3-trihydrobenzene (**77**) was initially assigned to be 2B_2 .^{294,295} However, later experimental and theoretical studies demonstrated that the ground state is actually 2A_1 , with the 2B_2 state lying about 1–2 kcal/mol higher in energy.^{297,298,299} The 2A_1 state was confirmed to be the ground state of the 1,2,3-isomer in most recent computational studies using recently developed theoretical methods, the improved virtual orbital-complete active space configuration interaction (IVO-CASCI) and complete active space based state specific multi-reference Møller–Plesset perturbation theory (SS-MRMPPT) methods.^{306–308} The IVO-CASCI method was found to be a good yet cheaper alternative for CASSCF methods.^{306,307}

Nguyen et al. examined the thermochemical properties of 1,3,5-trihydrobenzene (**79**; Chart 9).³⁰⁴ The gas-phase heat of formation ($\Delta H_{f,298,1atm}$) was calculated to be 179 ± 2 kcal/mol (CBS-QB3 method), which is in good agreement with the experimentally determined value.²⁹³ In addition, the electron affinity ($EA = 1.6 \pm 0.2$ eV), ionization energy ($IE_a = 7.2 \pm 0.2$ eV), and the proton affinity ($PA = 227 \pm 4$ kcal/mol) of **79** were calculated in this work.³⁰⁴ The consequences of the different bonding interactions between the radical centers in all three isomeric trihydrobenzenes were evaluated by Krylov and co-workers.^{294,295} Their electronic structure calculations for the ground state of 1,2,3-isomer (**77**; Chart 9) showed that the interactions between the *ortho*-positioned radical centers prevail over the *meta*-radical centers. However, the stabilizing interactions between the *ortho*-radical centers are weaker than in the *ortho*-benzyne due to electron delocalization over the three dehydrocarbons. In the case of the 1,2,4-isomer (**78**; Chart 9), interactions

between both the *ortho*- and *meta*-radical centers have a significant effect on its structure, with the *ortho*-radical centers being more important. In 1,3,5-tridehydrobenzene (**79**; Chart 9), the distance between two of the radical centers is much shorter than the others. This short distance is only slightly longer than for *meta*-benzyne, which suggests only weak interactions between this *meta*-benzyne moiety and the remaining radical center in the 1,3,5-isomer.²⁹⁴ The doublet-quartet (D-Q) splittings and, therefore, the strengths of the bonding interactions, decrease in the following order: 1,2,3 > 1,2,4 > 1,3,5.²⁹⁴ The calculated strengths of the interactions between each radical center and the remaining diradical moiety in each triradical are defined as triradical stabilization energies (TSE). The TSEs were derived from calculations of enthalpy changes of the isodesmic hydrogen atom transfer from the pyridinium cation to a triradical cation to give the corresponding biradical and monoradical cations. TSEs provide a direct indication of the stabilization (TSE > 0) or destabilization (TSE < 0) involved when all three radical sites are present in the same molecule. TSEs were found to vary from 0.4 to 30 kcal/mol. In benzyne, stabilizing interactions between the two radical centers (biradical stabilization energies or BSE) were found to be in a similar range (4–32 kcal/mol).²⁹⁴ However, as shown by Fishtik, the proper evaluation of stabilization/destabilization effects strongly depends on the approach for their determination.³⁰⁵ The most commonly used approach, a single reaction scheme, balances the number and types of groups or interactions in molecules involved in the reaction scheme and results in isodesmic stabilization/destabilization energies. An alternative method, the group additivity approach, evaluates stabilization/destabilization energies of molecules as the difference between the enthalpy of formation calculated via a group additivity scheme and the experimental enthalpy of formation. The results of the second method significantly depend on the proper selection of reference molecules and the group additivity scheme used for the calculations.³⁰⁵ Based on this approach, the tridehydrobenzenes exhibit a destabilizing effect,³⁰⁵ which is contrary to previous findings.²⁹⁴ The thermochemical properties of tridehydrobenzenes were recently reviewed elsewhere.^{301,303}

Theoretical studies on the effects of substitution on the geometries and energetics of tridehydrobenzenes are limited to three publications.^{300,309,310} Sander and co-workers found that fluorination of **79** (Chart 9) has only a marginal effect on the structures and relative stabilities of the two lowest energy doublet states, the 2A_1 ground state and the 2B_2 lowest-energy excited state.³⁰⁰ Recently, Krylov et al. reported that the doublet state ordering in 2,4,6-tridehydropyridine **85** (Chart 9) is reversed compared to **79**.³¹⁰ According to these calculations, the 2B_2 ground state of **85** is 14.1 kcal/mol more stable than the 2A_1 state, whereas the D-Q splitting (28.2 kcal/mol) is similar to that of 1,3,5-tridehydrobenzene.

Extensive quantum chemical characterization of the structures, some thermochemical properties and D-Q splittings of the tridehydropyridinium cations **86** - **91** (Chart 10) were reported recently by Nash et al.³⁰⁹ Some of the results of this work are summarized in Figure 21. The results are used in this section to rationalize the reactivities of triradicals **86** and **87**. Nash showed that for **86**, similarly to **85**,³⁰⁹ the 2B_2 state is lower in energy than the 2A_1 state by 11.1–16.6 kcal/mol, depending on the level of theory used, thus making 2B_2 the ground state. The ground state of the 3,4,5-tridehydropyridinium cation **87**, as opposed to the parent tridehydrobenzene **77**, is 2B_2 , which is better resonance stabilized than the 2A_1 state. The 2,3,4-isomer **89** was predicted to have $^2A'$ ground state, and it therefore differs significantly from **77** and **87**. The tridehydropyridinium cations **88**, **90** and **91** have been predicted to have $^2A'$ ground state, similarly to their parent tridehydrobenzene **78**.

The chemical properties of the tridehydrobenzenes are intriguing as they may be influenced by both their ground electronic state and the low-lying excited electronic states. Unfortunately, examination of the reactivity of tridehydroarenes in condensed phases is challenging due to the difficulty of cleanly generate them. Hence, for a long time,

experimental studies on the tridehydrobenzenes were limited to the above mentioned thermochemical measurements on 1,3,5-tridehydrobenzene by Wenthold et al.²⁹³ and matrix isolation and IR detection of 1,2,3-tridehydrobenzene and perfluoro-1,3,5-tridehydrobenzene by Sander and co-workers.^{299,300} In the latter study, the 1,2,3-tridehydrobenzene (**77**) and the trifluoro-1,3,5-tridehydrobenzene were generated in low temperature matrices and IR spectra were measured for them.^{299,300}

Considering such a small number of experimental studies, recently published reactivity studies on related σ,σ,σ -triradicals (Chart 10), the 2,4,6-tridehydropyridinium cation (**86**), 3,4,5-tridehydropyridinium cation (**87**), 3-hydroxy-2,4,6-tridehydropyridinium cation (**98**), 1,2,4-tridehydropyridinium cation (**93**), and 1,2,6-tridehydropyridinium cation (**95**), are of great importance since they provide first evidence for the influence of the electronic structures of these triradicals on their chemical properties.^{311–314} A review of these studies is provided below.

4.4.1 Ion-molecule Reactions—The tridehydropyridinium cations **86**, **87** and **98** (Chart 10) were generated in a similar manner as demonstrated for **86** in Scheme 22 and Figure 22. In the first step, a suitable iodoprecursor was protonated by using CI or self-CI. Two or three SORI-CAD events were employed to cleave off iodine atoms and/or HI molecules. In order to generate triradical **95** (Chart 10), 2,6-diiodopyridine was protonated and subjected to SORI-CAD. The loss of an iodine atom and HI were observed. The product of the iodine atom loss, a monoradical, was isolated and subjected to a second SORI-CAD event. Again, both an iodine atom and HI were lost. The desired triradical **95** was the product of the HI loss in the second SORI-CAD event (Scheme 23). Generation of triradical **93** (Chart 10) was carried out in similar manner.

The 2,4,6-tridehydropyridinium cation (**86**) was found³¹¹ to form products similar to those of the related mono- and diradicals. However, it reacts with most reagents more efficiently than related biradicals and at about equal efficiency as related monoradicals. The reactivity of **86** clearly indicates the presence of three radical sites as it abstracts three SCH₃ groups from dimethyl disulfide molecules, three hydrogen atoms from cyclohexane, and three iodine atoms from allyl iodide molecules. However, the three radical sites are not equally reactive. For example, the first and third iodine atom abstractions occur faster than the second. This finding suggests that the first iodine atom is abstracted by the radical at the 6-position to generate 2,4-didehydro-6-iodopyridinium cation (Scheme 24).

The most important ionic resonance structure of **86** (Scheme 25) and the TSE determined for **86** (Figure 23) explain the different rates observed for the three iodine atom abstractions. The ionic resonance structure of **86** indicates a stronger coupling between the radical sites at positions 2 and 4 (or 4 and 6) than at 2 and 6, which suggests that the radical site at position 6 (or 2) should be more reactive than the others. This suggestion is supported by calculated TSEs, which indicate that the removal of the radical site in position 6 (or 2) is preferred (Figure 23). The second iodine atom abstraction is slow as a result of the strong coupling between the remaining radical sites 2 and 4. However, once the second iodine atom is abstracted, a phenyl radical is produced that undergoes a fast iodine atom abstraction. Triradical **86** is calculated to have a doublet ground state (D-Q splitting: -29.3 kcal/mol; BD(T)/cc-pVTZ//BPW91/cc-pVDZ), which explains why it undergoes radical reactions faster than the analogous diradical whose reactions are hindered by its singlet ground state. The triradical reacts with allyl iodide and dimethyl disulfide at about the same efficiency as related monoradicals. Apparently, the large EA_v of **86** counterbalances most of the radical reactivity hindering effect caused by the coupling of the three unpaired electrons.³¹¹

The successful synthesis of the precursor for triradical **98** enabled the investigation of the effects of substitution on the reactivity of triradical **86**.³¹³ The results were surprising. Triradical **98** was found to behave just like related monoradicals and not like related triradicals. While three consecutive abstractions of iodine atom from allyl iodide and thiomethyl group from dimethyl disulfide molecules were observed for triradical **86**, only one iodine atom and one thiomethyl group abstraction was observed for triradical **98**. Triradical **98** reacts faster with allyl iodide and dimethyl disulfide than **86**. These results can be explained based on a specific ionic resonance structure and TSEs (for unsubstituted and hydroxyl substituted triradicals, see Scheme 26). Based on the ionic resonance structures of **86** and **98** shown in Schemes 25 and 26, respectively, the coupling between the radical sites at positions 2 and 4 (or 4 and 6) in **98** is expected to be even stronger than coupling in **86**. Therefore, the interactions between the radical sites at positions 2 and 6 in **98** should be weaker than in **86**, which suggest that the radical site at position 6 (or 2) should be more reactive than the radical site at position 6 (or 2) in **86**. These predictions are supported by TSEs calculated for triradical **98**.

Another tridehydropyridinium cation whose chemical properties have been characterized is the 3,4,5-tridehydropyridinium cation (**87**; Chart 10). The reactivity of **87** reveals the presence of three radical sites as it abstracts both SCH₃ and HSCH₃ groups from dimethyl disulfide molecules, three hydrogen atoms from cyclohexane, and an iodine atom as well as HI from allyl iodide molecules. Triradical **87** forms products similar to those of the related mono- and diradicals. However, it reacts with most reagents more efficiently than the related mono- and biradicals. The high reactivity of **87** can be attributed to its very high EA_v (7.20 and 6.70 eV for the ²B₂ and ²A₁ states, respectively; (U)BLYP/aug-cc-pVDZ//((U)BLYP/cc-pVDZ, Figure 21) compared to related mono- and diradicals.³⁰⁹ Thus, it appears that the very high electron affinity of **87** counterbalances the reduction in the radical reactivity caused by the coupling of the three unpaired electrons. This is also likely to be the case for triradical **86** discussed above (EA_vs for the (ground) ²B₂ and (excited) ²A₁ states: 7.01 and 7.18 eV, respectively; calculated at the (U)BLYP/aug-cc-pVDZ//((U)BLYP/cc-pVDZ level of theory).³⁰⁹

In principle, any one of the three radical sites in **87** may be involved in the first reaction. The diradical formed upon the abstraction of the first iodine atom from allyl iodide by **87** reacts with allyl iodide by abstraction of allyl-H, allyl, and HI (Scheme 27). The branching ratios for these three secondary products are nearly identical to those observed for the reaction of the 3,4-didehydropyridinium cation with allyl iodide. Moreover, abstraction of an iodine atom, a reaction characteristic of *meta*-benzyne analogues, was not observed. Hence, the diradical formed from **87** upon the first iodine atom abstraction is more likely to have the iodine atom attached to C5 than to C4 (Scheme 27).

Additional evidence for the regioselectivity of above radical reactions was obtained³⁰⁹ by isolating the primary product ions formed upon abstraction of one and two hydrogen atoms from cyclohexane by **87** and allowing them to react with cyclohexane (Scheme 28). The product ion formed upon abstraction of one hydrogen atom reacts with cyclohexane by abstraction of two hydrogen atoms with an efficiency of 2%. This reaction efficiency is close to the efficiency of an *ortho*-benzyne analogue, 3,4-didehydropyridinium cation. Thus, the first hydrogen atom abstraction by **87** appears to occur at C3, which is in agreement with the results obtained for allyl iodide. The product ion formed upon abstraction of two hydrogen atoms reacts with cyclohexane by one hydrogen atom abstraction with an efficiency of 20%. The comparison of this result with the efficiencies of related monoradicals suggests that the first two hydrogen atoms are abstracted by the radical sites C3 (first) and C4 (second), leaving the radical site at C5 as the last to react. The proposed regioselectivity for reactions of **87** with allyl iodide and cyclohexane is in good agreement

with calculated triradical stabilization energies for this triradical.³⁰⁹ All these findings indicate much stronger *ortho*-interactions than *meta*-interactions between the radical sites in **87**.

The most recent kinetic reactivity study on isomeric triradicals based on the pyridine skeleton focused on the chemical properties of the 2,4-didehydropyridine radical cation **93** and the 2,6-didehydropyridine radical cation **95** (Scheme 29).³¹⁴ The reactivity of **93** and **95** indicates the presence of three radical sites as they abstract an iodine atom and HI from allyl iodide molecules. The main reactions observed for **93** and **95** with cyclohexane, allyl iodide and dimethyl disulfide were hydride abstraction, C₃H₅ group abstraction and electron transfer, respectively. These reactions are either minor or do not occur at all for triradicals **86** and **87**. The electrophilic, carbocation-like reactivity observed for **96** and **98** can be rationalized based on their ionic resonance structures (Scheme 29). Triradicals **93** and **95** also undergo a radical reaction, iodine atom abstraction, due to the presence of a (formally) unpaired electron in all resonance structures.

One of the goals of the reactivity studies on σ,σ,σ -triradicals was to improve the understanding on the factors controlling their reactivity.^{311–314} However, the reported reactivity studies are limited only to five different tridehydropyridinium cations.^{311–314} This limited amount of data may not be sufficient to make definite conclusions on what factors control the reactivity of these reactive species. However, possible reactivity controlling factors are discussed below.

The D-Q splittings (calculated at the BD(T)/cc-pVTZ//BPW91/cc-pVDZ level of theory) for most studied tridehydropyridinium cations are similar to the D-Q splittings of the related tridehydrobenzenes (Figure 21).³⁰⁹ This trend does not hold for **88** and **91** (Chart 10), whose D-Q splittings are lower by 7–9 kcal/mol compared to the related 1,2,4-tridehydrobenzene (**78**; Chart 9). This is likely due to the proximity of the *ortho*-benzyne moiety to the NH⁺ group in these molecules. However, the D-Q splittings do not seem to have a major influence on the reactivity of the tridehydropyridinium ions, since triradicals with larger D-Q splittings can have higher reaction efficiencies. The observed trend of reaction efficiencies seems to be related to the EA_v values of the radical sites in the triradicals. This trend is similar to that observed for monoradicals, where a higher EA_v usually results in faster reactions.^{67,68}

The results of the above studies show a greater complexity for the triradical systems compared to related mono- and diradical systems. This is reflected in the reaction efficiencies and reaction products observed for the triradicals. However, more triradicals have to be investigated in order to better understand the chemical behavior of these reactive intermediates.

4.5 Charged Aryl Tetraradical: the 2,4,6-Tridehydropyridinium Radical Cation

Aromatic carbon-centered $\sigma,\sigma,\sigma,\sigma$ -tetraradicals are very complex and interesting systems, yet extremely difficult to study experimentally. Several tetradehydroaromatic systems have been explored computationally so far, including benzdiynes (**99**, **100** and **101** in Chart 11) and bisarynes with extended π -systems, such as anthracene (**103**) and naphthalenes **104** - **107** (Chart 11).³¹⁵ However, until now, the only successful generation of aromatic tetraradicals was achieved for derivatives of **100**.^{316,317}

The intermediacy of **99** and **100** (Chart 11) was first proposed in 1966 by Meyerson.³¹⁸ The pyrolyses of pyromellitic (1,2,3,4-benzenetetracarboxylic) **108** and mellophanic (1,2,4,5-benzenetetracarboxylic) **109** dianhydrides were found to yield products that were likely derived from the benzdiynes **99** and **100** (Chart 12). About 25 years later, McNaughton

obtained high-resolution IR spectra of the products of gas-phase flash vacuum photolysis of pyromellitic dianhydride.³¹⁹ Among the products, 1,3,5-hexatriyne **110** (Scheme 30) was detected. **110** was likely formed by ring opening of benzdiynes **99**, **100** or **101**. The photolysis of pyromellitic dianhydride **108** in a low-temperature argon matrix, as reported by Yabe et al., also yielded **110**.³²⁰ However, no direct evidence for the existence of 1,4-benzdiyne **100** was obtained by IR or UV-Vis spectroscopies. In the following years, Yabe and co-workers reported several successful and unsuccessful attempts to generate benzdiynes by photolysis of unsubstituted and substituted benzenetetracarboxylic anhydrides and trap them.^{316,317,321–323} In order to explain the formation of **110** upon photolyses of two different precursors, **108** and **109**, the reaction pathway shown on Scheme 30 was proposed.^{321,322} Based on this pathway, the initial formation of benzdiynes is followed by ring opening and rearrangements to yield diradical **111** and carbene **112**, which rearranged further to form the 1,3,5-hexatriyne **110**.

The first generation of benzdiynes, as confirmed by IR and UV-Vis spectroscopies, was carried out in a low temperature matrix by using the substituted dianhydrides **113** and **114**.^{316,317} The photolyses of **113** and **114** yielded the benzdiynes **115** and **116**, respectively, which upon further irradiation underwent ring opening and rearrangement to hexatriynes **117** and **118** (Scheme 31). In contrast, as mentioned above, no benzdiynes were observed for the unsubstituted dianhydrides. This was proposed to be due to facile hydrogen atom migration in the electron-rich benzdiyne **100** to form **110**.^{316,317} However, migration of a CF₃ group or F atom is less favorable than hydrogen atom migration, which explains the observation of the hexatriynes for the fluoro-compounds.

The experimental studies on benzdiynes were complemented by computational studies.^{292,324–329} Hoffmann calculated the electronic structures of the three tetrahydrobenzenes by using the extended Hückel theory.³²⁴ Radom et al. used ab initio molecular orbital theory to study didehydro- and tetrahydroarenes, among them the benzdiynes **99** and **100**.³²⁷ Zahradník et al. reported ab initio and semi-empirical quantum chemical calculations on several strained unsaturated molecules, including the benzdiynes **99** and **100**.³²⁹ Schaefer's group studied the three unsubstituted benzdiyne isomers and benzdiyne **115** by using both ab initio and density functional theory (DFT).²⁹² The nine isomers of C₆H₂, including the three unsubstituted benzdiyne isomers, were characterized by Sattelmeyer and Stanton by using high-level ab initio methods.³²⁸ Yabe and co-workers performed calculations of the structures and energetics of unsubstituted and substituted benzdiyne isomers by using both ab initio and DFT methodologies.^{325,326} Based on the most recent theoretical studies, **101** is the most stable among tetrahydrobenzenes, lying 14 kcal/mol below **100** and 7.8 kcal/mol below **99** (at G2 level of theory).³²⁵ In addition, it was suggested that **101** adopts a bicyclic geometry.^{292,328} The S-T splittings in **99**, **100** and **101** were calculated to be 26.3, 18.3 and 37.5 kcal/mol, respectively (at CCSD(T)TZ2P//CCSD/DZP level of theory). The S-T splitting in *ortho*-benzyne at the same level of theory is estimated to be 35.8 kcal/mol. The large heats of formation of benzdiynes predicted by Arulmozhiraja et al., are probably one of the reasons for the difficulty in generating and isolating these species.³²⁵ According to the theoretical studies of Yabe et al.,³²⁶ electron withdrawing substituents, such as NO₂, CN and CF₃, destabilize the benzdiyne more than they destabilize the corresponding benzenes, whereas electron-donating groups have the opposite effect.

Compared to benzdiynes, studies on tetrahydropyridines are very limited.³²⁷ Two pyridiynes, **119** and **120**, have been investigated computationally using an ab initio molecular orbital approach (Chart 13). Based on these studies, **120** is favored over **119** by 8 kJ/mol (at CASSCF/3-21G level of theory).

In spite of the above studies, the chemical properties of tetrahydrobenzenes have been unexplored, until recently. The first reactivity study on such a tetradical, the 1,2,4,6-tetrahydro-pyridinium cation **102** (Chart 11; a derivative of benzdiyne **101**), was reported recently.³¹⁴ Tetradical **102** was generated in a dual-cell FT-ICR mass spectrometer by using previously described methodology, as shown in Scheme 32.³³⁰ The precursor, 2,4,6-triiodopyridine,³¹¹ was introduced into the mass spectrometer and chemically ionized with ionized deuterated acetone to form a deuterated precursor cation. SORI-CAD was used to cleave two C-I bonds in the precursor cation, followed by elimination of DI to give the tetradical cation **102** (Scheme 32). The structure of **102** was confirmed by using structurally diagnostic ion-molecule reactions,^{311,330} including the abstraction of four hydrogen atoms from cyclohexane, which confirmed the presence of four radical sites in **102**.

The tetradical cation **102** was allowed to react with cyclohexane, allyl iodide and dimethyl disulfide, and its gas-phase reactivity was compared to that of related bi- and triradicals. Surprisingly, **102** was found not to undergo radical reactions. This was rationalized by an even-electron resonance structure (Scheme 33). The tetradical cation **102** is highly electrophilic and it rapidly reacts with many nucleophiles by quenching the C-N *ortho*-benzyne moiety, which results in the formation of a relatively unreactive *meta*-benzyne analogue (Scheme 33).

4. Summary

Many different experimental techniques now exist for generating aryl distonic ions and examining their structures and chemical properties in mass spectrometers. The experiments have been complemented by successful quantum chemical approaches for predicting electronic and thermochemical properties for these distonic ions. Numerous research groups are involved in this research, with interests ranging from fundamental properties of neutral and charged mono- and polyradicals to exploration of radical-induced dissociation of biopolymers. The properties and reactivity of a variety of small cations with aryl radical sites have been thoroughly examined both theoretically and experimentally. These distonic ions have been found to be stable molecules with very different properties from those of the isomeric conventional radical cations. Much of their reactivity closely resembles that of related neutral radicals as hydrogen atom abstraction and addition reactions are common. However, when the charge can delocalize into the benzene ring containing the radical site, also nucleophilic addition reactions to the radical site have been observed. The influence of the polarity and type of the charge site and its location with respect to the radical site, as well as additional substituents, has been investigated. In sharp contrast, the chemistry of related negative ions is almost entirely unexplored. Further, while several factors that control reactions of distonic *ortho*- and *meta*-benzyne analogs have been identified, including electron affinities of the radical sites, hydrogen bonding interactions in transition state, S-T splittings and dehydrocarbon atom separation for *meta*-benzyne analogs, the same is not true for related *para*-benzyne analogs. These species have proven to be particularly difficult to generate in mass spectrometers. Finally, much still remains to be learned on distonic aryl tri- and tetradicals. Especially those polyradicals containing the *meta*-benzyne moiety are expected to show interesting chemical properties since *meta*-benzyne analogs' reactivity is extremely sensitive to minor structural changes and can be readily "tuned" to be either nonradical or radical-like.

Acknowledgments

The National Science Foundation and the National Institutes of Health are acknowledged for generous financial support.

References

1. Gross ML, McLafferty FW. *J Am Chem Soc.* 1971; 93:126.
2. Gross ML. *J Am Chem Soc.* 1972; 94:3744.
3. Bouma WJ, MacLeod JK, Radom L. *J Chem Soc, Chem Comm.* 1978:724.
4. Yates BF, Bouma WJ, Radom L. *J Am Chem Soc.* 1984; 106:5805.
5. Yates BF, Bouma WJ, Radom L. *Tetrahedron.* 1986; 42:6225.
6. Tomazela DM, Sabino AA, Sparrapan R, Gozzo FC, Eberlin MN. *J Am Soc Mass Spec.* 2006; 17:1014.
7. Munsch TE, Slipchenko LV, Krylov AI, Wenthold PG. *J Org Chem.* 2004; 69(5735)
8. Hammerum S. *Mass Spectrom Rev.* 1988; 7:123.
9. Stirk KM, Kiminkinen LKM, Kenttämää HI. *Chem Rev.* 1992; 92:1649.
10. Kenttämää HI. *Org Mass Spectrom.* 1994; 29:1.
11. Bielanski, A.; Haber, J. *Oxygen in Catalysis.* Mercel Dekker; New York: 1991.
12. Love N, Parthasarathy N, Gollahallia R. *Int J Green Energy.* 2011; 8:113.
13. Denisov, E.; Afanas'ev, I. *Oxidation and Antioxidants in Organic Chemistry and Biology.* Taylor & Francis; Boca Raton: 2005.
14. Abdul-Rasoul FA, Ledwith A, Yagci Y. *Polymer.* 1978; 19:1219.
15. Crivello, J. *Initiators -- Poly-Reactions -- Optical Activity.* Springer-Verlag; Heidelberg, Germany: 1984. *Cationic Polymerization -- Iodonium and Sulfonium Salt Photoinitiators*; p. 1-48.
16. Odian, G. *Principles of Polymerization.* Wiley-Interscience; New York: 2004.
17. Gunz DW, Hoffmann MR. *Atmos Environ.* 1990; 24:1601.
18. Atkinson R, Arey J. *Atmos Environ.* 2003; 37:S197.
19. Herrmann H, Hoffmann D, Schaefer T, Bräuer P, Tilgner A. *Chem Phys Chem.* 2010; 11:3796. [PubMed: 21120981]
20. Greenberg M. *Astrophys Space Sci.* 1976; 39:9.
21. Kraka E, Cremer D. *J Am Chem Soc.* 2000; 122:8245.
22. Griffiths J, Murphy JA. *J Chem Soc, Chem Comm.* 1991:1422.
23. Griffiths J, Murphy JA. *J Chem Soc, Chem Comm.* 1992:24.
24. Wender PA, Jeon R. *Org Lett.* 1999; 1:2117. [PubMed: 10836065]
25. Wender PA, Jeon R. *Bioorg Med Chem Lett.* 2003; 13:1763. [PubMed: 12729660]
26. Hoffner J, Schottelius MJ, Feichtinger D, Chen P. *J Am Chem Soc.* 1998; 120:376.
27. Chen P. *Angew Chem, Int Ed.* 1996; 35:1478.
28. Greenley TL, Davies MJ. *Biochim Biophys Acta.* 1993; 1157:23. [PubMed: 8388732]
29. Hazlewood C, Davies MJ. *Arch Biochem Biophys.* 1996; 332:79. [PubMed: 8806712]
30. Meunier B, Pratviel G, Bernadou J. *Bull Soc Chim Fr.* 1994; 131:933.
31. Pratviel G, Bernadou J, Meunier B. *Angew Chem, Int Ed.* 1995; 34:746.
32. Nicolaou KC, Dai WM. *Angew Chem, Int Ed.* 1991; 30:1387.
33. Myers AG, Cohen SB, Kwon BM. *J Am Chem Soc.* 1994; 116:1255.
34. Bridger RF, Russell GA. *J Am Chem Soc.* 1963; 85:3754.
35. Pryor WA, Guard J. *J Am Chem Soc.* 1964; 86:1150.
36. Fu JLL, Bentrude WG. *J Am Chem Soc.* 1972; 94:7710.
37. Scaiano JC, Stewart LC. *J Am Chem Soc.* 1983; 105:3609.
38. Sommeling PM, Mulder P, Louw R, Avila DV, Luszyk J, Ingold KU. *J Phys Chem.* 1993; 97:8362.
39. Yu T, Lin MC, Melius CF. *Int J Chem Kinet.* 1994; 26:1095.
40. Fang W, Mertens R, von Sonntag CJ. *J Chem Soc Perk Trans 2.* 1995; 6:1033.
41. Parker VD. *Acta Chem Scand.* 1998; 52:154. [PubMed: 9487659]
42. Ingold KU. *Curr Med Chem.* 2003; 10:2631. [PubMed: 14529453]
43. Schepp NP. *J Org Chem.* 2004; 69:4931. [PubMed: 15255718]

44. Salamone M, DiLabio GA, Bietti M. *J Am Chem Soc.* 2011; 133:16625. [PubMed: 21895002]
45. Widjaja F, Jin Z, Nash JJ, Kenttämaa HI. *J Am Chem Soc.* 2012; 134:2085. [PubMed: 22148169]
46. Stirk KM, Orłowski JC, Leeck DT, Kenttämaa HI. *J Am Chem Soc.* 1992; 114:8604.
47. Chyall LJ, Kenttämaa HI. *J Am Chem Soc.* 1994; 116:3135.
48. Chyall LJ, Kenttämaa HI. *J Mass Spectrom.* 1995; 30:81.
49. Smith RL, Kenttämaa HI. *J Am Chem Soc.* 1995; 117:1393.
50. Li R, Smith RL, Kenttämaa HI. *J Am Chem Soc.* 1996; 118:5056.
51. Thoen KK, Beasley BJ, Smith RL, Kenttämaa HI. *J Am Soc Mass Spec.* 1996; 7:1245.
52. Thoen KK, Smith RL, Nousiainen JJ, Nelson ED, Kenttämaa HI. *J Am Chem Soc.* 1996; 118:8669.
53. Thoen KK, Perez J, Joseph J, Ferra J, Kenttämaa HI. *J Am Soc Mass Spec.* 1998; 9:1135.
54. Nelson ED, Li R, Kenttämaa HI. *Int J Mass Spectrom.* 1999; 7:1245.
55. Heidbrink JL, Thoen KK, Kenttämaa HI. *J Org Chem.* 2000; 65:645.
56. Heidbrink JL, Ramirez-Arizmendi LE, Thoen KK, Guler L, Kenttämaa HI. *J Phys Chem A.* 2001; 105:7875.
57. Ramirez-Arizmendi LE, Guler L, Joseph J, Ferra J, Thoen KK, Kenttämaa HI. *Int J Mass Spectrom.* 2001; 210/211:511.
58. Tichy SE, Thoen KK, Price JM, Ferra J, Petucci CJ, Kenttämaa HI. *J Org Chem.* 2001; 66:2726. [PubMed: 11304194]
59. Petzold CJ, Nelson ED, Lardin HA, Kenttämaa HI. *J Phys Chem A.* 2002; 106:9767.
60. Petzold CJ, Ramirez-Arizmendi LE, Heidbrink JL, Perez J, Kenttämaa HI. *J Am Soc Mass Spec.* 2002; 13:192.
61. Ramirez-Arizmendi LE, Heidbrink JL, Guler L, Kenttämaa HI. *J Am Chem Soc.* 2003; 125:2272. [PubMed: 12590557]
62. Jing L, Guler L, Nash JJ, Kenttämaa HI. *J Am Soc Mass Spec.* 2004; 15:913.
63. Huang Y, Guler L, Heidbrink JL, Kenttämaa HI. *J Am Chem Soc.* 2005; 127:3973. [PubMed: 15771534]
64. Huang Y, Kenttämaa HI. *J Am Chem Soc.* 2005; 127:7952. [PubMed: 15913386]
65. Liu, J-a; Petzold, CJ.; Ramirez-Arizmendi, LE.; Perez, J.; Kenttämaa, HI. *J Am Chem Soc.* 2005; 127:12758. [PubMed: 16159243]
66. Jing L, Guler L, Pates GO, Kenttämaa HI. *J Phys Chem A.* 2008; 112:9708. [PubMed: 18774790]
67. Jing L, Nash JJ, Kenttämaa HI. *J Am Chem Soc.* 2008; 130:17697. [PubMed: 19061320]
68. Adeuya A, Price JM, Jankiewicz BJ, Nash JJ, Kenttämaa HI. *J Phys Chem A.* 2009; 113:13663. [PubMed: 19902945]
69. Adeuya A, Nash JJ, Kenttämaa HI. *J Phys Chem A.* 2010; 114:12851. [PubMed: 21080694]
70. Jankiewicz BJ, Gao J, Reece JN, Vinueza NR, Narra P, Nash JJ, Kenttämaa HI. *J Phys Chem A.* 2012; 116:3089. [PubMed: 22352890]
71. Comisarow MB, Marshall AG. *Chem Phys Lett.* 1974; 25:282.
72. Dempster AJ. *Phys Rev.* 1918; 11:316.
73. Harrison, AG. *Chemical Ionization Mass Spectrometry.* CRC Press; Boca Raton: 1992.
74. Munson MSB, Field FH. *J Am Chem Soc.* 1966; 88:2621.
75. Whitehouse CM, Dreyer RN, Yamashita M, Fenn JB. *Anal Chem.* 1985; 57:675. [PubMed: 2581476]
76. Fenn JB, Mann M, Meng CK, Wong SF, Whitehouse CM. *Science.* 1989; 246:64. [PubMed: 2675315]
77. Fenn JB, Mann M, Meng CK, Wong SF, Whitehouse CM. *Mass Spectrom Rev.* 1990; 9:37.
78. Pramanik, BN.; Ganguly, AK.; Gross, ML. *Applied Electrospray Mass Spectrometry.* Marcel Dekker; New York: 2002.
79. Carroll DI, Dzidic I, Stillwell RN, Haegele KD, Horning EC. *Anal Chem.* 1975; 47:2369.
80. Thomson BA. *J Am Soc Mass Spectrom.* 1998; 9:187.

81. Tanaka K, Waki H, Ido Y, Akita S, Yoshida Y, Yoshida T. *Rapid Commun Mass Spectrom.* 1988; 2:151.
82. Karas M, Bachmann D, Bahr U, Hillenkamp F. *Int J Mass Spectrom Ion Proc.* 1987; 78:53.
83. Karas M, Krüger R. *Chem Rev.* 2003; 103:427. [PubMed: 12580637]
84. Knochenmuss R, Zenobi R. *Chem Rev.* 2003; 103:441. [PubMed: 12580638]
85. Marshall AG, Roe DC. *J Chem Phys.* 1980; 73:1581.
86. Wang M, Marshall AG. *Int J Mass Spectrom Ion Proc.* 1990; 100:323.
87. Marshall AG, Wang TCL, Ricca TL. *J Am Chem Soc.* 1985; 107:7893.
88. Gauthier JW, Trautman TR, Jacobson DB. *Anal Chim Acta.* 1991; 246:211.
89. Mirgorodskaya E, O'Connor PB, Costello CE. *J Am Soc Mass Spec.* 2002; 13:318.
90. Freiser, SB. *Techniques for the Study of Ion-Molecule Reactions.* Farrar, JM.; Saunders, WH., Jr, editors. John Wiley & Sons; New York: 1988.
91. Laude DA Jr, Johlman CL, Brown RS, Weil DA, Wilkins CL. *Mass Spectrom Rev.* 1986; 5:107.
92. Kofel P, Allemann M, Kellerhals H, Wanczek KP. *Int J Mass Spectrom Ion Proc.* 1985; 65:97.
93. McIver RT Jr, Hunter RL, Bowers WD. *Int J Mass Spectrom Ion Proc.* 1985; 64:67.
94. Hunt, dF; Shabanowitz, J.; McIver, RT., Jr; Hunter, RL.; Syka, JEP. *Anal Chem.* 1985; 57:765. [PubMed: 3993950]
95. Wysocki VH, Kenttämää HI. *J Am Chem Soc.* 1990; 112:5110.
96. Gozzo FC, Eberlin MN. *J Am Soc Mass Spec.* 1995; 6:554.
97. Eberlin MN. *Mass Spectrom Rev.* 1997; 16:113.
98. Carvalho M, Gozzo FC, Mendes MA, Sparrapan R, Kascheres C, Eberlin MN. *Chem Eur J.* 1998; 4:1161.
99. Gozzo FC, Eberlin MN. *J Org Chem.* 1999; 64:2188.
100. Sparrapan R, Mendes MA, Carvalho M, Eberlin MN. *Chem Eur J.* 2000; 6:321. [PubMed: 11931112]
101. Eberlin MN. *Int J Mass Spectrom.* 2004; 235:263.
102. Gronert S. *Chem Rev.* 2001; 101:329. [PubMed: 11712250]
103. Gronert S. *Mass Spectrom Rev.* 2005; 24:100. [PubMed: 15389862]
104. Gronert S. *J Am Soc Mass Spec.* 1998; 9:845.
105. Harman DG, Blanksby SJ. *Org Biomol Chem.* 2007; 5:3495. [PubMed: 17943209]
106. Squires RR. *Int J Mass Spectrom Ion Proc.* 1992; 118/119:503.
107. Levsen, K. *Fundamental Aspects of Organic Mass Spectrometry.* Budzikiewicz, HvH, editor. Vol. 4. Verlag Chemie; New York: 1978.
108. Burgers PC, Holmes JL. *Rapid Commun Mass Spectrom.* 1989; 3:279.
109. Holmes, JL.; Aubry, C.; Mayer, PM. *Assigning Structures to Ions in Mass Spectrometry.* CRC Press; Boca Raton: 2007.
110. Wenthold, PG. *Studies of the Thermochemical Properties of Reactive Intermediates by Mass Spectrometric Methods.* In: Platz, MS.; Moss, RA.; Maitland Jones, J., editors. *Reviews of Reactive Intermediate Chemistry.* John Wiley & Sons, Inc; Hoboken, New Jersey: 2007.
111. Wenthold PG, Squires RR. *J Am Chem Soc.* 1994; 116:6401.
112. Cooks, RG. *Collision Spectroscopy.* Cooks, RG., editor. Plenum Press; New York: 1978.
113. McLuckey SA. *J Am Soc Mass Spec.* 1992; 3:599.
114. Hoffman, Ed. *J Mass Spectrom.* 1996; 31:129.
115. Yu SJ, Holliman CL, Rempel DL, Gross ML. *J Am Chem Soc.* 1993; 115:9676.
116. Bjornholm T, Hammerum S, Kuck D. *J Am Chem Soc.* 1988; 110:3862.
117. Osburn S, Ryzhov V. *Anal Chem.* 2013; 85:769. [PubMed: 23077968]
118. Green MK, Lebrilla CB. *Mass Spectrom Rev.* 1997; 16:53. [PubMed: 9414490]
119. Brodbelt JS. *Mass Spectrom Rev.* 1997; 16:91.
120. Yem, ST. M S Thesis. Purdue University; IN: 2009.

121. Sorrilha AEP, Gozzo FC, Pimpim RS, Eberlin MN. *J Am Soc Mass Spectrom.* 1996; 7:1126. [PubMed: 24203075]
122. Salem L, Leforestier C, Segal G, Wetmore R. *J Am Chem Soc.* 1975; 97:479.
123. Salem L. *Science.* 1976; 191:822. [PubMed: 1251196]
124. Dauben WG, Salem L, Turro NJ. *Acc Chem Res.* 1975; 8:41.
125. Donahue NM. *Chem Rev.* 2003; 103:4593. [PubMed: 14664625]
126. Heitler W, London F. *Z Phys Chem.* 1927; 44:455.
127. Evans MG, Polanyi M. *Trans Faraday Soc.* 1938; 34:11.
128. Marcus RA. *J Chem Phys.* 1956; 24:966.
129. Marcus RA. *J Phys Chem.* 1968; 72:891.
130. Donahue NM, Clarke JS, Anderson JG. *J Phys Chem A.* 1998; 102:3923.
131. Shaik SS, Hiberty PC. *Adv Quantum Chem.* 1995; 26:99.
132. Pross A. *Adv Phys Org Chem.* 1985; 21:99.
133. Maitre P, Hiberty PC, Ohanessian G, Shaik SS. *J Phys Chem.* 1990; 94:4089.
134. Pross A, Yamataka H, Nagase S. *J Phys Org Chem.* 1991; 4:135.
135. Porter RM, Karplus M. *J Chem Phys.* 1964; 40:1105.
136. Pross, A. *Theoretical and Physical Principles of Organic Reactivity.* John Wiley and Sons; New York: 1995.
137. Donahue NM. *J Phys Chem A.* 2001; 105:1489.
138. Clarke JS, Kroll JH, Donahue NM, Anderson JG. *J Phys Chem A.* 1998; 102:9847.
139. Clarke JS, Rypkema HA, Kroll JH, Donahue NM, Anderson JG. *J Phys Chem A.* 2000; 104:4458.
140. Beasley BJ, Smith RL, Kenttämää HI. *J Mass Spectrom.* 1995; 30:384.
141. Pryor WA, Platt PK. *J Am Chem Soc.* 1963; 85:1496.
142. Pryor WA, Smith K. *J Am Chem Soc.* 1970; 92:2731.
143. Danen, WC. *Methods in Free-Radical Chemistry.* Huyser, ES., editor. Vol. 5. Marcel Dekker; New York: 1974.
144. Poutsma, M. *Free Radicals.* Kochi, JK., editor. Vol. II. Wiley; New York: 1973.
145. Danen WC, Winter RL. *J Am Chem Soc.* 1971; 93:716.
146. Migita T, Takayama K, Abe K, Kosugi M. *J Chem Soc Perk Trans 2.* 1979; 8:1137.
147. Takayama K, Kosugi M, Migita T. *Chem Lett.* 1973:193.
148. Petucci CJ, Nyman M, Guler L, Kenttämää HI. *J Am Chem Soc.* 2002; 124:4108. [PubMed: 11942850]
149. *CRC Handbook of Chemistry and Physics.* 78. CRC Press; Boca Raton: 1997.
150. Lam AKY, Li C, Khairallah G, Kirk BB, Blanksby SJ, Trevitt AJ, Wille U, O'Hair RAJ, da Silva G. *Phys Chem Chem Phys.* 2012; 14:2417. [PubMed: 22249506]
151. Hoffman, RW. *Dehydrobenzyne and Cycloalkynes.* Academic Press; New York: 1967.
152. Winkler, M.; Wenk, HH.; Sander, W. *Reactive Intermediate Chemistry.* Moss, RA.; Platz, MS.; Jones, JM., editors. Wiley-Interscience; New York: 2004.
153. Debbert SL, Cramer CJ. *Int J Mass Spectrom.* 2000; 201:1.
154. Salem L, Rowland C. *Angew Chem, Int Ed.* 1972; 11:92.
155. IUPAC. *Compendium of Chemical Terminology.* 2. Blackwell Scientific Publications; Oxford: 1997. (the "Gold Book")
156. Wittig G. *Naturwissenschaften.* 1942:696.
157. Wittig G, Harborth G. *Ber Dtsch Chem Ges.* 1944; 77:306.
158. Wittig G, Pohmer L. *Chem Ber.* 1956; 89:1334.
159. Roberts JD, Semenov DA, Simmons JHE, Carlsmith LA. *J Am Chem Soc.* 1956; 78:601.
160. Roberts JD, Simmons HE, Carlsmith LA, Vaughan CW. *J Am Chem Soc.* 1953; 75:3290.
161. Panar M, Roberts JD. *J Am Chem Soc.* 1960; 82:3692.
162. Huisgen R, Rist H. *Naturwissenschaften.* 1954; 42:358.
163. Huisgen R, Rist H. *Ann Chem.* 1955; 594:137.

164. Huisgen R, Knorr R. *Tetrahedron Lett.* 1963;1017.
165. Fisher IP, Lossing FP. *J Am Chem Soc.* 1963; 85:1018.
166. Chapman OL, Mattes K, McIntosh CL, Pacansky J, Calder GV, Orr G. *J Am Chem Soc.* 1973; 95:6134.
167. Chapman OL, Chang CC, Kolc J, Rosenquist NR, Tomioka H. *J Am Chem Soc.* 1975; 97:6586.
168. Wentrup C, Blanch R, Briehl H, Gross G. *J Am Chem Soc.* 1988; 110:1874.
169. Schweig A, Muenzel N, Meyer H, Heidenreich A. *Struct Chem.* 1990; 1:89.
170. Simon JGG, Muenzel N, Schweig A. *Chem Phys Lett.* 1990; 170:187.
171. Radziszewski JG, Hess BA, Zahradnik R. *J Am Chem Soc.* 1992; 114:52.
172. Berry RS, Clardy J, Schafer ME. *J Am Chem Soc.* 1964; 86:2738.
173. Berry RS, Spokes GN, Stiles M. *J Am Chem Soc.* 1962; 84:3570.
174. Kauffmann T. *Angew Chem, Int Ed.* 1965; 77:557.
175. den Hertog HJ, van der plas HC. *Adv Heterocycl Chem.* 1965; 4:121.
176. Kauffmann T, Wirthwein R. *Angew Chem, Int Ed.* 1971; 10:20.
177. Reinecke MG. *Tetrahedron.* 1982; 38:427.
178. Sander W. *Acc Chem Res.* 1999; 32:669.
179. Wenk HH, Winkler M, Sander W. *Angew Chem, Int Ed.* 2003; 42:502.
180. Moss, RA.; Platz, MS.; Jones, JM. *Reactive Intermediate Chemistry.* John Wiley & Sons; New York: 2004.
181. Platz, MS.; Moss, RA.; Jones, JM. *Reviews of Reactive Intermediate Chemistry.* John Wiley & Sons; New York: 2007.
182. Wenthold PG, Squires RR, Lineberger WC. *J Am Chem Soc.* 1998; 120:5279.
183. Hoffmann R, Imamura A, Hehre WJ. *J Am Chem Soc.* 1968; 90:1499.
184. Kauffmann T, Boettcher FP, Hansen J. *Angew Chem.* 1961; 73:341.
185. Kauffmann T, Boettcher FP. *Chem Ber.* 1962; 95:949.
186. Lewis ES, Cooper JE. *J Am Chem Soc.* 1962; 84:3847.
187. Kauffmann T, Marhan T. *Chem Ber.* 1963; 96:2519.
188. Abramovitch RA, Singer GM. *J Org Chem.* 1974; 39:1795.
189. Bunnett JF, Singh P. *J Org Chem.* 1981; 46:4567.
190. Ohkura K, Seki KI, Tershima M, Kanaoka Y. *Tetrahedron Lett.* 1989; 30:3433.
191. Turecek F, Wolken JK, Sadilek M. *Eur J Mass Spectrom.* 1998; 4:321.
192. Wolthuis E, Bouma B, Modderman J, Sytsma L. *Tetrahedron Lett.* 1970; 6:407.
193. Kauffmann T, Boettcher FP. *Angew Chem.* 1961; 73:65.
194. Martens RJ, den Hertog HJ. *Tetrahedron Lett.* 1962; 15:643.
195. Jankiewicz BJ, Reece JN, Vinuesa NR, Nash JJ, Kenttämäa HI. *Angew Chem, Int Ed Engl.* 2008; 47:9860. [PubMed: 19006159]
196. Jankiewicz, BJ. Ph D Dissertation. Purdue University; West Lafayette, IN: 2008.
197. Yamabe S, Minato T, Ishiwata A, Irinamihira O, Machiuguchi T. *J Org Chem.* 1997; 72:2832. [PubMed: 17367189]
198. Campbell, KM. Ph D Dissertation. Purdue University; West Lafayette, IN: 2007.
199. Kirkpatrick, LM. Ph D Dissertation. Purdue University; West Lafayette, IN: 2010.
200. Berry RS, Clardy J, Schafer ME. *Tetrahedron Lett.* 1965; 415:1011.
201. Washburn WN, Zahler R, Chen I. *J Am Chem Soc.* 1978; 100:5863.
202. Marquardt R, Sander W, Kraka E. *Angew Chem, Int Ed.* 1996; 108:825.
203. Winkler M, Sander W. *J Phys Chem A.* 2001; 105:10422.
204. Sander W, Exner M, Winkler M, Balster A, Hajerpe A, Kraka E, Cremer D. *J Am Chem Soc.* 2002; 124:13072. [PubMed: 12405834]
205. Sander W, Marquardt R, Bucher G, Wandel H. *Pure & Appl Chem.* 1996; 68:353.
206. Wenk HH, Sander W. *Chem Eur J.* 2001; 7:1837. [PubMed: 11405460]
207. Kraka E, Cremer D, Bucher G, Wandel H, Sander W. *Chem Phys Lett.* 1997; 348:115.

208. Bucher G, Sander W, Kraka E, Cremer D. *Angew Chem, Int Ed.* 1992; 31:1230.
209. Bucher G, Sander W. *J Org Chem.* 1992; 57:2185.
210. Sander W, Bucher G, Wandel H, Kraka E, Cremer D, Sheldrick WS. *J Am Chem Soc.* 1997; 119:10660.
211. Sander W, Exner M. *J Chem Soc Perk Trans 2.* 1999:2285.
212. Wenk HH, Sander W. *Eur J Org Chem.* 2002:3927.
213. Sander W, Exner M, Winkler M, Balster A, Hjerpe A, Kraka E, Cremer D. *J Am Chem Soc.* 2002; 124:13072. [PubMed: 12405834]
214. Noell JO, Newton MD. *J Am Chem Soc.* 1979; 101:51.
215. Thiel W. *J Am Chem Soc.* 1981; 103:1420.
216. Dewar MJ, Ford GP, Reynolds CH. *J Am Chem Soc.* 1983; 105:3162.
217. Wenthold PG, Paulino JA, Squires RR. *J Am Chem Soc.* 1991; 113:7414.
218. Wenthold PG, Squires RR. *J Am Chem Soc.* 1994; 116:6961.
219. Wenthold PG, Squires RR. *J Am Chem Soc.* 1996; 118:11865.
220. Kraka E, Cremer D. *Chem Phys Lett.* 1993; 216:233.
221. Nicolaides A, Borden WT. *J Am Chem Soc.* 1993; 115:11951.
222. Wierschke SG, Nash JJ, Squires RR. *J Am Chem Soc.* 1993; 115:11958.
223. Lindh R, Persson BJ. *J Am Chem Soc.* 1994; 116:4963.
224. Lindh R, Lee TJ, Bernhardsson BJ, Persson BJ, Karlström G. *J Am Chem Soc.* 1995; 117:7186.
225. Nash JJ, Squires RR. *J Am Chem Soc.* 1996; 118:11872.
226. Cramer CJ, Nash JJ, Squires RR. *Chem Phys Lett.* 1997; 277:311.
227. de Visser SP, Filatov M, Shaik S. *Phys Chem Chem Phys.* 2000; 105:10422.
228. Smith CE, Crawford TD, Cremer D. *J Chem Phys.* 2005; 122:174309. [PubMed: 15910034]
229. Evangelista FA, Allen WD, Schaefer HF III. *J Chem Phys.* 2007; 127:24102.
230. Li H, Yu S-Y, Huang M-B, Wange Z-X. *Chem Phys Lett.* 2007; 450:12.
231. Al-Saidi WA, Umrigar CJ. *J Chem Phys.* 2008; 128:154324. [PubMed: 18433226]
232. Li X, Paldus J. *J Chem Phys.* 2008; 129:174101. [PubMed: 19045327]
233. Shen J, Fang T, Hua W, Li S. *J Phys Chem A.* 2008; 112:4703. [PubMed: 18433169]
234. Karton A, Kaminker I, Martin JML. *J Phys Chem A.* 2009; 113:7610. [PubMed: 19265420]
235. Wei H, Hrovat DA, Mo Y, Hoffman R, Borden WT. *J Phys Chem A.* 2009; 113:10351. [PubMed: 19711935]
236. Chattopadhyay S, Chaudhuri RK, Mahapatra US. *Chem Phys Lett.* 2010; 491:102.
237. Gavina F, Luis SV, Safont VS, Ferrer P, Costero AM. *Tetrahedron Lett.* 1986; 27:4779.
238. Billups WE, Buynak JD, Butler D. *J Org Chem.* 1979; 44:4218.
239. Hess JBA. *Eur J Org Chem.* 2001:2185.
240. Kraka E, Anglada J, Hjerpe A, Filatov M, Cremer D. *Chem Phys Lett.* 2001; 348:115.
241. Schottelius MJ, Chen P. *J Am Chem Soc.* 1996; 118:4896.
242. Logan CF, Chen P. *J Am Chem Soc.* 1996; 118:2113.
243. Wenthold PG, Rau NJ. *J Phys Chem A.* 2011; 115:10353. [PubMed: 21812461]
244. Johnson WTG, Cramer CJ. *J Am Chem Soc.* 2001; 123:923. [PubMed: 11456626]
245. Wenthold PG, Hu J, Squires RR. *J Am Chem Soc.* 1994; 116:6961.
246. Wenthold PG, Hu J, Squires RR. *J Mass Spectrom.* 1998; 33:796.
247. Price JM, Nizzi KE, Campbell LJ, Kenttämää HI, Seierstad M, Cramer CJ. *J Am Chem Soc.* 2003; 125:131. [PubMed: 12515514]
248. Thoen KK, Kenttämää HI. *J Am Chem Soc.* 1997; 119:3832.
249. Thoen KK, Kenttämää HI. *J Am Chem Soc.* 1999; 121:800.
250. Nelson ED, Artau A, Price JM, Tichy SE, Jing L, Kenttämää HI. *J Phys Chem A.* 2001; 105:10155.
251. Nash JJ, Nizzi KE, Adeuya A, Yurkovich MJ, Cramer CJ, Kenttämää HI. *J Am Chem Soc.* 2005; 127:5760. [PubMed: 15839653]

252. Johnson WTG, Cramer CJ. *J Phys Org Chem*. 2001; 14:597.
253. Vinueza NR, Archibold EF, Jankiewicz BJ, Gallardo VA, Habicht SC, Aqueel MS, Nash JJ, Kenttämää HI. *Chem Eur J*. 2012; 18:8692. [PubMed: 22693151]
254. Rees CW, Storr RC. *Chem Commun*. 1965:193.
255. Nakayama J, Dan S, Hoshino M. *J Chem Soc Perk Trans 1*. 1981:413.
256. Benati B, Montevocchi PC. *J Org Chem*. 1976; 41:2639.
257. de Luca G, Renzi G, Pizzabiocca A. *Tetrahedron Lett*. 1983; 24:821.
258. Squires RR, Cramer CJ. *J Phys Chem A*. 1998; 102:9072.
259. Berry RS, Clardy J, Schafer ME. *Tetrahedron Lett*. 1965; 415:1003.
260. Jones RR, Bergman RG. *J Am Chem Soc*. 1972; 94:660.
261. Dewar MJS, Li WK. *J Am Chem Soc*. 1974; 96:5569.
262. Yamaguchi K, Nishio A, Yabushita S, Fueno T. *Chem Phys Lett*. 1978; 53:109.
263. Dewar MJS, Ford GP, Reynolds CH. *J Am Chem Soc*. 1983; 105:3162.
264. Chapman OL, Chang CC, Kolc J. *J Am Chem Soc*. 1976; 98:5703.
265. Wong HNC, Sondheimer F. *Tetrahedron Lett*. 1980; 21:217.
266. Nicolaou KC, Dai WM. *Angew Chem, Int Ed*. 1995; 34:746.
267. Marinelli PJ, Paulino JA, Sunderlin LS, Wenthold PG, Poutsma JC, Squires RR. *Int J Mass Spectrom Ion Processes*. 1994; 130:89.
268. Graul ST, Squires RR. *Mass Spectrom Rev*. 1988; 7:1.
269. Kirkpatrick LM, Vinueza NR, Jankiewicz BJ, Gallardo VA, Archibold EF, Nash JJ, Kenttämää HI. *Chem Eur J*. 2013; 19:9022. [PubMed: 23703949]
270. Hawkins CL, Davies MJ. *Biochim Biophys Acta*. 2001; 1504:196. [PubMed: 11245785]
271. Dean RT, Fu S, Stocker R, Davies MJ. *Biochem J*. 1997; 324:1. [PubMed: 9164834]
272. Blackburn, GM.; Gait, MJ. *Nucleic Acids in Chemistry and Biology*. IRL Press; Oxford: 1990.
273. von Sonntag, CJ. *The Chemical Basis of Radiation Biology*. Taylor and Francis; London: 1987.
274. Burrows CJ, Muller JG. *Chem Rev*. 1998; 98:1109. [PubMed: 11848927]
275. Hazlewood C, Davies MJ. *J Chem Soc Perk Trans 2*. 1995:895.
276. Carter N, Taverner T, Schiesser CH, Greenberg MM. *J Org Chem*. 2000; 65:8375. [PubMed: 11101400]
277. Stubbe J, Kozarich JW. *Chem Rev*. 1987; 87:1107.
278. Steenken S. *Chem Rev*. 1989; 89:503.
279. Petzold CJ, Ramirez-Arizmendi LE, Heidbrink JL, Perez J, Kenttämää HI. *J Am Soc Mass Spec*. 2001; 13:192.
280. Li S, Fu M, Habicht SC, Pates GO, Nash JJ, Kenttämää HI. *J Org Chem*. 2009; 74:7724. [PubMed: 19772324]
281. Pates GO, Guler LP, Nash JJ, Kenttämää HI. *J Am Chem Soc*. 2011; 133:9331. [PubMed: 21612203]
282. Adeuya A, Yang L, Amegayibor S, Nash JJ, Kenttämää HI. *J Am Soc Mass Spec*. 2006; 17:1325.
283. Yang L, Nash JJ, Kenttämää HI. *Org Lett*. 2008; 10:1889. [PubMed: 18429617]
284. Fu M, Li S, Archibold E, Yurkovich MJ, Nash JJ, Kenttämää HI. *J Am Soc Mass Spec*. 2010; 21:1737.
285. Hazlewood C, Davies MJ, Gilbert BC, Packer JE. *J Chem Soc Perk Trans 2*. 1995:2167.
286. Perez J, Ramirez-Arizmendi LE, Petzold CJ, Guler LP, Nelson ED, Kenttämää HI. *Int J Mass Spectrom*. 2000; 198:173.
287. Reid GE, Tichy SE, Perez J, O'Hair RAJ, Simpson RJ, Kenttämää HI. *J Am Chem Soc*. 2001; 123:1184. [PubMed: 11456672]
288. Dow AR, Wittrig AM, Kenttämää HI. *Eur J Mass Spectrom*. 2012; 18:77.
289. Sherman CL, Pierce SE, Brodbelt JS, Tuesuwan B, Kerwin SM. *J Am Soc Mass Spec*. 2006; 17:1342.
290. Salem L, Rowland C. *Angew Chem Int Ed Engl*. 1972; 11:92.

291. Slipchenko LV, Krylov AI. *J Chem Phys.* 2003; 118:9614.
292. Bettinger HF, Schleyer PvR, Schaefer HF III. *J Am Chem Soc.* 1999; 121:2829.
293. Lardin HA, Nash JJ, Wenthold PG. *J Am Chem Soc.* 2002; 124:12612. [PubMed: 12381207]
294. Cristian AMC, Shao Y, Krylov AI. *J Phys Chem A.* 2004; 108:6581.
295. Krylov AI. *J Phys Chem A.* 2005; 109:10638. [PubMed: 16863112]
296. Krylov AI. *Acc Chem Res.* 2006; 39:83. [PubMed: 16489727]
297. Koziol L, Winkler M, Houk KN, Venkataramani S, Sander W, Krylov AI. *J Phys Chem A.* 2007; 111:5071. [PubMed: 17518455]
298. Manohar PU, Krylov AI. *J Chem Phys.* 2008; 129:194105. [PubMed: 19026043]
299. Venkataramani S, Winkler M, Sander W. *Angew Chem Int Ed Engl.* 2005; 44:6306. [PubMed: 16196083]
300. Venkataramani S, Winkler M, Sander W. *Angew Chem Int Ed Engl.* 2007; 46:4888. [PubMed: 17568464]
301. Winkler M, Sander W. *Aust J Chem.* 2010; 53:1013.
302. Wenthold PG. *Angew Chem Int Ed Engl.* 2005; 44:7170. [PubMed: 16276554]
303. Wenthold PG. *Aust J Chem.* 2010; 63:1091.
304. Nguyen HMT, Hölzl T, Gopakumar G, Veszpremi T, Peeters J, Nguyen MT. *Chem Phys.* 2005; 316:125.
305. Fishtik I. *Tetrahedron Lett.* 2009; 50:5596.
306. Chattopadhyay S, Chaudhuri RK, Freed KF. *Phys Chem Chem Phys.* 2011; 13:7514. [PubMed: 21423953]
307. Chattopadhyay S, Chaudhuri RK, Freed KF. *J Phys Chem A.* 2011; 115:3665. [PubMed: 20586459]
308. Chattopadhyay S, Mahapatra US, Chaudhuri RK. *Chem Phys.* 2012; 401:15.
309. Nash JJ, Kenttämä HI, Cramer CJ. *J Phys Chem A.* 2008; 112:5542. [PubMed: 18503288]
310. Manohar PU, Koziol L, Krylov AI. *J Phys Chem A.* 2009; 113:2591. [PubMed: 19236028]
311. Jankiewicz BJ, Adeuya A, Yurkovich MJ, Vinueza NR, Gardner ISJ, Zhou M, Nash JJ, Kenttämä HI. *Angew Chem Int Ed Engl.* 2007; 46:9198. [PubMed: 17890659]
312. Jankiewicz BJ, Reece JN, Vinueza NR, Nash JJ, Kenttämä HI. *Angew Chem Int Ed Engl.* 2008; 47:9860. [PubMed: 19006159]
313. Jankiewicz BJ, Vinueza NR, Reece JN, Lee YC, Williams PE, Nash JJ, Kenttämä HI. *Chem Eur J.* 2012; 18:969. [PubMed: 22180095]
314. Gallardo VA, Jankiewicz BJ, Vinueza NR, Nash JJ, Kenttämä HI. *J Am Chem Soc.* 2012; 134:1926. [PubMed: 22211475]
315. Sato T, Niino H. *Aust J Chem.* 2010; 63:1048.
316. Moriyama M, Ohana T, Yabe A. *J Am Chem Soc.* 1997; 119:10229.
317. Sato T, Arulmozhiraja S, Niino H, Sasaki S, Matsuura T, Yabe A. *J Am Chem Soc.* 2002; 124:4512. [PubMed: 11960482]
318. Fields EK, Meyerson S. *J Org Chem.* 1966; 31:3307.
319. McNaughton D, Bruget DN. *J Mol Spectrosc.* 1991; 150:620.
320. Moriyama M, Ohana T, Yabe A. *Chem Lett.* 1995; 24:557.
321. Moriyama M, Ohana T, Yabe A. *Chem Lett.* 1998; 27:337.
322. Moriyama M, Sato T, Uchimaru T, Yabe A. *Phys Chem Chem Phys.* 1999; 1:2267.
323. Sato T, Niino H, Yabe A. *J Am Chem Soc.* 2003; 125:11936. [PubMed: 14505415]
324. Adam W, Grimison A, Hoffmann R. *J Am Chem Soc.* 1969; 91:2590.
325. Arulmozhiraja S, Sato T, Yabe A. *J Comp Chem.* 2001; 22:923.
326. Arulmozhiraja S, Sato T, Yabe A. *J Org Chem.* 2003; 68:5084. [PubMed: 12816461]
327. Radom L, Nobes RH, Underwood DJ, Li WK. *Pure & Appl Chem.* 1986; 58:75.
328. Sattelmeyer KW, Stanton JF. *J Am Chem Soc.* 2000; 122:8220.

329. Zahradnik R, Hobza P, Burcl R, Hess JBA, Radziszewski JG. J Mol Struct (THEOCHEM). 1994; 313:335.
330. Price JM, Kenttämä H. J Phys Chem A. 2003; 107:8985.
331. Linstrom, PJ.; Mallard, WG. NIST Chemistry WebBook, NIST Standard Reference Database Number 69. National Institute of Standards and Technology; Gaithersburg MD, 20899: <http://webbook.nist.gov>

Biographies

Peggy Williams

Peggy Williams received her B.S. in chemistry in 2009 from Indiana State University. Her undergraduate research focused on efficient synthesis of benzylic Grignard reagents. She began pursuing a Ph.D. in chemistry at Purdue University in 2009. Under the supervision of Prof. Kenttämä she is currently studying the reactivity of charged aryl monoradicals in the gas phase and solution as well as using ion-molecule reactions for the functional group identification of drug metabolites by mass spectrometry.

Bartłomiej Jankiewicz

Bartłomiej Jankiewicz was born in 1980 in Reszel, Poland. During his diploma thesis at the Military University of Technology in Warsaw, Poland he worked on the synthesis and characterization of liquid crystalline compounds. He received his Ph.D. in 2008 from Purdue University, where under supervision of Prof. Kenttämä he investigated gas phase reactivity of mono- and polyradicals. His studies have led to first characterization of triradicals' reactivity. Currently he works in the Institute of Optoelectronics at Military University of Technology in Warsaw, Poland. His research focuses mainly on the synthesis of plasmonic nanostructures for applications in sensing (SERS, SEF) and photovoltaics.

Linan Yang

Linan Yang obtained her Ph.D. in Professor Hilkka Kenttämää's group in the Chemistry Department at Purdue University in 2007. Her research was focused on gas-phase studies of reactivity of biradicals toward biomolecules in mass spectrometers. She is now a Research Associate working at Purdue University, Center for Direct Catalytic Conversion of Biomass to Biofuels on collaborative efforts with chemical engineers, catalysis scientists and plant biologists to understand and optimize processes to convert biomass to biofuels.

Hilkka Kenttämää

Professor Hilkka Inkeri Kenttämää received her Ph.D. in chemistry at Helsinki University, Finland, in 1986. Following post doctoral research at Purdue University under the guidance of Professor Graham Cooks, she accepted an Assistant Professor position at Purdue University in 1989. In 1999, she was promoted to a Full Professor. Her research interests include fundamental chemical properties of gaseous organic bi- and polyradicals, development of mass spectrometric methods for the identification of unknown organic compounds in mixtures, such as drug degradation products, crude oil, and degraded biomass, and the design and construction of novel tandem mass spectrometers to facilitate above studies.

Figure 1. MS³ sequential CAD product ion mass spectra of the ionized methylene transfer products of three isomeric ethyl pyridines, (a) **2b**, (b) **2c**, and (c) **2d**, showing characteristic CAD fragmentation. Reproduced from reference 96 with kind permission from Springer Science and Business Media.

Figure 2. CAD mass spectrum of the adduct ion of m/z 107 formed in the reaction of pyridine radical cation and ethene. Reproduced with permission from reference 115. Copyright (1993) American Chemical Society.

Figure 3. CAD mass spectrum of the product ion of m/z 107 formed in the reaction of the methylcyclopropane radical cation with pyridine. Reproduced with permission from reference 115. Copyright (1993) American Chemical Society.

Figure 4. CAD mass spectrum of ethylpyridine radical cations: (A) 4-ethyl-, (B) 3-ethyl-, and (C) 2-ethylpyridine radical cations formed by chemical ionization of the corresponding neutral pyridines. Reproduced with permission from reference 115. Copyright (1993) American Chemical Society.

Figure 5. Adduct of m/z 139 formed by reaction of the β -distonic ethylenepyridine cation with O₂. Reproduced with permission from reference 115. Reproduced with permission from reference 145. Copyright (1993) American Chemical Society.

Figure 6. Graphic representation of atom transfer barriers according to (a) Marcus theory, (b) covalent (Heitler-London) avoided curve crossing model, and (c) ionic avoided curve crossing model. Reproduced with permission from reference 130. Copyright (1998) American Chemical Society.

Figure 7. Energy-resolved CAD product ions for (a) 2-dehydroanilinium, (b) 3-dehydroanilinium, and (c) 4-dehydroanilinium cation, as well as (d) ionized aniline, as a function of collision energy. Reproduced with permission from reference 16. Copyright (1995) John Wiley & Sons, Ltd.

Figure 8. Reaction of the 4-dehydroanilinium cation with (a) dimethyl disulfide, (b) 2-iodopropane and (c) pyridine. Reproduced with permission from reference 16. Copyright (1995) John Wiley & Sons, Ltd.

Figure 9.

Natural logarithm of the reaction efficiencies for hydrogen atom abstraction from cyclohexane versus calculated vertical electron affinities (eV) for 23 aryl radicals with a fit to the linear trend line ($R^2 = 0.92$). Reproduced with permission from reference 67. Copyright (2008) American Chemical Society.

Figure 10.

Natural logarithm of the reaction efficiencies for hydrogen atom abstraction from 15 different hydrogen atom donors by N-(3-dehydrophenyl)pyridinium cation (**g**) versus the vertical ionization energies (eV) of the hydrogen atom donors. The data are fit to a linear trend line ($R^2 = 0.94$). The hydrogen-atom donors are tributyltin hydride (**Tri**), benzeneselenol (**Ben**), proline (**Pro**), 2-deoxy-D-ribose (**2-Deoxy**), 1-O-methyl-2-deoxy-D-ribose (**1-O-Meth**), ribose (**Rib**), valine (**Val**), 1,4-dioxane, tetrahydrofuran (**THF**), L-alanine (**Ala**), glycine (**Gly**), cyclohexane (**Cyclo**), isopropyl alcohol (**Iso**), tert-butanol (**t-Buta**), and ethanol (**Eth**). Reproduced with permission from reference 67. Copyright (2008) American Chemical Society.

Figure 11.

Natural logarithm of the reaction efficiencies for hydrogen atom abstraction from 18 different hydrogen-atom donors by 29 different aryl radicals versus IE - EA (eV). The data are fit to a linear trend line ($R^2 = 0.87$); the dashed lines represent the 95% prediction interval. The horizontal dotted line represents 100% reaction efficiency. Reproduced with permission from reference 67. Copyright (2008) American Chemical Society.

Figure 12.

Mulliken spin densities calculated at B3LYP/cc-pVDZ//B3LYP/6-31G(d) level of theory for the intermediate resulting from the addition of *tert*-butyl isocyanide to radicals **17**, **24** and **25**. Reproduced with permission from reference 68. Copyright (2009) American Chemical Society.

Figure 13. Natural logarithm of the reaction efficiencies for hydrogen atom abstraction from cyclohexane versus calculated vertical electron affinities (eV) for charged radicals **17**, **24**, **26** - **31**. The data are fit to a linear trend line ($R^2 = 0.83$). Reproduced with permission from reference 69. Copyright (2010) American Chemical Society.

Figure 14.

Natural logarithm of the reaction efficiencies for hydrogen atom abstraction from methanol versus calculated activation enthalpies (kcal/mol) for charged radicals **17**, **24**, **26** - **31**. MPW1K (circles) and G3MP2B3 (squares) data are fit to linear trend lines ($R^2 = 0.93$ and 0.96 , respectively). Reproduced with permission from reference 69. Copyright (2010) American Chemical Society.

Figure 15.

Calculated (MPW1K) transition-state structures for hydrogen atom abstraction from methanol by radicals **17** (g), **24** (b), **26 – 31** (a, c, d–f, h). Distances between the hydrogen atom being transferred and the α -carbon atom in methanol, distances between the hydrogen atom being transferred and the radical site, and hydrogen bond lengths (dashed lines), are also shown. Reproduced with permission from reference 69. Copyright (2010) American Chemical Society.

Figure 16.

Calculated heavy atom bond lengths and dehydrocarbon atom separations (in Ångstroms) for the singlet and triplet states (in parentheses) of diradicals **32** and **35** – **38** at the BPW91/cc-pVDZ and CAS(8,8)/cc-pVDZ (italics) levels of theory. Reproduced from reference 153 with permission from Elsevier. Copyright (2000) Elsevier Science B.V.

Figure 17. Calculated heavy atom bond lengths and dehydrocarbon atom separations (in Ångstroms) for singlet and triplet states (in parenthesis) of radicals **33** and **53-55** at the BLYP/cc-pVDZ level of theory. Reproduced with permission from reference 244. Copyright (2001) American Chemical Society.

Figure 18. Calculated heavy atom bond lengths and dehydrocarbon atom separations (in Ångstroms) for singlet and triplet states (in parenthesis) of benzyne and didehydropyridine isomers at the BPW01/cc-pVDZ and CAS(8,8)/cc-pVDZ (italics) levels of theory. Reproduced from reference 153 with permission from Elsevier. Copyright (2000) Elsevier Science B.V.

Figure 19. Relative energy versus dehydrocarbon atom separation for **51**, **68**, and **69** (**b**, **c** and **d**, respectively). Reproduced with permission from reference 251. Copyright (2005) American Chemical Society.

Figure 20. Relative energy versus dehydrocarbon atom separation for radicals **65**, **70** and **71** (**e**, **f**, and **g**, respectively). Reproduced with permission from reference 251. Copyright (2005) American Chemical Society.

	87	89	88	90	91	86
Ground State:	2B_2	$^2A'$	$^2A'$	$^2A'$	$^2A'$	2B_2
D-Q splittings (kcal/mol):	-48.8	-46.2	-32.4	-39.8	-30.2	-29.3
EA (eV):	7.20	6.63	7.57	7.71	7.73	7.01

	77		78		79	
Ground State:	2A_1		$^2A'$		2A_1	
D-Q splittings (kcal/mol):	-49.5		-39.3		-29.7	
EA (eV):	1.52		1.61		0.68	

Ground State and D-Q splittings (kcal/mol): BD(T)/cc-pVTZ//BPW91/cc-pVDZ;
EA (eV): UBLYP/aug-cc-pVDZ/UBLYP/cc-pVDZ

Figure 21.

Ground states, doublet-quartet (D-Q) splittings and electron affinities (EA) of parent trihydrobenzenes and the corresponding tridehydropyridinium cations.

Figure 22.

Generation of 2,4,6-tridehydropyridinium cation **86** in an FT-ICR mass spectrometer depicted as follows: (A) Mass spectrum of radical cations generated upon first SORI-CAD event. (B) Mass spectrum of isolated ion of m/z 331. (C) Mass spectrum of ions generated upon second SORI-CAD event.

Triradical stabilization energy (TSE) (kcal/mol) at BD(T)/cc-pVTZ//BPW91/cc-pVDZ level of theory

Figure 23.
Triradical stabilization energies calculated for **86**.

Scheme 1.

Scheme 2.

Scheme 3.

Scheme 4a. **Scheme 4b.** **Scheme 4.**

Scheme 5.

Scheme 6.

NIH-PA Author Manuscript

Scheme 8.

Scheme 9.

Scheme 10.

Scheme 11.

Scheme 12.

Scheme 13.

Scheme 14.

Scheme 15.

Scheme 16.

Scheme 17.

Scheme 18.

Scheme 19.

Scheme 20.

Scheme 21.

Scheme 22.

Scheme 23.

Scheme 24.

Scheme 25.

Scheme 26.

Scheme 27.

Scheme 28.

Scheme 29.

Scheme 30.

Scheme 31.

Scheme 32.

Scheme 33.

Chart 1.

Chart 2.

Chart 3.

Chart 4.

Chart 5.

Chart 6.

Chart 7.

Chart 8.

Chart 9.

Chart 11.

Chart 12.

119 **120****Chart 13.**

Table 1

S-T energy splittings (kcal/mol) of *ortho*-arynes **35** and **38** – **41**. Reproduced with permission from reference 153. Copyright © 2000 Elsevier Science B.V.

	Benzynes	Pyridinium cations		Pyridynes

	32	37	38	35	36
BPW91 level	-31.3	-24.1	-31.5	-17.9	-35.2
CASPT2 level	-30.4	-23.6	-30.9	-14.6	-32.9
CCSD(T) level	-32.6	-25.2	-32.9	-18.9	-34.4
Expt.^a	-37.5 ± 0.3				

^aReference 92.

Table 2

S-T energy splittings (kcal/mol) of *meta*-arynes **36**, **49** – **51** and **53** – **58**. Reproduced with permission from reference 244. Copyright (2001) American Chemical Society.

	Benzyne	Pyridynes

	33	46	47	48
BPW91 level	-19.4	-23.4	-12.9	2.1
CASPT2 level	-18.0	-23.3	-14.2	1.2
CCSD(T) level	-18.3	-25.1	-14.8	-0.3
Expt. ^a	-21.0 ± 0.3			
	Benzyne	Pyridinium cations

	33	50	51	52
BPW91 level	-19.4	-24.8	-24.0	-8.1
CASPT2 level	-18.0	-18.3	-17.5	-10.6
CCSD(T) level	-18.3	-21.5	-19.6	-12.1
Expt. ^a	-21.0 ± 0.3			
	Benzyne	HO-substituted pyridynes

	33	53	54	55
BPW91 level	-19.4	-18.5	-24.9	-28.0
CASPT2 level	-18.0	-18.5	-24.0	-26.8
CCSD(T) level	-18.3	-18.6	-24.2	-27.1
Expt. ^a	-21.0 ± 0.3			

^aReference 245.

Table 3

Ionic products from gas-phase reactions of the *meta*-benzyne anion **49** with selected compounds. Reproduced with permission from reference 246. Copyright (1998) John Wiley & Sons, Ltd.

Compound	<i>meta</i> -Benzyne anion

NH ₃	No reaction
ND ₃	C ₆ H _{4-n} D _n ^{-*} (<i>n</i> = 1,2)
H ₂ O	No signal
D ₂ O	C ₆ H _{4-n} D _n ^{-*} (<i>n</i> = 1,2)
CS ₂	C ₆ H ₄ S ^{-*} , C ₆ H ₄ S ₂ ^{-*}
N ₂ O	C ₆ H ₄ O ^{-*} , (C ₅ H ₄) ^{-*}
CH ₃ SSCH ₃	CH ₃ S ⁻ , CH ₃ SCH ₂ S ⁻
NO ₂	NO ₂ ⁻
NO	C ₆ H ₄ NO ⁻ , C ₆ H ₄ (NO) ₂ ^{-*}
O ₂	C ₆ H ₄ O ^{-*}
RX, X = Cl, Br, R = alkyl	X ⁻
CH ₃ OH	CH ₃ O ⁻
CH ₃ OCH ₃	No reaction
PhCH ₃	PhCH ₂ ⁻
Cyclohexa-1,3-diene	<i>c</i> -C ₆ H ₇ ⁻

Table 4

S-T energy splittings (kcal/mol) of various benzyne. Reproduced with permission from reference 247.
Copyright (2003) American Chemical Society.

Benzyne	Theory	S-T Energy splitting
BLYP	-26.2
	BD(T)	-24.4
	hfc ^a	-23.8
BLYP	-20.0
	BD(T)	-19.8
	hfc ^a	-19.4
X = pyridinium	BLYP	-28.0
	hfc ^a	-24.2
X = pyridinium	BLYP	-22.3
	hfc ^a	-20.2
X = pyridinium	BLYP	-22.0
	hfc ^a	-20.9
Exp. ^b	-21.0 ± 0.3

^a Hyperfine splitting constant method using UBPW91/cc-pvDZ-optimized geometry.

^b Reference 245.

Table 5

Electron affinities (eV), S-T energy splittings (kcal/mol), and dehydrocarbon atom separations (Å) of various benzyne. Reproduced with permission from reference 251. Copyright (2005) American Chemical Society.

	EA ^a (eV)	S-T Energy splitting ^b (kcal/mol)	Dehydrocarbon atom separation ^c (Å)
51	6.30	-16.3	1.6
68	6.50	-14.1	1.9
69	6.59	-14.4	1.6
65	5.26	-16.7	1.6
70	5.62	-11.1	1.6
71	5.48	-15.4	2.0

^a Calculated at the (U)BLYP/aug-cc-pVDZ//MCSCF/cc-pVDZ level of theory.

^b Calculated at the CASPT2/cc-pVDZ//(U)BLYP/cc-pVDZ level of theory.

^c Calculated at the (U)BLYP/cc-pVDZ//(U)BLYP/cc-pVDZ level of theory.

Table 6

S-T energy splittings (kcal/mol) of *para*-arynes **37**, **76** and **77**. Reproduced with permission from reference 153. Copyright © 2000 Elsevier Science B.V.

	Benzyne	Pyridinium cation	Pyridyne

	34	73	74
BPW91 level	2.0	-4.3	-14.0
CASPT2 level	-5.8	-5.2	-13.0
CCSD(T) level	-2.8	-5.6	-12.9
Expt.^a	-3.9 ± 0.4		

^aReference 92.

Table 7

Ionic products from gas-phase reactions of the *para*-benzyne anion with selected compounds. Reproduced with permission from reference 246. Reproduced with permission from reference 246. Copyright (1998) John Wiley & Sons, Ltd.

Compound	<i>para</i> -Benzyne anion

NH ₃	No reaction
ND ₃	C ₆ H _{4-n} D _n ^{-*} (<i>n</i> = 1,2)
H ₂ O	No signal
D ₂ O	OD ⁻ , C ₆ H _{4-n} D _n ^{-*} (<i>n</i> = 1,2)
CS ₂	C ₆ H ₄ S ^{-*} , C ₆ H ₄ S ₂ ^{-*}
N ₂ O	C ₆ H ₄ O ^{-*} , (C ₅ H ₄ ^{-*})
CH ₃ SSCH ₃	CH ₃ S ⁻ , CH ₃ SCH ₂ S ⁻
NO ₂	NO ₂ ⁻
NO	C ₆ H ₄ NO ⁻ , C ₆ H ₄ (NO) ₂ ^{-*}
O ₂	C ₆ H ₄ O ^{-*}
RX, X = Cl, Br, R = alkyl	X ⁻
CH ₃ OH	CH ₃ O ⁻
CH ₃ OCH ₃	No reaction
PhCH ₃	PhCH ₂ ⁻
Cyclohexa-1,3-diene	<i>c</i> -C ₆ H ₇ ⁻