Rudimentary signs of immunosenescence in *Cytomegalovirus*-seropositive healthy young adults

James E. Turner · John P. Campbell · Kate M. Edwards ·
Lauren J. Howarth · Graham Pawelec · Sarah Aldred ·
Paul Moss · Mark T. Drayson · Victoria E. Burns · Jos A. Bosch

Received: 2 October 2012 / Accepted: 18 June 2013 / Published online: 12 July 2013
© American Aging Association 2013

Abstract Ageing is associated with a decline in immune competence termed immunosenescence. In the elderly, this process results in an accumulation of differentiated ‘effector’ phenotype memory T cells, predominantly driven by *Cytomegalovirus* (CMV) infection. Here, we asked whether CMV also drives immunity towards a senescent profile in healthy young adults. One hundred and fifty-eight individuals (mean $\pm$ SD; age 21 ± 3 years, body mass index 22.7 ± 2.7 kg m²) were assessed for CMV serostatus, the numbers/proportions of CD4⁺ and CD8⁺ late differentiated/effector memory cells (i.e. CD27[−]CD28[−]/CD45RA⁺), plasma interleukin-6 (IL-6) and antibody

responses to an in vivo antigen challenge (half-dose influenza vaccine). Thirty percent (48/158) of participants were CMV⁺. A higher lymphocyte and CD8⁺ count (both $p < 0.01$) and a lower CD4/CD8 ratio ($p < 0.03$) were observed in CMV⁺ people. Eight percent (4/58) of CMV⁺ individuals exhibited a CD4/CD8 ratio < 1.0 , whereas no CMV[−] donor showed an inverted ratio ($p < 0.001$). The numbers of CD4⁺ and CD8⁺CD27[−]CD28[−]/CD45RA⁺ cells were ~fourfold higher in CMV⁺ people ($p < 0.001$). Plasma IL-6 was higher in CMV⁺ donors ($p < 0.05$) and showed a positive association with the numbers of CD8⁺CD28[−] cells ($p < 0.03$). Finally, there was a

J. E. Turner · J. P. Campbell · L. J. Howarth · S. Aldred ·
V. E. Burns
School of Sport and Exercise Sciences,
University of Birmingham,
Birmingham, UK

J. E. Turner (✉) · P. Moss
School of Cancer Sciences, University of Birmingham,
Edgbaston, Birmingham B15 2TT, UK
e-mail: j.e.turner@bham.ac.uk

J. E. Turner
e-mail: turnerje1984@gmail.com

K. M. Edwards
Exercise Health and Performance Research Group,
University of Sydney,
Lidcombe, NSW, UK

J. P. Campbell · M. T. Drayson
School of Immunity and Infection,
University of Birmingham,
Birmingham, UK

G. Pawelec
Department of Internal Medicine II, Centre for Medical
Research, University of Tübingen,
Tübingen, Germany

J. A. Bosch
Mannheim Institute of Public Health, Social and Preventive
Medicine (MIPH), Mannheim Medical Faculty,
University of Heidelberg,
Heidelberg, Germany

J. E. Turner
Department for Health,
University of Bath,
Bath BA2 7AY, UK

J. A. Bosch (✉)
School of Psychology,
University of Amsterdam,
Amsterdam, The Netherlands
e-mail: j.a.bosch@uva.nl

significant negative correlation between vaccine-induced antibody responses to the A/Brisbane influenza strain and CMV-specific immunoglobulin G titres ($p < 0.05$). This reduced vaccination response was associated with greater numbers of total CD8⁺ and CD4⁺ and CD8⁺CD27[−]CD28[−]/CD45RA⁺ cells ($p < 0.05$). This study observed marked changes in the immune profile of young adults infected with CMV, suggesting that this virus may underlie rudimentary aspects of immunosenescence even in a chronologically young population.

Keywords Immunosenescence · *Cytomegalovirus* · Interleukin-6 · CD8⁺ T cells · Vaccination efficacy

Introduction

Ageing is associated with a gradual decline in immune competence, termed immunosenescence, which has been associated with increased susceptibility to infection, accelerated cognitive decline, frailty and increased mortality (Wikby et al. 2005). A hallmark of immunosenescence is an accumulation of differentiated or ‘effector’ phenotype CD8⁺ T cells (i.e. CD27[−]CD28[−]/CD45RA⁺). Other characteristics include a decline in the frequency and number of naïve CD8⁺ T cells (i.e. CD27⁺CD28⁺/CD45RA⁺) and increased inflammatory activity (e.g. plasma interleukin-6 (IL-6) levels) (Chidrawar et al. 2009; Wikby et al. 2005; Derhovanessian et al. 2009). It has become clear that these features are to a significant extent determined by latent *Cytomegalovirus* (CMV) infection (Derhovanessian et al. 2009; Olsson et al. 2000). CMV is a beta-herpes virus that establishes lifelong latency after primary infection. Seroprevalence, as determined by the presence of CMV-specific immunoglobulin G antibodies (anti-CMV IgG), increases with age to reach levels of over 70 % in many elderly populations (Staras et al. 2006).

It has been shown that the magnitude of the CMV-specific immune response can also influence clinical outcomes in CMV-seropositive individuals. High titres of anti-CMV IgG and increased numbers of late differentiated CD8⁺ T cells (i.e. CD27[−]CD28[−]) have been associated with lower antibody responses to influenza vaccination and higher circulating levels of inflammatory markers (Saurwein-Teissl et al. 2002; Goronzy et al. 2001; Trzonkowski et al. 2003; Wikby et al. 2006;

Moro-Garcia et al. 2012). In contrast, a recent study observed comparable antibody responses to influenza vaccination, irrespective of CMV serostatus, in residents of long-term care facilities (den Elzen et al. 2011). Indeed, as with the latter study, most observations of this kind have been made in elderly individuals that might already exhibit immune impairment. Thus, it remains largely unexplored whether CMV infection also drives immunity towards a senescent immune profile in healthy young adults. Here, in a chronologically young population, we have examined the influence of CMV serostatus on immune parameters including the CD4/CD8 ratio, the number of late differentiated/effector memory T cells and plasma IL-6 levels, as well as the *in vivo* functional response to antigen challenge (a half-dose influenza vaccine).

Methods

Participants

One hundred and fifty-eight healthy university students were recruited by local campus advertisement (Edwards et al. 2010). Mean±SD age and body mass index (BMI) were 21±3 years and 22.7±2.7 kg m², respectively. An equal number of males and females were recruited, and 90 % were of White-British ethnicity. Exclusion criteria were smoking and self-reported history of inflammatory, autoimmune or cardiovascular disease, self-reported pregnancy or suspected pregnancy and use of prescription medication in the past month (excluding the contraceptive pill). Participants self-reported no influenza-like illness in the year preceding this investigation and no symptoms of acute infection at the time of vaccination and follow-up measurements. All participants provided written informed consent, and the study protocol was approved by the Black Country Local Research Ethics Committee.

Procedures

Participants visited the laboratory between 12:00 and 16:00 to provide a baseline pre-vaccine blood sample and to receive an influenza vaccination. Participants returned to provide additional blood samples 24 h and 28 days after vaccination. Before arrival, participants were instructed to abstain from vigorous exercise and over-the-counter medication for 24 h, alcohol for 12 h

and food or caffeine for 2 h prior to their visit. Immediately after the baseline blood sample and prior to vaccination, the majority of participants ($n=119$) undertook a 25-min exercise intervention involving a series of weightlifting exercises. The results of this manipulation are described elsewhere (Edwards et al. 2010) and did not influence the present results.

Vaccination

Influenza vaccination was used as a marker of in vivo immune function (Burns and Gallagher 2010). To increase the sensitivity of this marker, a half dose (0.25 ml/7.5 µg; 50 % adult recommended dose) of the 2008/2009 Northern Hemisphere influenza vaccine (Fluarix, GlaxoSmithKline, Inactivated Split Virion, Lot No. AFLUA384AB) was used. The vaccine contained A/Brisbane- (both H3N2 and H1N1 strains), A/Uruguay- (H1N1 strain) and B/Florida-like strains. A nurse administered the vaccine via intramuscular injection into the deltoid.

Blood sample processing

Serum was obtained by allowing blood to clot in plain Vacutainer tubes (Becton-Dickinson, UK), centrifuging at $3,400\times g$ for 5 min at 21 °C, and was stored at −20 °C. Plasma was obtained by centrifuging blood in potassium ethylene-diamine-tetra-acetic acid (K_3EDTA) Vacutainer tubes at $3,400\times g$ for 10 min at 1 °C and was stored at −80 °C.

Assays

Influenza antibody titre was determined in serum before vaccination (baseline) and at 28 days using a haemagglutination inhibition test as previously described (Edwards et al. 2010). An antibody titre represents the highest serum dilution to inhibit the agglutination of test erythrocytes which bind together into a lattice-like structure upon exposure to influenza virus particles (Burns and Gallagher 2010). Anti-CMV IgG and IL-6 were measured in plasma before vaccination (baseline). CMV seropositivity was defined as having an anti-CMV IgG titre >3 IU/ml by ELISA, according to manufacturer's instructions (Genesis Diagnostics, UK). IL-6 was measured using a high sensitivity ELISA (Quantikine HS Human IL-6 ELISA, R&D Systems, UK). Assay sensitivity was 0.1 IU/ml and

0.039 pg/ml for the CMV and IL-6 ELISAs, respectively. Only one individual fell below the sensitivity threshold for IL-6 (0.02 pg/ml) and was included in the analyses. Intra- and inter-assay precision (CV %) were <10 % for both assays.

Flow cytometry and immunophenotyping

Leukocyte differential was assessed in K_3EDTA blood 24 h post-vaccination and repeated 28 days later (Coulter ACT^{diff} haematology analyser; Beckman-Coulter, High Wycombe, UK). These samples were processed for flow cytometric measurements as previously described (Turner et al. 2010). Fixed cell preparations were read on a multi-parameter flow cytometer (BD FACS CANTO II, BD Biosciences). Lymphocytes were gated on the forward versus side scatter. Sub-populations of $CD3^+CD4^+$ and $CD3^+CD8^+$ T cells were identified using two analytical strategies. First, expression of CD27 in combination with CD45RA identified naïve ($CD27^+CD45RA^+$), central memory (CM; $CD27^+CD45RA^-$), effector memory (EM; $CD27^-CD45RA^-$) and 'revertant' effector memory cells which re-express CD45RA (EMRA; $CD27^-CD45RA^+$). Second, CD27 expression was analysed in combination with CD28 to identify early ($CD27^+CD28^+$), intermediate ($CD27^+CD28^-$) and late differentiated ($CD27^-CD28^-$) sub-populations (Hamann et al. 1997; Appay et al. 2002). Data were analysed with FlowJo (Treestar, Ashland, OR, USA).

Statistical analyses

Data were inspected for normal distribution using the Kolmogorov–Smirnov test. Non-normally distributed data were transformed logarithmically. Differences between CMV^+ and CMV^- participants were assessed with univariate analyses of variance (ANOVA) (continuous variables) or chi-squared tests (categorical variables). Relationships between IL-6 and lymphocyte sub-populations were established using stepwise regression analysis. Anti-influenza antibody titres were assessed between baseline and 28 days post-vaccination with repeated measures ANOVA. To identify factors moderating the vaccination response, key variables (e.g. anti-CMV IgG) were entered into ANOVAs as individual covariates and examined for significance (analysis of co-variance—ANCOVA). Significant covariates were further examined adjusting

for sex, CMV serostatus, baseline influenza antibody titre and the pre-vaccine exercise intervention. Pearson's correlations established the direction of relationships between key variables (e.g. anti-CMV IgG) and fold change of influenza antibody titre. Fold change was calculated by dividing the geometric mean antibody titre by the baseline titre. Effect sizes are reported as eta-squared (η^2). Conventionally, η^2 values of 0.01, 0.06 and 0.14 are considered small, medium and large effect sizes, respectively. For example, η^2 of 0.25 indicates that 25 % of the observed effects (i.e. dependent variable) are accounted for by the experimental manipulation (i.e. the independent variable). Data are presented as means $\pm$ standard error of the mean (SEM) unless otherwise stated. Data were analysed using SPSS statistical package version 18.0 for Windows (SPSS Inc., USA).

Results

CMV serostatus

Approximately 30 % of individuals ($n=48/158$) were CMV⁺ as determined by anti-CMV IgG titres > 3 IU/ml. CMV serostatus was not associated with age, sex, ethnicity or BMI ($p>0.1$).

CMV seropositivity is associated with a higher lymphocyte count and lower CD4/CD8 ratio

Lymphocyte and T cell sub-population data were averaged for the two collection days (i.e. 24 h and 28 days post-vaccination). Analyses showed that for both days, the proportions of cells were identical ($p>0.38$), except for CD8⁺CD27[−]CD45RA^{−/+} cells ($p<0.01$) and correlated between samples (average correlation $r=0.72$, $p<0.001$). Moreover, the results reported below remained the same when collection days were analysed separately.

CMV seropositivity was associated with a higher lymphocyte count and a greater number of CD8⁺ T cells ($F_{(1, 152)}>8.1$; $p<0.01$, $\eta^2>0.050$; data not shown). CMV seropositivity was also associated with a significantly lower CD4/CD8 ratio ($F_{(1, 152)}=5.1$, $p<0.03$, $\eta^2=0.032$; see Fig. 1a). It is notable that a CD4/CD8 ratio < 1.0 was observed in 4/48 CMV⁺ individuals (8 % of CMV⁺) and none of the CMV[−] individuals (independent samples t test; $t_{(108)}=3.8$, $p<0.001$).

CMV infection is associated with an expansion of late differentiated effector memory CD4⁺ and CD8⁺ T cells

The numbers and proportions of CD4⁺ and CD8⁺ EM (CD27[−]CD45RA[−]), EMRA (CD27[−]CD45RA⁺) and late differentiated (CD27[−]CD28[−]) cells were significantly increased in CMV⁺ individuals ($F_{(1, 152)}>30.6$; $p<0.001$, $\eta^2>0.167$, see Fig. 1b, c and Table 1). Further, individuals with higher anti-CMV IgG titres (in CMV⁺ participants) had increased numbers of total lymphocytes, CD8⁺ EMRA and CD4⁺ late stage differentiated cells (Pearson's correlations $r=0.32$ to 0.38 , $p<0.03$; CD8⁺ late differentiated cells $r=0.27$, $p=0.064$; see Fig. 2a–d).

Plasma IL-6 is increased with CMV infection and associated with the number of CD8⁺CD28[−] cells

Modestly elevated plasma IL-6 was observed in CMV⁺ individuals, as compared to CMV[−] donors ($F_{(1, 152)}=4.3$, $p<0.05$, $\eta^2=0.027$; see Fig. 3a). Individuals with higher IL-6 had increased numbers of both intermediate (CD27⁺CD28[−]) and late stage (CD27[−]CD28[−]) differentiated CD8⁺ cells (Pearson's correlation $r>0.18$, $p<0.03$; data not shown). The relationship between CD28[−] cells and IL-6 was largely an effect of CMV infection. After statistical adjustment for anti-CMV IgG, the association between IL-6 and late stage differentiated (CD27[−]CD28[−]) cells was lost. This association remained close to significance for intermediate stage (CD27⁺CD28[−]) cells ($r=0.19$; $p<0.05$).

Higher anti-CMV IgG titres are associated with weaker antibody responses to A/Brisbane antigen

CMV serostatus did not predict vaccination responses (defined as a >fourfold change in antibody titre 28 days post-vaccination) to any of the influenza strains ($p>0.1$). However, a higher concentration of anti-CMV IgG was associated with a smaller increase in antibody titres against the A/Brisbane antigen (anti-CMV IgG $\times$ time interaction $F_{(1, 152)}=4.1$, $p<0.05$, $\eta^2=0.026$; Pearson's correlation $r=0.16$, $p<0.05$; see Fig. 3b). This effect remained significant after statistical adjustment for sex, A/Brisbane baseline antibody titre and the pre-vaccine exercise intervention (adjusted anti-CMV IgG $\times$ time interactions; $F_{(1, 155)}>3.9$, $p<0.05$, $\eta^2>0.025$). There were no relationships between anti-CMV IgG and antibody responses to the other influenza strains.

Fig. 1 Composition of the T cell pool CMV infection. **a** Ratio of CD4⁺ to CD8⁺ T lymphocytes; 4/58 CMV⁺ individuals (8 %) exhibited a ratio <1.0. **b** CD4⁺ and CD8⁺ effector memory (EM; CD27⁺CD45RA⁺) and ‘revertant’ effector memory (EMRA; CD27⁺CD45RA⁺) T lymphocytes. **c** CD4⁺ and CD8⁺ T intermediate stage (CD27⁺CD28⁺) and late differentiated (CD27⁺CD28⁺) T lymphocytes. Data are means±SEM. ****p*<0.001, **p*<0.05 CMV⁺ compared to CMV⁻

High numbers of total lymphocytes and late differentiated T cells are associated with smaller antibody responses to A/Brisbane antigen

A weaker A/Brisbane antibody response was associated with a higher total lymphocyte and CD8⁺ T cell count, as well as increased numbers of late differentiated CD4⁺ and CD8⁺ T cells (CD27⁺CD28⁺) and CD8⁺ EM (CD27⁺CD45RA⁺) and EMRA (CD27⁺CD45RA⁺) cells (cell population×time interactions; $F_{(1, 156)} > 4.4$, $p < 0.05$, $\eta^2 > 0.041$; Pearson’s correlations $r = -0.16$ to -0.23 , $p < 0.05$). These effects remained significant after statistical adjustment for sex, A/Brisbane baseline antibody titre and the pre-vaccine exercise intervention (adjusted analyses for all; $F_{(1, 155)} > 4.2$, $p < 0.05$, $\eta^2 > 0.019$). However, the observation that individuals

with higher cell counts exhibited smaller A/Brisbane antibody responses was largely an effect of CMV infection. After statistical adjustment for anti-CMV IgG or CMV serostatus, most of these associations were lost, except for total lymphocytes and total CD8⁺ T cells (analyses adjusted for CMV infection; $F_{(1, 155)} > 4.2$, $p < 0.05$, $\eta^2 > 0.026$). There were no relationships between lymphocyte and lymphocyte sub-populations with antibody responses to the other influenza strains.

with higher cell counts exhibited smaller A/Brisbane antibody responses was largely an effect of CMV infection. After statistical adjustment for anti-CMV IgG or CMV serostatus, most of these associations were lost, except for total lymphocytes and total CD8⁺ T cells (analyses adjusted for CMV infection; $F_{(1, 155)} > 4.2$, $p < 0.05$, $\eta^2 > 0.026$). There were no relationships between lymphocyte and lymphocyte sub-populations with antibody responses to the other influenza strains.

Discussion

The results of this study show that young adults infected with CMV exhibit mild signs of immunosenescence, characterised by increased numbers of late

Table 1 Proportion of lymphocyte sub-populations in CMV⁻ and CMV⁺ participants (mean±SD)

Cells % ^a	CMV ⁻	CMV ⁺
CD4 ⁺ ^b	39.44±6.18	38.53±5.55
CD4 ⁺ naïve	46.26±10.27	44.81±10.96
CD4 ⁺ CM	48.61±9.23	46.16±9.54
CD4 ⁺ EM	4.90±2.31	7.50±4.22***
CD4 ⁺ EMRA	0.22±0.49	1.53±3.83***
CD4 ⁺ early	93.60±2.60	89.68±7.32*
CD4 ⁺ inter	1.27±0.81	1.28±1.03
CD4 ⁺ late	0.19±0.43	3.29±6.07***
CD8 ⁺ ^b	20.77±4.12	23.42±5.36**
CD8 ⁺ naïve	57.77±10.52	48.69±13.71***
CD8 ⁺ CM	34.38±9.35	30.10±11.97**
CD8 ⁺ EM	3.88±2.68	7.23±5.48***
CD8 ⁺ EMRA	3.96±5.32	13.98±11.12***
CD8 ⁺ early	79.38±9.14	65.98±14.88***
CD8 ⁺ inter	12.78±6.44	12.81±8.18
CD8 ⁺ late	5.39±5.54	15.90±12.87***

^a as a proportion of CD4⁺ or CD8⁺ T lymphocytes^b Cells as a proportion of lymphocytes****p*<0.001, ***p*<0.01, **p*<0.05, univariate ANOVA

differentiated/effector memory T cells and elevated inflammatory activity. The present study adds to the very sparse data on the relationship between CMV and T cell differentiation in healthy young adults and is the first to confirm that the magnitude of the humoral response to CMV (i.e. anti-CMV IgG) is associated with an accumulation of late differentiated/effector memory T cells. Until now, to the best of our knowledge, this phenomenon has only been shown in elderly subjects (Alonso Arias et al. 2013; Vescovini et al. 2010). Moreover, donors with the strongest CMV-specific immune response, characterised by high anti-CMV IgG titres and memory T cell numbers, exhibited a reduced antibody response to a half-dose influenza vaccination.

Fig. 2 Associations between anti-CMV IgG and lymphocyte populations. **a** Total lymphocytes. **b** CD8⁺ ‘revertant’ effector memory (EMRA; CD27⁻CD45RA⁺) T lymphocytes. **c** CD4⁺ late differentiated (CD27⁻CD28⁻) T lymphocytes. **d** CD8⁺ late differentiated (CD27⁻CD28⁻) T lymphocytes (**p*=0.064). Statistical analyses were restricted to CMV⁺ individuals (*n*=48) and were conducted on log10 data. β represents standardised regression coefficient

Although the majority of individuals exhibited a >fourfold change in antibody titre 28 days post-vaccination to each influenza strain (i.e. individuals were

Fig. 3 Elevated inflammatory activity and reduced antibody responses to antigen challenge with CMV infection. **a** Plasma IL-6 concentration. **b** Data depict the association between anti-CMV IgG titre and fold change in A/Brisbane antibody titre 28 days after influenza vaccination ($p < 0.05$). For illustrative purposes, anti-CMV IgG is presented in quartiles; CMV⁻ < 3.0 IU/ml, CMV⁺ low 3.0–5.1 IU/ml, CMV⁺ med 5.1–9.1 IU/ml, CMV⁺ high > 9.1 IU/ml. Statistical analyses were conducted on continuous data (Altman and Royston 2006). Data are means $\pm$ SEM. * $p < 0.05$ CMV⁺ compared to CMV⁻

defined by the World Health Organisation as ‘vaccine responders’), we found that an increased anti-CMV IgG titre was associated with a smaller antibody response to the A/Brisbane influenza antigen only. Anti-CMV IgG

was also associated with increased numbers of late differentiated/effector memory T cells. In turn, a higher number of these cells were also associated with lower vaccine responses to the A/Brisbane antigen, but this was a consequence of CMV infection: when controlling statistically for CMV serostatus, or anti-CMV IgG titre, most of these associations were lost.

It is unclear why our observations were limited to the A/Brisbane vaccine component although strain-specific differences in the magnitude of antibody response to influenza vaccination have commonly been reported in a variety of contexts. For example, obese individuals have been shown to exhibit larger antibody responses to single influenza vaccine components (Sheridan et al. 2012; Talbot et al. 2012). Further, sexual dimorphism and diurnal variation in vaccine efficacy have been observed with single influenza sub-types in trivalent vaccines (Phillips et al. 2008; Langlois et al. 1995). Finally, the age-associated decline in antibody response to vaccination is not uniform between strains. Differences in antibody response between young and old are largest with H1N1 and B antigens, compared to H3N2 antigens (Goodwin et al. 2006). Indeed, our laboratory has repeatedly shown that weaker vaccine antigens are the most sensitive to mild forms of immunomodulation, such as exercise or chronic stress (Edwards et al. 2006, 2007, 2008, 2010; Burns and Gallagher 2010). We speculate that strain-specific variation in the immunogenicity of vaccine components might explain contradictory findings in the literature (Burns and Gallagher 2010; Burns et al. 2003; den Elzen et al. 2011; Goodwin et al. 2006). In the present study, the A/Brisbane strain was the most immunogenic vaccine component (Edwards et al. 2010) consisting of both H1N1 and H3N2 subtypes, which show the greatest age-related differences in immunogenicity (Goodwin et al. 2006). Significantly, H3N2 was highly prevalent around the time of this investigation (CDC 2008) and is the strain associated with most influenza-related complications and deaths (Thompson et al. 2003). It is therefore fascinating to show that the immunosuppressive effects of CMV infection can be observed with highly immunogenic influenza vaccine components in such a young and healthy population. Thus, the present findings provide further support for the concept that CMV is a major driving force behind immunosenescence and might result in vaccine failure and increased infection risk in CMV-seropositive individuals (Derhovanessian et al. 2009).

It is currently unknown exactly how infection with CMV could promote lower antibody responses to vaccination. Immunosenescence and chronological ageing might be associated with impaired dendritic cell function (Panda et al. 2010). It is unclear whether this is a direct effect of CMV infection on antigen presenting cells, or a secondary downstream consequence of infection (e.g. indirect effects caused by the accumulation of memory T cells). Indeed, dendritic cell function is impaired by co-incubation with CD8⁺CD28⁻ cells, which promote expression of negative regulatory receptors such as immunoglobulin-like transcripts 3 and 4 (ILT3 and ILT4) (Chang et al. 2002). In addition, CD8⁺CD28⁻ cells cause a down-regulation of CD40 ligand on CD4⁺ T helper cells (Cortesini et al. 2001) and the ensuing impaired responsiveness of CD4⁺ T helper cells may in turn inhibit antibody production, which is a hallmark of B cell immunosenescence (Siegrist and Aspinall 2009).

It is notable that 8 % of CMV⁺ individuals (4/58) in the present study showed an inverted CD4/CD8 ratio of less than 1.0. This marker of immunosenescence is a hallmark of the ‘immune risk profile’ (IRP), an immune phenotype observed in around 15 % of people aged over 85 years in the Swedish OCTO/NONA studies. The IRP is a cluster of immunological markers, including the accumulation of CD8⁺ memory T cells (resulting in an inverted CD4/CD8 ratio) as well as CMV infection, which predicts all-cause mortality in the very old (Wikby et al. 2005, 2006; Derhovanessian et al. 2009). Further, a recent 10-year follow up of ~14,000 adults from the National Health and Nutrition Examination Survey provides evidence that CMV infection alone predicts all-cause mortality independently of inflammatory activity (CRP level) (Simanek et al. 2011). The results of the present study suggest that a mild IRP-like phenotype (which might be associated with mortality) can be observed in some adults much earlier in life than has been reported previously.

To the best of our knowledge, we have shown for the first time that healthy young adults latently infected with CMV exhibit increased levels of IL-6. Until now, interaction between IL-6 and pathogen burden has only been observed in the elderly (Nazmi et al. 2010; Schmaltz et al. 2005). This result seems mechanistically plausible considering the strong in vitro evidence that CMV infection in a variety of cell types, including monocytes, epithelial cells and

adipocytes, results in a marked increase of IL-6 gene and protein expression (Bouwman et al. 2009; Geist and Dai 1996; Iwamoto and Konicek 1997; Visseren et al. 1999). Increased IL-6 is thought to reflect viral reactivation (Bennett et al. 2011) due to a potential role for this cytokine in chronic viral control (Harker et al. 2011). The present manuscript also shows that elevated IL-6 level is associated with the numbers of CD8⁺CD28⁻ T cells. Although our data suggest that this relationship is largely a direct effect of CMV infection, a causal relationship between CD8⁺CD28⁻ T cells and IL-6 is biologically plausible. For example, CD8⁺CD28⁻ are efficient producers of inflammatory cytokines such as TNF- α , IFN- γ and IL-6 (Zanni et al. 2003; Lorre et al. 1994; Effros et al. 2005). In addition, an association between systemic inflammation and an accumulation of CD8⁺CD28⁻ cells has also been found in healthy older adults (Zanni et al. 2003; Wikby et al. 2006), as well as in a variety of autoimmune diseases including rheumatoid arthritis, multiple sclerosis and Graves’ disease (Sun et al. 2008; Schmidt et al. 1996; Markovic-Plese et al. 2001). Together, our results extend the existing literature that links the concept of ‘inflamm-ageing’ and T cell senescence (Franceschi et al. 2000) by showing that such associations may even be present in healthy young adults, likely driven by CMV infection.

In the present study, individuals infected with CMV exhibited plasma IL-6 levels approximately 20 % higher than CMV-negative individuals. This is comparable to the elevated level of IL-6 seen with men and women who develop coronary heart disease later in life, compared to those who remain healthy (Pai et al. 2004). The present finding may thus provide an additional mechanism by which CMV has been implicated in the development of cardiovascular disease (Soderberg-Naucler 2006; Strandberg et al. 2009). Individuals with a guanine>cytosine single nucleotide polymorphism (SNP) at position -174 in the promoter region for the IL-6 gene show ~60 % increase in IL-6 levels and are at greater risk of cardiovascular disease (Rafiq et al. 2007). It would be interesting to examine whether being CMV seropositive with this SNP carries a further increased risk of disease (Soderberg-Naucler 2006; Pai et al. 2004).

This study does not provide data on specific T cell responses to CMV infection (e.g. IFN- γ production to CMV antigens and MHC class I or II tetramer staining). However, in CMV-seropositive individuals, it is

well accepted that the majority of CD27⁺ and CD28⁺ T cells are CMV specific. Moreover, a substantial number of studies have shown positive correlations between the number of late differentiated/effector memory T cells and the CMV-specific immune response across a range of ages (Griffiths et al. 2013; Lachmann et al. 2012; Khan et al. 2002; Pita-Lopez et al. 2009). In future studies, such measurements of CMV-specific T cells could be correlated with the antibody response to influenza vaccination in young adults. While these data may not necessarily contribute to our understanding of the mechanisms underlying diminished vaccine responses, it would broaden our understanding of CMV infection in young adults and might provide a potential predictor of vaccine efficacy in CMV-infected people. Based on the small amount of data on this topic, it might be expected that the magnitude of IFN- γ production by CMV-specific T cells in young adults would be negatively associated with influenza vaccine responses, as has been shown in the elderly (Moro-Garcia et al. 2012).

The role of CMV in immunosenescence remains a topic of debate (Wills et al. 2011). For example, it has been shown that in residents of long-term care facilities, CMV does not impair the antibody response to influenza vaccination (den Elzen et al. 2011). In light of these and other inconsistent findings, it cannot be assumed that the effects of CMV shown in the elderly in some, but not all studies, will translate into young adults. In the present study, by measuring multiple biomarkers of immunosenescence, we confirm that the deleterious effects of CMV infection, sometimes shown in the elderly, do indeed translate to young adults (Saurwein-Teissl et al. 2002; Goronzy et al. 2001; Trzonkowski et al. 2003; Wikby et al. 2006). Our findings show that healthy university students, infected with CMV, exhibit an accumulation of late differentiated/effector memory T cells, increased inflammatory activity and poorer immune responses to novel antigens early in life (i.e. well before overt signs of immune impairment appear). It should be considered, however, that the population examined in this work is a socially homogenous group of young adults, the majority of whom were of White-British ethnicity and of high socio-economic status. Making identical measurements in a more heterogeneous group of individuals might result in larger and stronger effects of CMV on the results presented.

The present observations are relevant to influenza control and might also apply to other routine vaccination

programmes (e.g. hepatitis B). Likewise, these findings might warrant investigation of vaccine efficacy in other contexts, such as yellow fever, which is important considering the popularity of worldwide travel. Another implication is that the current findings suggest a different theoretical approach to T cell immunosenescence, which might be relevant to the type and timing of interventions that intend to ameliorate or prevent immunosenescence (Adler 2008; Pawelec et al. 2010). Traditional thinking is that senescence is primarily driven by thymic involution, whereby CMV infection could act as a secondary or amplifying factor. The current observations instead suggest a more primary role of CMV in establishing rudimentary features of immunosenescence very early in life, whereby subsequent thymic involution may potentially act as a secondary feature.

Acknowledgments The authors would like to thank Alison Whitelegg for assistance with optimising the flow cytometry assay used in this study. Acknowledgements also go to Riyad Suliman Khanfer, Charlotte Downes, Joanna Long, Josephine Lumb, Alex Merry and Nicola Paine for their assistance with participant recruitment. This study was funded by Action Medical Research. LJH was supported by a Wellcome Trust Vacation Scholarship. GP was supported by the Deutsche Forschungsgemeinschaft (DFG PA 361/14-1), the Bundesministerium für Bildung und Forschung (BMBF 0315890 F, “Gerontoshield” and 89718016 “BASE-II”) and the European Commission (EU-FP7 IDEAL 259679).

References

- Adler SP (2008) Human CMV vaccine trials: what if CMV caused a rash? *J Clin Virol* 41(3):231–236
- Alonso Arias R, Moro-Garcia MA, Echeverria A, Solano-Jaurieta JJ, Suarez-Garcia FM, Lopez-Larrea C (2013) Intensity of the humoral response to cytomegalovirus is associated with the phenotypic and functional status of the immune system. *J Virol* 87(8):4486–4495
- Altman DG, Royston P (2006) The cost of dichotomising continuous variables. *BMJ* 332(7549):1080
- Appay V, Dunbar PR, Callan M, Klenerman P, Gillespie GM, Papagno L, Ogg GS, King A, Lechner F, Spina CA, Little S, Havlir DV, Richman DD, Gruener N, Pape G, Waters A, Easterbrook P, Salio M, Cerundolo V, McMichael AJ, Rowland-Jones SL (2002) Memory CD8⁺ T cells vary in differentiation phenotype in different persistent virus infections. *Nat Med* 8(4):379–385
- Bennett JM, Glaser R, Malarkey WB, Beversdorf DQ, Peng J, Kiecolt-Glaser JK (2011) Inflammation and reactivation of latent herpesviruses in older adults. *Brain Behav Immun*
- Bouwman JJ, Diepersloot RJ, Visseren FL (2009) Intracellular infections enhance interleukin-6 and plasminogen activator

- inhibitor 1 production by cocultivated human adipocytes and THP-1 monocytes. *Clin Vaccine Immunol* 16(8):1222–1227
- Burns VE, Carroll D, Ring C, Drayson M (2003) Antibody response to vaccination and psychosocial stress in humans: relationships and mechanisms. *Vaccine* 21(19–20):2523–2534
- Burns VE, Gallagher S (2010) Antibody response to vaccination as a marker of in vivo immune function in psychophysiological research. *Neurosci Biobehav Rev* 35(1):122–126
- CDC (2008) Influenza activity—United States and worldwide, 2007–08 season. *MMWR Morb Mortal Wkly Rep* 57(25):692–697
- Chang CC, Ciubotariu R, Manavalan JS, Yuan J, Colovai AI, Piazza F, Lederman S, Colonna M, Cortesini R, Dalla-Favera R, Suciuc-Foca N (2002) Tolerization of dendritic cells by T(S) cells: the crucial role of inhibitory receptors ILT3 and ILT4. *Nat Immunol* 3(3):237–243
- Chidrawar S, Khan N, Wei W, McLarnon A, Smith N, Nayak L, Moss P (2009) Cytomegalovirus-seropositivity has a profound influence on the magnitude of major lymphoid subsets within healthy individuals. *Clin Exp Immunol* 155(3):423–432
- Cortesini R, LeMaout J, Ciubotariu R, Cortesini NS (2001) CD8+CD28– T suppressor cells and the induction of antigen-specific, antigen-presenting cell-mediated suppression of Th reactivity. *Immunol Rev* 182:201–206
- den Elzen WP, Vossen AC, Cools HJ, Westendorp RG, Kroes AC, Gussekloo J (2011) Cytomegalovirus infection and responsiveness to influenza vaccination in elderly residents of long-term care facilities. *Vaccine* 29(29–30):4869–4874
- Derhovanessian E, Larbi A, Pawelec G (2009) Biomarkers of human immunosenescence: impact of Cytomegalovirus infection. *Curr Opin Immunol* 21(4):440–445
- Edwards KM, Burns VE, Adkins AE, Carroll D, Drayson M, Ring C (2008) Meningococcal A vaccination response is enhanced by acute stress in men. *Psychosom Med* 70(2):147–151
- Edwards KM, Burns VE, Allen LM, McPhee JS, Bosch JA, Carroll D, Drayson M, Ring C (2007) Eccentric exercise as an adjuvant to influenza vaccination in humans. *Brain Behav Immun* 21(2):209–217
- Edwards KM, Burns VE, Reynolds T, Carroll D, Drayson M, Ring C (2006) Acute stress exposure prior to influenza vaccination enhances antibody response in women. *Brain Behav Immun* 20(2):159–168
- Edwards KM, Campbell JP, Ring C, Drayson MT, Bosch JA, Downes C, Long JE, Lumb JA, Merry A, Paine NJ, Burns VE (2010) Exercise intensity does not influence the efficacy of eccentric exercise as a behavioural adjuvant to vaccination. *Brain Behav Immun* 24(4):623–630
- Effros RB, Dagarag M, Spaulding C, Man J (2005) The role of CD8+ T-cell replicative senescence in human aging. *Immunol Rev* 205:147–157
- Franceschi C, Bonafe M, Valensin S, Olivieri F, De Luca M, Ottaviani E, De Benedictis G (2000) Inflamm-aging. An evolutionary perspective on immunosenescence. *Ann N Y Acad Sci* 908:244–254
- Geist LJ, Dai LY (1996) Cytomegalovirus modulates interleukin-6 gene expression. *Transplantation* 62(5):653–658
- Goodwin K, Viboud C, Simonsen L (2006) Antibody response to influenza vaccination in the elderly: a quantitative review. *Vaccine* 24(8):1159–1169
- Goronzy JJ, Fulbright JW, Crowson CS, Poland GA, O'Fallon WM, Weyand CM (2001) Value of immunological markers in predicting responsiveness to influenza vaccination in elderly individuals. *J Virol* 75(24):12182–12187
- Griffiths SJ, Riddell NE, Masters J, Libri V, Henson SM, Wertheimer A, Wallace D, Sims S, Rivino L, Larbi A, Kemeny DM, Nikolich-Zugich J, Kern F, Klennerman P, Emery VC, Akbar AN (2013) Age-associated increase of low-avidity Cytomegalovirus-specific CD8+ T cells that re-express CD45RA. *J Immunol* 190(11):5363–5372
- Hamann D, Baars PA, Rep MH, Hooibrink B, Kerkhof-Garde SR, Klein MR, van Lier RA (1997) Phenotypic and functional separation of memory and effector human CD8+ T cells. *J Exp Med* 186(9):1407–1418
- Harker JA, Lewis GM, Mack L, Zuniga EI (2011) Late interleukin-6 escalates T follicular helper cell responses and controls a chronic viral infection. *Science* 334(6057):825–829
- Iwamoto GK, Konicek SA (1997) Cytomegalovirus immediate early genes upregulate interleukin-6 gene expression. *J Invest Med* 45(4):175–182
- Khan N, Shariff N, Cobbold M, Bruton R, Ainsworth JA, Sinclair AJ, Nayak L, Moss PA (2002) Cytomegalovirus seropositivity drives the CD8 T cell repertoire toward greater clonality in healthy elderly individuals. *J Immunol* 169(4):1984–1992
- Lachmann R, Bajwa M, Vita S, Smith H, Cheek E, Akbar A, Kern F (2012) Polyfunctional T cells accumulate in large human cytomegalovirus-specific T cell responses. *J Virol* 86(2):1001–1009
- Langlois PH, Smolensky MH, Glezen WP, Keitel WA (1995) Diurnal variation in responses to influenza vaccine. *Chronobiol Int* 12(1):28–36
- Lorre K, Kasran A, Van Vaeck F, de Boer M, Ceuppens JL (1994) Interleukin-1 and B7/CD28 interaction regulate interleukin-6 production by human T cells. *Clin Immunol Immunopathol* 70(1):81–90
- Markovic-Plese S, Cortese I, Wandinger KP, McFarland HF, Martin R (2001) CD4+CD28– costimulation-independent T cells in multiple sclerosis. *J Clin Invest* 108(8):1185–1194
- Moro-Garcia MA, Alonso-Arias R, Lopez-Vazquez A, Suarez-Garcia FM, Solano-Jaurrieta JJ, Baltar J, Lopez-Larrea C (2012) Relationship between functional ability in older people, immune system status, and intensity of response to CMV. *Age (Dordr)* 34(2):479–495
- Nazmi A, Diez-Roux AV, Jenny NS, Tsai MY, Szklo M, Aiello AE (2010) The influence of persistent pathogens on circulating levels of inflammatory markers: a cross-sectional analysis from the Multi-Ethnic Study of Atherosclerosis. *BMC Publ Health* 10:706
- Olsson J, Wikby A, Johansson B, Lofgren S, Nilsson BO, Ferguson FG (2000) Age-related change in peripheral blood T-lymphocyte subpopulations and cytomegalovirus infection in the very old: the Swedish longitudinal OCTO immune study. *Mech Ageing Dev* 121(1–3):187–201
- Pai JK, Pischon T, Ma J, Manson JE, Hankinson SE, Joshipura K, Curhan GC, Rifai N, Cannuscio CC, Stampfer MJ, Rimm EB (2004) Inflammatory markers and the risk of coronary heart disease in men and women. *N Engl J Med* 351(25):2599–2610

- Panda A, Qian F, Mohanty S, van Duin D, Newman FK, Zhang L, Chen S, Towle V, Belshe RB, Fikrig E, Allore HG, Montgomery RR, Shaw AC (2010) Age-associated decrease in TLR function in primary human dendritic cells predicts influenza vaccine response. *J Immunol* 184(5):2518–2527
- Pawelec G, Akbar A, Beverley P, Caruso C, Derhovanessian E, Fulop T, Griffiths P, Grubeck-Loebenstein B, Hamprecht K, Jahn G, Kern F, Koch SD, Larbi A, Maier AB, Macallan D, Moss P, Samson S, Strindhall J, Trannoy E, Wills M (2010) Immunosenescence and Cytomegalovirus: where do we stand after a decade? *Immun Ageing* 7:13
- Phillips AC, Gallagher S, Carroll D, Drayson M (2008) Preliminary evidence that morning vaccination is associated with an enhanced antibody response in men. *Psychophysiology* 45(4):663–666
- Pita-Lopez ML, Gayoso I, Delarosa O, Casado JG, Alonso C, Munoz-Gomariz E, Tarazona R, Solana R (2009) Effect of ageing on CMV-specific CD8 T cells from CMV seropositive healthy donors. *Immun Ageing* 6:11
- Rafiq S, Frayling TM, Murray A, Hurst A, Stevens K, Weedon MN, Henley W, Ferrucci L, Bandinelli S, Corsi AM, Guralnik JM, Melzer D (2007) A common variant of the interleukin 6 receptor (IL-6r) gene increases IL-6r and IL-6 levels, without other inflammatory effects. *Genes Immun* 8(7):552–559
- Saurwein-Teissl M, Lung TL, Marx F, Gschosser C, Asch E, Blasko I, Parson W, Bock G, Schonitzer D, Trannoy E, Grubeck-Loebenstein B (2002) Lack of antibody production following immunization in old age: association with CD8(+)CD28(−) T cell clonal expansions and an imbalance in the production of Th1 and Th2 cytokines. *J Immunol* 168(11):5893–5899
- Schmaltz HN, Fried LP, Xue QL, Walston J, Leng SX, Semba RD (2005) Chronic cytomegalovirus infection and inflammation are associated with prevalent frailty in community-dwelling older women. *J Am Geriatr Soc* 53(5):747–754
- Schmidt D, Goronzy JJ, Weyand CM (1996) CD4+ CD7−CD28− T cells are expanded in rheumatoid arthritis and are characterized by autoreactivity. *J Clin Invest* 97(9):2027–2037
- Sheridan PA, Paich HA, Handy J, Karlsson EA, Hudgens MG, Sammon AB, Holland LA, Weir S, Noah TL, Beck MA (2012) Obesity is associated with impaired immune response to influenza vaccination in humans. *Int J Obes (Lond)* 36(8):1072–1077
- Siegrist CA, Aspinall R (2009) B-cell responses to vaccination at the extremes of age. *Nat Rev Immunol* 9(3):185–194
- Simanek AM, Dowd JB, Pawelec G, Melzer D, Dutta A, Aiello AE (2011) Seropositivity to cytomegalovirus, inflammation, all-cause and cardiovascular disease-related mortality in the United States. *PLoS One* 6(2):e16103
- Soderberg-Naucler C (2006) Does cytomegalovirus play a causative role in the development of various inflammatory diseases and cancer? *J Intern Med* 259(3):219–246
- Staras SA, Dollard SC, Radford KW, Flanders WD, Pass RF, Cannon MJ (2006) Seroprevalence of cytomegalovirus infection in the United States, 1988–1994. *Clin Infect Dis* 43(9):1143–1151
- Strandberg TE, Pitkala KH, Tilvis RS (2009) Cytomegalovirus antibody level and mortality among community-dwelling older adults with stable cardiovascular disease. *Jama* 301(4):380–382
- Sun Z, Zhong W, Lu X, Shi B, Zhu Y, Chen L, Zhang G, Zhang X (2008) Association of Graves' disease and prevalence of circulating IFN-gamma-producing CD28(−) T cells. *J Clin Immunol* 28(5):464–472
- Talbot HK, Coleman LA, Crimin K, Zhu Y, Rock MT, Meece J, Shay DK, Belongia EA, Griffin MR (2012) Association between obesity and vulnerability and serologic response to influenza vaccination in older adults. *Vaccine* 30(26):3937–3943
- Thompson WW, Shay DK, Weintraub E, Brammer L, Cox N, Anderson LJ, Fukuda K (2003) Mortality associated with influenza and respiratory syncytial virus in the United States. *Jama* 289(2):179–186
- Trzonkowski P, Mysliwska J, Szmít E, Wieckiewicz J, Lukaszuk K, Brydak LB, Machala M, Mysliwski A (2003) Association between cytomegalovirus infection, enhanced proinflammatory response and low level of anti-hemagglutinins during the anti-influenza vaccination—an impact of immunosenescence. *Vaccine* 21(25–26):3826–3836
- Turner JE, Aldred S, Witard O, Drayson MT, Moss PM, Bosch JA (2010) Latent Cytomegalovirus infection amplifies CD8 T-lymphocyte mobilisation and egress in response to exercise. *Brain Behav Immun* 24(8):1362–1370
- Vescovini R, Biasini C, Telera AR, Basaglia M, Stella A, Magalini F, Bucci L, Monti D, Lazzarotto T, Dal Monte P, Pedrazzoni M, Medici MC, Chezzi C, Franceschi C, Fagnoni FF, Sansoni P (2010) Intense antiextracellular adaptive immune response to human cytomegalovirus in very old subjects with impaired health and cognitive and functional status. *J Immunol* 184(6):3242–3249
- Visseren FL, Verkerk MS, Bouter KP, Diepersloot RJ, Erkelens DW (1999) Interleukin-6 production by endothelial cells after infection with influenza virus and cytomegalovirus. *J Lab Clin Med* 134(6):623–630
- Wikby A, Ferguson F, Forsey R, Thompson J, Strindhall J, Lofgren S, Nilsson BO, Emerudh J, Pawelec G, Johansson B (2005) An immune risk phenotype, cognitive impairment, and survival in very late life: impact of allostatic load in Swedish octogenarian and nonagenarian humans. *J Gerontol A Biol Sci Med Sci* 60(5):556–565
- Wikby A, Nilsson BO, Forsey R, Thompson J, Strindhall J, Lofgren S, Emerudh J, Pawelec G, Ferguson F, Johansson B (2006) The immune risk phenotype is associated with IL-6 in the terminal decline stage: findings from the Swedish Nona immune longitudinal study of very late life functioning. *Mech Ageing Dev* 127(8):695–704
- Wills M, Akbar A, Beswick M, Bosch JA, Caruso C, Colonna-Romano G, Dutta A, Franceschi C, Fulop T, Gkrania-Klotsas E, Goronzy J, Griffiths SJ, Henson S, Herndler-Brandstetter D, Hill A, Kern F, Klenerman P, Macallan D, Macalalay R, Maier AB, Mason G, Melzer D, Morgan M, Moss P, Nikolich-Zugich J, Pachnio A, Riddell N, Roberts R, Sansoni P, Sauce D, Sinclair J, Solana R, Strindhall J, Trzonkowski P, van Lier R, Vescovini R, Wang G, Westendorp R, Pawelec G (2011) Report from the Second Cytomegalovirus and Immunosenescence Workshop. *Immun Ageing* 8(1):10
- Zanni F, Vescovini R, Biasini C, Fagnoni F, Zanlari L, Telera A, Di Pede P, Passeri G, Pedrazzoni M, Passeri M, Franceschi C, Sansoni P (2003) Marked increase with age of type 1 cytokines within memory and effector/cytotoxic CD8+ T cells in humans: a contribution to understand the relationship between inflammation and immunosenescence. *Exp Gerontol* 38(9):981–987