Published in final edited form as:

Soc Sci Med. 2014 January ; 0: 62–71. doi:10.1016/j.socscimed.2013.10.037.

The risk of unintended pregnancy among young women with mental health symptoms

Kelli Stidham Hall, PhD MS [Research Investigator and “BIRCWH” NIH K-12 Scholar],
Department of Obstetrics and Gynecology, School of Medicine Institute for Social Research,
Population Studies Center, University of Michigan L4000 Women's Hospital, 1500 East Medical
Center Dr., Ann Arbor, MI 48109 (t) 734-930-5621 (c) 859-533-0762 (f) 734-930-5609

Yasamin Kusunoki, PhD MPH [Assistant Research Scientist],
Institute for Social Research, Population Studies and Survey Research Centers, University of
Michigan, 426 Thompson St. Ann Arbor, MI 48106-1248

Heather Gatny, MS [Senior Research Area Specialist], and
Institute for Social Research, Population Studies and Survey Research Centers, University of
Michigan, 426 Thompson St., Ann Arbor, MI 48106-1248

Jennifer Barber, PhD [Professor of Sociology]
Department of Sociology; Research Professor, Institute for Social Research Population Studies
and Survey Research Centers, University of Michigan, 426 Thompson St., Ann Arbor, MI
48106-1248

Kelli Stidham Hall: hkelli@umich.edu

Abstract

Depression and stress have been linked with poor contraceptive behavior, but whether existing mental health symptoms influence women's subsequent risk of unintended pregnancy is unclear. We prospectively examined the effect of depression and stress symptoms on young women's pregnancy risk over one year. We used panel data from a longitudinal study of 992 U.S. women ages 18-20 years who reported a strong desire to avoid pregnancy. Weekly journal surveys measured relationship, contraceptive use and pregnancy outcomes. We examined 27,572 journal surveys from 940 women over the first study year. Our outcome was self-reported pregnancy. At baseline, we assessed moderate/severe depression (CESD-5) and stress (PSS-4) symptoms. We estimated the effect of baseline mental health symptoms on pregnancy risk with discrete-time, mixed-effects, proportional hazard models using logistic regression. At baseline, 24% and 23% of women reported moderate/severe depression and stress symptoms, respectively. Ten percent of young women not intending pregnancy became pregnant during the study. Rates of pregnancy were higher among women with baseline depression (14% vs. 9%, $p=0.04$) and stress (15% vs. 9%, $p=0.03$) compared to women without symptoms. In multivariable models, the risk of pregnancy was 1.6 times higher among women with stress symptoms compared to those without stress (aRR 1.6, CI 1.1,2.7). Women with co-occurring stress and depression symptoms had over twice the risk of pregnancy (aRR 2.1, CI 1.1,3.8) compared to those without symptoms. Among women without a prior pregnancy, having co-occurring stress and depression symptoms was the

© 2013 Elsevier Ltd. All rights reserved.

Correspondence to: Kelli Stidham Hall, hkelli@umich.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

strongest predictor of subsequent pregnancy (aRR 2.3, CI 1.2,4.3), while stress alone was the strongest predictor among women with a prior pregnancy (aRR 3.0, CI 1.1,8.8). Depression symptoms were not independently associated with young women's pregnancy risk. In conclusion, stress, and especially co-occurring stress and depression symptoms, consistently and adversely influenced these young women's risk of unintended pregnancy over one year.

Keywords

United States; unintended pregnancy; mental health; depression; stress; adolescent women

Introduction

Unintended pregnancy among young women has been attributed to many diverse demographic, social, and cognitive behavioral factors. Young age, low educational attainment, low income, minority race/ethnicity, disrupted childhood family situation, having a mother who had a teenage pregnancy, earlier age at coitarche, more sexual partners, ambivalent pregnancy desire, lack of contraceptive knowledge, misperceptions of side effects, and low self-efficacy are just a few examples of the many factors shown to predict poor contraceptive behavior and unintended pregnancy among young women (Finer & Henshaw, 2006; Frost & Darroch, 2008; Rosenberg et al., 1995). While this research has isolated specific risk factors for unintended pregnancy, studies have often failed to account for the interrelatedness of different dimensions of young women's lives. Notably, the impact of mental health on young women's risk of adverse family planning outcomes has seldom been the focus of rigorous study (Chen et al., 1997; Farr et al., 2010; Hall, 2011).

Among the most common mental health problems worldwide, depression, anxiety, and related psychological stress disproportionately impact young and socially disadvantaged women (APA, 2000; NIMH, 2012; USDHHS, 2001). These conditions, which are highly comorbid, are a significant burden to society and contribute to negative biopsychosocial outcomes and reproductive sequelae, including maternal and infant morbidity and mortality (Alder et al., 2007; APA, 2000; Kelly et al., 2002; NIMH, 2012; USDHHS, 2001; Williams et al., 2007).

A considerable body of research has linked mental health symptomatology with risky *sexual* experiences. Depression and stress symptoms have been associated with having more sexual partners, earlier coitarche, higher sexually transmitted infections (STI) rates, sex while under the influence of alcohol and drugs, non-consensual sex and intimate partner violence (Brooks et al., 2002; Chen et al., 1997; Lehrer et al., 2006; Silverman et al., 2001).

More recent studies, using mostly clinic-based samples, have begun to identify associations between women's mental health status and risky contraceptive behaviors, including contraceptive nonuse, misuse, discontinuation, and less effective method use (Bennett et al., 2006; Farr, et al., 2010; Garbers et al., 2010; Ko et al., 2012; Lee et al., 2005; Hall et al., 2013a; Hall et al., 2013b; Hall et al., 2012; Zink et al., 2002). In a study of 354 young minority urban family planning patients, Hall and colleagues found an association between elevated baseline depression and stress symptoms and 6-month oral contraceptive discontinuation rates. Using clinical data from 2,476 urban, predominantly Black and Latina women, Garbers and colleagues found that women who screened positive for depression had 45% higher odds of selecting condoms and 39% lower odds of selecting hormonal methods at their clinic visit compared to women without depressive symptoms. Similarly, Farr and colleagues used cross-sectional national data from 53,255 women ages 18 and older in the Behavioral Risk Factor Surveillance System and found that low-income women with

frequent mental distress had a reduced likelihood of using long-acting/hormonal methods (OR 0.5) and condoms (OR 0.6) than other less effective methods.

Given that inadequate contraceptive use accounts for 90% of the estimated 3.2 million annual U.S. unintended pregnancies, contraceptive (and sexual risk) behavior is indeed a primary mediating factor of unintended pregnancy (Finer & Henshaw, 2006; Kost et al., 2008; Trussell & Vaughn, 1999). Thus, research describing a link between mental health symptoms and family planning behaviors has been an important scientific contribution, one that appears to have theoretical validity. Within a cognitive behavioral framework, mental health symptoms may interfere with a woman's cognitive capacities for decision-making, including risk assessment, planning, and social learning, as well as influence or distort their perceptions of benefits and threats of a preventive health therapy like contraception and their perceived susceptibility to an outcome like pregnancy (Hall, 2011; Maner, & Schmidt, 2006; Yuen & Lee, 2003). While this has not been directly tested to our knowledge, it is reasonable to hypothesize that symptoms of depression or stress, such as decreased motivation or distraction, could impair a woman's ability to use certain methods correctly (e.g. contraceptive pills) or to make suboptimal contraceptive choices (e.g. condom non-use or method discontinuation), for example.

A potentially more important question, though, is whether the influence of depression and stress symptoms on contraceptive use translates to an actual effect on women's risk of unintended pregnancy. The majority of studies focused on mental health and reproduction have described the prevalence, correlates, and treatment of perinatal and postpartum depression among women who have already experienced an unintended pregnancy (Alder et al., 2007; Dennis et al., 2007; Flynn et al., 2007; Grote et al., 2010; Vesga-Lopez et al., 2008). Other studies that have examined associations between mental health and unintended pregnancy (and related outcomes like abortion) have encountered temporality problems inherent in cross-sectional datasets and retrospective designs (Ko et al., 2012; Steinberg et al., 2012; Tenkku et al., 2009). One analysis of data from pregnant Japanese women in a birth cohort study found that women who had been diagnosed with depression were more likely to report that their current pregnancy was mistimed compared to those without depression; women with an anxiety diagnosis were more likely to report that their pregnancy as unwanted (Takahashi et al., 2012). Another study of 5,877 respondents aged 15–54 years in the National Comorbidity Survey found that men and women who reported having an early onset psychiatric disorder had a higher cumulative probability of also having experienced a teenage childbirth compared to those without a psychiatric history (Kessler et al., 1997). So, even some research that has attempted to account for temporal ordering of mental health diagnoses and pregnancy outcomes has been limited by historical data and retrospective reporting (Kessler et al., 1997; Takahashi et al., 2012).

Moreover, even less is known about unintended pregnancy and the influence of psychological stress, which is prevalent among young women and commonly co-occurs with depression and anxiety. As a distinct condition, stress has been defined as an emotional, psychological, behavioral or physiological response (and a subjective appraisal of this response) to situational demands (Larzelere & Jones, 2008; Selye, 1956). Stress is increasingly recognized as an important and independent contributor to morbidity (APA, 2013; Larzelere & Jones, 2008), including reproductive sequelae like preterm birth (Hogue et al., 2013). However, the distinct effect that stress may have on young women's risk of unintended pregnancy, as well as the effect of co-occurring stress and depression symptoms, has not been studied.

We sought to prospectively estimate the influence of baseline depression and stress symptoms on women's risk of unintended pregnancy over one year. We hypothesized that

the proportion of women who experienced a new pregnancy would be higher among women with elevated mental health symptoms at baseline than those without symptoms. We also hypothesized that after taking into account confounding effects of women's sociodemographic characteristics and reproductive histories, the risk of pregnancy would be higher among women with mental health symptoms than those without symptoms. Finally, we hypothesized that both depression and stress would exhibit similar but independent adverse effects on women's pregnancy risk.

Materials and Methods

Sample and Design

In other reports on young women's contraceptive behavior (Hall et al., 2013a; Hall et al., 2013b), we previously described the study sample and design. Briefly, we used data from a representative population-based cohort study of 992 young women aged 18-20 years residing in a racial/ethnically and socioeconomically diverse Michigan county. Names and contact information were randomly selected from state driver's license and personal identification card registries. Eligible women (ages 18-20 and a county resident) were contacted by mail or phone and asked to participate. Sampling occurred between March 2008 and March 2009. The Institutional Review Board of the University of Michigan approved this study.

After informed consent, a trained research assistant conducted a 60-minute in-person baseline survey interview to elicit information on sociodemographics, relationship characteristics, reproductive attitudes, beliefs and intentions, contraceptive histories, and mental health symptoms. Participants were then invited to participate in the weekly journal-based survey study for a period of 2.5 years. The weekly journal surveys (i.e. entries), which measured contraceptive use patterns, relationship status and pregnancy outcomes, were completed online for participants with Internet access or by phone if Internet access was unavailable on a particular week. Participants were compensated \$1 per weekly survey with \$5 bonuses for on-time completion of five consecutive surveys. Ninety-eight percent of women agreed to participate in the study. The response rate for the full baseline interview was 88%; 78% of women completed 12 months or more of weekly surveys.

Here, we focused on the first 12 months of data given the high response rates during this time and because our mental health symptoms measures were included only at baseline. Given that our unit of analysis was woman weeks, we included all women who were not pregnant at baseline and who completed more than one weekly journal survey (so that we had an assessment of mental health symptoms and then at least one subsequent assessment of pregnancy status). Our final sample included 940 women who completed 27,572 journal surveys (i.e. women-weeks) over the first study year.

Measures

Baseline Mental Health Symptoms—We administered standardized depression and stress instruments during the baseline survey interview. The Center for Epidemiologic Studies – Depression Scale-5 (CES-D-5) assesses depressive symptoms over the previous week (Radloff, 1977). Women were asked how often they felt the following five symptoms over the past 7 days: “like you could not shake off the blues,” “depressed,” “sad,” “life was not worth living,” and “happy.” Responses were scored on a 4-point scale (0=rarely or none of the time, 1=some or little of the time, 2=occasionally or moderate amount of time, 3=most or all of the time); the positively worded item was reverse coded. Items were then summed for a total depression score. Scores range from 0-15; a higher score indicates a higher degree of symptoms. We examined depression symptoms as a continuous score as

well as a binary variable using a standardized cut-off of 4 points to denote moderate/severe depression symptoms. We also conducted sensitivity tests of different score cut-off points for moderate/severe depression symptoms. Overall, findings in relation to pregnancy were similar, so we present results from the binary 4-point moderate/severe depression symptom variable (which captured one standard deviation above our population mean). The CES-D has demonstrated good sensitivity, specificity, and high validity in identifying patients classified as depressed by the full 20-item scale (Bohannon et al., 2003). The abbreviated CES-D has also demonstrated strong content, concurrent, and discriminant validity and internal consistency (Radloff).

The Perceived Stress Scale (PSS-4) assesses the degree to which one appraises their life situations as stressful, unpredictable, uncontrollable, and overloading over the previous month (Cohen, 1983). Women were asked how often (0=never, 1=almost never, 2=sometimes, 3=fairly often, 4=very often) they felt the following four symptoms: “unable to control important things in life,” “confident about ability to handle personal problems,” “things were going your way,” and “difficulties were piling up so high that you could not overcome them.” Positively worded items were reverse coded. Items were summed for a total score, which could range from 0-16 with higher scores indicating greater stress. Like the depression measure, we examined continuous PSS scores as well as different cut-off points for binary moderate/severe stress symptoms. We present results from the binary stress symptom variable in which we used the standardized cut-off of 9 points (one standard deviation above population mean) to denote moderate/severe stress symptoms. The PSS has been correlated with depression, anxiety and physical symptomatology measures in clinical populations but has been shown to measure a distinct, independent construct of appraised stress (Cohen, 1983). High construct validity, internal consistency, sensitivity and specificity have been previously demonstrated (Cohen, 1983).

First, we used the CES-D and PSS to examine depression and stress symptoms separately (but not exclusively). We also created a 4-point categorical variable to examine the independent and combined effects of having each or both mental health symptoms with the following categories: no mental health symptoms, depression symptoms only, stress symptoms only, or co-occurring depression and stress symptoms.

Unintended Pregnancy—In each weekly journal, women were asked if it was possible that she was pregnant and if so, whether a pregnancy test had indicated a positive pregnancy. We operationalized a pregnancy as a newly reported positive pregnancy test during the 12-month study period. We refer to pregnancy among these women during the first study year as “unintended” in our discussion given that this sample is comprised of adolescent women (a demographic for which pregnancies are overwhelmingly unintended), 98% of whom explicitly stated at baseline that they had no intentions and, in fact, strong desires to avoid pregnancy (reported a 4 or 5 on an avoidance scale of 1-5).

Sociodemographic, Relationship, and Reproductive Characteristics—We examined the following sociodemographic, relationship and reproductive history characteristics at baseline and each week: age, race/ethnicity (non-Hispanic White, non-Hispanic Black, Hispanic, or other), educational attainment (not enrolled in school, still in high school, enrolled in 2 year program, enrolled in 4 year college, or dropped out of high school), employment status (employed or unemployed), parental income level (<\$15,000, \$15,000-44,999, \$45,000-74,999, or \$75,000), public assistance recipient (including WIC, FIP, cash welfare or food stamps, yes or no), childhood household/family structure (lived with 2 biological/step parents, 1 parent only, or other), mother's education level (graduated high school or less than high school education), mother's age at first birth (less than 20 years old or 20 years old), frequency of religious service attendance (never, <weekly, or

weekly), relationship status (married, engaged, in special romantic relationship, having physical/emotional contact with someone or none), cohabitation with marital or non-marital partner (yes or no), sexual intercourse experience (yes or no), age at coitarche, lifetime number of sexual partners, histories of pregnancy, birth control use, unprotected sex, and current birth control use (all yes or no).

Statistical Analysis—We used univariate statistics to describe women's baseline sociodemographic and reproductive characteristics and to estimate the proportions (%) of women with baseline moderate/severe depression and stress symptoms. We also described the proportion of women who experienced a new pregnancy during the one-year study period (i.e. pregnancy rate). We conducted unadjusted bivariate X^2 tests to compare sociodemographic and reproductive characteristics among: 1) women with moderate/severe depression and stress symptoms versus those without depression or stress symptoms, 2) women with a new pregnancy versus those without, and 3) women with a new pregnancy, stratified by history of pregnancy (without or with a prior pregnancy).

We further examined the effect of baseline mental health symptoms on women's risk of pregnancy over one year while adjusting for sociodemographic and reproductive confounders. We used event history methods to model the risk of pregnancy each week and discrete-time methods to estimate the proportional hazard models. Person-weeks of exposure are the unit of analysis. We considered women to be at risk of pregnancy during all weeks that she reported not being currently pregnant. Our mixed effects multivariable logistic regression models included covariate fixed effects and a random effect for the woman, while also controlling for the number of weekly journals completed, number of pregnancy months, and number of pregnancy months squared as indicators of time (i.e. effects of repeated exposure to the questionnaire, continued participation in the study and being pregnant). We first estimated the effect of mental health symptoms on pregnancy risk in univariate models, then in multivariable models (adding sociodemographic first, followed by reproductive history variables), and then in final reduced models controlling only for significant covariates. We also stratified models by history of prior pregnancy. We examined depression and stress symptoms as independent variables in models first and then using our combined 4-point categorical mental health symptom variable.

Variables were considered for inclusion in regression models if their p-value in bivariate models was 0.25 or less. For variables that appeared to be collinear (e.g. reproductive and sexual history variables), we retained variables with the strongest associations with our outcome in final models. While we also examined time-variant sociodemographic characteristics, their effects on pregnancy risk were similar to baseline characteristics, so we present baseline models given that mental health symptoms were measured at baseline. Finally, we repeated all analyses using the first 18 months of data in order to estimate the effect of mental health on pregnancy risk with a larger sample of weeks at risk of pregnancy. Results were similar, so we present the more conservative one-year estimates.

We present exponentiated coefficients from regression models as adjusted relative risks (aRR) and corresponding 95% confidence intervals (CI). Two-tailed alphas of $p < 0.05^*$, $p < 0.01^{**}$, and $p < 0.001^{***}$ were considered significant. All data were analyzed using Stata 12.0 (StataCorp LP, College Station, TX).

Results

Baseline Mental Health Symptoms

Women's sociodemographic and reproductive history background characteristics are presented in Table 1. For baseline mental health symptoms, 24% of women met criteria for

moderate/severe depression symptoms (4pts CESD) and 23% met criteria for moderate/severe stress symptoms (9pts PSS-4); 15% of women had co-occurring depression and stress symptoms. Among women with moderate/severe depression symptoms, 65% also reported stress symptoms; among those with stress, 61% also reported depression symptoms (p 's<0.001, Pearson's $r=0.51$).

Having moderate/severe depression symptoms was significantly associated with nearly all sociodemographic and reproductive background characteristics in unadjusted bivariate tests (Table 1). Proportions of depression symptoms were higher among Black women, women who dropped out of high school, women with a low parental income, women receiving public assistance, women without two parents in their childhood household, women with a mother who had less than a high school education and who had given birth as a teen, and women with an early age at coitarche, higher numbers of sexual partners, and histories of pregnancy, contraceptive use and unprotected sex. Proportions of moderate/severe stress symptoms were higher among women who were not enrolled in school or had dropped out of high school, women without two parents in their childhood household, and women with an early coitarche, higher numbers of sexual partners, and histories of pregnancy, contraceptive use and unprotected sex.

Pregnancy Rates Over One Year

During the first year of study, 98 women (10%) reported a new pregnancy; 93 women experienced 1 pregnancy and five women experienced 2 pregnancies. Rates of pregnancy varied by nearly all women's background characteristics (Table 1). Pregnancy rates were higher among Black women, women who were still in high school or not enrolled in school, unemployed women, women with low parental incomes and those receiving public assistance, women without two parents in their childhood household, women with a mother who had given birth as a teen, women who were engaged or cohabiting, and women with an early age at coitarche, higher numbers of sexual partners, and histories of pregnancy, contraceptive use and unprotected sex. These associations were similar among women with no history of prior pregnancy ($n=734$, 78%). Among women who had previously been pregnant ($n=206$, 22%), only cohabitation status was significantly associated with experiencing a new pregnancy during the study ($p<0.001$).

Associations Between Baseline Mental Health Symptoms and Pregnancy Rates

In the unadjusted bivariate analysis (Table 1), rates of pregnancy over one year were higher among women with moderate/severe depression (14% versus 9%, $p=0.03$) and stress (15% versus 9%, $p=0.03$) symptoms than those without symptoms. This finding was also true of women without a pregnancy history, for both depression (13% versus 6%, $p=0.005$) and stress (11% versus 6%, $p=0.04$). Among women with a prior pregnancy, however, pregnancy rates were similar by depression ($p=0.60$) and stress ($p=0.54$) symptoms.

Results from multivariable logistic regression models are presented in Tables 2-4. Among all women, depression symptoms did not independently predict women's risk of pregnancy over one year (aRR 1.2, $p=0.52$). Women with moderate/severe stress symptoms, however, had a 1.6 times higher risk of pregnancy (aRR 1.6, $p=0.04$) than those without stress symptoms. The risk of pregnancy was highest among women who had co-occurring stress and depression symptoms (aRR 2.1, $p=0.03$) and not significant among the few women who had stress symptoms alone (aRR 1.3, $p=0.61$) or depression symptoms alone (aRR 0.7, $p=0.48$), compared to women without any mental health symptoms.

We further examined the effect of mental health symptoms on pregnancy risk in models stratified by pregnancy history (Table 4). For women without prior pregnancy, the combined

effect of having both stress and depression symptoms was the strongest predictor of pregnancy risk (aRR 2.3, $p=0.01$) compared to those with no symptoms. Among women with a prior pregnancy, the effect of having stress alone was the strongest predictor of pregnancy risk (aRR 3.0, $p=0.04$).

Other sociodemographic and reproductive characteristics associated with women's one year pregnancy risk included receipt of public assistance, having a mother who had given birth as a teen, cohabiting with a partner, and experiencing an early age at coitarche (Tables 2 & 3). In models stratified by pregnancy history (Tables 4), having a teen mother and an early age at coitarche were most strongly associated with pregnancy risk among women without a prior pregnancy, while cohabiting with a partner was most strongly associated among women with a prior pregnancy.

Discussion

In our previous analyses of young women's contraceptive behavior using these same data, we found that young women with moderate/severe baseline depression and stress symptoms were at increased risk of a wide range of risky contraceptive behaviors each week compared to women without depression and stress symptoms (Hall et al, 2013a; Hall et al, 2013b). More specifically, 10% of weeks during the first study year were not covered by contraceptive use and women with moderate/severe stress symptom had over twice the odds of not using contraception each week (including any family planning methods, ranging from rhythm and withdrawal to intrauterine devices and implants) than women without stress symptoms. The odds of consistent contraceptive use were 47% and 69% lower among women with depression and stress than those without symptoms, respectively. Finally, the relative risks of using long-acting methods (rather than oral contraceptives) were 48% and 25% lower among women with depression and stress symptoms than those without symptoms. In addition to contraceptive use, we are also currently examining the role of sexual behavior, including frequency of sexual activity and sexual partnerships, as additional mechanisms that may link mental health symptoms to pregnancy risk. While young women, including those wishing to avoid pregnancy, may have a variety of reasons for having more or less sex or for not using contraception (and indeed contraceptive nonuse is not uncommon (Frost & Darroch, 2008; Kost et al, 2008; Mosher & Jones, 2010; Trussell & Vaughn, 1999)), our collective research suggests that mental health may be one factor which shapes women's family planning behaviors and outcomes. Among our adolescent women wishing to avoid pregnancy, those with moderate/severe mental health symptoms were at increased risk of behaviors that can lead to unintended pregnancy and for unintended pregnancy itself. Furthermore, the effect of stress on young women's pregnancy risk appears to be more consistently present and stronger than that of depression symptoms.

Here, we build upon these findings to show the impact of stress on women's risk of unintended pregnancy over one year. Women with baseline stress symptoms had a 1.6 times higher risk of becoming pregnant over time than women without stress. Moreover, women who had co-occurring stress *and* depression symptoms had twice the risk of pregnancy than women without symptoms. In stratified models, in which we attempted to control for the confounding effects of a previous pregnancy (and potentially related post-pregnancy mental health status) and sociodemographic factors (e.g. Black race/ethnicity, receipt of public assistance, and cohabitation), stress was consistently associated with risk of pregnancy among women without and with a pregnancy history. Additionally, having concurrent stress and depression symptoms was even more strongly related to pregnancy risk among women who had never been pregnant.

Collectively, our findings extend the research on mental and reproductive health and help clarify the temporal order effect of mental health symptoms, not only risky contraceptive behavior but also on actual unintended pregnancy risk, which is arguably the more meaningful outcome. A major contribution of this study is our robust, longitudinal design with weekly measures of pregnancy-related behaviors and outcomes, which we believe advances the methods required to adequately examine complex relationships between mental and reproductive health. This work also illuminates areas for further research to understand the dynamic, causal relationships between the two. Specifically, findings point to the adverse role of stress, and especially co-occurring stress and depression symptoms, in influencing young women's family planning behaviors and outcomes. Additional research is warranted to further disentangle interrelated mental and reproductive health experiences during adolescence and young adulthood (Hall, 2011; Maner, & Schmidt, 2006; Yuen & Lee, 2003).

The independent effect that depression may have on young women's unintended pregnancy risk remains unclear, with statistically insignificant and variable point estimates noted across our analytical models. It is plausible that for some women, especially those with a history of unintended pregnancy, experiencing depression may be a protective factor against subsequent pregnancy (which would be supported by a trend noted in our stratified models, with a 0.5 RR for depression alone among previously pregnant women). Women who have experienced an unintended pregnancy and/or postpartum mental health conditions may exhibit adaptive coping strategies to avoid another pregnancy. This "active behavioral coping" hypothesis has been suggested by Steinberg and colleagues (2013), who conducted a study of contraceptive method selection among post-abortion patients and found that women with mental distress were more likely to choose more effective methods following their unintended pregnancy than those without distress. Effective contraceptive use may reflect a motivation to prevent rapid repeat pregnancy among depressed women. Unfortunately, small subsample sizes in our stratified regression models resulted in reduced statistical power and limited our ability to adequately analyze these relationships.

Additionally, we were unable to determine the onset of the mental health symptoms, which we measured only at baseline, and whether they may have been attributable to women's prior pregnancy and postpartum status. Finally, as our own previous research and that of others has shown, stress and depression symptoms (just as with depression, anxiety and other mental health symptomatology) are all highly correlated, which was indeed the case in this sample. While we attempted to measure distinct constructs using validated scales for depression and stress, and while our findings appear to point to subtle differences in the effects of experiencing "sadness" or "depressed mood" versus "feeling out of control" or "burdened by life situations" on pregnancy risk, the independent (and interrelated) effects of depression and stress on family planning behaviors and outcomes are not fully clear and require further investigation. Nonetheless, this work is the first of which we are aware to prospectively examine the relationships between mental health symptoms, including co-occurring depression and stress symptoms, and subsequent unintended pregnancy among young women, especially by pregnancy history.

Another limitation of our study is the potential for underreporting of unintended pregnancies, particularly those that may have ended in abortion. We had a relatively wide range of the number of journal surveys completed per woman and it is possible that women did not complete surveys in weeks that they experienced a pregnancy or abortion. Such biased reporting could have underestimated our pregnancy rate and undermined our ability to detect associations between mental health symptoms and pregnancy risk. Indeed, self-report bias with unintended pregnancy and abortion is a documented problem in population-based survey research (Finer & Henshaw, 2006) and although we believe our weekly

assessments of pregnancy are more rigorous than most survey studies are able to achieve, our design was not without limitations. Other previous research has described an effect of preexisting mental health morbidity on women's risk of having an abortion (Steinberg & Finer, 2011). Given that 90% of abortions are the result of unintended pregnancies and women who have abortions are at risk for subsequent unintended pregnancies (Jones et al., 2006; Upadhyay et al., 2012), this may also have led to an underestimate of unintended pregnancy, overall and especially among the women with underlying depression and stress. While pregnancy outcomes are not the focus of this analysis, our results warrant further study and suggest a more rigorous measurement of mental health status and family planning outcomes in population-based samples is needed.

Additionally, we were unable to assess potential time-varying effects of depression and stress on pregnancy risk over time since our depression and stress measures were administered to all women only at baseline and not each week. It is unclear how changing mental health symptoms may impact changing pregnancy risk (as well as pregnancy intentions) over time. Other researchers have suggested that mental health symptoms are relatively across adolescence and young adulthood and over short periods (e.g. 5-6 years) (Holsen et al., 2000; Prenoveau et al., 2011), but unfortunately we were unable to assess whether this was true of our sample and whether and how dynamic mental health symptoms impact pregnancy risk over time. Finally, our sample, which was young (18-20 years at enrollment) and which was reflective of the demographic composition of a diverse county in Michigan, was not representative of the greater U.S. population, which limits generalizability.

Despite these limitations, our work has highlighted the adverse impact of mental health symptoms, especially stress and co-occurring mental health symptoms, on young women's risk of experiencing an unintended pregnancy over one year. It has also provided preliminary insight into a potential differential effect that depression and stress may have on subsequent pregnancy risk among women who have and have not had a previous pregnancy. Additional research is needed to disentangle the effects of depression and stress symptoms on women's pregnancy risk, especially among women with different reproductive backgrounds and pregnancy-related mental health experiences. Our ongoing and future work aims to further investigate psychological determinants of women's family planning outcomes, including birth, abortion and postpartum mental health, as well as the behavioral and biological mechanisms that may link mental health with unintended pregnancy. We are also studying the role of adverse social circumstances, which we believe are intrinsically tied to psychological and physical health outcomes and which may further help explain the relationships between mental and reproductive health among young women.

Acknowledgments

Funding source: This work was supported by an NIH NICHD Building Interdisciplinary Careers in Women's Health K-12 Career Development Award #K12HD001438 (Hall), by NICHD grant #R01-HDHD050329 (Barber), by NICHD grant #R24HD041028 (Smock).

References

- Alder J, Fink N, Bitzer J, Hosli I, Holzgreve W. Depression and anxiety during pregnancy: A risk factor the obstetric, fetal and neonatal outcome? A critical review of the literature. *Journal of Maternal-Fetal Neonatal Medicine*. 2007; 20:189–209. [PubMed: 17437220]
- American Psychiatric Association (APA). *Diagnostic and Statistical Manual of Mental Disorders*. 4th, Text Revised. Washington, DC: Author; 2000.
- American Psychiatric Association (APA). *Stress in America*. 2013. Retrieved May 12, 2013 from: <http://www.apa.org/news/press/releases/stress/index.aspx>

- Bennett IM, Culhane JF, McCollum KF, Elo IT. Unintended rapid repeat pregnancy and low education status: Any role for depression and contraceptive use? *Obstetrics & Gynecology*. 2006; 194:749–54.
- Bohannon RW, Maljanian R, Goethe J. Screening for depression in clinical practice: reliability and validity of a five-item subset of the CES-Depression. *Perceptual & Motivational Skills*. 2003; 97:855–61.
- Brooks TL, Harris SK, Thrall JS, Woods ER. Association of adolescent risk behaviors with mental health symptoms in high school students. *Journal of Adolescent Health*. 2002; 31:240–46. [PubMed: 12225736]
- Chen YW, Stiffman AR, Cheng LC, Dore P. Mental health, social environment and sexual risk behaviors of adolescent service users: a gender comparison. *Journal of Child & Family Studies*. 1997; 6:9–25.
- Cohen S, Kamarck T, Mermelstein R. A global measure of perceived stress. *Journal of Health & Social Behavior*. 1983; 24:385–96. [PubMed: 6668417]
- Dennis, CL.; Ross, LE.; Grigoriadis, S. Cochrane Database of Systematic Reviews. 2007. Psychosocial and psychological interventions for treating antenatal depression. Art. No.: CD006309
- Farr SL, Bitsko RH, Hayes DK, Dietz PM. Mental health and access to services among US women of reproductive aged. *American Journal of Obstetrics & Gynecology*. 2010; 203:e1–9. [PubMed: 20817143]
- Flynn HA, Walton MA, Chermack ST, Cunningham RM, Marcus SM. Brief detection and co-occurrence of violence, depression and alcohol risk in prenatal care settings. *Archives Women's Mental Health*. 2007; 10:155–161.
- Finer LB, Henshaw SK. Disparities in rates of unintended pregnancy in the United States, 1994 and 2001. *Perspectives in Sexual & Reproductive Health*. 2006; 38:90–6.
- Frost JJ, Darroch JE. Factors associated with contraceptive choice and inconsistent method use, United States, 2004. *Perspectives in Sexual & Reproductive Health*. 2008; 40:94–104.
- Garbers S, Correa N, Tobier N, Blust S, Chiasson MA. Associations between symptoms of depression and contraceptive method choices among low-income women at urban reproductive health centers. *Maternal Child Health*. 2010; 14:102–109.
- Grote NK, Bridge JA, Gavin AR, Melville JL, Iyengar S, Katon WJ. A meta-analysis of depression during pregnancy and the risk of pre-term birth, low birthweight and intrauterine growth restriction. *Archives of General Psychiatry*. 2010; 67:1012–1024. [PubMed: 20921117]
- Hall K. The Health Belief Model can guide modern contraceptive research. *Journal Midwifery & Women's Health*. 2011 Jan-Feb;57:74–81.
- Hall K, Moreau C, Trussell J, Barber J. Role of young women's depression and stress symptoms in their weekly use and nonuse of contraceptive methods. *Journal of Adolescent Health*. 2013; 53(2): 241–248. [PubMed: 23582524]
- Hall, K.; Moreau, C.; Trussell, J.; Barber, J. Contraception. 2013. Young women's consistency of contraceptive use – do depression and stress matter. Published online ahead of print 2013 Jun 12
- Hall K, Reame N, O'Connell K, Rickert V, Weshoff C. Influence of depressed mood and psychological stress symptoms on perceived oral contraceptive side effects and discontinuation in young minority women. *Contraception*. 2012; 86:518–25. [PubMed: 22673038]
- Holsen I, Kraft P, Vittersø J. Stability in depressed mood in adolescence: Results from a 6-year longitudinal panel study. *Journal of Youth Adolescence*. 2000; 29:61–78.
- Hogue CJR, Parker CB, Willinger M, et al. A Population-based Case-Control Study of Stillbirth: The Relationship of Significant Life Events to the Racial Disparity for African Americans. *American Journal of Epidemiology*. 2013 Published online ahead of print. 10.1093/aje/kws381
- Jones, RK.; Singh, S.; Finer, LB.; Frohwirth, LF. Guttmacher Institute Occasional Report. Vol. 29. New York, NY: Guttmacher Institute; 2006. Repeat abortion in the United States.
- Kelly RH, Russo J, Holt V, et al. Psychiatric and Substance Use Disorders as Risk Factors for Low Birth Weight and Preterm Delivery. *Obstetrics & Gynecology*. 2002; 100:297–304. [PubMed: 12151153]

- Kessler RC, Berglund PA, Foster CL, Saunders WB, Stang PE, Walters EE. Social consequences of psychiatric disorders, II: Teenage parenthood. *American Journal of Psychiatry*. 1997; 154(10): 1405–1411. [PubMed: 9326823]
- Ko JY, Farr SL, Dietz PM, Robbins CL. Depression and treatment among US pregnant and non-pregnant women of reproductive age, 2005–2009. *Journal of Women's Health*. 2012; 21:830–6.
- Kost K, Singh S, Vaughan B, Trussell J, Bankole A. Estimates of contraceptive failure from the 2002 National Survey of Family Growth. *Contraception*. 2008; 77:10–21. [PubMed: 18082661]
- Larzelere MM, Jones GN. Stress and health. *Primary Care*. 2008; 35:839–856. [PubMed: 18928833]
- Lee, Li-Ching; Casanueva, CE.; Martin, SL. Depression among female family planning patients: prevalence, risk factors and use of mental health services. *Journal of Women's Health*. 2005; 14:225–232.
- Lehrer JA, Shrier LA, Gortmaker S, Buka S. Depressive symptoms as longitudinal predictors of sexual risk behaviors among US middle and high school students. *Pediatrics*. 2006; 118:189–200. [PubMed: 16818565]
- Maner JK, Schmidt NB. The role of risk avoidance in anxiety. *Behavior Therapy*. 2006; 37(2):181–189. [PubMed: 16942970]
- Mosher WD, Jones J. Use of contraception in the United States: 1982–2008. *Vital Health Statistics*. 2010; 23:1–43. [PubMed: 20928970]
- National Institute of Mental Health (NIMH). Major depressive disorder among adults. Retrieved October 24, 2012 from: http://www.nimh.nih.gov/statistics/1MDD_ADULT.shtml
- Prenoveau JM, Craske MG, Zinbarg RE, Mineka S, Rose RD, Griffith JW. Are anxiety and depression just as stable as personality during late adolescence? Results from a three-year longitudinal latent variable study. *Journal of Abnormal Psychology*. 2011; 120:832–43. [PubMed: 21604827]
- Radloff LS. The CES-D scale: A self-report depression scale for research in the general population. *Applied Psychological Measures*. 1977; 1:385–401.
- Rosenberg MJ, Waugh MS, Long S. Unintended pregnancies and use, misuse and discontinuation of oral contraceptives. *Journal of Reproductive Medicine*. 1995; 40:355–60. [PubMed: 7608875]
- Selye, H. *The stress of life*. New York: McGraw-Hill; 1956.
- Silverman JG, Raj A, Mucci LA, Hathaway JE. Dating Violence Against Adolescent Girls and Associated Substance Use, Unhealthy Weight Control, Sexual Risk Behavior, Pregnancy, and Suicidality. *JAMA*. 2001; 286:572–579. [PubMed: 11476659]
- Steinberg JR, Finer LB. Examining the association of abortion history and current mental health: A reanalysis of the National Comorbidity Survey using a common-risk-factors model. *Social Science & Medicine*. 2011; 72:72–82. [PubMed: 21122964]
- Steinberg, JR.; Tschann, JM.; Henderson, JT.; Drey, EA.; Steinauer, JE.; Harper, CC. *Contraception*. 2013. Psychological distress and post-abortion contraceptive choice at an urban clinic. PMC In Press
- Steinberg JR, Trussell J, Hall KS, Guthrie K. Fatal flaws in a recent meta-analysis on abortion and mental health. *Contraception*. 2012; 86(5):430–7. [PubMed: 22579105]
- Takahashi S, Tsuchiya KJ, Matsumoto K, Suzuki K, Mori N, Takei N. Psychosocial determinants of mistimed and unwanted pregnancy: The Hamamatsu Birth Cohort Study. *Journal of Maternal and Child Health*. 2012; 16(5):947–955.
- Tenkku LE, Flick LH, Homan S, Loveland Cook CA, Campbell C, McSweeney M. Psychiatric disorders among low-income women and unintended pregnancies. *Women's Health Issues*. 2009; 19(5):313–324. [PubMed: 19733801]
- Trussell J, Vaughn B. Contraceptive failure, method-related discontinuation and resumption of use: results from the 1995 NSFG. *Family Planning Perspectives*. 1999; 31(93):64–72. [PubMed: 10224544]
- Upadhyay UD, Brown BA, Sokoloff A, Raine TR. Contraceptive discontinuation and repeat unintended pregnancy within 1 year after an abortion. *Contraception*. 2012; 85(1):56–62. [PubMed: 22067792]
- U.S. Department of Health and Human Services (USDHHS). *A Report of the Surgeon General*. Rockville, Md: US Dept of Health and Human Services, Public Health Service, Office of the

Surgeon General; 2001. Mental Health: Culture, Race, and Ethnicity: A Supplement to Mental Health.

Vesga-Lopez O, Blanco C, Keyes K, Olfson M, Grant BF, Hasin DS. Psychiatric disorders in pregnant and postpartum women in the United States. *Archives of General Psychiatry*. 2008; 65:805–815. [PubMed: 18606953]

Williams KE, Marsh WK, Rasgon NL. Mood disorders and fertility in women: a critical review of the literature and implications for future research. *Human Reproduction Update*. 2007; 13:607–616. [PubMed: 17895237]

Yuen KSL, Lee TMC. Could mood state affect risk-taking decisions? *Journal of Affective Disorders*. 2003; 75(1):11–18. [PubMed: 12781345]

Zink TM, Shireman TI, Ho M, Buchanan T. High-risk teen compliance with prescription contraception: An analysis of Ohio Medicaid Claims. *Journal of Pediatric & Adolescent Gynecology*. 2002; 15:15–21. [PubMed: 11888805]

Research highlights

- Women with depression and stress symptoms had high one-year unintended pregnancy rates
- Stress was consistently, adversely associated with an increased risk of pregnancy over one year
- Combined effect of co-occurring stress and depression symptoms most strongly predicted pregnancy
- The independent effect of depression on subsequent unintended pregnancy remains unclear

Table 1

Baseline Mental Health Symptoms					Pregnancy Rates at One Year, Overall and by Pregnancy History				
Baseline characteristics	Depression 4pts CESD ^a n=224 24%		Stress 9pts PSS-n=213 23%		All women n=98 10%	Women without a prior pregnancy n=734 78%		Women with a prior pregnancy n=206 22%	
	%	%	p ¹	%		p ²	%		p ³
Age			0.17		0.05	0.12		0.13	0.57
18 years	41	23		22			9		24
19 years	50	26		25			7		22
20 years	9	16		12			2		12
Race/ethnicity			0.002		0.37	0.008		0.02	0.61
Non-Hispanic White	59	20		22			6		24
Non-Hispanic Black	31	31		25			12		21
Hispanic	8	31		22			7		16
Asian/Other	2	25		10			0		0
Educational enrollment			<0.001		0.004	0.005		0.01	0.61
Not enrolled	22	29		30			11		26
High school	14	23		20			13		22
2 year college	29	24		22			7		17
4 year college	28	16		16			3		28
High school drop-out	8	40		30			6		16
Employment status			0.19		0.84	0.02		0.008	0.66
Employed	50	22		22			5		23
Unemployed	50	26		23			10		20
Parental income			<0.001		0.06	<0.001		0.004	0.37
<\$15,000	14	34		27			16		27
\$15,000-44,999	28	25		23			8		14
\$45,000-74,999	19	23		26			4		17
\$75,000	19	13		15			4		29

Soc Sci Med. Author manuscript; available in PMC 2015 January 01.

Baseline Mental Health Symptoms										Pregnancy Rates at One Year, Overall and by Pregnancy History							
Depression 4pts CESD ^d n=224 24%		Stress 9pts PSS-n=213 23%		All women n=98 10%		Women without a prior pregnancy n=734 78%		Women with a prior pregnancy n=206 22%									
%	p ¹	%	p ²	%	p ³	%	p ⁴	%	p ⁴								
83	24	23	<0.001	8	<0.001	6	<0.001	15	0.47								
Not cohabitating																	
Age at first vaginal intercourse																	
24	15	15															
16	38	35		22		18		27									
35	29	24		15		13		19									
25	17	20		6		4		19									
Lifetime # sexual partners																	
24	15	15	<0.001		<0.001		<0.001		0.46								
17	16	16		11		6		30									
13	23	19		11		7		21									
46	31	31		16		14		20									
History of pregnancy																	
22	31	28	0.005	21													
78	22	21		7					0.53								
Ever used birth control																	
75	26	25	0.003	13	<0.001	10		22									
25	17	16		3		1		16									
Ever had unprotected sex																	
42	32	30	<0.001	17	0.006	14	0.03	22	0.98								
28	19	20		10		8		22									
30																	
Currently using birth control																	
50	24	24	0.24		0.07	8	0.21	19	0.23								
20	29	26		16		12		27									
30									0.60								
Moderate/severe depression ^d																	

Baseline characteristics	Baseline Mental Health Symptoms				Pregnancy Rates at One Year, Overall and by Pregnancy History			
	n=940 100%	Depression 4pts CESD ^a n=224 24%	Stress 9pts PSS n=213 23%	All women n=98 10%	Women without a prior pregnancy n=734 78%		Women with a prior pregnancy n=206 22%	
		%	%		%	%	p ⁴	%
Yes (4pts CESD)	24		61	14	13		28	
No (<4pts CESD)	76		11	9	6		32	
Moderate/severe stress ^b			<0.001			0.03		0.54
Yes (9pts PSS-4)	23	64		15	11		32	
No (<9pts PSS-4)	77	12		9	6		27	
Mental health symptoms						0.05		0.68
None	67			8	6		21	
Depression only	10			10	9		14	
Stress only	8			8	5		31	
Both depression and stress	15			23	15		21	

¹ Results are proportions (%) of all women in sample, women with moderate/severe depression and stress symptoms, and women who experienced a new pregnancy during the first year of study overall and by pregnancy history across sociodemographic and reproductive characteristics. P-values are from unadjusted X² analysis comparing sociodemographic and reproductive characteristics among women with and without moderate/severe depression symptoms,

² among women with and without moderate/severe stress symptoms,

³ among women who did and did not experience a new pregnancy during the study period,

⁴ among women who did and did not experience a new pregnancy during the study period stratified by history of previous pregnancy.

^a Center for Epidemiologic Studies – Depression Scale (CES-D-5) – 4 point cut-off for moderate/severe depression symptoms.

^b Perceived Stress Scale - 4 (PSS-4) - 9-point cut-off for moderate/severe stress symptoms. Analyses were also conducted to examine associations between mental health symptoms and pregnancy outcomes over an 18-month study period (n=176 pregnancies) and point estimates were similar to those from the one-year data (not shown).

Table 2
The effect of baseline depression symptoms on young women's risk of pregnancy over one year

N=940	Model 1 (Univariate)				Model 2 (Sociodemographics)				Model 3 (Full)				Model 4 (Final reduced)			
	RR	CI	aRR	CI	aRR	CI	aRR	CI	aRR	CI	aRR	CI	aRR	CI	aRR	CI
Depression symptoms ^a																
No (<4pts CESD)	1		1		1		1		1		1		1		1	
Yes (4pts CESD)	1.6	1.0,2.7	1.2	0.7,2.0	1.2	0.7,2.0	1.1	0.6,1.7	1.2	0.7,1.9	1.2	0.7,1.9	1.2	0.7,1.9	1.2	0.7,1.9
Age																
18 years			1		1		1		1		1		1		1	
19 years			1.0	0.6,1.6	1.0	0.6,1.6	1.0	0.6,1.6	1.0	0.6,1.6	1.0	0.6,1.6	1.0	0.6,1.6	1.0	0.6,1.6
20 years			0.5	0.1,1.4	0.4	0.1,1.3	0.4	0.1,1.3	0.4	0.1,1.3	0.4	0.1,1.3	0.4	0.1,1.3	0.4	0.1,1.3
Race/ethnicity																
Non-Black			1		1		1		1		1		1		1	
Black			1.5	0.8,2.6	1.2	0.7,2.0	1.2	0.7,2.0	1.2	0.7,2.0	1.2	0.7,2.0	1.2	0.7,2.0	1.2	0.7,2.0
Educational enrollment																
Not enrolled			1		1		1		1		1		1		1	
High school			1.0	0.5,2.0	1.1	0.6,2.2	1.1	0.6,2.2	1.1	0.6,2.2	1.1	0.6,2.2	1.1	0.6,2.2	1.1	0.6,2.2
2 year college			0.7	0.4,1.3	0.7	0.4,1.3	0.7	0.4,1.3	0.7	0.4,1.3	0.7	0.4,1.3	0.7	0.4,1.3	0.7	0.4,1.3
4 year college			0.5	0.3,1.1	0.8	0.4,1.5	0.8	0.4,1.5	0.8	0.4,1.5	0.8	0.4,1.5	0.8	0.4,1.5	0.8	0.4,1.5
High school drop-out			0.4	0.2,1.0	0.4*	0.2,0.9	0.4*	0.2,0.9	0.4*	0.2,0.9	0.4*	0.2,0.9	0.4*	0.2,0.9	0.4*	0.2,0.9
Employment status																
Unemployed			1		1		1		1		1		1		1	
Employed			0.7	0.4,1.2	0.7	0.4,1.1	0.7	0.4,1.1	0.7	0.4,1.1	0.7	0.4,1.1	0.7	0.4,1.1	0.7	0.4,1.1
Receiving public assistance																
No			1		1		1		1		1		1		1	
Yes			1.8*	1.1,3.0	1.3	0.8,2.3	1.3	0.8,2.3	1.9*	1.2,3.0	1.9*	1.2,3.0	1.9*	1.2,3.0	1.9*	1.2,3.0
Childhood family structure																
2 parents (biological/step)			1		1		1		1		1		1		1	
1 parent only			1.5	0.9,2.7	1.5	0.9,2.5	1.5	0.9,2.5	1.5	0.9,2.5	1.5	0.9,2.5	1.5	0.9,2.5	1.5	0.9,2.5
Other			1.2	0.5,2.7	1.1	0.5,2.4	1.1	0.5,2.4	1.1	0.5,2.4	1.1	0.5,2.4	1.1	0.5,2.4	1.1	0.5,2.4

N=940	Model 1 (Univariate)		Model 2 (Sociodemographics)		Model 3 (Full)		Model 4 (Final reduced)	
	RR	CI	aRR	CI	aRR	CI	aRR	CI
Mother's age at first birth								
20 years old			1		1		1	
<20 years old			1.9**	1.2,3.1	1.7*	1.1,2.7	2.0**	1.3,3.2
Religious service attendance								
Never			1		1			
< weekly			1.5	0.8,2.8	1.5	0.8,2.7		
weekly			1.3	0.6,2.9	1.7	0.7,3.7		
Relationship status								
Married			1		1			
Engaged			2.6	0.3,26.0	2.8	0.3,25.0		
Romantic relationship			1.6	0.2,14.0	1.7	0.2,14.0		
Physical/emotional			1.2	0.1,11.0	1.4	0.2,12.0		
None			0.6	0.1,6.3	1.2	0.1,10.8		
Cohabitation status								
Not cohabitating			1		1		1	
Cohabitating			2.3*	1.2,4.2	1.8	1.0,3.1	2.2**	1.3,3.7
Age at coitarche								
16 years					1		1	
<16 years					2.2*	1.1,4.4	4.2***	2.2,7.8
History of pregnancy								
No					1			
Yes					1.5	0.9,2.5		
Ever had unprotected sex								
No					1			
Yes					1.1	0.6,1.9		

Results are adjusted relative risk ratios (aRR) and 95% confidence intervals (CI) from discrete-time, mixed-effects, proportional hazard models using multivariable logistic regression with a random effect for the woman and controlling for number of pregnancy months, pregnancy months squared and number of journals completed.

* p-values (p) significant for two-tailed alpha at <0.05,

** <0.01,

and <0.001.

Outcome is the risk of experiencing a new pregnancy during the 1-year study period among all women (n=940). Self-reported pregnancy status was assessed each week. Depression, stress and covariates were measured at baseline.

^a Center for Epidemiologic Studies – Depression Scale (CES-D-5) – 4 point cut-off for moderate/severe depression symptoms. Regression modeling was also conducted to examine the effect of mental health symptoms on pregnancy outcomes over an 18-month study period (n=176 pregnancies) and point estimates were similar to those from the one-year models (not shown).

Table 3
The effect of baseline stress symptoms on young women's risk of pregnancy over one year

N=940	Model 1 (Univariate)			Model 2 (Sociodemographics)			Model 3 (Full)			Model 4 (Final reduced)		
	RR	CI		aRR	CI		aRR	CI		aRR	CI	
Stress symptoms ^d												
No (<9pts PSS)	1			1			1			1		
Yes (≥ 9pts PSS)	1.8*	1.1,3.0		1.6	1.0,2.7		1.4	0.8,2.3		1.6*	1.1,2.7	
Age												
18 years				1			1					
19 years				1.0	0.6,1.6		1.0	0.6,1.6				
20 years				0.5	0.2,1.5		0.4	0.1,1.4				
Race/ethnicity												
Non-Black				1			1					
Black				1.4	0.8,2.5		1.2	0.7,2.1				
Educational enrollment												
Not enrolled				1			1					
High school				1.0	0.5,2.0		1.1	0.6,2.2				
2 year college				0.7	0.4,1.3		0.7	0.4,1.3				
4 year college				0.6	0.3,1.1		0.7	0.4,1.4				
High school drop-out				0.4	0.2,1.0		0.4*	0.2,0.9				
Employment status												
Unemployed				1			1					
Employed				0.7	0.5,1.2		0.7	0.5,1.2				
Receiving public assistance												
No				1			1			1		
Yes				2.0*	1.1,3.3		1.4	0.8,2.3		2.0**	1.3,3.2	
Childhood family structure												
2 parents (biological/step)				1			1					
1 parent only				1.5	0.9,2.6		1.4	0.8,2.5				
Other				1.1	0.5,2.5		1.0	0.5,2.3				

N=940	Model 1 (Univariate)		Model 2 (Sociodemographics)		Model 3 (Full)		Model 4 (Final reduced)	
	RR	CI	aRR	CI	aRR	CI	aRR	CI
Mother's age at first birth								
<20 years old			1		1		1	
20 years old			1.9**	1.2,3.0	1.7*	1.1,2.8	1.9**	1.2,3.0
Religious service attendance								
Never			1		1			
< weekly			1.5	0.8,2.8	1.4	0.8,2.7		
weekly			1.4	0.6,3.0	1.6	0.7,3.7		
Relationship status								
Married			1		1			
Engaged			2.6	0.3,25.0	2.6	0.3,24.0		
Romantic relationship			1.6	0.2,14.0	1.7	0.2,14.0		
Physical/emotional			1.2	0.1,11.0	1.2	0.1,11.0		
None			0.7	0.1,6.6	1.0	0.1,8.9		
Cohabitation status								
Not cohabitating			1		1		1	
Cohabitating			2.4**	1.3,4.5	1.9*	1.0,3.5	2.3**	1.4,3.9
Age at coitarche								
16 years					1		1	
< 16 years					2.9**	1.5,5.6	3.7***	2.0,6.7
History of pregnancy								
Yes					1			
No					1.6	0.9,2.8		
Ever had unprotected sex								
No					1			
Yes					1.3	0.8,2.3		

Results are adjusted relative risk ratios (aRR) and 95% confidence intervals (CI) from discrete-time, mixed-effects, proportional hazard models using multivariable logistic regression with a random effect for the woman and controlling for number of pregnancy months, pregnancy months squared and number of journals completed.

* P-values (P) significant for two-tailed alpha at <0.05,

** <0.01,

and <0.001.

Outcome is the risk of experiencing a new pregnancy during the 1-year study period among all women (n=940). Self-reported pregnancy status was measured each week. Stress and covariates measured at baseline.

^a Perceived Stress Scale - 4 (PSS-4) - 9-point cut-off for moderate/severe stress symptoms. Regression modeling was also conducted to examine the effect of mental health symptoms on pregnancy outcomes over an 18-month study period (n=176 pregnancies) and point estimates were similar to those from the one-year models (not shown).

Table 4
The effect of baseline mental health symptoms on pregnancy risk over one year among women without and with a history of pregnancy

	Women without a prior pregnancy (n =734)		Women with (a prior pregnancy n=206)	
	aRR	CI	aRR	CI
Mental health symptoms ^{ab}				
None	1		1	
Depression symptoms only	1.0	0.4,2.4	0.5	0.1,1.8
Stress symptoms only	0.7	0.2,2.5	3.0*	1.1,8.8
Both depression and stress symptoms	2.3*	1.2,4.3	1.4	0.5,3.7
Receiving public assistance				
No	1		1	
Yes	1.9*	1.1,3.6	1.3	0.6,3.1
Mother's age at first birth				
20 years old	1		1	
<20 years old	2.9***	1.7,5.2	1.0	0.5,2.1
Cohabitation status				
Not cohabitating	1		1	
Cohabitating	1.7	0.9,3.2	2.9*	1.3,6.6
Age at coitarche				
16 years	1		1	
<16 years	6.0***	2.8,12.8	1.1	0.4,2.8

Results are adjusted relative risk ratios (aRR) and 95% confidence intervals (CI) from discrete-time, mixed-effects, proportional hazard models using multivariable logistic regression with a random effect for the woman and controlling for number of pregnancy months, pregnancy months squared and number of journals completed and significant covariates.

* P-values (p) significant for two-tailed alpha at <0.05,

** <0.01,

*** and <0.001.

Outcome is the risk of experiencing a new pregnancy during the one-year study period among women without a history of previous pregnancy (n=734) and among women with a previous pregnancy (n=206). Self-reported pregnancy status was measured each week. Depression, stress and covariates were measured at baseline.

^a Center for Epidemiologic Studies – Depression Scale (CES-D-5) – 4 point cut-off for moderate/severe depression symptoms.

^b Perceived Stress Scale - 4 (PSS-4) - 9-point cutoff for moderate/severe stress symptoms. Regression modeling was also conducted to examine the effect of mental health symptoms on pregnancy outcomes over an 18-month study period (n=176 pregnancies) and all point estimates were similar to those from the one-year models (not shown).