Published in final edited form as:

J Comp Physiol A Neuroethol Sens Neural Behav Physiol. 2013 December ; 199(12): 1105–1116. doi: 10.1007/s00359-013-0853-3.

Removing sensory input disrupts spinal locomotor activity in the early postnatal period

JeanMarie Acevedo and Manuel Díaz-Ríos

Department of Anatomy and Neurobiology and Institute of Neurobiology, University of Puerto Rico, Medical Sciences Campus, San Juan, PR 00936

Abstract

Motor patterns driving rhythmic movements of our lower limbs during walking are generated by groups of neurons within the spinal cord, called central pattern generators (CPGs). After suffering a spinal cord injury (SCI), many descending fibers from our brain are severed or become nonfunctional which leaves the spinal CPG network without its initiating drive. Recent studies are focusing on the importance of maintaining sensory stimulation to the legs on SCI patients as a way to initiate and control the CPG locomotor network. We began assessing the role of sensory feedback to the locomotor CPG network by using a neonatal mouse spinal cord preparation where the lower limbs are still attached. Removing sensory feedback coming from the hindlimbs by way of a lower lumbar transection or by ventral root denervation revealed a positive correlation in the ability of sensory input deprivation to disrupt ongoing locomotor activity on older versus younger animals. The differences in the motor responses as a function of age could be correlated with the loss of excitatory activity from sensory afferents. Continued studies on this field could eventually provide key information that translates in the design of novel therapeutic strategies to treat patients that have suffered a SCI.

Keywords

central pattern generator; locomotion; mouse; spinal cord; sensory input

INTRODUCTION

For decades it has been known that networks of interneurons located within the spinal cord are responsible for generating locomotion. These networks interact with specific sensory information to allow for the reinforcement as well as modifications in the organization of locomotor behaviors (Grillner 1981). The connected ensembles of spinal neurons are defined as central pattern generators (CPG) and are capable of generating locomotor-like neural activity independently of supraspinal (descending) and afferent (peripheral) input (Grillner and Zangger 1971). The basic principles of CPG function have been extensively studied in invertebrate model systems and locomotion has been detailed in a primitive fish – the lamprey (Grillner 1981; Grillner et al. 1995; see Marder 1998). In terrestrial mammalian systems of overground locomotion such as cat it has been shown that a spinal cord transection does not hinder the capability of relearning or re-developing hindlimb stepping in the absence of input from the brain (Lovely et al. 1986; Barbeau and Rossignol 1987; De Leon et al. 1998).

While sensory afferents are not essential for producing the patterned locomotor output in animals deprived of descending (brain) inputs, afferent input is an important source of feedback from the ongoing movements. Afferent inputs are involved in the modulation and correction of such parameters as speed, step cycle, burst duration and interlimb coupling in an ongoing locomotor pattern (see Rossignol et al. 2008). Moreover, in the absence of descending input, locomotion can be recruited and controlled by ongoing sensory afferent input from the lower limbs (see Edgerton et al. 2008). Studies conducted by these groups and others have demonstrated that combinational mechanical (treadmill and robotic training), pharmacological (agonists of excitatory pathways and/or antagonists of inhibitory pathways; neurotrophic factor application) and electrical (muscle and/or epidural) strategies re-instate near normal locomotion in animals, and have provided much optimism on the improvement of locomotor function in humans with a spinal cord injury (Rossignol 2006; Harkema 2007; Rossignol et al. 2008; Edgerton et al. 2008; Gerasimenko et al. 2008; Courtine et al. 2009; Harkema 2011).

Most previously and presently published scientific work regarding the understanding of the organization of the mammalian spinal CPG network for locomotion at the cellular level has involved studies using the rodent isolated spinal cord preparation (for recent reviews see Dougherty and Kiehn 2010; Guertin 2009). Physiological and molecular genetics experiments in this preparation have started to shed light to the basic circuit organization responsible for walking in mammals. Even though these experiments have been essential in the development of our present understanding of the organization of the mammalian neuronal network controlling locomotion, the effects of sensory information to spinal locomotor network function have been largely unexplored. We began assessing the role of sensory feedback to the locomotor CPG network by using a recently novel neonatal mouse spinal cord preparation where the lower limbs are still attached (Hayes et al. 2009) where the loss of descending brain and brainstem inputs was mimicked by cervical decapitation and the loss of sensory input was induced by either a upper sacral transection or by ventral root denervation. We show that removing sensory feedback coming from the hindlimbs by way of a lower lumbar transection or by ventral root denervation revealed subtle and, in some cases, drastic changes in motoneuron burst amplitude and even cessation of the locomotor rhythm itself. There was a positive correlation of age with the ability of sensory input deprivation to disrupt ongoing locomotor activity, with the majority younger animals (P0 - early P1) not showing significant effects after sensory input deprivation as compared to older animals (late P1 - P2) where over 80% showed significant reduction in motor activity including complete cessation. The disruption of locomotor activity in older animals was correlated to the loss of excitatory sensory drive as recorded from dorsal nerve roots. These results suggest a developmental relationship between spinal motor function and its dependence on sensory input in the absence of supraspinal inputs.

MATERIALS AND METHODS

Experiments were performed using spinal cords of 0- to 2-day-old (P0 -P2) ICR mice (Charles River, Wilmington, MA). Animals were killed by rapid decapitation. The spinal cord was exposed by ventral laminectomy under ice-cold (4°C) oxygenated (95% O₂-5% CO₂) low-calcium Ringer solution composed of (in mM): 128 NaCl, 4.7 KCl, 1.2 KH₂PO₄, 0.25 CaCl₂, 1.3 MgCl₂, 3.25 MgSO₄, 25 NaHCO₃, and 22 D-glucose). The spinal cord from segments C5 to S3 were pinned ventral-side up using part of the dorsal part of the rib cage and superfused with oxygenated normal Ringer solution composed of (in mM): 111 NaCl, 3.08 KCl, 25 NaHCO₃, 1.18 KH₂PO₄, 1.25 MgSO₄, 2.52 CaCl₂, and 11 D-glucose. In order to assess the effects of sensory input on locomotor activity, a whole spinal cord preparation still connected to the hindlimbs was used (Fig. 1). The preparation used in this

proposed work was adapted from an isolated in vitro neonatal rat spinal cord oriented dorsal-up with intact hindlimbs developed recently by Hayes et al. (2009).

Locomotor-like activity was evoked by perfusion with mouse Ringer solution containing a combination of 5-HT (9–12 M), NMDA (6–9 M) and dopamine (18 M). Small-diameter suction electrodes were placed either: (1) on both L2 ventral roots to monitor alternating flexor activity in the motor pattern from the right and left sides, (2) in both L5 ventral roots to monitor alternating extensor activity or (3) in the L2 and L5 ventral roots to record alternating motor activity between flexor and extensor motorneuron pools as previously done (Kiehn and Kjaerulff 1996). Sensory input deprivation was induced by either performing a lower lumbar-upper sacral transection with an ultra-sharp scissor; this cut will include the whole spinal cord and partial removal of surrounding ribcage. We additionally deprived sensory input by cutting all lumbar and upper sacral ventral roots with ultra-sharp micro-scissors; this will automatically disrupt any noticeable lower limb muscle contractility. All experiments include at least 5 preparations ($n=5$ or greater) at each time point (15, 30 and 60 minutes) after performing either (sacral transection or ventral nerve root denervation) sensory deprivation method. Ventral and dorsal root recordings were band-pass-filtered (100 Hz to 1 kHz) and recorded using an AC amplifier (Model 1600 from A-M systems). The locomotor rhythm was allowed to stabilize over 20 min, after which the sensory deprivation cutting methods were used. Serotonin (5-HT), N-methyl-d-aspartate (NMDA) and dopamine were purchased from Sigma and diluted in regular mouse ringer.

Data analysis

Locomotor-like activity was recorded in the intact spinal cord preparation during bath application of 5-HT, NMDA and dopamine. Clampfit 9.0 (Molecular Devices), Excel (Microsoft, Seattle, WA), and Spike 2 (Cambridge Electronic Design, Cambridge, UK) were used for data analysis. A cycle of motor nerve activity started at the onset of either an L2 or L5 ventral root burst and ended at the onset of the next L2 or L5 ventral root burst recorded from the same nerve; these onsets were determined by a custom-made program in Spike2 (courtesy of Dr. Thomas Cleland, Cornell University) to detect when the rectified signal exceeded the average noise level between bursts by a preset amount. Measurements of burst amplitude (measured from trough to crest) were determined by analysis of rectified and normalized L2 or L5 activity using Spike 2 software. Averages of burst amplitude were determined from all locomotor bursts that occurred once a stable pattern of locomotor-like activity had been established (after a minimum of 15–20 minutes of stable locomotor activity was recorded). Circular statistics (Zar 1974) were used to determine the coupling strength between opposing L2 and L5 ventral roots. Left or right L2 bursts occurring over a continuous 5 min interval were selected, and their phase values were calculated in reference to either the onsets of each left or right L5 burst, respectively (they were always located in the same side in order to have alternation). Phase values were determined by dividing the latency between the onset of the first L2 burst and the following burst in L5 by the step cycle period (time between the reference L2 burst and the next L2 burst). Locomotor steps in which the ipsilateral L2 and L5 roots were completely out of phase (i.e., appropriate flexor-extensor alternation) had phase values of approximately 0.5. Those completely in phase (co-bursting) had phase values of around 0. The r values are a measure of the concentration of phase values around the mean value for alternation (0.5). An r value of 1 indicates all the phase values are 0.5, whereas an r value of 0 indicates the phase values are distributed randomly.

Statistical comparisons between experimental conditions were made using one-way repeated measures ANOVA followed by a Fisher LSD post-hoc test if the data were normally

distributed and had equal variance. Otherwise, the data were compared with a Mann-Whitney rank sum test. Results were considered statistically significant at $p < 0.05$. Data are expressed as mean $\pm$ SD. Figures were compiled using Sigma Plot 10, Photoshop and Corel Draw.

RESULTS

We performed our experiments using mice that were between 0 and 2 days postnatal (P0-P2). A motor pattern was obtained neurochemically using serotonin (5-HT; 9–12 M), the glutamate analog NMDA (6–9 M) and dopamine (18 M) as previously published (Kiehn et al. 1996). Alternating locomotor-like activity was assessed with extracellular nerve recordings and visual confirmation of alternating hindlimb movements. We performed sensory deprivation experiments in only those spinal cord-leg attached preparations which showed stable rhythmic hindlimb activity (over 20 minutes) after adding the drug cocktail.

Limb deafferentation was achieved via a complete transection at the upper sacral level (around S1-S2) just above the hip bones of P0 animals. The rationale for using a sacral transection stemmed from previous studies in which a spinal cord injury was produced by either a compression or a cut of the spinal cord at a specific spinal level in cats and rodents (see Edgerton et al. 2008; Rossignol and Frigon 2011). We expected that by severing the spinal cord, as well as all ventral and dorsal roots, we will induce an almost immediate stoppage of the flow of afferent (sensory) and efferent (motor) information thus perturbing the ongoing (drug-induced) locomotor rhythm. We checked for any significant changes in the either burst amplitude or the phase of the locomotor pattern (alternating versus non-alternating) at 15, 30 and 60 minutes after performing the sacral transection. After 1 hour of deafferentation, no significant changes were seen in burst amplitude (Fig. 2a1; $n=5$) or locomotor coordination between the left and right L2 or between the right L2 and the right L5 ventral roots (Fig. 2a2; $n=5$). The average burst amplitude in control P0 preparations was 4.1 ± 2.8 mV. After 60 minutes of sensory deprivation the average burst amplitude in these P0 preparations changed to 3.8 ± 1.5 , which was not statistically significant. However, the same deafferentation protocol on P1 mice induced a progressive decrease in burst amplitude (Fig. 2b1; $n=5$) coupled with a loss in the alternating phase of the pattern (Fig. 2b2; $n=5$). A locomotor-like rhythm could not be seen in 3 of 5 preparations within an hour after the cut (Fig. 2b; $n=5$). The average time that it took to initiate the progressive loss of the locomotor pattern was 37 ± 8.5 minutes. The average burst amplitude in P1 control preparations was 3.3 ± 1.5 mV. After 60 minutes of sensory deprivation the average burst amplitude in these P1 preparations changed to 0.5 ± 1.3 , which was statistically significant. It is important to mention that even though 2 preparations did not completely lose rhythmic motor activity after sensory deprivation, the denervation still had significant effects on burst amplitude and in the alternating phase of the pattern (data not shown) on these preparations. In P2 preparations, deafferentation led to an even more rapid dissolution of rhythmic motor activity (within 20 ± 9.5 minutes) with one preparation losing all rhythmic activity within 15 minutes following deafferentation (Fig. 2c1,2; $n=5$) including a complete cessation of rhythmic motor activity in 4 of 5 preparations with 1 of 5 preparations not losing alternating activity but still showing a significant reduction in burst amplitude (data not shown). The average burst amplitude in control P2 preparations was 5.5 ± 3.3 mV, which was significantly reduced to 1.5 ± 1.7 mV after 60 minutes of sensory deprivation.

All preparations were left with ongoing perfusion of drugs (5-HT, NMDA and DA) for at least an hour with no recovery of rhythmic activity. In order to eliminate the possibility that the loss of the locomotor rhythm could be due to over-excitation caused by prolonged exposure to the drugs, control experiments were performed after obtaining a stable rhythm.

No sacral transection was made and the preparations were left untouched for at least 90 minutes to two hours with continuous drug perfusion. None of the preparations ($n=3$) displayed a loss of the locomotor rhythm although a small but significant decrease in burst amplitude and the speed of alternating motor activity (the hindlimbs never stopped alternating in all preparations tested) was seen after 1 hour of continuous perfusion in two of the preparations (data not shown) which could be attributed to preparation fatigue (lack of oxygenation to more internal areas of the cord).

Since the sacral transection method included cutting the spinal cord and ventral/dorsal nerve roots (including some muscle and bone from the rib cage), we repeated these experiments using a less invasive sensory deprivation method to eliminate the possibility that the observed effects were due to an injury-related effect rather than from sensory input deprivation. Instead of a sacral cut (transection) we repeated sensory deprivation experiments on P0 to P2 spinal cord lower limb-attached preparations by cutting all lumbar and sacral ventral nerve roots as the point where they exit the spinal cord. This manipulation will interrupt the flow of efferent (motor) drive from the spinal CPG network to the lower limbs which stops muscle contraction (including cessation of rhythmic alternating hindlimb movements) and lead to a cessation of afferent (sensory) input through the still intact dorsal nerve roots. The rationale for cutting just the ventral roots was to interrupt the flow of efferent (motor) control to the lower limb muscles thus reducing and eventually stopping the flow of afferent (sensory) information flowing back in the spinal CPG network through the dorsal nerve roots due to the loss of muscle contraction. Even though we did not interrupt the flow of cutaneous (skin)-related information since the afferent fibers (dorsal nerve roots) were left intact, we expected to obtain similar results as with the sacral transection method. This prediction is based on our theory that the disruption of locomotor activity is due to the deprivation of muscle spindle and Golgi tendon-related sensory information (not cutaneous-related sensory input) through to the cessation of motor output by cutting the ventral nerve roots thus interrupting the motor-sensory information loop.

Following 20 minutes of stable drug-induced (5-HT, NMDA and DA) alternating locomotor activity, all lumbar and sacral ventral roots were severed, eliminating alternating locomotor activity in the hindlimbs of the animals due to the loss of muscle contractions. Sensory deprivation via ventral nerve root cuts had no significant effects on 4 of 5 P0 preparations with just 1 of 5 preparations losing rhythmic motor activity within 30 minutes after cutting (Fig. 3a1,2). The average burst amplitude in control P0 preparations was 2.0 ± 1.7 mV which was reduced to 1.9 ± 1.3 mV after 60 minutes of sensory deprivation but this result was not statistically significant. Sensory deprivation via ventral nerve root cutting had significant effects on 3 of 5 preparations in P1 mice with 2 of 5 preparations losing locomotor-like activity within 60 minutes after sensory deprivation (Fig. 3b1,2). The average burst amplitude in P1 preparations was 3.1 ± 1.4 mV which was significantly reduced to 1.7 ± 1.2 mV after 60 minutes of sensory deprivation. Finally, we tested the effects of ventral nerve root cutting and subsequent sensory input loss on the drug-induced rhythmic motor patterns of P2 spinal cord lower limb-attached preparations. Sensory deprivation induced a cessation of locomotor activity in 4 of 5 preparations. Loss of activity was seen as early as 15 minutes after sensory input deprivation with an average loss of discernible rhythmic motor activity at 30.2 ± 8.1 minutes after ventral nerve root cutting (Fig. 3c1,2). The average burst amplitude in P2 preparations was 2.8 ± 1.3 mV which was significantly reduced to 0.6 ± 0.9 mV, after 60 minutes of sensory deprivation. A comparison of the magnitude of the effect of depriving sensory input via sacral transection and ventral nerve root cuts reveals that, after 60 minutes, a sacral transection decreased burst amplitude by about 85% and 73% in P1 and P2 animals, respectively. Although the sacral transection method decreased burst amplitude by a higher percent in P1 than in P2 animals,

the effect produced by this sensory input deprivation method was always faster in P2 than in P1 mice (37 ± 8.5 min in P1 mice versus 20 ± 9.5 min in P2 mice; Fig. 2). Depriving sensory input in P1 and P2 animals via ventral nerve root cuts revealed that this method reduced burst amplitude by 45% and 79% in P1 and P2 animals, respectively. As in the case of the sacral transection method, cutting all lumbar and sacral ventral nerve roots decreased burst amplitude and eventually disrupted locomotor activity faster in P2 than in P1 mice (47.8 ± 7.9 in P1 mice versus 30.2 ± 8.1 min in P2 mice; Fig. 3). Although there were qualitative differences regarding the effects of sensory deprivation on P0-P2 spinal cord lower limb-attached preparations using a sacral transection versus ventral nerve root cutting, there was still a positive correlation between animal age and robustness of the effects on sensory input loss on an ongoing rhythmic locomotor pattern of activity. These results suggest that the spinal locomotion CPG neuronal network is developing toward a more sensory-dependent state with age.

In order to elucidate potential mechanisms that could explain this age dependent loss of spinal locomotor activity after losing sensory inputs in P2 animals, we explored the possibility that the disruption of motor activity could be attributed to a loss of excitatory drive from sensory afferents. To assess this possibility we recorded motor activity from ventral roots of P2 animals as previously done but additionally recorded activity from both (left and right) dorsal nerve roots at the L4 level and measured any changes in the frequency of the activity measured from both types (ventral/dorsal) of nerve roots. Previous studies have shown that electrical stimulation of dorsal nerve roots from the mid to lower lumbar levels can elicit bouts of spontaneous locomotor like activity lasting up to several minutes (Gabbay et al. 2002; Lev-Tov et al. 2010) which prompted us to use the L4 dorsal nerve root to monitor afferent (sensory) activity. We measured the frequency of the firing activity recorded from the dorsal root and simultaneously recorded drug-induced (5-HT/NMDA/DA) locomotor-like activity from the ventral nerve roots before, during, and after cutting all lumbar and sacral ventral roots. We found that the loss of afferent activity always preceded the disruption of locomotor-like activity as seen by a significant reduction in burst amplitude as early as 5 minutes after the cut (Fig 4; $n=5$). We additionally attempted to regain locomotor-like activity in P2 animals after ventral nerve root denervation by increasing the concentration of the drug cocktail (5-HT/NDMA/DA). The rationale behind augmenting the concentration of locomotor-inducing drugs was that if the loss of alternating locomotor activity was due to the interruption of afferent (sensory) input, the raised concentration of the drug cocktail should compensate for this apparent void in excitatory drive left after stopping the flow of motor and sensory information through ventral root denervation. We sequentially changed the concentration of each of the drugs by 1 M increments up to 5 M (from 9 M 5-HT / 6 M NMDA / 18 M DA up to 14 M 5-HT / 11 M NMDA / 23 M DA) in 20 minutes time periods (each new concentration of drugs was maintained in the bath for at least 20 minutes). Although there was an increase in overall nerve root activity, increasing the concentration of drugs failed to recover alternating locomotor-like activity in all P2 animals tested (Fig. 5, $n=5$). This result suggests that this apparent loss of excitatory drive to the CPG network after denervation could be due to the progressive loss of a sensory-related neuromodulatory transmitter and not just from less flow of locomotor-related excitatory neurotransmitters, as those present in our drug cocktail, through the motor-sensory loop.

Overall our results support the idea that when the spinal locomotion CPG network has to interact only with sensory feedback due to the lost of supraspinal inputs (through injury for example), the network develops a more sensory-dependent state of spinal motor activity which could be correlated to the developmental maturity of the sensory and motor spinal networks (Fig. 6). Potential mechanisms explaining this apparent loss of sensory drive to the spinal locomotor network are now discussed.

DISCUSSION

It has been well studied and documented that networks of interneurons located within the spinal cord are responsible for generating locomotion and that these networks interact with specific sensory information to allow for the reinforcement as well as modifications in the organization of locomotor behaviors (Grillner 1981). After suffering a spinal cord injury, many (and in some cases all) descending fibers (axons) from supraspinal centers are severed or become nonfunctional which leaves the CPG network controlling locomotion without its initiating and controlling drive (input). Recent studies have focused on the importance of maintaining sensory stimulation to the lower limbs, via physical therapy for example, on spinal cord injury patients as a way to initiate and control the CPG locomotor network (Harkema et al. 2011; Courtine et al. 2009; Lavrov et al. 2008; Edgerton et al. 2008) but studies looking at the importance of sensory input to locomotor activity in the absence of supraspinal inputs are lacking. Here we studied the effects of sensory deprivation on spinal locomotor CPG network function using a recently developed spinal cord lower limb-attached preparation (Hayes et al. 2009). We sought to understand the role of sensory input to spinal locomotor function as the animal begins to actively interact with its environment during its first days after birth by disrupting sensory input via a sacral cord transection or through ventral nerve root denervation.

Our previous work with the isolated spinal cord (Díaz-Ríos et al. 2007; Zhong et al. 2006a,b) has shown that fictive locomotion can be elicited with a combination of 5-HT, NMDA and dopamine, and that this rhythm can last for many hours as long as the concentration of drugs in the bath is kept constant with perfusion. So then why would a drug-elicited locomotor rhythm stop its alternating activity after depriving it of sensory feedback from the hindlimbs? We would like to discuss two possible explanations: (1) locomotor activity induced pharmacologically in a spinal cord preparation with the hindlimbs attached could result in a motor pattern produced by a spinal CPG network that is more dependent on sensory feedback than a CPG network which relies on the synaptic properties of its component neurons to produce a “fictive” locomotor pattern (isolated spinal cord); (2) the deprivation of sensory information via a cut of all afferent (dorsal) roots “feeding” back to the CPG network could be leading to a “spinal shock” like state (seen previously after a spinal cord injury; see Dietz 2010 and 2011 for recent reviews) which could explain the “shutdown” of network activity after a lower sacral transection.

Dorsal root injuries, which can result in the partial or complete loss of connections between the sensory system and the spinal cord, can lead to sensory impairments, including both the loss of sensation and the development of neuropathic pain in the affected limb (Cragg et al. 2010). Our data does not support that the observed effects on locomotor activity were due to an injury-related effect of cutting the spinal cord due to the following facts: (1) we never saw any immediate effects on motor activity after cutting (data not included), these including any hyper-excitability on the motor output immediately after cutting or any other noticeable effect; (2) preparations were deprived of supraspinal input via cervical decapitation in ice-cold low calcium Ringer and the preparation was dissected and prepared in this Ringer before being moved to our recording chamber and perfused with regular Ringer solution (see Materials and Methods) in the presence of the drug cocktail. This procedure will take around 45 minutes to 1 hour and then the preparation is continuously perfused with these drugs, even during and after the sacral transection, which should counteract any “spinal shock-like” state induced by the decapitation procedure or spinal cord cutting. Still, in order to assess the possibility of an injury-mediated effect due to cutting the spinal cord we began using a ventral nerve root cutting procedure thus maintaining the spinal cord and all dorsal nerve roots intact. By not severing the spinal cord or any dorsal

roots we avoided any type of peripherally injury-mediated effects. Effects from ventral nerve cutting on locomotor activity were similar to our sacral transection experiments thus supporting our observations.

We believe that by removing sensory feedback to an ongoing locomotor pattern, the deprivation of sensory information (via sacral transection or ventral root denervation) could result in the disruption of a locomotor CPG network which had “learned” to produce locomotion in a sensory feedback-dependent manner. This concept will support the idea that in situations where there is no descending input to the spinal cord, the control of locomotion could be directed from specialized spinal networks directed by sensory afferent input from the lower limbs (Edgerton et al. 2008; Lavrov et al. 2008; Courtine et al. 2009; Rossignol and Frigon 2011; Rossignol et al. 2011). Additionally, our results confirmed a positive correlation between loss of afferent activity, as recorded from dorsal nerve roots using suction electrodes, and the subsequent disruption of locomotor activity in all P2 animals tested (Fig. 4). This further suggests that by severing the ventral roots, which caused a subsequent loss of muscle contraction, we disrupted the afferent (sensory) and efferent (motor) activity loop which lead to an apparent loss of sensory drive to the CPG network resulting in a disruption of locomotor-like activity. One possible source of this sensory drive could be via a yet to be identified neuromodulator such as a neuropeptide.

The tachykinin neuropeptide Substance P (SP) is abundant both in the periphery and in the central nervous system (CNS), where it is usually co-localized with one of the classical neurotransmitters, most commonly serotonin (5-HT; Marson, 1989; Thor et al. 1988; Arvidsson et al. 1994). Earlier and more recent work has linked Substance P to a centrally and peripherally located modulator of locomotor behavior in model systems such as the lamprey and rat (Piercey et al. 1981; Barthe and Clarac 1997; Ulström et al. 1999; Svensson et al. 2001; Kozlov et al. 2001; Parker and Bevan 2007; Thorn Pérez et al. 2009). Application of SP to the isolated lumbosacral spinal cord of newborn rats induced slow rhythmic activity in quiescent preparations (no other drugs present in the perfusate) and during drug-induced fictive locomotion, SP application increased locomotor cycle frequency and burst duration including a stabilizing effect by the neuropeptide on the locomotor rhythm (Barthe and Clarac 1997). Other neuropeptides which have been linked to the dorsal horn and sensory neurons include Neuropeptide Y and GABA (Parker and Grillner 1996; Ulström et al. 1999). NPY has the opposite effect of tachykinins in that it reduces the excitability of spinobulbar interneurons and presynaptically reduces dorsal cell-mediated glutamatergic synaptic transmission (Parker et al. 1998). GABA is colocalized with NPY in bipolar neurons in the dorsal column (Parker et al. 1998). The GABA_B receptor agonist baclofen has complementary effects to NPY in depressing sensory synaptic transmission (Christenson et al. 1991; Parker et al. 1998). Based on the observed inhibitory effects of Neuropeptide Y and GABA on sensory synaptic transmission, these are unlikely candidates for explaining the loss of rhythmic motor activity due to sensory input deprivation.

In summary, these experiments suggest that the effects of sensory deprivation on an ongoing drug-induced locomotor pattern could be related to a disruption of what seems to be a “hybrid” network comprised of the locomotor CPG network whose component neurons are located within the lumbar spinal cord and sensory feedback coming from the hindlimbs. The disruption of motor output in older animals could be attributed to a spinal CPG network which is more dependent in sensory input (Fig. 6) due to a maturing motor and sensory circuitry. We suggest that these effects are enhanced in the spinal cord lower limb-attached preparation used in this study since the spinal CPG circuitry can only access sensory input and not supraspinal inputs as could be the case in severe cases of cervical or thoracic spinal cord injury. These results are supported by recent studies that have shown that after inducing

a spinal cord injury via a complete transection of the nerve cord in cats and rodents, combinations of serotonergic agonists and epidural electrical stimulation were able to acutely transform spinal networks from nonfunctional to highly functional and adaptive states as early as 1 week after injury (Dorofeev et al. 2008; Courtine et al. 2009). The variability in the motor responses as a function of age recorded after the loss of sensory feedback could be correlated with the developmental maturity of the locomotor CPG network and a use-dependent relationship with sensory information. We are currently gathering data on slightly older animals (P3-P5) than those included in this present study (P0-P2) to contrast and compare any potential differences regarding the effects of sensory deprivation on locomotor behavior in the absence of supraspinal inputs. Other studies are also ongoing, including the use of electromyographic recordings (EMGs) to correlate muscle activity to CPG motor output (Tysseling et al. 2013), in order to elucidate the potential mechanism mediating the effects apparently induced by the disruption of sensory input to an ongoing centrally-initiated motor output.

Acknowledgments

We thank Thomas Cleland for Spike 2 scripts used in data analysis and to Alex Kwan for Spike 2 scripts used in the generation of the circular plots. We would also like to thank Ernesto Cabezas, Jeidiel De Leon, Marla Rivera Oliver and Nikol Matos Vergara for their helpful comments on the manuscript. We would additionally like to thank the University of Puerto Rico Puerto Rico Institutional Animal Care and Use Committee for the approval of the animal protocols used for the experiments included in this work in accordance with National Institutes of Health guidelines. The present study was supported by the Craig Neilsen Foundation grant 124554, the Research Centers in Minority Institution at the University of Puerto Rico-Medical Sciences Campus (RCMI-UPR-MS) grant G12RR03051 and a Minority Biomedical Research Support-Research Initiative for Scientific Enhancement (MBRS-RISE) pre-doctoral grant R25GM061838 to JeanMarie Acevedo-Rosario.

References

- Barbeau H, Rossignol S. Recovery of locomotion after chronic spinalization in the adult cat. *Brain Res.* 1987; 412:84–95. [PubMed: 3607464]
- Barthe JY, Clarac F. Modulation of the spinal network for locomotion by substance P in the neonatal rat. *Exp Brain Res.* 1997; 115:485–492. [PubMed: 9262203]
- Courtine G, Gerasimenko Y, van den Brand R, Yew A, Musienko P, Zhong H, Song B, Ao Y, Ichiyama RM, Lavrov I, Roy RR, Sofroniew MV, Edgerton VR. Transformation of nonfunctional spinal circuits into functional states after the loss of brain input. *Nat Neurosci.* 2009; 12:1333–1342. [PubMed: 19767747]
- Christenson J, Borman A, Lagerback PA, Grillner S. The dorsal cell, one class of primary sensory neurone in the lamprey spinal cord. I. Touch, pressure but no nociception—a physiological study. *Brain Res.* 1998; 440:1–8. [PubMed: 3359201]
- Cragg JJ, Scott AL, Ramer MS. Depletion of spinal 5-HT accelerates mechanosensory recovery in the deafferented rat spinal cord. *Exp Neurol.* 2010; 222:277–284. [PubMed: 20079735]
- De Leon RD, Hodgson JA, Roy RR, Edgerton VR. Locomotor capacity attributable to step training versus spontaneous recovery after spinalization in adult cats. *J Neurophysiol.* 1998; 79:1329–1340. [PubMed: 9497414]
- Díaz-Ríos M, Dombeck D, Webb WW, Harris-Warrick RM. Serotonin modulates dendritic calcium influx in commissural interneurons in the mouse spinal locomotor network. *J Neurophysiol.* 2007; 98:2157–2167. [PubMed: 17581844]
- Dietz V. Behavior of spinal neurons deprived of supraspinal input. *Nat Rev Neurol.* 2010; 6:167–174. [PubMed: 20101254]
- Dietz V. Neuronal plasticity after a human spinal cord injury: Positive and negative effects. *Exp Neurol.* 2011; 235:110–115. [PubMed: 21530507]

- Dorofeev IY, Avelev VD, Shcherbakova NA, Gerasimenko YP. The role of cutaneous afferents in controlling locomotion evoked by epidural stimulation of the spinal cord in decerebrate cats. *Neurosci Behav Physiol.* 2008; 38:695–701. [PubMed: 18720012]
- Dougherty KJ, Kiehn O. Firing and cellular properties of V2a interneurons in the rodent spinal cord. *J Neurosci.* 2010; 30:24–37. [PubMed: 20053884]
- Edgerton VR, Courtine G, Gerasimenko YP, Lavrov I, Ichiyama RM, Fong AJ, Cai LL, Otsoshi CK, Tillakaratne NJ, Burdick JW, Roy RR. Training locomotor networks. *Brain Res Rev.* 2008; 57:241–254. [PubMed: 18022244]
- Gabbay H, Delvolvé I, Lev-Tov A. Pattern generation in caudal-lumbar and sacrococcygeal segments of the neonatal rat spinal cord. *J Neurophysiol.* 2002; 2:732–739. [PubMed: 12163525]
- Gerasimenko Y, Roy RR, Edgerton VR. Epidural stimulation: comparison of the spinal circuits that generate and control locomotion in rats, cats and humans. *Exp Neurol.* 2008; 209:417–425. [PubMed: 17850791]
- Grillner, S. Handbook of physiology. The nervous system. Motor control. Vol. II. Am. Physiol. Soc./Wiley-Blackwell; Maryland: 1981. Control of locomotion in bipeds, tetrapods, and fish; p. 1179-1236. Sect. 1
- Grillner S, Deliagina T, Ekeberg O, El Manira A, Hill RH, Lansner A, Orlovsky GN, Wallen P. Neural networks that co-ordinate locomotion and body orientation in lamprey. *Trends Neurosci.* 1995; 18:270–279. [PubMed: 7571002]
- Grillner S, Zangger P. How detailed is the central pattern generation for locomotion? *Brain Res.* 1975; 88:367–371. [PubMed: 1148835]
- Guertin PA. The mammalian central pattern generator for locomotion. *Brain Res Rev.* 2009; 62:45–56. [PubMed: 19720083]
- Harkema SJ. Plasticity of interneuronal networks of the functionally isolated human spinal cord. *Brain Res Rev.* 2007; 57:255–264. [PubMed: 18042493]
- Harkema S, Gerasimenko Y, Hodes J, Burdick J, Angeli C, Chen Y, Ferreira C, Willhite A, Rejc E, Grossman RG, Edgerton VR. Effect of epidural stimulation of the lumbosacral spinal cord on voluntary movement, standing, and assisted stepping after motor complete paraplegia: a case study. *Lancet.* 2011; 377:1938–1947. [PubMed: 21601270]
- Hayes HB, Chang YH, Hochman S. An in vitro spinal cord-hindlimb preparation for studying behaviorally relevant rat locomotor function. *J Neurophysiol.* 2009; 101:1114–1122. [PubMed: 19073815]
- Kiehn O, Johnson BR, Raastad M. Plateau properties in mammalian spinal interneurons during transmitter-induced locomotor activity. *Neuroscience.* 1996; 75:263–273. [PubMed: 8923540]
- Kiehn O, Kjaerulff O. Spatiotemporal characteristics of 5-HT and dopamine-induced rhythmic hindlimb activity in the in vitro neonatal rat. *J Neurophysiol.* 1996; 75:1472–1482. [PubMed: 8727391]
- Kozlov A, Kotaleski JH, Aurell E, Grillner S, Lansner A. Modeling of substance P and 5-HT induced synaptic plasticity in the lamprey spinal CPG: consequences for network pattern generation. *J Comput Neurosci.* 2001; 11:183–200. [PubMed: 11717534]
- Lev-Tov A, Etlin A, Blivis D. Sensory-induced activation of pattern generators in the absence of supraspinal control. *Ann N Y Acad Sci.* 2010; 1198:54–62. [PubMed: 20536920]
- Lovely RG, Gregor R, Roy RR, Edgerton VR. Effects of training on the recovery of fullweight-bearing stepping in the adult spinal cat. *Exp Neurol.* 1986; 92:421–435. [PubMed: 3956672]
- Marder E. From biophysics to models of network function. *Annu Rev Neurosci.* 1998; 21:25–45. [PubMed: 9530490]
- Parker D, Bevan S. Modulation of cellular and synaptic variability in the lamprey spinal cord. *J Neurophysiol.* 2007; 97:44–56. [PubMed: 17021027]
- Parker D, Grillner S. Tachykinin-mediated modulation of sensory neurons, interneurons, and synaptic transmission in the lamprey spinal cord. *J Neurophysiol.* 1996; 76:4031–4039. [PubMed: 8985898]

- Parker D, Söderberg C, Zotova E, Shupliakov O, Langel U, Bartfai T, Larhammar D, Brodin L, Grillner S. Co-localized neuropeptide Y and GABA have complementary presynaptic effects on sensory synaptic transmission. *Eur J Neurosci*. 1998; 10:2856–2870. [PubMed: 9758155]
- Rossignol S. Plasticity of connections underlying locomotor recovery after central and/or peripheral lesions in the adult mammals. *Phil Trans R Soc Lond B Biol Sci*. 2006; 361:1647–1671. [PubMed: 16939980]
- Rossignol S, Barrière G, Frigon A, Barthélemy D, Bouyer L, Provencher J, Leblond H, Bernard G. Plasticity of locomotor sensorimotor interactions after peripheral and/or spinal lesions. *Brain Res Rev*. 2008; 57:228–240. [PubMed: 17822774]
- Rossignol S, Frigon A. Recovery of locomotion after spinal cord injury: some facts and mechanisms. *Annu Rev Neurosci*. 2011; 34:413–440. [PubMed: 21469957]
- Rossignol S, Frigon A, Barrière G, Martinez M, Barthélemy D, Bouyer L, Bélanger M, Provencher J, Chau C, Brustein E, Barbeau H, Giroux N, Marcoux J, Langlet C, Alluin O. Chapter 16--Spinal plasticity in the recovery of locomotion. *Prog Brain Res*. 2011; 188:229–241. [PubMed: 21333814]
- Svensson E, Grillner S, Parker D. Gating and braking of short- and long-term modulatory effects by interactions between colocalized neuromodulators. *J Neurosci*. 2001; 21:5984–5992. [PubMed: 11487621]
- Thörn Pérez C, Hill RH, El Manira A, Grillner S. Endocannabinoids mediate tachykinin-induced effects in the lamprey locomotor network. *J Neurophysiol*. 2009; 102:1358–1365. [PubMed: 19571197]
- Tysseling VM, Janes L, Imhoff R, Quinlan KA, Lookabaugh B, Ramalingam S, Heckman CJ, Tresch MC. Design and evaluation of a chronic EMG multichannel detection system for long-term recordings of hindlimb muscles in behaving mice. *J Electromyogr Kinesiol*. 2013; 23:531–539. [PubMed: 23369875]
- Ullström M, Parker D, Svensson E, Grillner S. Neuropeptide-mediated facilitation and inhibition of sensory inputs and spinal cord reflexes in the lamprey. *J Neurophysiol*. 1999; 81:1730–1740. [PubMed: 10200208]
- Zar, JH. Biostatistical analysis. Englewood Cliffs, New Jersey: Prentice-Hall; 1974.
- Zhong G, Díaz-Ríos M, Harris-Warrick RM. Serotonin modulates the properties of ascending commissural interneurons in the neonatal mouse spinal cord. *J Neurophysiol*. 2006a; 95:1545–1555. [PubMed: 16338993]
- Zhong G, Díaz-Ríos M, Harris-Warrick RM. Intrinsic and functional differences among commissural interneurons in the central pattern generator for locomotion in the neonatal mouse. *J Neurosci*. 2006b; 26:6509–6517. [PubMed: 16775138]

Fig. 1.

Spinal Cord lower limb-attached preparation. **a** Suction recording electrodes were placed to monitor motor activity from ventral root nerve before, during and after perfusion of drugs. **b** Actual preparation showing simultaneous extracellular recordings **c** Extracellular recordings showing rhythmic locomotor-like activity from lumbar nerve 2 and the left and right 5th lumbar nerves (scale bar in **b** panel: 3 mm)

Fig. 2. Effects of sensory deprivation on P0, P1 and P2 mouse spinal cord lower limb-attached preparations. All measurements were performed 15, 30 and 60 minutes after sensory deprivation via sacral transection. **a1** Mean data on the effects of the loss of sensory input on the burst amplitude of the ventral L2 and L5 nerve recordings from P0 mouse spinal cord preparations ($n = 5$). **a2** Circular plots showing the effects of sensory feedback to the phasing between the rL2 and rL5 ventral roots (Note: cycles of motor burst activity located at 0.5 are considered in alternation while 0 is considered synchronous activity). **b** Similar recording as in **a** panels but from a P1 mouse spinal cord lower limb-attached preparation. **c** Similar recording as in **a** panels but from a P2 mouse spinal cord lower limb-attached preparation ($n = 5$ for each age group)

Fig. 3.

Effects of sensory deprivation on P0, P1 and P2 mouse spinal cord preparations. All measurements were performed 15, 30 and 60 minutes after sensory deprivation after cutting all lumbar ventral nerve roots. **a1** Mean data on the effects of the loss of sensory input on the burst amplitude of the ventral L2 and L5 nerve recordings from P0 mouse spinal cord preparations ($n = 5$). **a2** Circular plots showing the effects of sensory feedback to the phasing between the rL2 and rL5 ventral roots (Note: cycles of motor burst activity located at 0.5 are considered in alternation while 0 is considered synchronous activity). **b** Similar recording as in **a** panels but from a P1 mouse spinal cord lower limb-attached preparation. **c** Similar recording as in **a** panels but from a P2 mouse spinal cord lower limb-attached preparation ($n = 5$ for each age group)

Fig. 4.

Comparing ventral nerve activity to dorsal root activity before and after sensory deprivation via ventral nerve root cuts on P2 mice. **a** Rectified ventral root activity from the right 2nd lumbar nerve root and raw dorsal root activity from the right 4th lumbar dorsal nerve root before and, 1 and 15 minutes after cutting all ventral nerve roots ($n = 5$). **b** panels: Mean data on the effects of loss of sensory input on burst amplitude of P2 mouse spinal cord lower limb-attached preparations (**b1**) and the firing activity as recorded from both L4 dorsal roots (**b2**) ($n = 5$)

Fig. 5.

Attempting to regain alternating locomotor-like activity after sensory deprivation by increasing the concentration of 5-HT, NMDA and dopamine (locomotion-inducing drug cocktail) in spinal cord-leg attached preparations of P2 mice. After the disruption of the recorded locomotor-like rhythm was achieved via sensory deprivation (ventral root denervation) we proceeded to raise the initial drug cocktail concentrations (5-HT: 9 M / NMDA: 6 M / Dopamine: 18 M) by 1 M increments up to 5 M (from 9 M 5-HT / 6 M NMDA / 18 M DA up to 14 M 5-HT / 11 M NMDA / 23 M DA) in 20 minutes time periods and check (via extracellular ventral root recordings) if the alternating motor pattern was recovered with a higher concentration of the drugs ($n = 5$)

Fig. 6.

Diagram of the potential mechanism by which stopping the flow of sensory input disrupts an ongoing pharmacologically-induced locomotor pattern. The included diagram is intended to suggest a potential mechanism by which depriving sensory input to a spinal CPG network which was activated pharmacologically in the presence of limb afferent feedback and no supraspinal inputs. We suggest that the developmental maturity of the sensory network could explain why depriving sensory input in a P0-early P1 animal did not produce a significant perturbation of motor output since apparently this motor pattern was mostly produced by the intrinsic circuitry of the spinal CPG network (arrows). In contrast, in late P1-P2 animals the spinal CPG network could have produced a motor pattern which depended in part on afferent feedback from the limbs (arrows). We suggest that this potential mechanism is enhanced in the absence of supraspinal inputs and the source of this afferent signal which is lost after sensory input deprivation is yet to be elucidated