Published in final edited form as:

Pediatr Obes. 2013 October ; 8(5): e59–e63. doi:10.1111/j.2047-6310.2013.00162.x.

Lifestyle intervention improves lipoprotein particle size and distribution without weight-loss in obese Latino adolescents

Justin R. Ryder¹, Sonia Vega-López^{1,2}, Rocio Ortega^{2,3}, Yolanda Konopken⁴, and Gabriel Q. Shaibi^{1,2,3}

¹School of Nutrition and Health Promotion, Arizona State University, Phoenix, AZ

²Southwest Interdisciplinary Research Center, Arizona State University, Phoenix, AZ

³College of Nursing and Health Innovation, Arizona State University, Phoenix, AZ

⁴St. Vincent de Paul Virginia G. Piper Medical and Dental Clinic, Phoenix, AZ

Abstract

Childhood obesity is associated with a pro-atherogenic phenotype contributing to increased cardiovascular disease (CVD) risk. This single-arm pilot study examined the effects of a lifestyle intervention on lipoprotein particle size and cholesterol distribution in obese Latino adolescents. Fifteen obese Latino adolescents (15.0±1.0 years) completed a 12-week nutrition education and exercise intervention. LDL particle size and distribution of cholesterol in lipoprotein subclasses were determined via polyacrylamide gel electrophoresis. The intervention resulted in increases in mean LDL-particle size (269.3±3.4 to 271.6±2.9Å, p=0.0003) and cholesterol in large HDL subfractions (22.4±11.2 to 26.8±10.6% area, p=0.007) along with decreases of cholesterol in small LDL (1.6±2.0 to 0.6±1.2% area, p<0.01) and HDL subfractions (23.2±9.4 to 19.0±6.7% area, p=0.05). These improvements were observed independent of changes in weight (90.7±26.2 to 89.9±27.8 kg, p>0.05) and suggest that lifestyle modification in obese youth may reduce cardiovascular risk by shifting lipoprotein particle size and cholesterol distribution to a less atherogenic phenotype.

Keywords

Obesity; LDL-size; cholesterol distribution; atherosclerosis; cholesterol

Introduction

Obesity places youth at higher risk for premature morbidity and mortality.(1) Obese youth exhibit cardiovascular disease (CVD) risk factors including an atherogenic lipoprotein distribution(2-4) that contributes to atherosclerosis and overt CVD.(5) Although the atherosclerotic process begins in childhood,(6) traditional markers do not account for the majority of CVD risk observed in adulthood.(7) In adults, atherogenic phenotyping that includes classification of size and distribution of LDL-cholesterol (LDL-c) and HDL-cholesterol (HDL-c) predicts incident CVD independent of total LDL-c and HDL-c.(8-11). Therefore, including a more comprehensive assessment of lipid markers in youth may improve identification of those at greatest risk for CVD.(12)

Corresponding Author: Gabriel Q. Shaibi, PhD, 500 N. 3rd St, Phoenix, AZ 85004, Phone: (602)496-0909, Gabriel.Shaibi@asu.edu.

Conflict of interest: The Authors have no conflicts of interest.

To date, few pediatric studies incorporate lipoprotein size and distribution measures and whether these markers can be improved in obese youth through lifestyle modification independent of weight-loss is not clear. Therefore, the purpose of this study was to determine the effects of a 12-week lifestyle intervention on traditional lipid profiles as well as LDL particle size and distribution of cholesterol in LDL-c and HDL-c subfractions in obese youth.

Methods

Fifteen (8 female; 7 male) obese (CDC based BMI-percentile=96.3±4.4) Latino adolescents (15.0±1.0 years) completed a 12-week lifestyle intervention (91% attendance, 3/18 withdrawals), described in detail elsewhere.(13) Briefly, weekly nutrition education classes were delivered to adolescents and their families along with 180-minutes (3 days/week for 60-minutes) of structured and unstructured physical activity at a target heart rate of 150 beats/min.

In addition to standard anthropometric measures, fasting (>10hrs) serum was collected at baseline and post-intervention (within 24-48 hours of last session) to evaluate changes in total cholesterol (TC), LDL-c, HDL-c, triglycerides (TG), and LDL particle size and distribution of cholesterol in LDL and HDL subfractions. TC, LDL-c, HDL-c, and TG concentrations were determined using an automated analyzer (Cobas c 111; Roche Diagnostics Corporation; Indianapolis, IN). Lipoprotein particles (LDL or HDL) were separated by polyacrylamide tube gel electrophoresis using the Lipoprint system (Quantimetrix Co., Redondo Beach, CA). The amount of cholesterol in each of the bands containing lipoprotein subfractions of different size was quantified by densitometry using the Lipoware software (Quantimetrix Co., Redondo Beach, CA) to estimate the proportion of cholesterol in each of the subfractions. Data were analyzed using SPSS 20.0 and are presented as means±SD. Baseline to post-intervention changes were evaluated by paired sample t-test (p 0.05). The Arizona State University Institutional Review Board approved the study, and all participants and a parent/guardian provided written informed consent prior to enrollment.

Results

Baseline and post-intervention anthropometric measurements along with fasting serum lipid concentrations are presented in Table 1. Improvements were observed for TC (-20 mg/dL; 13.1% decrease), LDL-c (-15 mg/dL; 15.5% decrease), and TG (-43 mg/dL; 30.5% decrease). Table 2 presents changes in the distribution of cholesterol in different lipoprotein subfractions in response to the intervention. Significant improvements were observed in mean LDL-particle size (2.4Å, 0.9% increase), the amount of TC in small LDL subfractions (62.5% decrease), and the amount of HDL-c in large HDL subfractions (19.6% increase) and small HDL subfractions (18.1% decrease).

Discussion

This is the first study to demonstrate that lifestyle intervention can improve LDL particle size and distribution of cholesterol in LDL and HDL subfractions independent of weight-loss in obese Latino youth. Coupled with changes in traditional lipoprotein measurements, these improvements represent a favorable shift in the atherogenic phenotype and suggest that lifestyle interventions (i.e. changes in dietary patterns and increases in physical activity) can improve early markers of atherosclerosis in youth. Furthermore, weight-loss may not be necessary for improvements in these novel markers of cardiovascular health.

Evidence suggests that lipoprotein particle size and the distribution of cholesterol in LDL and HDL subfractions may be better predictors of CVD than traditional lipid profiles.(8, 9, 11, 14, 15) Since, traditional lipoprotein levels were in the normal range at baseline (Table 1), it is possible that improvements in lipoprotein particle size and distribution following diet and exercise may contribute to CVD risk reduction above and beyond levels of and changes in traditional lipoprotein levels.(10, 11) In adults, the cardioprotective properties of both exercise and diet are related, in part, to favorable shifts in the distribution and size of cholesterol but may be weight-loss dependent(16). We did not observe changes in bodyweight and relatively small decreases in BMI-percentile and waist circumference. Moreover, neither change in BMI-percentile nor waist circumference was correlated with changes in any CVD risk factor (data not shown). Regardless, it is plausible that changes in body composition and / or fat distribution may contribute to the observed improvements independent of weight loss. Alternatively, the observed improvements may be related to the decreases in dietary fat consumption and sedentary time along with increases in cardiorespiratory fitness that were observed.(13) These findings have important implications for prevention programs for obese adolescents and are supported by previous work among obese youth showing that improvements in insulin sensitivity following exercise are independent of changes in adiposity.(17) It is noteworthy that, weight loss in obese youth has also been associated with improvements in cholesterol distribution among lipoprotein subclasses.(18) Therefore, obesity-dependent and independent mechanisms underlying improvements in lipoprotein particle size and distribution may be operative.

In children, LDL particle size is associated with traditional markers of CVD risk, including TG and HDL-c.(19) In the present investigation, changes in LDL particle size were observed in the absence of changes in total HDL-c (although redistribution of cholesterol in HDL was noted) and unrelated to changes in TG. In support of our data, Azadbakht et al,(20) found that reductions in LDL particle size using an NCEP II diet in dyslipidemic adolescents were unrelated to changes in TG or HDL-c. However, Ferguson et al,(21) reported that changes in LDL particle size following exercise in obese prepubertal children were correlated with changes in TG. Interestingly, a subsequent study by this same group suggested that changes in both LDL size and TG following exercise training are intensity dependent.(22) More work is needed to dissect how intervention parameters such as dietary modification and/or exercise intensity may influence lipid and lipoprotein metabolism in obese youth. In addition, studies should attempt to elucidate the mechanisms underlying the heterogeneity in response to lifestyle intervention as not all youth respond similarly. These studies will help to optimize lifestyle intervention delivery within targeted populations.

Our study has several strengths including a translational approach focused on a vulnerable population during a critical developmental period. Despite these strengths, there are limitations that are worthy of comment. The pilot nature of the intervention necessitated a relatively small sample that did not incorporate a control group. However, given the compelling evidence highlighting cardiovascular risk among obese Latino youth,(23) the relatively few intervention studies targeting this population, and the ethical considerations for enrolling obese youth into studies that are likely to improve health, we believe our findings extend the available science and represent an important contribution to the current literature. Additional limitations include the relative short-duration of the lifestyle intervention, lack of information on pubertal status, and the inability to determine the mechanisms underlying the observed improvements. Further, since this was a comprehensive lifestyle intervention, we were unable to determine the individual impact of and degree to which diet and/or exercise changes influenced lipoprotein profiles. Lastly, although participants did not lose weight, changes in BMI, BMI percentile, and waist circumference were observed.(13) These changes could reflect significant improvements in

body composition and/or fat distribution that were not assessed. These limitations need to be addressed in future studies.

Conclusion

These findings suggest lifestyle modification alters the lipoprotein profile in obese Latino youth to a less atherogenic phenotype. Randomized controlled trials that incorporate long-term outcome data are needed to determine the lasting effects of lifestyle intervention on future cardiovascular risk.

Acknowledgments

We would like to thank all of the children who participated in this study. Additionally, we acknowledge Ginger Hook, RN for her assistance with the measurement of lipid concentrations and thank Dr. C. Stephen Djedjos, MD for his critical review of our manuscript. The project was funded by the ASU Southwest Interdisciplinary Research Center, an Exploratory Center of Excellence for Health Disparities Research and Training through a grant from the National Institutes of Health, National Center on Minority Health and Health Disparities (P20MD002316).

References

1. Franks PW, Hanson RL, Knowler WC, Sievers ML, Bennett PH, Looker HC. Childhood Obesity, Other Cardiovascular Risk Factors, and Premature Death. *New England Journal of Medicine*. 2010; 362:485–93. [PubMed: 20147714]
2. Lee S, Gungor N, Bacha F, Arslanian S. Insulin resistance: link to the components of the metabolic syndrome and biomarkers of endothelial dysfunction in youth. *Diabetes Care*. 2007; 30:2091–7. [PubMed: 17475936]
3. Burns SF, Lee S, Arslanian SA. In vivo insulin sensitivity and lipoprotein particle size and concentration in black and white children. *Diabetes Care*. 2009; 32:2087–93. [PubMed: 19675203]
4. Perez-Mendez O, Torres-Tamayo M, Posadas-Romero C, et al. Abnormal HDL subclasses distribution in overweight children with insulin resistance or type 2 diabetes mellitus. *Clin Chim Acta*. 2007; 376:17–22. [PubMed: 16934792]
5. Van Gaal LF, Mertens IL, De Block CE. Mechanisms linking obesity with cardiovascular disease. *Nature*. 2006; 444:875–80. [PubMed: 17167476]
6. Newman WP, Wattigney W, Berenson GS. Autopsy Studies in United States Children and Adolescents. *Ann N Y Acad Sci*. 1991; 623:16–25. [PubMed: 2042824]
7. Green DJ. Exercise training as vascular medicine: direct impacts on the vasculature in humans. *Exerc Sport Sci Rev*. 2009; 37:196–202. [PubMed: 19955869]
8. El Harchaoui K, Arsenault BJ, Franssen R, et al. High-Density Lipoprotein Particle Size and Concentration and Coronary Risk. *Annals of Internal Medicine*. 2009; 150:84–93. [PubMed: 19153411]
9. Lamarche B, Tchernof A, Moorjani S, et al. Small, Dense Low-Density Lipoprotein Particles as a Predictor of the Risk of Ischemic Heart Disease in Men: Prospective Results From the Quebec Cardiovascular Study. *Circulation*. 1997; 95:69–75. [PubMed: 8994419]
10. Cromwell WC, Otvos JD, Keyes MJ, et al. LDL Particle Number and Risk of Future Cardiovascular Disease in the Framingham Offspring Study - Implications for LDL Management. *J Clin Lipidol*. 2007; 1:583–92. [PubMed: 19657464]
11. St-Pierre AC, Cantin B, Dagenais GR, et al. Low-Density Lipoprotein Subfractions and the Long-Term Risk of Ischemic Heart Disease in Men. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2005; 25:553–9.
12. Balagopal P, de Ferranti SD, Cook S, et al. Nontraditional Risk Factors and Biomarkers for Cardiovascular Disease: Mechanistic, Research, and Clinical Considerations for Youth: A Scientific Statement From the American Heart Association. *Circulation*. 2011; 123:2749–69. [PubMed: 21555711]

13. Shaibi GQ, Konopken Y, Hoppin E, Keller CS, Ortega R, Castro FG. Effects of a Culturally Grounded Community-Based Diabetes Prevention Program for Obese Latino Adolescents. *Diabetes Educ.* 2012; 38:504–12. [PubMed: 22585870]
14. Pascot A, Lemieux I, Prud'homme D, et al. Reduced HDL particle size as an additional feature of the atherogenic dyslipidemia of abdominal obesity. *Journal of Lipid Research.* 2001; 42:2007–14. [PubMed: 11734573]
15. Vakkilainen J, Mäkimattila S, Seppälä-Lindroos A, et al. Endothelial Dysfunction in Men With Small LDL Particles. *Circulation.* 2000; 102:716–21. [PubMed: 10942737]
16. Williams PT, Krauss RM, Vranizan KM, Wood PD. Changes in lipoprotein subfractions during diet-induced and exercise-induced weight loss in moderately overweight men. *Circulation.* 1990; 81:1293–304. [PubMed: 2317911]
17. Bell LM, Watts K, Siafarikas A, et al. Exercise Alone Reduces Insulin Resistance in Obese Children Independently of Changes in Body Composition. *J Clin Endocrinol Metab.* 2007; 92:4230–5. [PubMed: 17698905]
18. King F, Hobkirk J, Cooke B, Radley D, Gately J. Low-Density Lipoprotein Sub-Fraction Profiles in Obese Children Before and After Attending a Residential Weight Loss Intervention. *Journal of Atherosclerosis and Thrombosis.* 2008; 15:100–7. [PubMed: 18391473]
19. Kojima M, Kanno H, Yamazaki Y, Koyama S, Kanazawa S, Arisaka O. Association of low-density lipoprotein particle size distribution and cardiovascular risk factors in children. *Acta Pædiatrica.* 2005; 94:281–6.
20. Azadbakht L, Mirmiran P, Hedayati M, Esmailzadeh A, Shiva N, Azizi F. Particle size of LDL is affected by the National Cholesterol Education Program (NCEP) step II diet in dyslipidaemic adolescents. *British Journal of Nutrition.* 2007; 98:134–9. [PubMed: 17445337]
21. Ferguson MA, Gutin B, Le NA, et al. Effects of exercise training and its cessation on components of the insulin resistance syndrome in obese children. *Int J Obes Relat Metab Disord.* 1999; 23:889–95. [PubMed: 10490792]
22. Kang HS, Gutin B, Barbeau P, et al. Physical training improves insulin resistance syndrome markers in obese adolescents. *Med Sci Sports Exerc.* 2002; 34:1920–7. [PubMed: 12471297]
23. Johnson WD, Kroon JJM, Greenway FL, Bouchard C, Ryan D, Katzmarzyk PT. Prevalence of Risk Factors for Metabolic Syndrome in Adolescents: National Health and Nutrition Examination Survey (NHANES), 2001–2006. *Arch Pediatr Adolesc Med.* 2009; 163:371–7. [PubMed: 19349567]

Anthropometric measurements, blood pressure, fasting glucose and insulin, and fasting plasma lipids among youth participating in a lifestyle intervention.

Table 1

	Baseline	Range	Post-intervention	Range	P
Height (cm)	165.8 (12.1)	151.5-187.6	166.2 (12.2)	151.6-188.5	0.05
Weight (kg)	90.6 (26.2)	57.7-155.0	89.9 (27.8)	55.2-155.7	0.44
Waist Circumference (cm)	107.0 (16.7)	84.0-138.7	103.1 (19.5)	76.3-140.5	0.02
BMI (kg/m ²)	32.5 (6.3)	25.0-47.4	32.0 (6.7)	24.0-46.9	0.08
BMI percentile	96.3 (4.3)	85.9-99.9	95.0 (6.0)	82.2-99.9	0.02
SBP (mmHg)	122 (10)	108-138	118 (9)	100-136	0.03
DBP (mmHg)	71 (9)	51-88	68 (9)	55-89	0.18
Fasting Glucose (mg/dL)	89 (8)	76-105	90 (8)	77-110	0.60
Fasting Insulin (μU/L)	19 (10)	7-48	15 (6)	7-29	0.06
TC (mg/dl)	153 (27)	110-197	133 (31)	72-194	< 0.001
LDL-C (mg/dl)	90 (24)	55-128	76 (25)	40-131	< 0.001
HDL-C (mg/dl)	40 (7)	31-55	39 (7)	23-48	0.61
TG (mg/dl)	141 (55)	58-257	98 (38)	46-159	< 0.01

Data are presented as Mean (SD)

Abbreviations: BMI, body mass index; TG, triglycerides; TC, total cholesterol; HDL, high-density lipoprotein; LDL, low-density lipoprotein; SBP, systolic blood pressure; DBP, diastolic blood pressure

Table 2

Baseline and post-intervention LDL particle size and cholesterol distribution in lipoprotein subfractions among youth participating in a lifestyle intervention

	Baseline	Range	Post-intervention	Range	P
Mean LDL-C particle size (Å)	269.2 (3.4)	262-274	271.6 (2.9)	264-275	<0.001
Large HDL-C (%area)	22.4 (11.2)	10.3-46.2	26.8 (10.6)	13.3-46.4	<0.01
Small HDL-C (%area)	23.2 (9.4)	9.2-36.8	19.0 (6.8)	6.4-27.8	0.05
Large LDL-C (%area)*	25.8 (3.7)	17.8-31.4	24.3 (3.4)	19-31.1	0.11
Small LDL-C (%area) [†]	1.6 (2.0)	0-6.7	0.6 (1.2)	0-4.6	<0.01

Data presented as Means (SD)

Abbreviations: LDL-C, low-density lipoprotein cholesterol; HDL-C, high-density lipoprotein cholesterol; %area represents % of total cholesterol in each subfraction

* Large LDL-C is a combination of the 2 largest subfraction

[†] Small LDL-C is a combination of the 5 smallest subfraction