Published in final edited form as:

Biol Psychiatry. 2014 April 1; 75(7): 534–541. doi:10.1016/j.biopsych.2013.06.003.

Genetics of Alzheimer's disease in Caribbean Hispanic and African American Populations

Christiane Reitz^{1,2,3} and Richard Mayeux^{1,2,3,4,5}

¹Taub Institute for Research on Alzheimer's Disease and the Aging Brain, College of Physicians and Surgeons, Columbia University, New York, NY

²Department of Neurology, College of Physicians and Surgeons, Columbia University, New York, NY

³Gertrude H. Sergievsky Center, College of Physicians and Surgeons, Columbia University, New York, NY

⁴Department of Epidemiology, Joseph P. Mailman School of Public Health, Columbia University, New York, NY

⁵Department of Psychiatry, College of Physicians and Surgeons, Columbia University, New York, NY

Abstract

Late-onset Alzheimer's disease, which is characterized by progressive deterioration in cognition, function and behavior, is the most common cause of dementia and the sixth leading cause of all deaths placing a considerable burden on western societies. Most studies aiming to identify genetic susceptibility factors for LOAD have focused on non-Hispanic white populations. This is, in part related to differences in linkage disequilibrium and allele frequencies between ethnic groups that could lead to confounding. However, in addition, non-Hispanic white populations are simply more widely studied. As a consequence, minorities are genetically under-represented despite the fact that in several minority populations living in the same community as Whites (including African American and Caribbean Hispanics) LOAD incidence is higher. This review summarizes the current knowledge on genetic risk factors associated with LOAD risk in Caribbean Hispanics and African Americans and provides suggestions for future research. We focus on Caribbean Hispanics and African Americans as they have a high LOAD incidence and a body of genetic studies on LOAD that is based on samples with GWAS data and reasonable large effect sizes to yield generalizable results.

Keywords

Alzheimer's disease; genetics; gene; Caribbean Hispanic; African American; minorities

© 2013 Society of Biological Psychiatry. Published by Elsevier Inc. All rights reserved.

Address correspondence to: Richard Mayeux, MD, Gertrude H. Sergievsky Center, 630 West 168th Street, Columbia University, New York, NY 10032., Phone: 212-305-3192, Fax: 212-305-2518, rpm2@columbia.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Disclosure Statement

None of the authors have any actual or potential conflict of interest.

INTRODUCTION

Late-onset Alzheimer's disease (LOAD) places a considerable burden on western societies. LOAD is the most common cause of dementia increasing in frequency from 1% at age 65 years to more than 30% for people older than 80 years (1), and the fifth leading cause of death in persons aged 65 years. To date, an estimated 5.4 million Americans have LOAD but the prevalence in 2050 is expected to amount 11–16 million patients (2).

Senile plaques (SPs) and neurofibrillary tangles (NFTs) are considered the key pathological hallmarks of AD. The identification of β -amyloid ($A\beta$) in SPs, and genetic studies that identified mutations in the Amyloid Precursor Protein (*APP*)(3, 4), Presenilin 1 (*PSEN1*)(5) and Presenilin 1 (*PSEN2*) genes (5, 6) leading to the accumulation of $A\beta$ and early-onset familial dementia, resulted in the formulation of the "Amyloid Cascade Hypothesis". According to this hypothesis, the deposition of $A\beta$ is the initial pathological trigger in the disease, which subsequently leads to the formation of NFTs, neuronal cell death and dementia. While there is considerable evidence supporting this hypothesis, there are observations that seem to be inconsistent. First, SPs and NFTs may be reactive products resulting from neurodegeneration in AD rather than being its cause and, second, it remains unclear whether and how the deposition of $A\beta$ leads to the formation of NFTs.

It is clear that in non-Hispanic whites of European ancestry as much as 20% of the population attributable risk of LOAD is related to the $\epsilon 4$ variant in *APOE* (7–9). A series of large genome wide association studies (GWAS) identified several additional variants that affect disease susceptibility in non-Hispanic whites including *CRI1*, *CLU*, *PICALM*, *BINI1*, *CD2AP*, *CD33*, *EPHA1*, *MS4A6A/MS4E4*, *SORL1* and *ABCA7* (10–13). In addition, *SORCS1* was identified as a susceptibility gene in candidate gene and functional studies (14), and a rare variant in *TREM2* was identified in two recent sequencing studies (15, 16). In summary, these variants point to three distinct pathways, namely lipid metabolism, inflammation and endocytosis/intracellular trafficking. However, LOAD heritability estimates are high ($h^2 \approx 60$ –80%) and a large part of the genetic contribution to LOAD in this ethnic group remains unexplained (17–20).

Most genetic association studies have focused on non-Hispanic white populations because there are differences in linkage disequilibrium (LD) and allele frequencies between ethnic groups, which lead to genetic background noise and the likelihood of false-positive findings due to confounding. In addition, there is a paucity of datasets with appropriate genotyping and phenotyping in minority groups. As a consequence, ethnic groups other than non-Hispanic Whites are genetically understudied despite the fact that in several minority populations living in the same community as Whites LOAD incidence is higher (21). In addition, the reported LOAD risk associated with *APOE*- $\epsilon 4$ heterozygosity is inconsistent in most of these ethnic groups (22). This review summarizes the current knowledge on genetic risk factors associated with LOAD risk in Caribbean Hispanics and African Americans and provides suggestions for future research. We are focusing on these two ethnic groups as they are the best studied minority groups with high LOAD incidence that have GWAS data and large enough sample sizes to reliably detect risk loci. We first discuss the epidemiology of LOAD and role of *APOE* genotype in both ethnic groups followed by a separate discussion on genetic studies performed in either ethnic group outside the *APOE* locus.

EPIDEMIOLOGY OF LOAD IN AFRICAN AMERICANS AND CARIBBEAN HISPANICS

African Americans are 2–4 times and Caribbean Hispanics twice as likely as non-Hispanic whites to have LOAD (21, 23). Although differences in LOAD etiology across populations

have been widely studied, they are still poorly understood. The occurrence of multiple demented individuals in African American and Caribbean Hispanic families is significantly higher than in white families, although the genetic risk of LOAD is similar (24). While comparisons of risk across ethnic groups are complicated by differences in assessment of cognitive decline across studies and by population differences in willingness to participate in medical research, the increased risk in these specific ethnic groups may be a result of higher rates of risk factors such as poor education, cardio-and cerebrovascular disease and the metabolic syndrome (23).

APOE REGION AND RISK OF LOAD IN AFRICAN AMERICANS AND CARIBBEAN HISPANICS

In non-Hispanic Whites, the strongest susceptibility gene for LOAD is *APOE*, a lipid-binding protein expressed in humans as three common isoforms coded for by three alleles, *APOE*ε2, ε3, and ε4. The first reports linking *APOE* with LOAD found a significant increase in the *APOE*ε4 allele frequency in White patients with the disease. The large body of epidemiologic data that subsequently accumulated in cohorts of Whites supported this notion by demonstrating that *APOE*ε4 decreases the age-at-onset of LOAD in this ethnic group in a gene dosage-dependent manner (25–34), and that *APOE*ε4 is associated with lower cognitive performance, in particular the memory domain. It is thought that in non-Hispanic Whites *APOE* may account for as much as 20–50% of LOAD risk (7, 35). It is important to note that calculation of population attributable risk (PAR) is specific for a genetic factor and does not allow conclusions for other genetic variants meaning that the sum of all other PAR's can exceed 100%.

In vitro studies have indicated that the *APOE*-ε4 isoform binds Aβ peptides with a higher avidity compared to *APOE*-ε3 (36). Furthermore, there is a strong correlation between the presence of an *APOE*-ε4 allele and a higher Aβ burden in the brains of LOAD patients (37, 38), suggesting that *APOE* interacts with Aβ in enhancing its deposition in plaques. This is supported by the observation that homozygous *APOE* knockout (*APOE* -/-) mice develop fewer and more diffuse, non-fibrillar Aβ deposits (39–41). Some but not all studies assessing the effect of different *APOE* isoforms on Aβ fibrillization showed that the ε4 isoform leads to increased Aβ aggregation in vitro (42, 43). Similarly, in vivo studies in *APOE* -/- mice indicated that Aβ fibrillization and plaques formation was increased in mice expressing human *APOE*-ε4 (APPV717F+/-, apo E-/-) compared to mice not expressing human *APOE* (44, 45). Still, it is possible that *APOE* exerts its effects through different mechanisms, e.g. *APOE* is a major cholesterol transporter and high cholesterol levels have been associated with an increased Aβ load in animal models (46, 47) and changes in APP processing (48, 49). Thus *APOE* isoform-specific changes in cholesterol binding and transport in brain might also affect plaque formation in LOAD brains.

As described above, in African Americans and Caribbean Hispanics the reported LOAD risk associated with the *APOE*-ε4 allele is inconsistent (18–20, 22). While in several studies number of copies of the ε4 allele were not associated with risk or age-of onset of LOAD (18, 20, 22, 50), other studies observed such effect (19). The disparity could be due to recruitment bias, differences in age distribution, sample size, population stratification or differences in residual confounding through environmental or cultural factors. The largest GWAS performed to date in African Americans strongly suggests an increased risk of LOAD for *APOE*ε4 carriers (51). Roses et al.(52) previously reported an association between a variable length poly-T polymorphism ("poly-T") at rs10524523 in the gene encoding the channel-forming subunit of the translocase of the mitochondrial outer membrane (*TOMM40*) and risk for LOAD and age of onset (AAO) of LOAD in a small sample of non-Hispanic Whites (n=34). Subsequently, the same group assessed both the '523' allele frequencies of

this polymorphism and their linkage pattern with *APOE* (which resides in the same region on chromosome 19) and reported associations in non-Hispanic Whites and other ethnic groups. However, a more recent study of this polymorphism in a much larger sample of non-Hispanic Whites failed to confirm the original findings after adjusting for the effect of *APOE* $\epsilon 4$ (53). In addition, the Alzheimer's Disease Genetics Consortium (ADGC) showed in a large sample of over 22,000 White subjects that *APOE* alleles $\epsilon 2$, $\epsilon 3$, and $\epsilon 4$ account for essentially all the inherited risk of LOAD associated with the *APOE* region and that other variants including the poly-T track in *TOMM40* are not independent risk or AAO loci (54). While no additional large-scale studies have reassessed this issue in other ethnic groups, it is likely that, due to the lesser extent of LD in African Americans and Caribbean Hispanics compared to Whites, this is also true for these ethnic groups.

GENETIC STUDIES IN CARIBBEAN HISPANICS OUTSIDE THE *APOE* REGION

Family-based linkage studies

Multiple genome wide linkage studies (GWLS) for LOAD were published between 1997 and 2006 and most were performed on white populations. While some chromosomal regions have been studied and replicated extensively using linkage (most notably chromosomes 9, 10, and 12) (55–58), no consistently replicated LOAD gene has yet been identified using this method. There are several reasons for these limited results including the generally small datasets, the inability of the then available molecular genotyping technologies to capture all the segregation information in the families and the sensitivity of linkage studies to underlying locus heterogeneity when using datasets consisting of a large number of small families. However, the inability to conclusively identify causal genes within these regions supports the possibility that multiple rare variants could be involved in AD risk in these families.

In linkage analyses of in 79 Caribbean Hispanic multiplex LOAD families from the participating in the Estudio Familiar de Influencia Genetica de Alzheimer (EFIGA) study using 35 microsatellite markers near the centromere of chromosome 12 Mayeux et al (59), observed modest evidence of linkage with support for D12S1623 and D12S1042. Linkage varied by age at onset of LOAD and by the presence or absence of the *APOE*-epsilon 4 allele. In larger follow-up studies, first in 490 individuals from 96 Caribbean Hispanic families using 340 microsatellite markers (60), and then in 1,075 individuals from 209 families (61, 62), Lee et al. obtained support for linkage on 3q28, 10q26, 12p12–13, and 18q21, some of which had also been repeatedly reported by linkage or case-control studies in Whites (in particular 10q and 12p) or Amish (18q) on LOAD (55, 60, 63–70). All these regions include candidate genes that may be biologically plausible but still remain to be confirmed by sequencing and functional studies. Finally, the same group observed a small effect of the alpha-2 macroglobulin deletion/insertion polymorphism on familial LOAD risk. alpha-2 macroglobulin is a proteinase inhibitor which binds β -amyloid peptide and prevents fibril formation (71).

Heritability of LOAD endophenotypes

Johnson et al. and Lee et al. explored the heritability of several cognitive endophenotypes of LOAD in Caribbean Hispanics and observed high heritability for memory performance (delayed recall (h^2) = 0.60); delayed recognition (h^2) = 0.41)) and abstract reasoning (h^2) = 32.6%)(72, 73).

Genome-wide association (GWAS) and candidate gene studies

Table 1 summarizes the GWAS and candidate gene studies performed. In a community-based case-control candidate-gene study on several ethnic groups which included 372 Caribbean Hispanic individuals, Lee et al.(74) identified multiple LOAD-associated SNPs and haplotypes in the 5' and 3' ends of the L(DLR class) A repeats-containing sortilin-related receptor (*SORL1*) that were associated in African Americans. *SORL1* is a cargo molecule of the retromer complex and involved in trafficking of APP, and underexpression of *SORL1* leads to overproduction of A β (75). The variants identified include SNPs 12 and 26 (rs12285364 and rs1784933, respectively) and the TTC haplotype at SNPs 23 through 25 (rs3824968, rs2282649, rs1010159), which was significantly associated with LOAD in the North European white individuals in previous reports (75, 76).

Lee et al.(77) also performed the largest GWAS to date in Caribbean Hispanics. The study included 549 cases and 544 controls originating from the Dominican Republic and Puerto Rico who are part of the Washington Heights-Inwood Columbia Aging Project (WHICAP) and the Estudio Familiar de Influencia Genetica de Alzheimer (EFIGA) family study. While the strongest support was observed for rs9945493 on 18q23, the study identified 5 SNPs that could subsequently be replicated in an independent Caucasian validation dataset and are located near genes or regions that could be biologically relevant to LOAD and/or have also been reported other studies performed in Caucasians including SNPs on 2p25.1, 3q25.2, 7p21.1, 10q23.1 containing *HPCAL1*, *DGKB*, *GHITM*, *C10orf99*, *PCDH21*, *LRIT2*, *LRIT1* and *RGR*. In addition, the study replicated *CLU*, *PICALM*, and *BINI*. The effect sizes for all observed variants were modest ($0.33 < OR < .87$).

In the same Caribbean Hispanic dataset, Reitz et al. explored the association between variants in the Fat and Obesity Associated (*FTO*) gene and risk of LOAD (78). The authors identified three SNPs (rs17219084, rs11075996, rs11075997) that were associated with LOAD, consistent with independent studies in non-Hispanic Whites showing that polymorphisms in the *FTO* gene have robust effects on obesity, obesity-related traits and endophenotypes associated with LOAD(79–82). In the same cohort, the authors explored the association between SNPs in leucine-rich repeat transmembrane 3 (*LRRTM3*) gene and LOAD (83). *LRRTM3* is a neuronal gene-promoting amyloid precursor protein (APP) processing by β -secretase 1 thereby modulating the levels of A β 40 and A β 42. In addition, *LRRTM3* is nested in the alpha-3 catenin gene (*CTNNA3*) on chromosome 10q22.2 that in turn binds presenilin 1. Four SNPs belonging to two distinct LD blocks and including one promoter SNP were associated (rs16923760, rs1925608, rs7082306, rs1925609) as were their corresponding haplotypes. In functional analyses *LRRTM3* knockdown with small-hairpin RNAs caused a significant decrease in amyloid precursor protein processing compared with the scrambled small-hairpin RNA condition consistent with the notion that that *LRRTM3* may modulate γ -secretase processing of APP(83).

A study by Ghani et al.(84) that conducted a genome-wide scan for large copy number variation (CNV) in this dataset observed a nominal association between LOAD and a ~470 kb duplication on chromosome 15q11.2 which encompasses up to five genes (*TUBGCP5*, *CYFIP1*, *NIPA2*, *NIPA1*, and *WHAMML1*) and was present in 10 cases and 3 controls. The dosage increase of the *CYFIP1* and *NIPA1* genes was further confirmed by quantitative PCR. Both genes are interesting LOAD candidate genes. *NIPA1* encodes a magnesium transporter associated with early endosomes in neuronal and epithelial cells; *CYFIP1* forms a complex at synapses with the fragile X mental retardation protein (FMRP) and eIF4E (FMRP-CYFIP1-eIF4E complex). FMRP acts as an APP translation repressor releasing CYFIP1 from the FMRP-CYFIP1-eIF4E complex in response to synaptic stimulation and unbalanced dosage of CYFIP1 might result in altered APP turnover in AD patients. The study did not detect CNVs (including common variants) affecting the well-confirmed LOAD

loci reported by large GWAS in Non-Hispanic Whites (*CLU*, *PICALM*, *BIN1*, *CRI*, *MS4A4/MS4A6E*, *CD2AP*, *CD33*, *EPHA1*, and *ABCA7*). However, this could be explained by analytical challenges in the detection of common CNVs from SNPintensity data. In general, a case-control setting can only test common CNVs that cluster and are well-tagged by common SNPs. These CNVs could be of a multiallelic or complex nature (*e.g.* a small deletion within a large CNV duplication) and can only be accurately genotyped using a combination of custom arrays and deep sequencing. Finally, in a subset of this dataset that had information on plasma A β 40 and 42 levels (160 cases, 294 controls), Reitz et al.(85) explored the association of genetic variation in the *IDE-KIF11-HHEX* complex with LOAD. Out of 32 SNPs in this region, 3 SNPs (rs2421943, rs12264682, rs11187060) were associated with plasma A β 40 or A β 42 levels in single marker and haplotype analyses after correction for multiple testing. All these SNPs lie within the same LD block, and are in LD with haplotypes previously reported in Whites(86, 87). IDE binds and degrades A β 40 and A β 42 and this A β degrading activity has been shown to be lower in LOAD brains than in controls (88). Consistent with this notion, in *IDE* knock-out mice, brain A β levels are elevated(89). Polymorphisms in *IDE* may also contribute to the risk of type 2 diabetes(90) which itself is associated with LOAD. Taken together the findings reported here support the possibility that the *IDE-KIF11-HHEX* region on chromosome 10q may contain genetic variants modifying A β 40 and 42 levels.

Sequencing studies

While a few sequencing studies of individual polymorphisms associated with the early-onset form of the disease have been performed in individual Caribbean Hispanic families(91, 92), no targeted candidate gene sequencing or whole exome or genome sequencing studies have been conducted. Reasons for this include both the previous lack of high-throughput sequencing technology that was only developed over the past five years, and the paucity of appropriate datasets.

GENETIC STUDIES IN AFRICAN AMERICANS OUTSIDE THE APOE REGION

GWAS and candidate gene studies

While no family-based linkage or sequencing studies on LOAD have been performed in African Americans outside the *APOE* region, several GWAS and candidate gene studies were conducted (Table 1). Logue et al.(93) analyzed a genome-wide set of 2.5 million imputed markers in 513 well-characterized African American LOAD cases and 496 cognitively normal controls collected from multiple sites as part of the Multi-Institutional Research on Alzheimer Genetic Epidemiology (MIRAGE) Study and the Henry Ford Health System as part of the Genetic and Environmental Risk Factors for Alzheimer Disease Among African Americans (GenerAAtions) Study. The analyses identified rs6859 in *PVRL2* as a novel susceptibility locus and replicated *CLU*, *PICALM*, *BIN1*, *EPHA1*, *MS4A*, *ABCA7*, and *CD33* as susceptibility genes, although the effect direction for some SNPs and the most significant SNPs in these genes differed from findings of the White datasets(11–13).

The largest GWAS to date on LOAD in African Americans(51) was performed by the Alzheimer's Disease Genetics consortium (ADGC) and included 5,896 subjects (1,968 cases and 3,928 controls) that were collected from multiple sites within the US. The top-ranked SNP observed in this study is located in *ABCA7* (rs115550680) and notably has an effect size that is nearly as strong as the effect size of *APOE* ϵ 4 (70–80% increase in risk). This observation clearly differs from the GWAS performed in Whites in which the reported *ABCA7* SNPs (rs3752246; rs3764650) but also the SNPs in all other reported genes (*CRI*, *BIN1*, *PICALM*, *CLU*, *EPHA1*, *MS4A* cluster, *CD2AP*, *CD33*) have significantly lower effect sizes (~10–20% increase in risk)(11–13). It remains possible that this could be due to

population differences in the frequencies of the causative variant(s) tagged by the associated SNPs or the result of a bias in the estimated effect of a newly identified allele on disease (also termed “winner’s curse”). However, it is also possible that the large difference in the effect size of the *ABCA7* locus on the risk of LOAD is explained by population-specific causative variants with variable impact on protein structure or function. The LD block in which rs115550680 is located spans across several introns and exons which implies that rs115550680 is in LD with exonic variants that could be potentially causative. The study by Reitz et al. also replicated *CRI*, *BINI*, *EPHA1* and *CD33* with significance in gene-based analyses and effect sizes similar to those in Whites although with different disease-associated top SNPs(51). These differences in associated SNPs not allowing direct comparison between specific SNPs were expected due to the differences in minor allele frequencies or linkage disequilibrium patterns between the ethnic groups.

In addition to these GWAS studies, several studies on specific LOAD candidate genes were performed in African Americans (table 1). In their candidate-gene study on *SOLRL1* Lee et al.(74) also explored the associations of *SORL1* SNPs and haplotypes with LOAD in 246 African American individuals and observed several disease-associated SNPs and haplotypes in the 5’ and 3’ ends. Burgess et al.(94) genotyped 119 cases and 252 controls for 15 SNPs in the kidney and brain expressed protein (*KIBRA*) which included rs17070145 previously reported to be associated with better episodic memory performance in Whites(95). Consistent with these earlier reports, the authors found a significantly reduced risk associated with the T allele of rs17070145 in the older African-American subjects ($p = 0.007$) but there was no association with episodic memory in control subjects. *KIBRA* interacts with a multitude of proteins involved in synaptic function, cell polarity, vesicular transport, and neuronal plasticity. It is expressed in memory-related structures of the brain and has increased expression in laser-capture microdissected neurons from the hippocampus, middle temporal gyrus, and posterior cingulate of LOAD cases in comparison with controls. Akomolafe et al. assessed the effect of the Glu298Asp variant of the endothelial nitric oxide synthase (*NOS3*) gene, which is involved in oxidative stress, on LOAD in 467 sibships and unrelated controls in the MIRAGE African American dataset, and observed an increased risk for carriers of the GG genotype. Oxidative stress accelerates degenerative changes including those leading to LOAD via β -amyloid/lipid interactions and can also lead to hypertension, ischemic heart disease and other cardiovascular diseases that indirectly contribute to progression of LOAD. In the same dataset, several SNPs and haplotypes in the Paraoxonase (*PON*) gene cluster (*PON1*, *PON2*, *PON3*) were associated with LOAD(96). Paraoxonase is an arylesterase enzyme that is expressed in the liver and found in the circulation in association with apoA1 and the high-density lipoprotein, and prevents the accumulation of oxidized lipids in low-density lipoproteins in vitro. Common polymorphisms in genes encoding paraoxonase are established risk factors in a variety of vascular disorders including coronary artery disease and carotid artery stenosis. As described above, genetic, epidemiological, autopsy and neuroimaging studies suggest that vascular disease increases risk of LOAD.

Genetics of LOAD endophenotypes

Benke et al.(97) explored the association of genetic variants in interleukin-1 genes with cognition in the Cardiovascular Health Study and observed significant association of SNPs in the *IL1B* gene with baseline performance on the 3MS. Cuenco et al.(98) explored the effect of 16 SNPs in transthyretin (*TTR*), which inhibits the production of the amyloid β protein, with LOAD risk and measures of neurodegeneration and cerebrovascular disease defined by magnetic resonance imaging in African American sibships. They observed a marginal effect of two SNPs significant in Caucasian sibships with hippocampal atrophy. Melville et al.(99) conducted a 2-stage GWAS in non-Hispanic White and African

Americans for measures of hippocampal volume, total cerebral volume and white matter hyperintensities. They attained genome-wide significant associations for hippocampal volume with SNPs in the APOE, F5/SELP, LHFP and GCFC2 gene regions in both ethnic groups. Except for APOE all these genes remain to be confirmed by independent African American datasets or functional studies.

CONCLUSIONS

Genetic differences in LOAD risk alleles across populations have not been studied sufficiently. In general, there is a paucity of datasets with appropriate phenotyping and genotyping, and most of the few studies that have these measures available are due to small sample sizes limited in their ability to detect the small effect sizes expected for this complex disease. Studies aiming to replicate the specific loci identified in non-Hispanic whites are furthermore often hampered by cross-population differences in allele frequencies of the identified SNPs and differences in linkage disequilibrium patterns.

As a consequence, loci recently identified in non-Hispanic whites may not modify LOAD risk in other ethnic groups; similarly it remains unclear whether there are population-specific causative variants. The few studies performed in African Americans and Caribbean Hispanics suggest that these ethnic groups share some but not all LOAD susceptibility loci with whites. The linkage studies performed in Caribbean Hispanics further support the possibility that multiple rare variants could be involved in LOAD risk in multiplex LOAD families in this ethnic group. Whether the effect of these variants is modified by epistatic effects or environmental factors remains unclear. Although essential for further elucidating the pathogenic mechanism underlying LOAD these questions can – due to limited statistical power – likely not sufficiently be answered by the currently available datasets.

Currently, large-scale whole exome and whole genome sequencing efforts are under way aiming to identify additional common and rare variants associated with LOAD. These efforts include Whites and Caribbean Hispanics, and while variants identified by this effort will need to be functionally confirmed, these studies hold the promise to lead to a more accurate understanding of the genetic risk factors in these ethnic groups, which in turn can be incorporated in diagnostic and predictive testing protocols and help to identify novel targets for prevention and treatment. Similar efforts are needed for African Americans and additional ethnic groups that have a high prevalence of LOAD but have been widely neglected by genomic LOAD research.

Acknowledgments

This work was supported by grants from the National Institute of Health and the National Institute on Aging: R37-AG15473, P01-AG07232, The Blanchett Hooker Rockefeller Foundation, The Charles S. Robertson Gift from the Banbury Fund, and The Merrill Lynch Foundation. Dr. Reitz was further supported by a Paul B. Beeson Career Development Award (K23AG034550).

REFERENCES

1. Fratiglioni L, De Ronchi D, Aguero-Torres H. Worldwide prevalence and incidence of dementia. *Drugs Aging*. 1999; 15(5):365–375. [PubMed: 10600044]
2. Hebert LE, Weuve J, Scherr PA, Evans DA. Alzheimer disease in the United States (2010–2050) estimated using the 2010 census. *Neurology*. 2013; 80(19):1778–1783. [PubMed: 23390181]
3. Goate A. Segregation of a missense mutation in the amyloid beta-protein precursor gene with familial Alzheimer's disease. *J Alzheimers Dis*. 2006; 9(3 Suppl):341–347. [PubMed: 16914872]

4. Goate A, Chartier-Harlin MC, Mullan M, Brown J, Crawford F, Fidani L, et al. Segregation of a missense mutation in the amyloid precursor protein gene with familial Alzheimer's disease. *Nature*. 1991; 349(6311):704–706. [PubMed: 1671712]
5. Sherrington R, Rogaev EI, Liang Y, Rogaeva EA, Levesque G, Ikeda M, et al. Cloning of a gene bearing missense mutations in early-onset familial Alzheimer's disease. *Nature*. 1995; 375(6534):754–760. [PubMed: 7596406]
6. Levy-Lahad E, Wasco W, Poorkaj P, Romano DM, Oshima J, Pettingell WH, et al. Candidate gene for the chromosome 1 familial Alzheimer's disease locus. *Science*. 1995; 269(5226):973–977. [PubMed: 7638622]
7. Slioter AJ, Cruts M, Kalmijn S, Hofman A, Breteler MM, Van Broeckhoven C, et al. Risk estimates of dementia by apolipoprotein E genotypes from a population-based incidence study: the Rotterdam Study. *Arch Neurol*. 1998; 55(7):964–968. [PubMed: 9678314]
8. Farrer LA, Cupples LA, Haines JL, Hyman B, Kukull WA, Mayeux R, et al. Effects of age, sex, and ethnicity on the association between apolipoprotein E genotype and Alzheimer disease. A meta-analysis. APOE and Alzheimer Disease Meta Analysis Consortium. *JAMA*. 1997; 278(16):1349–1356. [PubMed: 9343467]
9. Ward A, Crean S, Mercaldi CJ, Collins JM, Boyd D, Cook MN, et al. Prevalence of apolipoprotein E4 genotype and homozygotes (APOE e4/4) among patients diagnosed with Alzheimer's disease: a systematic review and meta-analysis. *Neuroepidemiology*. 2012; 38(1):1–17. [PubMed: 22179327]
10. Lambert JC, Heath S, Even G, Campion D, Sleegers K, Hiltunen M, et al. Genome-wide association study identifies variants at CLU and CR1 associated with Alzheimer's disease. *Nat Genet*. 2009; 41(10):1094–1099. [PubMed: 19734903]
11. Hollingworth P, Harold D, Sims R, Gerrish A, Lambert JC, Carrasquillo MM, et al. Common variants at ABCA7, MS4A6A/MS4A4E, EPHA1, CD33 and CD2AP are associated with Alzheimer's disease. *Nat Genet*. 2011; 43(5):429–435. [PubMed: 21460840]
12. Seshadri S, Fitzpatrick AL, Ikram MA, DeStefano AL, Gudnason V, Boada M, et al. Genome-wide analysis of genetic loci associated with Alzheimer disease. *JAMA*. 2010; 303(18):1832–1840. [PubMed: 20460622]
13. Naj AC, Jun G, Beecham GW, Wang LS, Vardarajan BN, Buross J, et al. Common variants at MS4A4/MS4A6E, CD2AP, CD33 and EPHA1 are associated with late-onset Alzheimer's disease. *Nat Genet*. 2011; 43(5):436–441. [PubMed: 21460841]
14. Reitz C, Tokuhiro S, Clark LN, Conrad C, Vonsattel JP, Hazrati LN, et al. SORCS1 alters amyloid precursor protein processing and variants may increase Alzheimer's disease risk. *Ann Neurol*. 2011; 69(1):47–64. [PubMed: 21280075]
15. Guerreiro R, Wojtas A, Bras J, Carrasquillo M, Rogaeva E, Majounie E, et al. TREM2 variants in Alzheimer's disease. *N Engl J Med*. 2013; 368(2):117–127. [PubMed: 23150934]
16. Jonsson T, Stefansson H, Steinberg S, Jonsdottir I, Jonsson PV, Snaedal J, et al. Variant of TREM2 associated with the risk of Alzheimer's disease. *N Engl J Med*. 2013; 368(2):107–116. [PubMed: 23150908]
17. Gatz M, Pedersen NL, Berg S, Johansson B, Johansson K, Mortimer JA, et al. Heritability for Alzheimer's disease: The study of dementia in Swedish twins. *Journals of Gerontology Series A - Biological Sciences and Medical Sciences*. 1997; 52(2):M117–M125.
18. Evans DA, Bennett DA, Wilson RS, Bienias JL, Morris MC, Scherr PA, et al. Incidence of Alzheimer disease in a biracial urban community: relation to apolipoprotein E allele status. *Arch Neurol*. 2003; 60(2):185–189. [PubMed: 12580702]
19. Graff-Radford NR, Green RC, Go RC, Hutton ML, Edeki T, Bachman D, et al. Association between apolipoprotein E genotype and Alzheimer disease in African American subjects. *Arch Neurol*. 2002; 59(4):594–600. [PubMed: 11939894]
20. Murrell JR, Price B, Lane KA, Baiyewu O, Gureje O, Ogunniyi A, et al. Association of apolipoprotein E genotype and Alzheimer disease in African Americans. *Arch Neurol*. 2006; 63(3):431–434. [PubMed: 16533971]
21. Tang MX, Cross P, Andrews H, Jacobs DM, Small S, Bell K, et al. Incidence of AD in African-Americans, Caribbean Hispanics, and Caucasians in northern Manhattan. *Neurology*. 2001; 56(1):49–56. [PubMed: 11148235]

22. Tang MX, Stern Y, Marder K, Bell K, Gurland B, Lantigua R, et al. The APOE-epsilon4 allele and the risk of Alzheimer disease among African Americans, whites, and Hispanics. *JAMA*. 1998; 279(10):751–755. [PubMed: 9508150]
23. 2012 Alzheimer's disease facts and figures. *Alzheimers Dement*. 2012; 8(2):131–168. [PubMed: 22404854]
24. Green RC, Cupples LA, Go R, Benke KS, Edeki T, Griffith PA, et al. Risk of dementia among white and African American relatives of patients with Alzheimer disease. *JAMA*. 2002; 287(3): 329–336. [PubMed: 11790212]
25. Breitner JC, Wyse BW, Anthony JC, Welsh-Bohmer KA, Steffens DC, Norton MC, et al. APOE-epsilon4 count predicts age when prevalence of AD increases, then declines: the Cache County Study. *Neurology*. 1999; 53(2):321–331. [PubMed: 10430421]
26. Corder EH, Saunders AM, Strittmatter WJ, Schmechel DE, Gaskell PC, Small GW, et al. Gene dose of apolipoprotein E type 4 allele and the risk of Alzheimer's disease in late onset families. *Science*. 1993; 261(5123):921–923. [PubMed: 8346443]
27. Gomez-Isla T, West HL, Rebeck GW, Harr SD, Growdon JH, Locascio JJ, et al. Clinical and pathological correlates of apolipoprotein E epsilon 4 in Alzheimer's disease. *Ann Neurol*. 1996; 39(1):62–70. [PubMed: 8572669]
28. Holmes C, Levy R, McLoughlin DM, Powell JF, Lovestone S. Apolipoprotein E: non-cognitive symptoms and cognitive decline in late onset Alzheimer's disease. *J Neurol Neurosurg Psychiatry*. 1996; 61(6):580–583. [PubMed: 8971103]
29. Hyman BT, Gomez-Isla T, Rebeck GW, Briggs M, Chung H, West HL, et al. Epidemiological, clinical, and neuropathological study of apolipoprotein E genotype in Alzheimer's disease. *Ann N Y Acad Sci*. 1996; 802:1–5. [PubMed: 8993478]
30. Kurz A, Altland K, Lautenschlager N, Zimmer R, Busch R, Gerundt I, et al. Apolipoprotein E type 4 allele and Alzheimer's disease: effect on age at onset and relative risk in different age groups. *J Neurol*. 1996; 243(6):452–456. [PubMed: 8803817]
31. Murman DL, Foster NL, Kilgore SP, McDonagh CA, Fink JK. Apolipoprotein E and Alzheimer's disease: strength of association is related to age at onset. *Dementia*. 1996; 7(5):251–255. [PubMed: 8872415]
32. Poirier J, Davignon J, Bouthillier D, Kogan S, Bertrand P, Gauthier S. Apolipoprotein E polymorphism and Alzheimer's disease. *Lancet*. 1993; 342(8873):697–699. [PubMed: 8103819]
33. Roses AD. Alzheimer's disease: the genetics of risk. *Hosp Pract (Minneap)*. 1997; 32(7):51–55. 58–63, 67–59.
34. Tang MX, Maestre G, Tsai WY, Liu XH, Feng L, Chung WY, et al. Relative risk of Alzheimer disease and age-at-onset distributions, based on APOE genotypes among elderly African Americans, Caucasians, and Hispanics in New York City. *Am J Hum Genet*. 1996; 58(3):574–584. [PubMed: 8644717]
35. Ashford JW, Mortimer JA. Non-familial Alzheimer's disease is mainly due to genetic factors. *J Alzheimers Dis*. 2002; 4(3):169–177. [PubMed: 12226536]
36. Strittmatter WJ, Weisgraber KH, Huang DY, Dong LM, Salvesen GS, Pericak-Vance M, et al. Binding of human apolipoprotein E to synthetic amyloid beta peptide: isoform-specific effects and implications for late-onset Alzheimer disease. *Proc Natl Acad Sci U S A*. 1993; 90(17):8098–8102. [PubMed: 8367470]
37. Schmechel DE, Saunders AM, Strittmatter WJ, Crain BJ, Hulette CM, Joo SH, et al. Increased amyloid beta-peptide deposition in cerebral cortex as a consequence of apolipoprotein E genotype in late-onset Alzheimer disease. *Proc Natl Acad Sci U S A*. 1993; 90(20):9649–9653. [PubMed: 8415756]
38. Rebeck GW, Reiter JS, Strickland DK, Hyman BT. Apolipoprotein E in sporadic Alzheimer's disease: allelic variation and receptor interactions. *Neuron*. 1993; 11(4):575–580. [PubMed: 8398148]
39. Bales KR, Verina T, Cummins DJ, Du Y, Dodel RC, Saura J, et al. Apolipoprotein E is essential for amyloid deposition in the APP(V717F) transgenic mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A*. 1999; 96(26):15233–15238. [PubMed: 10611368]

40. Bales KR, Verina T, Dodel RC, Du Y, Altstiel L, Bender M, et al. Lack of apolipoprotein E dramatically reduces amyloid beta-peptide deposition. *Nat Genet.* 1997; 17(3):263–264. [PubMed: 9354781]
41. Kindy MS, Rader DJ. Reduction in amyloid A amyloid formation in apolipoprotein-E-deficient mice. *Am J Pathol.* 1998; 152(5):1387–1395. [PubMed: 9588907]
42. Ma J, Yee A, Brewer HB Jr, Das S, Potter H. Amyloid-associated proteins alpha 1-antichymotrypsin and apolipoprotein E promote assembly of Alzheimer beta-protein into filaments. *Nature.* 1994; 372(6501):92–94. [PubMed: 7969426]
43. Sanan DA, Weisgraber KH, Russell SJ, Mahley RW, Huang D, Saunders A, et al. Apolipoprotein E associates with beta amyloid peptide of Alzheimer's disease to form novel monofibrils. Isoform apoE4 associates more efficiently than apoE3. *J Clin Invest.* 1994; 94(2):860–869. [PubMed: 8040342]
44. Holtzman DM, Bales KR, Tenkova T, Fagan AM, Parsadanian M, Sartorius LJ, et al. Apolipoprotein E isoform-dependent amyloid deposition and neuritic degeneration in a mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A.* 2000; 97(6):2892–2897. [PubMed: 10694577]
45. Holtzman DM, Bales KR, Wu S, Bhat P, Parsadanian M, Fagan AM, et al. Expression of human apolipoprotein E reduces amyloid-beta deposition in a mouse model of Alzheimer's disease. *J Clin Invest.* 1999; 103(6):R15–R21. [PubMed: 10079115]
46. Sparks DL, Scheff SW, Hunsaker JC 3rd, Liu H, Landers T, Gross DR. Induction of Alzheimer-like beta-amyloid immunoreactivity in the brains of rabbits with dietary cholesterol. *Exp Neurol.* 1994; 126(1):88–94. [PubMed: 8157129]
47. Refolo LM, Malester B, LaFrancois J, Bryant-Thomas T, Wang R, Tint GS, et al. Hypercholesterolemia accelerates the Alzheimer's amyloid pathology in a transgenic mouse model. *Neurobiol Dis.* 2000; 7(4):321–331. [PubMed: 10964604]
48. Bodovitz S, Klein WL. Cholesterol modulates alpha-secretase cleavage of amyloid precursor protein. *J Biol Chem.* 1996; 271(8):4436–4440. [PubMed: 8626795]
49. Howland DS, Trusko SP, Savage MJ, Reaume AG, Lang DM, Hirsch JD, et al. Modulation of secreted beta-amyloid precursor protein and amyloid beta-peptide in brain by cholesterol. *J Biol Chem.* 1998; 273(26):16576–16582. [PubMed: 9632729]
50. Tycko B, Lee JH, Ciappa A, Saxena A, Li CM, Feng L, et al. APOE and APOC1 promoter polymorphisms and the risk of Alzheimer disease in African American and Caribbean Hispanic individuals. *Arch Neurol.* 2004; 61(9):1434–1439. [PubMed: 15364690]
51. Reitz C, Jun G, Naj A, Rajbhandary R, Vardarajan BN, Wang LS, et al. Variants in the ATP-binding cassette transporter (ABCA7), apolipoprotein E 4, and the risk of late-onset Alzheimer disease in African Americans. *JAMA.* 2013; 309(14):1483–1492. [PubMed: 23571587]
52. Roses AD, Lutz MW, Amrine-Madsen H, Saunders AM, Crenshaw DG, Sundseth SS, et al. A TOMM40 variable-length polymorphism predicts the age of late-onset Alzheimer's disease. *Pharmacogenomics J.* 2010; 10(5):375–384. [PubMed: 20029386]
53. Cruchaga C, Nowotny P, Kauwe JS, Ridge PG, Mayo K, Bertelsen S, et al. Association and expression analyses with single-nucleotide polymorphisms in TOMM40 in Alzheimer disease. *Arch Neurol.* 2011; 68(8):1013–1019. [PubMed: 21825236]
54. Jun G, Vardarajan BN, Buross J, Yu CE, Hawk MV, Dombroski BA, et al. Comprehensive search for Alzheimer disease susceptibility loci in the APOE region. *Arch Neurol.* 2012; 69(10):1270–1279. [PubMed: 22869155]
55. Pericak-Vance MA, Bass MP, Yamaoka LH, Gaskell PC, Scott WK, Terwedow HA, et al. Complete genomic screen in late-onset familial Alzheimer disease. Evidence for a new locus on chromosome 12. *JAMA.* 1997; 278(15):1237–1241. [PubMed: 9333264]
56. Rogava E, Premkumar S, Song Y, Sorbi S, Brindle N, Paterson A, et al. Evidence for an Alzheimer disease susceptibility locus on chromosome 12 and for further locus heterogeneity. *JAMA.* 1998; 280(7):614–618. [PubMed: 9718052]
57. Scott WK, Grubber JM, Conneally PM, Small GW, Hulette CM, Rosenberg CK, et al. Fine mapping of the chromosome 12 late-onset Alzheimer disease locus: potential genetic and phenotypic heterogeneity. *Am J Hum Genet.* 2000; 66(3):922–932. [PubMed: 10712207]

58. Scott WK, Grubber JM, Abou-Donia SM, Church TD, Saunders AM, Roses AD, et al. Further evidence linking late-onset Alzheimer disease with chromosome 12. *JAMA*. 1999; 281(6):513–514. [PubMed: 10022104]
59. Mayeux R, Lee JH, Romas SN, Mayo D, Santana V, Williamson J, et al. Chromosome-12 mapping of late-onset Alzheimer disease among Caribbean Hispanics. *Am J Hum Genet*. 2002; 70(1):237–243. [PubMed: 11715112]
60. Lee JH, Mayeux R, Mayo D, Mo J, Santana V, Williamson J, et al. Fine mapping of 10q and 18q for familial Alzheimer's disease in Caribbean Hispanics. *Mol Psychiatry*. 2004; 9(11):1042–1051. [PubMed: 15241431]
61. Lee JH, Cheng R, Rogaeva E, Meng Y, Stern Y, Santana V, et al. Further examination of the candidate genes in chromosome 12p13 locus for late-onset Alzheimer disease. *Neurogenetics*. 2008; 9(2):127–138. [PubMed: 18340469]
62. Lee JH, Cheng R, Santana V, Williamson J, Lantigua R, Medrano M, et al. Expanded genomewide scan implicates a novel locus at 3q28 among Caribbean hispanics with familial Alzheimer disease. *Arch Neurol*. 2006; 63(11):1591–1598. [PubMed: 17101828]
63. Hahs DW, McCauley JL, Crunk AE, McFarland LL, Gaskell PC, Jiang L, et al. A genome-wide linkage analysis of dementia in the Amish. *Am J Med Genet B Neuropsychiatr Genet*. 2006; 141B(2):160–166. [PubMed: 16389594]
64. Hiltunen M, Mannerman A, Thompson D, Easton D, Pirskanen M, Helisalmi S, et al. Genome-wide linkage disequilibrium mapping of late-onset Alzheimer's disease in Finland. *Neurology*. 2001; 57(9):1663–1668. [PubMed: 11706108]
65. Kehoe P, Wavrant-De Vrieze F, Crook R, Wu WS, Holmans P, Fenton I, et al. A full genome scan for late onset Alzheimer's disease. *Hum Mol Genet*. 1999; 8(2):237–245. [PubMed: 9931331]
66. Pericak-Vance MA, Grubber J, Bailey LR, Hedges D, West S, Santoro L, et al. Identification of novel genes in late-onset Alzheimer's disease. *Exp Gerontol*. 2000; 35(9-10):1343–1352. [PubMed: 11113612]
67. Wu WS, Holmans P, Wavrant-DeVrieze F, Shears S, Kehoe P, Crook R, et al. Genetic studies on chromosome 12 in late-onset Alzheimer disease. *JAMA*. 1998; 280(7):619–622. [PubMed: 9718053]
68. Myers A, Holmans P, Marshall H, Kwon J, Meyer D, Ramic D, et al. Susceptibility locus for Alzheimer's disease on chromosome 10. *Science*. 2000; 290(5500):2304–2305. [PubMed: 11125144]
69. Ertekin-Taner N, Graff-Radford N, Younkin LH, Eckman C, Baker M, Adamson J, et al. Linkage of plasma Aβ₄₂ to a quantitative locus on chromosome 10 in late-onset Alzheimer's disease pedigrees. *Science*. 2000; 290(5500):2303–2304. [PubMed: 11125143]
70. Bertram L, Blacker D, Mullin K, Keeney D, Jones J, Basu S, et al. Evidence for genetic linkage of Alzheimer's disease to chromosome 10q. *Science*. 2000; 290(5500):2302–2303. [PubMed: 11125142]
71. Hughes SR, Khorkova O, Goyal S, Knaeblein J, Heroux J, Riedel NG, et al. Alpha2-macroglobulin associates with beta-amyloid peptide and prevents fibril formation. *Proc Natl Acad Sci U S A*. 1998; 95(6):3275–3280. [PubMed: 9501253]
72. Johnson B, Santana V, Schupf N, Tang MX, Stern Y, Mayeux R, et al. The heritability of abstract reasoning in Caribbean Latinos with familial Alzheimer disease. *Dement Geriatr Cogn Disord*. 2007; 24(6):411–417. [PubMed: 17938569]
73. Lee JH, Flaquer A, Stern Y, Tycko B, Mayeux R. Genetic influences on memory performance in familial Alzheimer disease. *Neurology*. 2004; 62(3):414–421. [PubMed: 14872023]
74. Lee JH, Cheng R, Schupf N, Manly J, Lantigua R, Stern Y, et al. The association between genetic variants in SORL1 and Alzheimer disease in an urban, multiethnic, community-based cohort. *Arch Neurol*. 2007; 64(4):501–506. [PubMed: 17420311]
75. Rogaeva E, Meng Y, Lee JH, Gu Y, Kawarai T, Zou F, et al. The neuronal sortilin-related receptor SORL1 is genetically associated with Alzheimer disease. *Nat Genet*. 2007; 39(2):168–177. [PubMed: 17220890]

76. Reitz C, Cheng R, Rogaeva E, Lee JH, Tokuhira S, Zou F, et al. Meta-analysis of the association between variants in SORL1 and Alzheimer disease. *Arch Neurol*. 2011; 68(1):99–106. [PubMed: 21220680]
77. Lee JH, Cheng R, Barral S, Reitz C, Medrano M, Lantigua R, et al. Identification of novel loci for Alzheimer disease and replication of CLU, PICALM, and BIN1 in Caribbean Hispanic individuals. *Arch Neurol*. 2011; 68(3):320–328. [PubMed: 21059989]
78. Reitz C, Tosto G, Mayeux R, Luchsinger JA. Genetic variants in the Fat and Obesity Associated (FTO) gene and risk of Alzheimer's disease. *PLoS One*. 2012; 7(12):e50354. [PubMed: 23251365]
79. Dina C, Meyre D, Gallina S, Durand E, Korner A, Jacobson P, et al. Variation in FTO contributes to childhood obesity and severe adult obesity. *Nat Genet*. 2007; 39(6):724–726. [PubMed: 17496892]
80. Hertel JK, Johansson S, Sonestedt E, Jonsson A, Lie RT, Platou CG, et al. FTO, type 2 diabetes, and weight gain throughout adult life: a meta-analysis of 41,504 subjects from the Scandinavian HUNT, MDC, MPP studies. *Diabetes*. 2011; 60(5):1637–1644. [PubMed: 21398525]
81. Wang K, Li WD, Zhang CK, Wang Z, Glessner JT, Grant SF, et al. A genome-wide association study on obesity and obesity-related traits. *PLoS One*. 2011; 6(4):e18939. [PubMed: 21552555]
82. Ho AJ, Stein JL, Hua X, Lee S, Hibar DP, Leow AD, et al. A commonly carried allele of the obesity-related FTO gene is associated with reduced brain volume in the healthy elderly. *Proc Natl Acad Sci U S A*. 2010; 107(18):8404–8409. [PubMed: 20404173]
83. Reitz C, Conrad C, Roszkowski K, Rogers RS, Mayeux R. Effect of genetic variation in LRRTM3 on risk of Alzheimer disease. *Arch Neurol*. 2012; 69(7):894–900. [PubMed: 22393166]
84. Ghani M, Pinto D, Lee JH, Grinberg Y, Sato C, Moreno D, et al. Genome-wide survey of large rare copy number variants in Alzheimer's disease among Caribbean hispanics. *G3 (Bethesda)*. 2012; 2(1):71–78. [PubMed: 22384383]
85. Reitz C, Cheng R, Schupf N, Lee JH, Mehta PD, Rogaeva E, et al. Association between variants in IDE-KIF11-HHEX and plasma amyloid beta levels. *Neurobiol Aging*. 2012; 33(1):199, e113–e197. [PubMed: 20724036]
86. Prince JA, Feuk L, Gu HF, Johansson B, Gatz M, Blennow K, et al. Genetic variation in a haplotype block spanning IDE influences Alzheimer disease. *Hum Mutat*. 2003; 22(5):363–371. [PubMed: 14517947]
87. Ertekin-Taner N, Allen M, Fadale D, Scanlin L, Younkin L, Petersen RC, et al. Genetic variants in a haplotype block spanning IDE are significantly associated with plasma Abeta42 levels and risk for Alzheimer disease. *Hum Mutat*. 2004; 23(4):334–342. [PubMed: 15024728]
88. Perez A, Morelli L, Cresto JC, Castano EM. Degradation of soluble amyloid beta-peptides 1–40, 1–42, and the Dutch variant 1–40Q by insulin degrading enzyme from Alzheimer disease and control brains. *Neurochem Res*. 2000; 25(2):247–255. [PubMed: 10786709]
89. Farris W, Mansourian S, Chang Y, Lindsley L, Eckman EA, Frosch MP, et al. Insulin-degrading enzyme regulates the levels of insulin, amyloid beta-protein, and the beta-amyloid precursor protein intracellular domain in vivo. *Proc Natl Acad Sci U S A*. 2003; 100(7):4162–4167. [PubMed: 12634421]
90. Rudovich N, Pivovarova O, Fisher E, Fischer-Rosinsky A, Spranger J, Mohlig M, et al. Polymorphisms within insulin-degrading enzyme (IDE) gene determine insulin metabolism and risk of type 2 diabetes. *J Mol Med (Berl)*. 2009; 87(11):1145–1151. [PubMed: 19809796]
91. Athan ES, Williamson J, Ciappa A, Santana V, Romas SN, Lee JH, et al. A founder mutation in presenilin 1 causing early-onset Alzheimer disease in unrelated Caribbean Hispanic families. *JAMA*. 2001; 286(18):2257–2263. [PubMed: 11710891]
92. Bertoli Avella AM, Marcheco Teruel B, Llibre Rodriguez JJ, Gomez Viera N, Borrajerio Martinez I, Severijnen EA, et al. A novel presenilin 1 mutation (L174 M) in a large Cuban family with early onset Alzheimer disease. *Neurogenetics*. 2002; 4(2):97–104. [PubMed: 12484344]
93. Logue MW, Schu M, Vardarajan BN, Buross J, Green RC, Go RC, et al. A comprehensive genetic association study of Alzheimer disease in African Americans. *Arch Neurol*. 2011; 68(12):1569–1579. [PubMed: 22159054]

94. Burgess JD, Pedraza O, Graff-Radford NR, Hirpa M, Zou F, Miles R, et al. Association of common KIBRA variants with episodic memory and AD risk. *Neurobiol Aging*. 2011; 32(3):557, e551–e559. [PubMed: 21185624]
95. Papassotiropoulos A, Stephan DA, Huentelman MJ, Hoernkli FJ, Craig DW, Pearson JV, et al. Common Kibra alleles are associated with human memory performance. *Science*. 2006; 314(5798):475–478. [PubMed: 17053149]
96. Erlich PM, Lunetta KL, Cupples LA, Huyck M, Green RC, Baldwin CT, et al. Polymorphisms in the PON gene cluster are associated with Alzheimer disease. *Hum Mol Genet*. 2006; 15(1):77–85. [PubMed: 16319130]
97. Benke KS, Carlson MC, Doan BQ, Walston JD, Xue QL, Reiner AP, et al. The association of genetic variants in interleukin-1 genes with cognition: findings from the cardiovascular health study. *Exp Gerontol*. 2011; 46(12):1010–1019. [PubMed: 21968104]
98. Cuenco KT, Friedland R, Baldwin CT, Guo J, Vardarajan B, Lunetta KL, et al. Association of TTR polymorphisms with hippocampal atrophy in Alzheimer disease families. *Neurobiol Aging*. 2011; 32(2):249–256. [PubMed: 19328595]
99. Melville SA, Buross J, Parrado AR, Vardarajan B, Logue MW, Shen L, et al. Multiple loci influencing hippocampal degeneration identified by genome scan. *Ann Neurol*. 2012; 72(1):65–75. [PubMed: 22745009]

Table 1
Genome-wide association and candidate gene studies on LOAD performed in Caribbean Hispanics and African Americans

Study	Source Population(s)	case s (n)	contr ols (n)	Age rang e	Phenotype	Genotyping	Finding (disease-associated SNP, Allele)
Caribbean Hispanics							
Lee et al. (2007)	WHICAP	178	194	> 65 years	LOAD (NINCDS-ARDRA criteria)	29 <i>SORL1</i> SNPs	rs12285364 (T allele, ↑) associated with LOAD
Lee et al. (2011)	WHICAP, EFICA	549	544	> 65 years	LOAD (NINCDS-ARDRA criteria)	Illumina 650Y	rs9945493 on 18q23; rs4669573 and rs10197851 on 2p25.1; rs11711889 on 3q25.2; rs117750 on 7p21.1; and rs7908652 on 10q23.1; CLU (rs881146), PICALM (rs17159904), BIN1 (rs7561528)
Reitz et al. (2012)	WHICAP, EFICA	549	544	> 65 years	LOAD (NINCDS-ARDRA criteria)	Illumina 650Y	3 SNPs (rs17219084 (G allele, ↑), rs11075996 (T allele, ↑), rs11075997 (T allele, ↑)) and corresponding haplotypes in FTO associated with LOAD
Reitz et al. (2012)	WHICAP, EFICA	549	544	> 65 years	LOAD (NINCDS-ARDRA criteria)	Illumina 650Y	4 SNPs (rs16923760 (C allele, ↓), rs1925608 (C allele, ↓), rs7082306 (A allele, ↓), rs1925609 (T allele, ↑) and corresponding haplotypes in LRRTM3 associated with LOAD
Ghani et al. (2012)	WHICAP, EFICA	549	544	> 65 years	LOAD (NINCDS-ARDRA criteria)	Illumina 650Y	470 kb duplication on chromosome 15q11.2 with dosage increase of CYFIP1 and NIPA1 associated with LOAD
Reitz et al. (2012)	WHICAP, EFICA	160	294	> 65 years	plasma Aβ40 and Aβ42 levels	Illumina 650Y	3 SNPs in IDE (rs2421943 (A allele, ↑), rs12264682 (A allele, ↓), rs1187060 (T allele, ↓)) associated with plasma Aβ40 or Aβ42 levels
African Americans							
Lee et al. (2007)	WHICAP	88	158	> 65 years	LOAD (NINCDS-ARDRA criteria)	29 <i>SORL1</i> SNPs	rs12285364 (T allele, ↓), rs1784933(G allele, ↑) associated with LOAD
Logue et al. (2011)	MIRAGE, GenerAAtion s	513	496	> 65 years	LOAD (NINCDS-ARDRA criteria)	Illumina 610 Quad, Illumina 370 Duo	Various SNPs in APOE, PVRL2, CLU, PICALM, BIN1, EPHA1, MS4A, ABCA7, CD33 associated with LOAD **
Reitz et al. (2013)	multiple sites	1,968	3,928	> 65 years	LOAD (NINCDS-ARDRA criteria)	multiple Illumina chips	APOE, ABCA7, CRI1, BIN1, EPHA1, CD33 associated with LOAD **
Burgess et al. (2011)	Mayo Clinic, Jacksonville	119	252	> 60 years	LOAD (NINCDS-ARDRA criteria)	15 <i>KIBRA</i> SNPs	significantly reduced LOAD risk for rs17070145 T allele in the older subjects
Akomolafe et al. (2006)	MIRAGE	241	226	>65 years	LOAD (NINCDS-ARDRA criteria)*	11 <i>NOS3</i> SNPs	Glu298 allele associated with higher LOAD risk

Study	Source Population(s)	case s (n)	contr ols (n)	Age rang e	Phenotype	Genotyping	Finding (disease-associated SNP, Allele)
Erich et al. (2006)	MIRAGE	241	226	>65 years	LOAD (NINCDS-ARDRA criteria)*	29 SNPs in <i>PON1</i> gene cluster	8 SNPs and several haplotypes associated with increased LOAD including <i>PON1</i> promoter SNP-161[C/T]

* control status based on modified Telephone Interview of Cognitive Status

** disease-associated alleles differ compared to Whites