Published in final edited form as:

Vaccine. 2009 February 5; 27(6): 833–845. doi:10.1016/j.vaccine.2008.11.109.

A discrete computer model of the Immune System reveals competitive interactions between the humoral and cellular branch and between cross-reacting memory and naïve responses

Yiming Cheng¹, Dario Gherzi¹, Claudia Calcagno¹, Liisa K. Selin², Roberto Puzone³, and Franco Celada^{1,4,*}

¹Division of Rheumatology, Department of Medicine, New York University School of Medicine, New York, NY, USA

²Department of Pathology, University of Massachusetts Medical School, Worcester, MA

³IST, National Institute for Cancer Research, Genova, Italy

⁴DOBIG Department, University of Genova, Genova, Italy

Abstract

In an agent-based computer model, we simulate the formation and recall of anti-virus immunological memory. Specifically we try to predict what will happen, both to the response and to memory, when the second infecting virus is partly different from the first one, and when the cross-reactivity of the two branches of the immune system, humoral and cellular, is asymmetrical, or “split”. The simulations explore systematically epitope distances, and measure all changes in affinity, cellularity and efficiency in clearing the infection. Besides obvious cooperations, they reveal powerful competitions between the branches, and more intriguing, between cross-reacting and new responses when the latter suffer the competition by preformed cell-rich but inefficient clones, as memory, usually an asset, becomes a liability.

Keywords

Viral Cross-reactivity; Cellular/Humoral Response; Agent-based Computer Model

1. Introduction

The wealth of data, ranging from the molecular to the organism level gathered by the very successful “reductionist” approach of modern immunologists has resulted in accrued knowledge of all components of the immune system (IS), whose complexity at all levels appears to have grown concomitantly. Thus, to translate the progress into understanding of the functioning principles and the logic of the interactions of parts, a multidisciplinary effort may be helpful. Systems biologists and immunologists agree in integrating advanced

© 2013 Published by Elsevier Ltd.

*Corresponding author at: Hospital for Joint Disease, Room 1611, 301 East 17th street, New York, NY 10003 USA. Tel. +1 212 598 6507, Fax: +1 212 598 7604. fc15@nyu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

technologies (such as imaging) and computational modelling with experimental data, and their efforts on defined sections of molecular activation and cellular interactions have yielded understandable images and interpretations [1].

Many models of the immune system have been proposed in the last fifteen years, each understandably focused on sections of the system and most falling into one of the two categories: *continuous* or *discrete*. Continuous models usually represent the dynamic of the several entities involved in the immune response in the form of differential equations: their solution yields an “average” or “typical” response, and its faithfulness to the real biologic reality depends on the chosen parameter values, and the simplifications and assumptions introduced to make the simulated phenomenon mathematically solvable. On the other hand, discrete models seldom reach general quantitative conclusions; however, being intrinsically apt to handle local dynamics, they fill an important complementary area of modelling. They are very dependent on computing power, but what appeared to be a severe limitation has been overcome by the rapidly increasing availability of computing capacity at low cost, which favours the applications of these kinds of models in any scientific field.

“Agent-based” discrete models are inspired by Von Neumann “cellular automata” [2] which in turn may be considered equivalent to mathematical calculations. They consist of discrete dimensional entities (two/three dimensional space and time in discrete steps), where the agents are the relevant cells (or molecules) equipped with virtual receptors and capabilities, which reflect the experimental observations. Both continuous and discrete models can be useful in unravelling complexity by representing and simulating hypotheses; their virtual results may stimulate new theories that will have to be tested via new experiments *in vivo*. Both kinds face the danger of becoming unreal and self-referential. One way to dodge this risk is to systematically confront the evolving biological consensus and - avoiding falling in love with one’s creature - be willing to modify the equation or the code to accommodate any new data or emerging hypothesis.

IMMSIM [3] has been conceived to allow the dynamic representation of hypotheses and their preliminary testing. This in turn may elicit new ideas and hypotheses to be eventually tested *in vivo*. In several applications over recent years, the model has generated emergent and sometimes surprising data, which allowed shedding light on the mechanisms and interactions of the model itself and on their counterparts in the biological immune system. For example, during the simulation of the affinity maturation of the humoral response, the varying density of cells and availability of antigen were shown to configure the shift from the severest bottleneck of the primary response, obtaining the T help, to the secondary bottleneck, winning the competition for antigen [4]. The model also offers the possibility to manipulate the elements of virtual runs like experimental biologists do, by using the computational equivalent of knock out mice or gene transfer. Stratagems of this kind were applied in parallel experiments comparing the response of the humoral branch *only*, the cellular branch *only*, and *both* branches, in order to relate the efficiency of responses to different viral features [5]. In a study about cross-reactive memory, the silencing of one or the other of two suspected kinds of attrition (passive vs active) was used recently, and revealed interesting cooperative effects of the combined mechanisms [6]. In the present study, selective freezing of humoral cross-reactive responses is obtained by increasing the bit distance in epitopes but not in peptides, while to reveal antibody mediated competition against cellular responses, the antibody lifetime was artificially shortened or extended over a 50-fold range.

The aim of this article is to present a systematic study of homologous and heterologous responses against viruses, representing the variable molecular complementarity by using the mathematical concept of binary string space, and tackling the recently found asymmetry of

the humoral vs cellular cross-reactivity. The obvious disclaimer is that any emergent findings and novel relations must await biological validation before being accepted and applied. A more general aim is to confirm that IMMSIM modelling complements rather than competes with the continuous approaches, exemplified by De Boer et al. [7], and carry a distinct intuitive appeal.

2. Materials and Methods

2.1. The IMMSIM Model

The IMMSIM model considers discrete events (e.g., in the immune response, the encounters between single cell and molecules) inside a quantized “lattice” space. It works from these events to delineate the whole simulation, which evolves in discrete time steps. In its simple form, the model consists of a 15×16 bidimensional grid hosting different entities in variable amounts. These entities are cells, e.g., T and B lymphocytes, antigen presenting cell (APC), epithelial cells (EC) etc., and molecules such as antibodies (Ab), antigens (Ag) and cytokines; because of the emphasis on interactions, it is easiest to consider the IMMSIM space as a virtual IS organ, such as a lymph node. The entities are selected depending on the aspect of the immune response to be simulated and their quantity can be decided at the beginning of each simulation. Entities are represented by data structures storing tuneable working properties (e.g., interaction strength, clonal expansion, death, diffusion) and present individual status (e.g., position in the grid, age, activation state). Most agents can diffuse within the grid; interactions between individual agents obey probabilistic rules, and are allowed only if they “meet” within a site in the grid. Any interaction between two entities is limited to a single successful one (if any) at each time step of simulation; the time step is the common pacer in all the interactions. Since the fastest cell division take one time step, the latter can be considered equivalent to 6 hours. Antigen and antibody specific properties, and the clonal characteristics of the cells, are represented by assigning to each single entity one or more individual “receptor” in the form of an N-bit binary string. All specific interaction rules depend upon the “matching” of the two interacting actors’ receptors; the rule is a function of the binary distance of the respective N-bit binary string. Any single interaction usually results in a discrete alternative (to die or not, activate or not, divide or not, diffuse or not, etc) that is stochastically decided from the probability calculated by the interaction rule. What happens at each step depends stochastically on the contents and conditions of each site, yielding a system that can be described as a stochastic agent-based discrete model.

The model incorporates many of the consensus facts of today’s immunological knowledge, e.g., the diversity of specific elements [8], MHC restriction [9], clonal selection by antigen affinity [10], thymic education of T cells [11], antigen processing and presentation [12], cell-cell cooperation [13], maturation of the response and memory [14], all tuned in the model to balance the responses in the minimalistic scale (compared to biology) of the simulations. As the stochastic behaviours in the model is based on a random number generator, when repeating a run with the same set-up but different initial random “seed”, it will result in a similar but stochastically different response, which also simulates the repeating of a biological experiment in different syngeneic mice.

The IMMSIM simulator was conceived in late 80’s in the collaboration between the immunologist F.Celada and the astrophysicist P.Seiden; the latter wrote the first code using the IBM’s APL language. A large IBM host was needed to run the simulator, which was limited to humoral responses with small specificity repertoires; its description and applications were published in the 90’s [3]. The simulator was then expanded by adding a basic representation of the cellular responses to viruses (IMMSIM3) [5]; it could run on a “big” PC through proprietary IBM libraries. The APL’s handicap was its limited diffusion among modellers. Thus the physicist R. Puzone, starting in 1999 developed a new version of

IMMSIM in C language (IMMSIM-C), and aligned it with Seiden's. When Seiden died prematurely in 2001, IMMSIM-C, completed with an input/output interface (designed for didactic and research) and additional specific interfaces (immscinput, immplot), became the only official simulator and is maintained at Puzone's workplace IST [15]. The code, all the interfaces and also the former IMMSIM3 APL and its manuals are available at www.immsim.org.

Several independent IMMSIM versions in C language exist: among those, a parallel version (c-immsim) by P. Castiglione [16], mostly based on the original IMMSIM project, and recently applied on AIDS [17], and one developed by S. Valensin [18], which aimed at simulating aging of the Immune System. The stochastic characteristics originally introduced by the IMMSIM model, are adopted in many IS models, even the analytical ones [19]. The present simulator displays many original and enhanced features: a new memory model for all lymphocytes with particular attention to T cells, a new T cell memory model with attrition, compact and effective B hypermutation and T thymus selection, support for flexible receptor and affinity function representations, optimized diffusion, interactive run mode (also network controlled), improved random number generation, support for lymphocyte extraction during the response and adoptive transfer into new mice, with possibility of manipulating adoptive transfer before transfer, optimized fast code. The code structures allow IMMSIM-C to run rather fast and to be handled by any standard PC with minimum system resources. They can also take advantage of any parallel computational power in networked hosts, thereby facilitating the analysis of large number of virtual mice (VM).

2.2. Agents

All agents in the model, including cells and molecules, are listed as following:

2.2.1. Cells—APC --- has major histocompatibility complex class I (MHC1) and class II (MHC2) receptors. Not specific. Has low affinity for all antigens, viruses. When in proximity of a source of DANGER signal, becomes activated and is able to activate any T helper cell (Th) and any T cell receptor (TCR) specifically. B cells act as APC, since they process and present antigen reacting with its B cell receptor (BCR), but are not able to activate Th *ex novo*.

B --- has specific BCR (recognizing antigen epitope) and MHC2. When it binds antigen the latter is internalized and processed, and the hidden peptide is mounted on MHC2 receptor. The combination of a part of peptide and MHC constitutes a T-epitope that may be recognized and bound by a Th2 cell's TCR. For this interaction to be effective, the Th2 must have been previously activated by an active APC.

Th1 (T helper cell type 1) --- has specific TCR capable to bind to a APC's MHC presenting a peptide. This causes the APC to activate a Tc if the two T cells bind to the APC at the same time or within a given time lapse.

Th2 (T helper cell type 2) --- has specific TCR capable to bind to a B cell's MHC presenting a peptide. This constitutes the signal for triggering the B and Th cell's activation process (simulating the lymphokine secretion and activation effect).

Tc (cytotoxic T lymphocyte) --- has specific TCR. When naïve, it will be activated if the TCR binds specifically to a APC's MHC1 presenting a peptide. When activated or in memory state, it will be able to bind by its TCR to any specific loaded MHC on virus-infected EC. This TCR binding causes the death of the EC and the elimination of all virus particles it contains.

EC --- is the example of body cell not belonging to the IS. As such, it has only MHC1. If infected by virus, signals the presence of invader by displaying a viral peptide on the MHC1. If the virus completes the proliferation inside the cell and reaches its critical multiplicity, EC burst open, frees the viral particles in the environment and diffuses a DANGER signal.

2.2.2. Molecules—Ab --- antibody, is identical to the BCR of the B cells that secretes it when it becomes a Plasma cell. Diffuses. Binds epitope by affinity proportional to bit complementarities. If it binds to a free virus, the latter is killed. Cannot penetrate into cells, not even if they are infected.

Ag --- antigen, is composed by two binary strings (more epitopes, or more peptides can be added for *ad hoc* experiments). One string (Epitope) is accessible by specific BCR and Antibodies. The other (Peptide) is not visible, and becomes accessible only when a B cell or an APC bind the antigen, internalizes it and displays it on MHC2 after processing.

V --- virus, is an antigen with the added capabilities to penetrate ECs and begin duplication when inside. Each virus has specific Epitopes and Peptides. In addition, it has one degree, Low, Moderate, High or Very High, of the three properties that define its lifestyle: a) I, infectivity (probability of penetrating into a EC in the same site); b) S, speed (number of duplications/time step); and c), L, lethality (multiplicity of virus particles that will cause the host EC to burst).

IFN --- interferon, represents a cytokine and is the effector of active attrition. A molecule of non-defined structure is selected in controlled amounts and has controlled life time. Diffuses. Its presence causes controlled apoptosis in memory cells present in the site.

D --- Danger, is a virtual cytokine and the effector of non-specific APC activation. Selected in controlled amounts in the site where virus-caused cell death occurs, and has controlled life time and can diffuse.

2.3. Flowchart of events

All actions and events that may occur during a simulation are here described in the form of flowcharts (Fig. 1–3). The flowcharts also help placing all cells and molecules in their context within the simulation. Fig. 1 describes the virus cycle; Fig. 2 describes the humoral response; Fig. 3 describes the cellular response.

2.4. Cure of infection

Whether the infection is cured or becomes persistent or even kills the virtual mouse depends on the virus dose and the quality (determined by its combination of infectivity, multiplicity and lethal load levels). These also determine whether the IS success requires the cooperation of both the cellular and the humoral branch, as has been shown in several occasions [5].

The parameter values in Table 1 are used whenever the program applies the “sequence of events” described above. The result is the viral infection of a VM followed by a full immune response. Runs are governed by random numbers generated by the computer, since all events, such as cell/cell and cell/molecule encounters, are probabilistic. A rerun with a different set of random numbers results in a different VM and a different response, even if the same list of default values is used. The default values are based on consensus among immunologists, but are not “real”, as they reflect the model’s simplification/minimisation of the real response. They cannot be singled out, since they have been selected in relation to each other and balanced by *in machina* experiments.

2.5. Matrix

For a comprehensive study of anti-virus cross-reactive memory, we have adopted a “matrix” approach that relies on the distinctive properties of IMMSIM, to allow total access to the data regarding the kinetics of the invader and of each population of lymphoid cells during an infection. The matrix is a design by which VM are infected with a virus (such as v1 in Fig. 4(a)), then subdivided in five groups and challenged either with the original infectant or with one of the 4 new viruses differing from the original by 1, 3, 4, 6 bits in either the Epitope (which will be recognized by the B cell receptor) or the Peptide (which will be processed and then presented on MHC to the Tc receptor). Then another virus (v2) is used for the role of primary infectant, and the other 4 viruses, now differing from the first by 1, 2, 3, 5 (Fig. 4(a)) are used as challenges. The process is repeated until all combinations are run, fulfilling the checkerboard in Fig. 4(b), and the results of all combinations are averaged according to nine binary string differences (0 to 8) between first and second antigen. This design allows a balanced analysis of the effect of increasing distance among antigens on cross-reactivity. The weakest possible cross-reaction is expected to happen at distance 6 (Fig. 5), while antigens at distance 7 and 8 will act as non-related agents.

The experiment is at first run with all viral peptides differing from each other by 0, 1, 2 ... up to 8 bit mismatches associated with an identical B-epitope. Then it is re-run with B-epitopes differing from each other by 0, 1, 2 ... up to 8 bit mismatches with an identical peptide. Finally, it is run with both the B-epitopes and the peptides changing. It is repeated 5 times, each time with a different set of antigens in one virtual mouse, i.e., with the same random number. Then by changing random numbers, a total of 5 virtual mice are repeated. So for each run with two specified injections, the statistical data are based on 25 runs.

Thanks to the complete transparency afforded by the model, a large number of data derived by the matrix runs can be gathered, compared with each other, and their significance in statistical or biological terms evaluated. The default matrix is characterized by a large set of parameters (e.g. the cell number, the antigen dose, the density of elements per site). If we modify one single parameter, a new matrix is created and the subsequent run constitutes a new experiment, whose result will be compared to those of the default matrix.

The matrix approach, albeit labour intensive for the computer (each matrix is run over-night and take several hours to complete) has allowed probing the following phenomena and how they are affected by the bit distance between challenging and priming virus:

- a. The efficiency of the response
- b. The weighted average affinity, also called strength
- c. The relative recall (in terms of cell and clone numbers of B and Tc)
- d. The variation (+ or -) of repertoires (B, Tc)
- e. The simulated Tc stimulation in the absence of B memory.

2.6. Statistical significance

Genetically identical animals differ in their immune response to the same antigen because the somatic recombination of immunoglobulin genes includes random processes. The statistical treatment of data from groups of syngeneic mice is necessary to separate the fluctuations from regularities of the phenomenon. In the IMMSIM model, the individual diversity is simulated effectively by introducing a random seed that influences every interaction of the response and classic statistics is applied on repeated *in machina* simulations for the responses mentioned above.

In the case of the matrix setup, we have calculated the standard error of the mean $SE_{\bar{x}} = \delta / \sqrt{n}$, where δ is the standard deviation and n is the sample size) and have determined how many repetitions were sufficient to reach significance. The curves in Fig. 6 pertain to B cell strength towards the original antigen and show that 25 runs are proper for different setup. The data also confirms the expectation that the variation is larger in intermediate cross-reactions. In addition, an ANOVA analysis for the significance of mean differences has been conducted using F value cut-off of 3.35 or alpha value cut-off of 0.05.

3. Results

3.1. The homologous response

Repeating infections by the same virus generates a “homologous response”, which is expected to use the feature “memory” at its highest level, i.e., the characteristic ability of the Immune System to make use of information acquired during the first infection in order to improve the quality and the efficiency of the response to the second. The mechanisms involved are: a) specific clonal expansion of B and T cells; b) affinity maturation brought about, for all cell lines, by competitive selection of high affinity clones, and in the case of B cells preceded by a repertoire expansion introduced by hypermutation of the V-genes. Several detailed features are illustrated as following.

3.1.1 Kinetics and clonal composition of a humoral and cellular response—

Fig. 7 shows the events and changes occurring in a representative virtual mouse injected 3 times with the same type of virus, and the 3 successive responses mounted by the humoral and cellular defences, which in the set-up are able to rapidly cure the infection. The virus enjoys its greatest success after the first injection when there is no preformed memory in the VM and the primary response takes some time to build up. In contrast, the second and third injections are conquered rapidly. The limited production of new virus is also the cause of the limited “size” of the secondary and tertiary response; the time to reach the peak is progressively shortened both for the humoral and cellular responses. During these events, there is partial recall of formed memory and selection of high affinity clones.

3.1.2. Variation of B and Tc repertoire during three infections by the same virus—We call repertoire of the primary humoral and cellular responses (Fig. 8) the B or Tc cells from all clones stimulated by the antigen and in addition, an unspecified minority of naïve cells and cells newly created by the “bone marrow”. Thus, not all of the primary repertoire will be re-called by the homologous challenge. The secondary repertoire will contain all primary memory clones that are re-stimulated by the challenging virus, plus new clones specifically elicited. The trend continues during the third challenge and the net result is a consistent decrease of the clonal memory repertoire from the first to the third response, which can be quantified in ~45% in the humoral and ~33% in the cellular response.

3.1.3. Strength of the anti-virus homologous response—Fig. 9 shows that during

three successive infections, the weighted average affinity ($\text{strength} = \sum_i c_i 2^{a_i} / \sum_i c_i$, where i represents different clones, c_i is the peak cellularity of clone i , a_i is the affinity of clone i and defined by 4 minus the mismatch, i.e., the affinity is 1 for the maximal mismatch 3) of both cellular response increases significantly even if in the humoral response a peak may be reached at the secondary response. The availabilities of the memory repertoire decreases in both responses (Fig. 8). The opposite tendencies of affinity maturation and available repertoires suggest a progressive competitive selection of memory clones where those with higher affinity gain dominance and afford a more efficient and streamlined response.

3.2. The cross-reactive responses

Viral antigens can be recognized inside and outside infected cells, and elicit a single type of response or both humoral and cellular responses. Consequently, B and T memory cells build up and the secondary response will be composed of re-called clones and new clones in different proportions, depending on the bit distance separating the challenge virus from the original immunogen. By using the matrix organization (Fig. 4), the cross-reactive response can be defined systematically by a number of parameters.

3.2.1. Recall efficiency—Even if the secondary infection is homologous, only a significant fraction of the B clones in the memory is able to bind the antigen, find T cell help and multiply. With infecting viruses progressively distant from the immunogen, the relative recall is predictably smaller. When the distance is 7 or more, there is no structural possibility of cross-reaction (Fig. 5). Fig. 10 shows the B and Tc clonal recall (the number of re-stimulated memory clones relative to the total clonal types at the peak response after the second challenge) and the cellular recall (the number of re-stimulated memory cells relative to the total cellularity at the peak response after the second challenge).

3.2.2. The Affinity Build-up—The “strength ratio” of a response represents the average affinity of the cells weighted by the cellularities in the secondary response, relative to the primary. In the homologous response, a “strength ratio” greater than 1 represents the affinity maturation due to the competitive selection of higher affinity clones after the second challenge, where 1 is the level of primary response (Fig. 11). In cross-reactive responses, the general trend is an expected decrease of strength toward the original antigen with increasing bit distance between first immunogen and challenging antigen and this happens in both humoral and cellular responses. When strength ratio is measured toward the second challenge, the humoral response shows a progressive decrease from the peak of the homologous challenge to the level of the primary response, which is reached after distance 6. In the cellular response, the strength ratio falls under 1 at distance 2, 3 and 4 (Fig. 11 (b)), but after this dip the curve picks up, to progressively reach an asymptote of 1 with antigens 7 and 8. “1” means that these responses reach the same average affinity that the primary response had. 7 and 8 are sufficiently distant from the original immunogen as not to entertain any cross-reaction with the memory clones, and in fact, constitute “new” primary responses. The lower strength ratio observed in the intermediate cross-reactants may be caused by competition between memory cross-reacting clones and new virgin clones, hindering the development of the latter. Thus the cross-reaction does not help but hinders the strength of the response against these secondary infectants, at least in terms of “quality” of the response. This situation can be described as an “antigenic sin” effect [20] where the response to an invader is pre-empted or hindered by a cross-reacting memory.

3.2.3. The repertoire size changes during cross-reaction—After a cross-reacting challenge, the re-called memory clones are expected to decrease with the increasing antigenic distance, while the fraction of newly stimulated clones will increase, until with challenging distance 7 and 8 (differing 7 and 8 bits from the immunogen) where the entire “secondary” response will be constituted by new clones. These trends are confirmed by the results of matrix experiments. However, the quantitative data reveal a biphasic variation that could not be predicted, even if recent *in vivo* experiments of the UMass group [21–23] had hinted at a restriction of the memory repertoire after challenge with “near” or relatively similar cross-reactive epitope, in contrast with an enlargement of the repertoire after contact with a “far” or more dissimilar cross-reactive epitope [22]. Fig. 12 shows the number of primary and secondary clones in B and Tc responses, while Fig. 13 separates two phases, of relative decrease and of relative increase of the number of clones in the secondary peak, compared to the primary. The two phases show remarkably similar trends in the humoral

and the cellular memory, even if the number of clones involved differs. While it is easy to explain the shrinking on the basis of dominance by few cross-reacting memory clones, which would selectively trim the repertoire and provide a better defence, the observation that in far responses the number of clones surpasses the level of a totally new primary is unexpected. A verifiable hypothesis is the presence of leftover inactive clonotypes.

3.2.4. Efficiency of the cross-reacting response—The ultimate test of efficiency is the capacity to clear the infection. As for the other functions examined, we need to evaluate the relative effect of cross-reacting memory in opposing the progressively distant invaders, and compare it to the effect seen in the homologous secondary response, which is obviously the most efficient in fighting the infection. Since the measure of the infection is defined as “viral success” (VS) and is represented by the sum of virus during the response, the efficiency of the response is inversely proportioned to the viral success. In the present set-up, thwarting viral success emerged as a very sensitive measure. By taking VS during the first infection as the asymptote, VS is about 20 times lower in the homologous secondary infection and increases only moderately from distance 1 to distance 4 (Fig. 14). There is no significant difference of VS between the primary and secondary responses at distance 6, 7 and 8.

3.3. Split Cross-reactivity

To model the situation of many heterologous responses, we simulated primary responses in a large group of virtual mice who either were assigned 500 B cells instead of 2500, or had incorporated a code variant by which the minimal affinity (maximal mismatch between B paratope and epitope) was increased from the default level of efficient affinity =1 (3 mismatches) to 2 (2 mismatches). As expected, both methods produced a significant number of VM with “cellular only” response while a slight majority responded with both Tc and B.

The cartoon (Fig. 15) illustrates three theoretical relations between humoral and cellular cross-reactivity, represented respectively by upward and downward diagonal hatch marked areas. The *classic* situation on the *left*, assumes that cross-reactions are represented equally in the humoral and the cellular branch of the IS, yielding a complete overlap of the two areas. The *central* and the *right* side drawings propose two arrangements allowing cross-reactions to take place in only one of the branches; therefore, in contrast with the *left*, they both could accommodate the findings underpinning heterologous immunity [6], where serologically unrelated viruses cross-react at the T cell level.

We have simulated the split situation by constructing a matrix cross-reactive scheme in which the epitopes had been modified to be 8 bit distant from the epitope of the primary infections, while the peptides followed the scheme of step by step increments of the distance. The results obtained by running this matrix are compared with the default situation and allow by contrast pinpointing some interferences between the two branches of IS.

3.3.1. Comparative effects on Tc memory recall—Tc memory is established as the final step of the response to the primary infection. In this simulation the primary response has been humoral and cellular, but only the cellular is effectively cross-reacting. In “split” regime, the homologous response necessarily elicits a humoral and cellular secondary response. All the cross-reacting infections in “split” can only elicit the cellular response. The relative efficiency of the recall is compared with the response to the stimulation of both the humoral and cellular branch. The comparative study of memory recall in default and split regimes reveals that Tc recall is highest in the split cross-reactive experiments, after challenge with homologous (distance 0) and near cross-reactive (distance 1, 2, 3) antigens, while there is a relative depression of recall in the default setting, corresponding to the same

challenges. There is no difference at more distant challenges (distance 4 and 5) and there is no recall at all at distances 6, 7 and 8 (Fig. 16). This unexpected finding signals a competition between the two branches of the immune system, hampering the secondary Tc response. Since preformed antibodies could be involved, we ran “what if?” experiments on virtual mice whose antibody half lifetime, normally set at 100 time steps (ts), has been shortened (to 10, or 50) or extended (to 200 or 500). The results are shown in Fig. 17 and compared with the “split” response, where there is no secondary antibodies produced.

3.3.2. Strength of the Tc response—The second feature of the cross-reactive response examined by comparing the split and standard set-up is the strength ratio of the second response, which is measured against the first and also against the second antigen. All responses are relative to the strength of the primary response. As expected, the strength ratio against the first antigen decreases progressively from a level higher than one to very low while the strength against the second antigen is supposed to reach 1, the level of a primary response also at antigenic distances beyond cross-reaction. However, when compared with the “split” response (Fig. 18 (b)), the standard H+C curves (Fig. 18 (a)) shows a consistent depression in correspondence of the homologous and near cross-reacting challenge. To strengthen the evidence that the Tc response is disturbed by the competing presence of the humoral response, we have run a series of “what if?” tests, introducing manipulated Ab half lifetime that had already been used in the recall study. The curves obtained (Fig. 19 upper panel) support the role of humoral competition. A different scenario appears when the strength against the second antigen is measured: the standard H+C and the split set-up (Fig. 18) show similar Tc affinity curves, both featuring a small consistent depression at distance 2, 3, 4, and then reaching up to strength approximate to 1 at 5, 6 and farther. The cause of this depression originates in the cellular memory itself, as hinted by Fig. 19 lower panel. The mechanism could involve the presence of large number of cross-reacting memory Tc cells, ready for action but necessarily low in affinity: they could hinder by resource competition, the triggering of new higher affinity clones.

4. Discussion

The general aim of this study is to better understand the structural and functional interplay of the branches of the Immune System, humoral and cellular, a task where the IMMSIM model finds its congenial place and has already proven useful [5]. We have shown that the model can both simulate and be predictive of biological experiments in a series of collaborations with the UMass group, which is studying the dynamics of the memory T cell responses under sequences of heterologous viral infections. That group has shown that virus-induced type 1 interferon causes an apoptotic loss of memory cells early in viral infections, and after resolution of an infection there is a permanent loss of memory cells specific to previously encountered viruses. This mechanism is labelled Active Attrition in contrast to the space and resource competition between memory cells (Passive Attrition). IMMSIM modelling showed that the long term loss in memory could be accounted exclusively by active attrition or by a combination of active and passive attrition, but not by passive attrition alone [6]. In the case of viruses sharing cross-reactive epitopes, the UMass group noted that cross-reactive memory cells were not inherently resistant to the early active attrition, and we questioned with the model what advantage susceptibility to active attrition of memory cells could provide to a host [24]. The model predicted that without active attrition, cross-reactive memory cells would cause dramatic immunodomination of the T cell response, but with active attrition there could be a far more diverse response, generally considered to be beneficial in the fight against viruses. Recent studies with T cells from older mice, which resist active attrition, validated this prediction by showing greater immunodomination by cross-reactive T cells in these older hosts (manuscript in preparation). Even with active attrition, however, cross-reactive T cells can still dominate immune responses. Again,

IMMSIM modelling predicted that cross-reactive responses could result in extreme narrowing of the T cell repertoire that varied between mice and that repertoire narrowing could be a function of the avidity of cross-reactive T cells. Very similar narrowing of the repertoire was mimicked with biological experiments using the cross-reactive viruses LCMV and PV [23].

The present paper, all based on computer experiments, focuses on the mechanisms governing cross-reactivity in heterologous anti virus responses. The very label “heterologous”, used by Selin & Welsh [6] in describing their findings, reveals the novelty of viruses which are serologically foreign to each other, but nevertheless cross-react at the level of cellular responses. This means that the humoral and cellular recognition of the universe of foreign antigens follows different criteria, and that it becomes interesting and important to find out the how, when and why of the phenomenon, that we like to re-name Split Cross-reactivity (SC) remembering Brent’s description of cases where the state of tolerance affected only one branch of the Immune System [25]. SC is interesting on several counts: a) it can help understanding the obviously different mechanics underpinning cross-reaction in the two branches, until recently considered similar; b) it sheds light on evolutionary aspects of the IS. For example, even if SC occurs only in a fraction of infections and depends on the viral traits [5], it shakes the assumption that the coexistence of the branches is justified by their obligatory cooperation in the anti-virus defence; c) the deployment of one or both responses can be considered as another example of the cognitive properties of IS [26], the capacity to select the most favourable and economical setup.

We believe that a computational approach such an agent-based discrete model is the right if not the only tool to represent hypotheses of this breadth. However, the model’s reliability and its limitations must be displayed up front to the reader. This applies particularly to IMMSIM, since it has undergone extensive upgrading and re-writing in C language after the last comprehensive study on viral immunology [5]. Therefore, before tackling the split situation we have applied the model to homologous memory responses and on standard (humoral and cellular) cross-reactivity. For the latter study, we needed a systematic assessment of the effect of stepwise increase of the bit distance between primary and secondary infecting virus.

The first group of *in machina* experiments shows the responses of VM to repeated infections of a non-lethal virus. The results are predictable and can be considered reassuring. Repeated infections are met by secondary and tertiary responses which show improvement relative to the primary, in terms of speed and quality obtained through selection of higher affinity clones both in humoral and cellular responses. Only in “size” (number of antibodies and effector cells), the response differs from the traditional expectation. This is because with the virus used, the cure is successful and its very success against repeated infections limits the availability of antigen and thus the cell population engaged.

A more detailed and complex picture is provided by the second series of *in machina* experiments, which is concerned with cross-reaction. Here we have introduced a matrix design, which consists in programming a network of interactions and in grouping the results solely according to the structural distance of primary and secondary antigen, whereas all other differences are randomized. The results are consistent, and the matrix system has passed the test. The impact of increasing distance of the second antigen, in the hypothetical case of parallel effect on humoral and cellular responses is tested by focusing on a number of sensitive features of the memory response: the “recall” efficiency, the affinity (strength) maturation, the evolution of repertoires and the effectiveness of the defence. The key findings regard the clonal and cellular recall, and the strength of the responses. Curves

illustrating different variables show repeated features, suggesting high connectivity between responses, and the existence of different types of competition.

It is when we introduced the “split cross-reactivity” (obtained in the model by the stratagem of extending the bit distance between B epitopes over the range of cross-reaction) that we found a tool able to dissect and reveal some important connections. First, it turns out that the “split” situation is the most favourable for the full display of the cellular response: while default homologous and “near” cross-reacting T cell responses are depressed in the presence of the B machinery (a key factor being the persistence of circulating antibodies), split responses are not affected. Second, another depression, affecting the strength of Tc responses is characteristically localized in the intermediate distance cross-reactions. Here split and standard responses do not differ, suggesting that this competition is within the cellular response, and precisely between cross-reacting memory and new clones. The negative effect of memory may be surprising at first but can easily be explained by considering that far cross-reactivity is carried out by clones that display low affinity to the challenging virus, but can compete out new and possibly better clones by the sheer power of cell numbers.

From these observations, we may conclude that memory can in certain cases become a hindrance to the defence. Does this also happen in the real immune system? If it does, a case in point would be, e.g. the flu vaccine: the model predicts that it is useful if the challenging infection is identical or very similar to the vaccine while if it is far cross-reactive, the effect would be damaging. In the interplay of modellers and wet lab experimenters, this is one experiment that we would like to see repeated in the laboratory.

An interesting by-product of these studies is the potential use of the concept that preformed antibody block the initiation of the secondary response. While T memory triggering is efficient with split cross-reactivity, it is hampered in “non-split” if the lifetime of antibody is extended over the physiological range. The absence of Ab barrier in split cross-reactive responses lends itself to an interesting application “on the field”: to use, instead of the homologous vaccine challenge, a cross-reacting antigen that will reach directly the memory cells, dodging the antibodies.

Acknowledgments

Research in this publication was supported by NIH/NIAID through Grant R01-AI054455 awarded to F. Celada. The copyright on the code of IMMSIMM-C utilized throughout the research belonging to R. Puzone and his employer, IST (Italian Institute of Cancer Research). Also, the secretarial assistance by Francesca Priarone and Elisa Tedone is gratefully acknowledged.

References

1. Kitano H. Systems biology: a brief overview. *Science*. 2002; 295(5560):1662–4. [PubMed: 11872829]
2. Neumann, JV. *Theory of Self-Reproducing Automata*. University of Illinois Press; 1966.
3. Celada F, Seiden PE. A computer model of cellular interactions in the immune system. *Immunol Today*. 1992; 13(2):56–62. [PubMed: 1575893]
4. Seiden PE, Celada F. A model for simulating cognate recognition and response in the immune system. *J Theor Biol*. 1992; 158(3):329–57. [PubMed: 1287364]
5. Kohler B, Puzone R, Seiden PE, Celada F. A systematic approach to vaccine complexity using an automaton model of the cellular and humoral immune system. I. Viral characteristics and polarized responses. *Vaccine*. 2000; 19(7–8):862–76. [PubMed: 11115710]

6. Selin LK, Cornberg M, Brehm MA, Kim SK, Calcagno C, Ghersi D, et al. CD8 memory T cells: cross-reactivity and heterologous immunity. *Semin Immunol.* 2004; 16(5):335–47. [PubMed: 15528078]
7. De Boer RJ, Mohri H, Ho DD, Perelson AS. Turnover rates of B cells, T cells, and NK cells in simian immunodeficiency virus-infected and uninfected rhesus macaques. *J Immunol.* 2003; 170(5): 2479–87. [PubMed: 12594273]
8. Fanning LJ, Connor AM, Wu GE. Development of the immunoglobulin repertoire. *Clin Immunol Immunopathol.* 1996; 79(1):1–14. [PubMed: 8612345]
9. Fink PJ, Bevan MJ. H-2 antigens of the thymus determine lymphocyte specificity. *J Exp Med.* 1978; 148(3):766–75. [PubMed: 308986]
10. Burnet, FM. The clonal selection theory of acquired immunity. London: Cambridge University Press; 1959.
11. Ashton-Rickardt PG, Bandeira A, Delaney JR, Van Kaer L, Pircher HP, Zinkernagel RM, et al. Evidence for a differential avidity model of T cell selection in the thymus. *Cell.* 1994; 76(4):651–63. [PubMed: 8124708]
12. Germain RN. MHC-dependent antigen processing and peptide presentation: providing ligands for T lymphocyte activation. *Cell.* 1994; 76(2):287–99. [PubMed: 8293464]
13. Lanzavecchia A. Antigen-specific interaction between T and B cells. *Nature.* 1985; 314(6011): 537–9. [PubMed: 3157869]
14. Celada F. Quantitative studies of the adoptive immunological memory in mice. II. Linear transmission of cellular memory. *J Exp Med.* 1967; 125(2):199–211. [PubMed: 6019132]
15. Puzone R, Kohler B, Seiden PE, Celada F. IMMSIM, a flexible model for in machina experiments on immune system responses. *Future Generation Computer Systems.* 2002; 18(7):961–72.
16. Castiglione F, Bernaschi M, Succi S. Simulating the immune response on a distributed parallel computer. *Int J Mod Phys C.* 1997; 8(3):527–45.
17. Bernaschi M, Castiglione F. Design and implementation of an immune system simulator. *Comput Biol Med.* 2001; 31(5):303–31. [PubMed: 11535199]
18. Valensin S, di Caro G. A theoretical model for “in machina” experiments on immunosenescence. *Ann N Y Acad Sci.* 2000; 908:344–7. [PubMed: 10911982]
19. Chao DL, Davenport MP, Forrest S, Perelson AS. A stochastic model of cytotoxic T cell responses. *J Theor Biol.* 2004; 228(2):227–40. [PubMed: 15094017]
20. Fazekas de St G, Webster RG. Disquisitions on Original Antigenic Sin. II. Proof in lower creatures. *J Exp Med.* 1966; 124(3):347–61. [PubMed: 5926092]
21. Clute SC, Watkin LB, Cornberg M, Naumov YN, Sullivan JL, Luzuriaga K, et al. Cross-reactive influenza virus-specific CD8+ T cells contribute to lymphoproliferation in Epstein-Barr virus-associated infectious mononucleosis. *J Clin Invest.* 2005; 115(12):3602–12. [PubMed: 16308574]
22. Clute, SC.; Naumov, YN.; Watkin, LB.; Sullivan, JL.; Luzuriaga, K.; Welsh, RM., et al. Increased breadth of cross-reactive T cell receptor repertoires recognizing relatively dissimilar Epstein-Barr and influenza A virus epitopes. In manuscript
23. Cornberg M, Chen AT, Wilkinson LA, Brehm MA, Kim SK, Calcagno C, et al. Narrowed TCR repertoire and viral escape as a consequence of heterologous immunity. *J Clin Invest.* 2006; 116(5):1443–56. [PubMed: 16614754]
24. Bahl K, Kim SK, Calcagno C, Ghersi D, Puzone R, Celada F, et al. IFN-induced attrition of CD8 T cells in the presence or absence of cognate antigen during the early stages of viral infections. *J Immunol.* 2006; 176(7):4284–95. [PubMed: 16547266]
25. Brent, L.; Courtenay, TH. On the induction of split tolerance. Czechoslovak Academy of Sciences; Prague: 1962. p. 113
26. Celada, F. Does the human mind use a logic of signs developed by lymphocytes 10⁸ years ago. Berlin and Heidelberg: Springer Verlag; 1988.

Appendix

Grid size effect

To mimic the “close encounters” that require actual contact, the cells and molecules of “the body” are distributed randomly in sites whose number and encompassed area is defined by the grid. The grid is called “hexagonal” because each site has one edge in common with six other sites. The default conformation of the grid is 15×16 , and defines 240 sites. Decreasing or increasing the number of sites will have an effect on the population of agents eligible for immediate contacts without an intermediate phase of cells and molecules diffusing from sites to sites. A greater difficulty of interaction will also have a selective effect, favouring among interaction within the site, those of a higher affinity. We have compared matrix experiments carried out with different grids, and the results may shed some light on the functioning of the immune system.

The comparison of the data shown in Fig. A1–2, reveals some similarities and some differences between the humoral and the cell-mediated responses. For heterologous response, the affinity maturation of the humoral and the cell-mediated responses is hampered by “small” size grid (i.e. large size sites) as there is a tendency to reach maximal strength at 19×20 grid. However, for homologous response, it is the opposite. The relative recall of B and T cells after the second viral challenge is also affected by the grid size: a larger grid size (e.g., 19×20) correlates with efficiency of maturation and takes place in larger grid sizes (Fig. A3).

Figure A1.

Strength of B response after challenge with decreasing antigen distance 8→0, as affected by grid size. Diamonds: 9×10 grid; squares: 15×16 grid; triangles: 19×20 grid. ANOVA analysis is conducted at distance 0, 1, 2, 3, 4, F-value > 3.35.

Figure A2.

Strength of Tc responses after challenge with decreasing antigen distance 8→0, as affected by grid size. Diamonds: 9×10 grid; squares: 15×16 grid; triangles: 19×20 grid. ANOVA analysis is conducted at distance 0, 1, 2, 3, 4, F-value > 3.35.

Figure A3.

Recall of cross-reactive clones after challenge with decreasing antigen distance 80, as affected by grid size. Diamonds: 9×10 grid; squares: 15×16 grid; triangles: 19×20 grid. ANOVA analysis is conducted at distance 0, 1, 2, 3, 4, F-value > 3.35.

Figure 1.
IMMSIM flowchart of the virus cycle. I: infectivity; P: probability; S: multiplicity; L: lethal load.

Figure 2.
IMMSIM flowchart of the humoral response.

Figure 3.
IMMSIM flowchart of the cellular response.

Figure 4. Scheme of a 5-virus cross-reactivity matrix. (a). Circles v1 to v5 represent viral antigens (epitopes or peptides) differing from each other by the number of bits indicated by integer numbers on the joining segments. (b). Each of the 25 experiments features secondary challenges by antigens differing by 0, 1, 2, 3, 4, 5, 6, 7, 8 bits from the primary immunogen. Distance 0 represents the homologous challenge.

Figure 5.

Scheme showing the maximal possible cross-interaction among antigens, when default receptor-epitope interaction takes place at 3 mismatches or fewer. The same receptor can interact with two antigens differing at most by 6 bits.

Figure 6.

Statistical significance of humoral response (strength). Upper panel: primary response; lower panel: cross-reacting secondary response. Ordinate: standard error of the mean; abscissa: number of matrix runs. The results indicate that 25 runs are sufficient to approach a low asymptote in all curves.

Figure 7.

Repeated homologous infections and responses of one virtual mouse. 20 viral particles (with coordinates $I = 0.001$, $S = 1.05$, $L=80$) were injected at three different time steps (ts) $ts = 0$, $ts = 1000$, and $ts = 2000$. The top and middle panel show the kinetics of B and Tc responses; the bottom panel shows the kinetics of the virus. The virus multiplicity peaks 200 ts after the first injection and 110 ts after the second injection and almost immediate after the third injection. The humoral response peaks at almost 200, 150 and 70 ts after the injections while the Tc peaks are 220, 180 and 60 Ts after the injections.

Figure 8.

Trimming of repertoire during successive responses against the same virus. The panels show numbers of B clones (left) and Tc clones (right) involved in primary, secondary and tertiary response. ANOVA analysis is conducted for the B and Tc repertoire and the F value > 3.35.

Figure 9. Affinity maturation of the humoral response (left panel) and of the cellular response (right panel) is shown in terms of strength. ANOVA analysis is conducted for the B and Tc repertoire and the F value > 3.35.

Figure 10.

Clonal and cellular recall ratio (see definition in text). X-axis is the bit distance of the first antigen and the second antigen, y-axis is the recall ratio. Clear squares: Tc clone; clear diamonds: Tc cellularity; black squares: B clone; black diamonds: B cellularity.

Figure 11. Relative affinity of secondary responses of B (a) and Tc (b) cells after challenge with homologous and cross-reactive viruses. Clear diamonds: affinity against the first antigen; black squares: affinity against the second antigen.

Figure 12.

B and Tc clones active in primary and secondary responses, as affected by the antigenic distance of the challenges. Clear columns: number of clones present in the primary response, dark columns: number of clones present in the secondary response.

Figure 13.

Relative repertoire variation after secondary challenge showing (based on data in Figure 12). Decrease or increase depends on distance between primary/challenging antigens and there is a dramatic inversion of tendency between distance 4 and 5. Black square: Tc repertoire; black diamond: B cell repertoire. X-axis is the distance between two antigens, including epitopes and peptides; y-axis is the increase/decrease of the secondary repertoire relative to the primary repertoire.

Figure 14.

Thwarting of “viral success” during the secondary infection decreases as a function of the antigenic distance of the invaders. Black squares: secondary viral success; black diamond: primary viral success. Viral success is the summation of the virus level at each time step.

Theoretical overlap among Humoral and Cellular repertoire*

Figure 15.

Cartoon to represent hypothetical overlaps of humoral and cellular cross-reacting repertoires. Only the central and the right drawing are compatible with split cross-reactivity. Upward diagonal hatch mark: humoral response; downward diagonal hatch mark: cellular response.

Figure 16.

The recall ratio is the number of re-stimulated memory cells relative to the total cellularity at the peak response after the second challenge. The split situation where there is no competition by the humoral response, is compared with H+C responses. The recall rate is significantly depressed in homologous and near cross-reactive challenges when both humoral and cellular responses are active. Black square: split; black diamond: default parameter setting. For split, x-axis is the distance between two peptides with identical epitopes; for default, x-axis is the distance between two antigens, including epitopes and peptides.

Figure 17.

This figure shows the results of a “what if?” experiment, where in addition of the two curves of Fig. 16, the effect of artificially reducing (to 10, 50) and extending (to 200, 500) of Ab half lifetime are added. The procedure demonstrates that the depression of recall in homologous and near cross-reacting Tc responses is caused by competition with the humoral response. ANOVA analysis is conducted for distance 0, 1, 2, 3, 4 and the F-value > 3.35. Clear squares: split; black squares: Ab lifetime 10; black triangle: Ab lifetime 50; black diamond: the default Ab lifetime 100; crosses: Ab lifetime 200; stars: Ab lifetime 500.

Figure 18.

Strength of the secondary Tc response after challenge with viruses when both H and C responses are active. Panel (a) shows that strength directed at first antigen decreases with antigen bit distance, and strength directed at the second antigen has a shallow dip from distance 1 to 4, before approaching unit. T-test is applied at distance 2 and 8, the p-value < 0.05. Panel (b) shows Tc strength towards first and towards second antigen in the split set-up (no cross-reacting responses in the humoral response). T-test is applied at distance 4 and 8, the p-value < 0.05. Black diamond: against 1st antigen; black squares: against 2nd antigen.

Figure 19.

Upper panel is the Tc strength towards first antigen, comparing SC with 4 “what if?” cases of H+C responses with artificially shortened (10, 50) or extended (200, 500) Ab half lifetime. The vertical arrow (“Effect of humoral response”) indicates the focus of inter-branch competition with the presence of humoral response thwarting the affinity of the Tc response. Lower panel is the Tc strength towards second antigen, comparing SC with 4 “what if?” cases of H+C responses with artificially shortened (10, 50) or extended (200, 500) Ab half lifetime. The vertical long arrow indicates the focus of the competition extended by the humoral response while the short vertical arrow shows the focus of the depression of Tc memory of the affinity build up by new primary cells. White square: split; black square: Ab lifetime 10; black triangle: Ab lifetime 50; black diamonds: default (Ab lifetime 100); crosses: Ab lifetime 200; stars: Ab lifetime 500.

Table 1

Default parameters of IMMSIM used in the present study, also values different from those listed have been used in order to perform specific experiments.

Parameter	Default values
String length (bit)	16
Mismatches	3
Grid size	15×16
Ep death level	0.5
Death load	200000
B, Tc, Th population	2500
APC population	1500
Ep population	20000
B, Th multiplication	3
Tc multiplication	7
Plasma cell, Ab life time	10
APC, B, Th, Tc life time	50
Ep life time	100
Memory cell life time	20000
IFN, IL4, Damage life time	3
Binding probability APC+Ag	0.001
Binding probability APC+Fc	0.02
Binding probability Ab+Ag	1
Mutation parameter	0.2,1.0,3.0
Affinity level and enhancer	0.03, 1.1
Antigen 1 lethality (mult, infect, lethal load)	1.05, 0.001, 80
Antigen 2 lethality	1.06, 0.001, 80