Published in final edited form as:

Inhal Toxicol. 2013 December ; 25(14): 766–773. doi:10.3109/08958378.2013.847993.

Comparison of how ambient PM_c and PM_{2.5} influence the inflammatory potential

Matthew D. Ferguson¹, Chris Migliaccio¹, and Tony Ward^{1,*}

¹ University of Montana, Center for Environmental Health Sciences, Missoula, Montana.

Abstract

Airborne particulate matter (PM) is one of six criteria air pollutants currently regulated by the U.S. Environmental Protection Agency (EPA), with existing ambient standards for PM_{2.5} and PM₁₀. Currently there are no health-based regulations for the size fraction between 2.5 and 10 μ m, commonly known as the coarse fraction (PM_c). The present study investigates current gaps in knowledge for PM_c including exposure toxicity and PM ratios (PM_c:PM_{2.5}) in PM₁₀. Throughout the world, all three PM size fractions have been shown to be associated with adverse impacts. Recent studies have shown that PM_c can be more detrimental to susceptible populations when directly compared to PM_{2.5}, and that the PM_c fraction in PM₁₀ can account for the majority of the inflammatory response from PM₁₀ exposure.

In our studies we utilized a bone marrow-derived mouse macrophage *in vitro* system to compare the inflammatory potential of PM_c, PM_{2.5}, and mixtures of the two. The result was a linear increase in interleukin(IL) –1 β with increasing levels of exposure to winter and summer PM_c, as compared to PM_{2.5}, which exhibited logarithmic growth. Also, exposure to PM₁₀ as a function of PM_{2.5} and PM_c mass ratios showed that IL-1 β and TNF- α levels increased synergistically with a greater burden of PM_c. Endotoxin content in the PM did not correlate with these results, suggesting that other activators in PM_c are likely responsible for activating the NF- κ B pathway and the inflammasome.

Keywords

inflammation; pulmonary macrophage; coarse fraction; ambient particulate matter; mixtures; NAAQS; IL-1 β ; TNF- α ; endotoxin; NF- κ B

Introduction

Airborne particulate matter (PM) is a known causative agent in inducing or exacerbating a variety of respiratory and cardiovascular diseases (Pelucchi et al. 2009, Brook et al. 2010, Dvonch et al. 2009, Samet et al. 2009, Weinmayr et al. 2010, Zanobetti et al. 2010, Xu et al. 2011). PM is also one of six criteria air pollutants currently regulated by the United States Environmental Protection Agency (EPA), with existing ambient standards for both PM_{2.5} and PM₁₀ (i.e. particulate matter (PM) with mean aerodynamic diameters <2.5 and <10 micrometers (μ m), respectively). Currently there are no health-based regulations for the size fraction between 2.5 and 10 μ m, commonly known as the coarse fraction (PM_c). Compared to ambient PM_{2.5} and PM₁₀ (which are routinely measured across the US), much less is known about PM_c seasonal concentrations and health effects. In multiple research studies conducted throughout the world, all three PM size fractions have been shown to be

* author for correspondence, Phone: (406) 243-4092, Fax: (406) 243-2807, tony.ward@umontana.edu.

associated with adverse impacts (Gilmour et al. 2007, Gordian et al. 1996, Host et al. 2008, Mar et al. 2004, Monn and Becker 1999, Ostro et al. 2000, Pelucchi et al. 2009 2007).

PM_c has been shown to cause a more significant pulmonary inflammatory response than PM_{2.5} in animal models (Tong et al. 2010, Wegesser and Last 2009), as well as increased inflammatory potential in multiple respiratory cell lines (Gualtieri et al. 2010). Becker et al. (Becker et al. 2005) reported that compared to PM_{2.5}, PM_c induced higher levels of the inflammation markers IL-6 and Cox-2 mRNA in alveolar macrophages, and higher levels of IL-8 and Cox-2 mRNA in normal human bronchial epithelial cells. These studies suggest that the coarse fraction in PM₁₀ accounts for 90-95% of the resulting inflammatory response (Becker et al. 2005). These findings may be explained by the biological composition of PM_c, as the coarse fraction can contain up to 10 times the amount of biological content compared to PM_{2.5} (Heinrich et al. 2003). This biological content includes pathogen associated molecular patterns (PAMPS), such as endotoxin, which prime the activation of the inflammasome (Tschopp and Schroder 2010, Yazdi et al. 2010). The inflammasome is a protein complex that once formed can stimulate maturation of pro-IL-1 β to its active form, IL-1 β . However, a pool of pro-IL-1 β will need to be maintained allowing the inflammasome to continue its function. Endotoxin has also been shown to trigger the NF- κ B pathway and downstream pro-IL-1 β production via TLR4 (Schins et al. , Bauernfeind et al. 2009), suggesting that ambient PM-induced adverse health outcomes may be heavily influenced by the PM_c fraction.

A tenet of toxicology is that exposure to environmental toxicants will likely occur as a mixture, with the outcome being a function of the joint effects of this mixture. A measure of how a mixture can maintain, lower, or increase a toxic outcome is termed additivity. For example, laboratory studies exposing humans to both carbon black particles and O₃ resulted in greater lung inflammation and suppression of alveolar macrophage phagocytosis than would be predicted using the sum of both toxicities alone (Jakab and Hemenway 1994). These results suggest a *synergism* in the lung, or a *greater than additive* toxicity between carbon black and O₃.

Toxic consequences from mixture exposures have two other outcomes other than synergism. An inverse relationship to synergism may also be observed where *antagonism* takes place, i.e. a decrease in toxic potential, *less than additive*. Finally, when two toxicants are introduced in a mixture and if one toxic does not affect the other, and visa versa, the mixture effect is termed *additive*.

Although multiple studies have been conducted linking either PM_{2.5} or PM₁₀ to respiratory disease throughout the world, little research has been conducted investigating the role of these two PM size fractions in mixture, specifically how the PM_c size class influences overall PM₁₀ inducible toxicity. In the real world, exposure to PM₁₀ occurs as a function of PM_c and PM_{2.5} mass ratios. These mixtures can change as a function of geography, temporality, and seasonal conditions (e.g. wildfires, winter inversions, etc.). In this manuscript, we describe the results of studies that evaluated the additive effect of ambient PM_{2.5} and PM_c mixtures. Following the harvesting of ambient PM using a particle concentrator, an *in vitro* model of inflammation was used to determine the contributions of PM_c within a mixture. The overall goal of these studies was to not only investigate the inflammation potential of PM_c compared to PM_{2.5}, but also to elucidate the relative contributions of these two size fractions to observed health effects.

Methods

Collection of Ambient Particles

Seasonal PM was collected in Missoula, Montana, a valley airshed located in west-central Montana. With a population of 70 000, Missoula is the 2nd largest city in the state of Montana. Particles were collected using a versatile aerosol concentration enrichment system particle concentrator (VACES-PC, University of Southern California). The concentrator has three parallel sampling lines that simultaneously collect PM_{2.5} and PM_c, respectively, at a set flow rate of 110 L/min (Kim et al. 2001a, Kim et al. 2001b). Following each sample run, the product (i.e. particles in ultrapure water) was stored at -20° and later lyophilized to concentrate particles for further studies. The particle concentrator is housed on the roof of a four story building on the University of Montana campus. Particles were harvested during the winter (February 2012), spring (May 2012), and summer months (July and August 2012). These collection periods yielded from 100 µg to about 1 mg of each PM type, depending on conditions.

Ambient PM Levels in Missoula

Although PM_c sampling is not currently conducted in Missoula, continuous PM_{2.5} and PM₁₀ are routinely measured as part of Missoula's EPA compliance monitoring program. This program uses these continuous PM data to calculate 24-hour ambient PM averages, which are then reported. Using these continuous PM_{2.5} and PM₁₀ data (collected with a MetOne BAM at the Missoula Boyd Park site), we were able to calculate, by difference, the seasonal ambient PM_c mass concentrations during 2012 ($[PM_{10}] - [PM_{2.5}] = [PM_c]$). We also calculated the 24-hour average mass concentrations on days when PM was collected with the particle concentrator and used in this work. The unique seasonal patterns in Missoula, MT are accounted for in this study, as high latitudes and high elevations stimulate longer cold seasons. For the purpose of this manuscript, winter months are categorized as the months of November, December, January, and February; spring are the months of March, April, and May; summer months are the months of June, July, and August; and autumn is September and October.

Mice and Bone Marrow-Derived Macrophages

A major component of respiratory immunity is the pulmonary macrophage, which plays a key role in the regulation of inflammation as well as subsequent adaptive responses. Pulmonary macrophages are perpetual residents of the lung and act as sentinels or first responders for particle uptake. These macrophages have the necessary receptors for particle uptake/phagocytosis (Hirayama et al. 2011, Geiser 2010, Goodridge et al. 2012, Underhill and Goodridge 2012). Our laboratory has published multiple *in vivo* particulate exposure studies using the Balb/c mouse model (Migliaccio et al. 2005, Migliaccio et al. 2008, Lacher et al. 2010, Migliaccio et al. 2013). In addition to these studies, our laboratory has also utilized an *in vitro* macrophage system using bone marrow-derived macrophages (BMdMs) to evaluate the effects of particles on cell functions (Migliaccio et al. 2005, Migliaccio et al. 2008). Therefore, by exposing mouse BMdMs to PM, an inflammatory response can be measured.

A variety of studies have described the key role of the pulmonary macrophage in particle clearance (Geiser 2010), and its link between the innate and adaptive immune systems (Gordon 2003, Mosser 2003). Bone marrow-derived macrophages utilized in this study to evaluate the inflammatory response of relevant doses of different PM fractions (PM_{2.5}, PM₁₀, and PM_c). Methods for BMdM cultures are previously described (Pfau et al. 2004, Migliaccio et al. 2008). Briefly, hind legs of euthanized Balb/c mice (Jackson Laboratory, Bar Harbor, ME, USA) were flushed with media (RPMI 1640 with 10% FBS) to release the

marrow. After overnight stromal elimination, cells were stimulated with M-CSF for an additional 7-10 days of culturing. On day 10, cells were aliquoted (10^5 cells/well) in 96-well plates for culturing with particles. All animal procedures were approved by the University of Montana Institutional Animal Care and Use Committee (Missoula, MT, USA). Method of euthanasia was by lethal dose of sodium pentobarbital via interperitoneal injection.

PM_c vs. PM_{2.5} Comparison Studies

During the summer (July, 2011) and winter (February, 2012) months, PM_{2.5} and PM_c, were simultaneously harvested in Missoula using our VACES-PC. Stock solutions of PM were made by diluting pre-determined masses of PM into a small volume of PBS (< 0.5 mL) followed by sonication. The volume of PBS necessary is dependent on the amount of PM harvested. Cells were incubated with particles for 1 hour (mixing) in 1.5 mL eppendorf tubes, then aliquoted to 96-well plates and cultured overnight. Final cultures contained 150 μ L per well of media, with each condition in triplicate exposures at 100 000 cells/well. A media negative control was also used that contained no particles. Particle exposure levels included varying masses of PM_{2.5} or PM_c at 5 μ g, 20 μ g, and 40 μ g/well. Collection of supernatants occurred after 24 hours.

PM Mixture Studies

PM_{2.5} and PM_c used in the mixture exposures were harvested during May 2012. BMdMs were exposed in a similar fashion as the comparison studies described above, but at one level of exposure at 20 μ g/well. To represent the varying conditions of PM₁₀, 20 μ g mixtures of PM_{2.5} and PM_c varied from 0%, 20%, 40%, 60%, 80%, and 100% PM_c by mass. For these studies, 20 μ g was used, as it was the lowest mass of exposure that induced the most significant differences between PM size and season with both IL-1 β and TNF- α . In similar studies using *in vivo* models, particulate exposures ranged from 5-50 μ g of particles per instillation (Porter et al. 2013, Stoeger et al. 2006). Considering that there are roughly 200-300 thousand alveolar macrophages in the mouse lung, if 50 μ g is distributed evenly, each cell will theoretically receive approximately 0.17-0.25 ng of particles. In our model using 20 μ g and 100 000 cells, if particles were distributed evenly, each cell would receive roughly 0.2 ng of particles. Therefore, the levels of particulates used in this research are representative of *in vivo* studies.

Collection of supernatants occurred after 24 hours of incubation. The resulting inflammatory cytokine levels were used to determine the additivity potential of the mixtures. To ensure that these particles were undergoing phagocytosis, bright field illumination microscopy of BMdMs following 24-hours of exposure to PM₁₀, was performed (Nikon Eclipse E800, Tokyo, Japan; Nuance Multispectral Imaging System, Hopkinton, Massachusetts).

Endotoxin Analysis

Levels of endotoxin were evaluated using methods adapted by Thorne (Thorne 2000). Briefly, the harvested PM samples were analyzed for endotoxin levels using a kinetic chromogenic *Limulus* amebocyte lysate (LAL) assay (Charles River Endosafe® - Endochrome-K™, Charles River Laboratories, Charleston, SC, USA), where a 12-point standard curve ranged from 50 to 0.005 EU/ml. The absorbance was measured at 405 nm every 30 seconds for 90 minutes.

Mixture Effect Calculation

Additivity was modeled using equation 1, a modified version of the toxic unit model (PapeLindstrom and Lydy 1997). Modification was necessary, as endpoints were not based on concentration/dose response thresholds, but were based on the resulting levels of the

expressed cytokines from 20 µg of PM exposure. Where additivity is > 1 , $= 1$, or < 1 the effect of the mixture is deemed synergistic, additive, or antagonistic, respectively. The fraction of $PM_{2.5}$ or PM_c in PM_{10} is represented with f and can range anywhere from 0 to 1. For example, if PM_{10} contained 25% $PM_{2.5}$ and 75% PM_c , f would equal 0.25 and 0.75, respectively.

$$f_{PM_{2.5}} \frac{[PM_{10}]}{[PM_{2.5}]} + f_{PM_c} \frac{[PM_{10}]}{[PM_c]} = \text{additivity} \quad (1)$$

As the ratios of $PM_{2.5}$ and PM_c change within total PM_{10} , a departure from an effect level of 1 explains whether a larger burden of one PM size fraction is either affecting a synergistic (additivity > 1) or an antagonistic (additivity < 1) effect. One other alternative is that the mixture will not cause a departure from 1, and that the increased presence of one or another PM type will result in an additive response.

Cytokine and Statistical Analysis

Collection of pro-inflammatory cytokines was completed following the 24-hour exposure. The collected media/supernatant was then stored at -20°C prior to analysis via enzyme-linked immunosorbent assay (ELISA; R&D Systems, Minneapolis, MN, USA). Each well's concentration was measured in duplicate. Levels of IL-1 β and TNF- α released from BMdMs were then plotted using Prism, version 5.0a. Student t-tests were also performed in Prism to determine significant differences between two PM types within each mass of exposure (5 µg, 20 µg, and 40 µg).

Results

Ambient air data in Missoula, MT

Table 1 summarizes the average concentrations of $PM_{2.5}$, PM_c , and PM_{10} measured throughout 2012, illustrating that PM_c is not only a substantial component of PM_{10} throughout the year, but also is present in elevated concentrations (compared to $PM_{2.5}$) throughout all four seasons in this northern Rocky Mountain airshed. The data presented in Table 1 indicate that ambient PM concentrations ($PM_{2.5}$, PM_c , and PM_{10}) vary as a function of season. Throughout the year, seasonal PM_c concentrations measured in Missoula, MT were consistently higher than $PM_{2.5}$, even in winter when there were more combustion-related particles in the Missoula airshed due to residential wood stove use (Ward and Lange 2010). Unlike $PM_{2.5}$, PM_c average concentrations were fairly consistent through winter and spring (9.3-9.6 µg/m³), with a marked increase in PM_c from summer to fall. $PM_{2.5}$ concentrations in fall were at least twice as high as all other seasons.

PM concentrations on days that ambient particles were harvested for the controlled laboratory exposure studies (PM_c vs. $PM_{2.5}$ comparison and mixture studies) are presented in Table 2. During the days that PM was harvested during the winter months, the PM_c concentrations were more than twice as high compared to the winter-long average presented in Table 1, though $PM_{2.5}$ concentrations were similar. The authors are unsure why PM_c was elevated on these sample days. When comparing PM_c concentrations measured during harvest days compared to overall summer averages, concentrations are comparable. The same is true for collected Spring PM. Also during these collection days (Table 2), $PM_{2.5}$ concentrations were comparable to the winter, spring, and summer averages presented in Table 1.

PM non-mixture exposures, in vitro

In side-by-side controlled exposures within the laboratory, winter PM_C induced the highest levels of IL-1 β and TNF- α at all three doses (Figure 1), supporting our hypothesis that PM_C is more biologically active compared to PM_{2.5}. Summer PM_C was the second most bioactive PM, followed by summer PM_{2.5} and winter PM_{2.5}. In each season (winter and summer), the most bioactive size fraction of PM was PM_C, with winter PM_C inducing more than twice the levels of IL-1 β than summer PM_C. Similar results were observed for TNF- α . IL-1 β increased in a linear fashion in response to both winter and summer PM_C. However, IL-1 β from winter and summer PM_{2.5} increased logarithmically, and leveled out at around 25 pg/mL. This logarithmic growth was not observed for TNF- α , as all four PM types generated linear increases in cytokine expression.

The data on the significant differences between PM_C and PM_{2.5} are listed in Table 3. Two-tailed, student *t*-tests were used to determine if the two PM size fractions induced significantly different responses at a single dose. Both summer and winter PM_C induced significantly higher IL-1 β levels compared to their PM_{2.5} counterparts (at 20 μ g, IL-1 β between summer PM_C v. PM_{2.5} *p*=0.02, and winter PM_C v. PM_{2.5} *p*=0.03). TNF- α levels were only significantly different between PM_C and PM_{2.5} for winter, but not for summer (at 20 μ g, TNF- α between summer PM_C v. PM_{2.5} *p*=0.09, and winter PM_C v. PM_{2.5} *p*=0.0002). IL-1 β levels from winter PM_C were triple that of summer PM_C. Differences in IL-1 β levels were mostly non-significant when comparing low doses of PM exposure, except between both summer and winter PM_{2.5} where there was no significant differences. These results suggest that PM_C is an important mediator in inducing an inflammatory response.

Cellular uptake of ambient PM₁₀

Most of the particles used in these exposures were phagocytosed by the BMdMs, as the majority of the particles were intracellular (Figure 2). Using BFI microscopy, we determined that the cells exposed to PM₁₀ had phagocytosed the majority of the particles in solution. In other images where PM_C and PM_{2.5} burden in PM₁₀ varied, similar uptake was observed (images not shown). Particle size (both PM_{2.5} and PM_C) did not seem to play a major role in the uptake of these particles. Small particles and larger sized particulates, such as the fungi, *Torula herbarum* (Figure 2), with diameters of approximately 3-4 μ m, appear to be equally phagocytosed. This confirms that both PM_C and PM_{2.5} are being taken up by the BMdMs.

PM endotoxin burden

Endotoxin levels were determined as a function of PM mass (Figure 3). Results show that overall endotoxin concentrations were highest during the summer months compared to the winter months for both fractions. As predicted, there was more endotoxin in the winter PM_C compared to the winter PM_{2.5}. Endotoxin concentrations during the summer were comparable between the two size fractions. These results suggest that endotoxin concentrations composing ambient PM can vary as a function of PM size and season, and not primarily found in the coarse fraction (compared to the fine fraction).

Mixture effects

As noted earlier, PM_{2.5} and PM_C used in the mixture exposures were harvested during May 2012. Results from the mixture studies show that varying the combinations of PM_{2.5} and PM_C in PM₁₀ influence cytokine output, with Figure 4 illustrating the effect of PM_C in a PM₁₀ mixture. All cytokine concentrations (i.e. [IL-1 β] and [TNF- α]) varied as a function of exposure to 20 μ g of each PM type (PM₁₀, PM_{2.5}, or PM_C). Shown in Figure 4, as PM₁₀ gains a greater proportion of PM_C, a synergistic release of IL-1 β and TNF- α occurs. Conversely, as the fraction of PM₁₀ contains more PM_{2.5} than PM_C, antagonistic effects are

observed. These results support our hypothesis that the PM_c fraction is an important mediator in inducing an inflammatory response with PM_{10} exposure.

Discussion

Understanding the mass ratio mixtures of ambient $PM_{2.5}$ and PM_c composing PM_{10} can help determine additivity of the inflammatory response. As shown in our controlled laboratory studies, PM_c was constantly more bioactive than $PM_{2.5}$ throughout each side-by-side exposure in comparing both summer and winter months, and with its increased burden in PM_{10} . We were successful in showing that there are certainly additive effects from these different mass ratio exposures. Furthermore, our results suggest that the outcome from exposure to PM_{10} can be partly understood as a function of $PM_{2.5}$ and PM_c burden in PM_{10} .

A source apportionment study has yet to be completed for PM_c in Missoula, MT. However, $PM_{2.5}$ source apportionment modeling has shown that the largest $PM_{2.5}$ source throughout the year is wood combustion, contributing an average of 41% to the fine fraction. This includes residential wood combustion during the winter, slash and prescribed burns from the surrounding areas during the spring and fall, and smoke from wildfires during the summer months. Other sources of $PM_{2.5}$ throughout the year include diesel exhaust (19%), secondary ammonium nitrate (17%), local industry (20%), and street sand (5%) (Ward and Smith 2005, Ward and Lange 2010). Though these studies did not focus specifically on apportioning the sources of PM_c , they do provide an idea of the types of exposure conditions that exist throughout the year in Missoula, MT.

The present study is the first attempt at determining ambient concentrations of PM_c in the state of Montana within the last 15 years. As illustrated in Table 1, ambient concentrations of PM_c were consistently greater than $PM_{2.5}$ throughout the year, with the exception of the winter months where the average PM_c and $PM_{2.5}$ concentrations were not significantly different. It should be noted, however, that harvested particles used in our laboratory studies were collected on days when the average PM_c mass concentrations were about 2 - 10 $\mu\text{g}/\text{m}^3$ greater than $PM_{2.5}$. In addition, particles were harvested during sample days in which there was higher winter PM_c concentrations ($22.6 \pm 9.2 \mu\text{g}/\text{m}^3$) compared to the winter 2012 average PM_c concentrations presented in Table 1 ($9.3 \pm 7.7 \mu\text{g}/\text{m}^3$). $PM_{2.5}$ concentrations (both throughout the winter of 2012 and during the dates of sample collection) were consistent ($9.1 \pm 4.9 \mu\text{g}/\text{m}^3$ vs. $12.3 \pm 3.0 \mu\text{g}/\text{m}^3$). During the *PM_c vs. $PM_{2.5}$ Comparison studies*, particle exposure levels ($PM_{2.5}$ and PM_c) were normalized at 5 μg , 20 μg , and 40 $\mu\text{g}/\text{well}$ in the 96-well plates. Therefore, it is unknown how the overall ambient concentrations during the days of particle harvesting influenced the results of the comparison studies (if at all). Additional studies within the laboratory across multiple seasons/years will be able to address this issue.

As evidenced in Figure 3, the differences in endotoxin content on the PM cannot explain the differences in the resulting IL-1 β expression. Something other than endotoxin is potentially stimulating the production of pro-IL-1 β upon exposure to PM_c , as the levels of IL-1 β do not seem to be limited with an increase in PM_c exposure. In addition, increasing concentrations of $PM_{2.5}$ did not induce a linear increase in IL-1 β expression (Figure 1). Further studies can evaluate the levels of pro-IL-1 β in cells, via western blot, to determine if a decreased IL-1 β release is due to a lowered level of expression of the pro form of IL-1 β or if the inflammasome is simply not active. A large pool of pro-IL-1 β remaining in the cells would suggest that there is likely a decrease in inflammasome activation, which catalyzes the maturation of pro-IL-1 β to its active IL-1 β form (Tschopp 2008).

Figure 1 supports the hypothesis that ambient PM_c is more bioactive than $PM_{2.5}$. However, these exposures were done on a side-by-side comparison using controlled laboratory conditions, and did not take into account that real-world exposures to these two size fractions occur concomitantly. The mixture exposures take into account possible mixing conditions of $PM_{2.5}$ and PM_c in PM_{10} . The results of the *in vitro* mixture exposures showed that $PM_{2.5}$ is not as potent an activator of the inflammasome as PM_c , even in the presence of another PM type. In order to stimulate these levels of IL-1 β production, the inflammasome complex must be activated by a receptor (e.g. binding of receptor to PAMP) or lysosomal stress (e.g. the release of cathepsin-B) (Tschopp 2008, Yazdi et al. 2010). Potentially, there are components in PM_c that are responsible for priming the formation of the inflammasome, such as endotoxin, an activator of the NF- κ B pathway via TLR4, and another component affecting lysosomal membrane integrity. NF- κ B activation is a precursor to transcription induction of pro-IL-1 β (Yazdi et al. 2010, De Nardo and Latz 2011). Activation of the TLR or priming of the NF- κ B pathway has been shown to stimulate the production of pro-IL-1 β and further production of NLRP3 inflammasome precursors (De Nardo and Latz 2011). It is likely that not only is the inflammasome being formed, but also more pro-IL-1 β is being produced following PM_c exposure. Also, reactive oxygen species (ROS) may play a significant role in inflammation as generation of ROS from particulates has been shown to exacerbate the inflammatory response (Fubini and Hubbard 2003). Furthermore, exposure to biogenic substances, such as pollen, largely found in PM_c , can increase ROS in the lung (Boldogh et al. 2005).

Conclusion

In this study, we utilized harvested ambient particles and biological *in vitro* models to investigate potential adverse health effects following exposures to different size fractions found in the Rocky Mountains in North America. The results of these studies conducted under controlled laboratory conditions suggest that there are properties (chemical, physical, or biological) of PM_c that are responsible for elevated levels of inflammation activity compared to ambient $PM_{2.5}$. An important step to further this research would be to investigate PM_c harvested throughout the year. For example, winter PM_c contained about two-thirds the endotoxin burden as compared to summer PM_c , yet winter PM_c induced more than twice the levels of IL-1 β than summer PM_c . Other methods of *in vitro* exposure could also be employed, such as an air-liquid interface model. Also, an alternative method of collecting particles that does not include the removal of water (via lyophilization) could be explored to minimize the loss of volatile and semi-volatile compounds. Such methods could include employing the use of a cyclone or another dry method of harvesting ambient air particles.

The mixture results presented here suggest that if the percentage of PM_c is known, the relative bioactivity of the ambient PM_{10} might be predicted. This study supports our hypothesis that PM_c adversely affects macrophage activity with increasing ratios of $PM_c:PM_{2.5}$ in PM_{10} . It is also important to note that there are currently no National Ambient Air Quality Standards (NAAQS) for the coarse fraction, though our results suggest that a new PM_c ambient standard should perhaps be considered in addition to the existing daily and annual $PM_{2.5}$ NAAQS. Including additional chemical and biogenic characterizations will facilitate our future source apportionment studies for different PM fractions. Also, evaluation of reactive oxygen species activity in both PM_c and $PM_{2.5}$ should be considered in future research. Further seasonal and dose-response relationship studies will aid in determining real-world mixture effects, as well as provide an understanding of the causal health effects from exposure to PM_c .

Acknowledgments

We would like to thank the Montana Department of Environmental Quality (Mr. Hoby Rash and Mr. Dave Simonson) and the Missoula City-County Health Department (Mr. Ben Schmidt) for supplying Missoula's ambient PM data, and also Lou Herritt (University of Montana) for imaging assistance. Support was provided by NCRR (COBRE P2ORR017670).

Declaration of Interest

The authors have no financial or other conflicts of interest.

References

- Bauernfeind FG, Horvath G, Stutz A, Alnemri ES, MacDonald K, Speert D, Fernandes-Alnemri T, Wu JH, Monks BG, Fitzgerald KA, Hornung V, Latz E. 'Cutting Edge: NF-kappa B Activating Pattern Recognition and Cytokine Receptors License NLRP3 Inflammasome Activation by Regulating NLRP3 Expression'. *Journal of Immunology*. 2009; 183(2):787–791.
- Becker S, Mundandhara S, Devlin RB, Madden M. 'Regulation of cytokine production in human alveolar macrophages and airway epithelial cells in response to ambient air pollution particles: further mechanistic studies'. *Toxicol Appl Pharmacol*. 2005; 207(2 Suppl):269–75. [PubMed: 15993911]
- Boldogh I, Bacsí A, Choudhury BK, Dharajiya N, Alam R, Hazra TK, Mitra S, Goldblum RM, Sur S. 'ROS generated by pollen NADPH oxidase provide a signal that augments antigen-induced allergic airway inflammation'. *Journal of Clinical Investigation*. 2005; 115(8):2169–2179. [PubMed: 16075057]
- Brook RD, Rajagopalan S, Pope CA 3rd, Brook JR, Bhatnagar A, Diez-Roux AV, Holguin F, Hong Y, Luepker RV, Mittleman MA, Peters A, Siscovick D, Smith SC Jr, Whitsel L, Kaufman JD. 'Particulate matter air pollution and cardiovascular disease: An update to the scientific statement from the American Heart Association'. *Circulation*. 2010; 121(21):2331–78. [PubMed: 20458016]
- De Nardo D, Latz E. 'NLRP3 inflammasomes link inflammation and metabolic disease'. *Trends in Immunology*. 2011; 32(8):373–379. [PubMed: 21733753]
- Dvornch JT, Kannan S, Schulz AJ, Keeler GJ, Mentz G, House J, Benjamin A, Max P, Bard RL, Brook RD. 'Acute Effects of Ambient Particulate Matter on Blood Pressure Differential Effects Across Urban Communities'. *Hypertension*. 2009; 53(5):853–859. [PubMed: 19273743]
- Fubini B, Hubbard A. 'Reactive oxygen species (ROS) and reactive nitrogen species (RNS) generation by silica in inflammation and fibrosis'. *Free Radical Biology and Medicine*. 2003; 34(12):1507–1516. [PubMed: 12788471]
- Geiser M. 'Update on Macrophage Clearance of Inhaled Micro- and Nanoparticles'. *Journal of Aerosol Medicine and Pulmonary Drug Delivery*. 2010; 23(4):207–217. [PubMed: 20109124]
- Gilmour MI, McGee J, Duvall RM, Dailey L, Daniels M, Boykin E, Cho SH, Doerfler D, Gordon T, Devlin RB. 'Comparative toxicity of size-fractionated airborne particulate matter obtained from different cities in the United States'. *Inhalation Toxicology*. 2007; 19:7–16. [PubMed: 17886044]
- Goodridge HS, Underhill DM, Touret N. 'Mechanisms of Fc Receptor and Dectin-1 Activation for Phagocytosis'. *Traffic*. 2012; 13(8):1062–1071. [PubMed: 22624959]
- Gordian ME, Ozkaynak H, Xue JP, Morris SS, Spengler JD. 'Particulate air pollution and respiratory disease in Anchorage, Alaska'. *Environmental Health Perspectives*. 1996; 104(3):290–297. [PubMed: 8919767]
- Gordon S. 'Alternative activation of macrophages'. *Nat Rev Immunol*. 2003; 3(1):23–35. [PubMed: 12511873]
- Gualtieri M, Ovrevik J, Holme JA, Perrone MG, Bolzacchini E, Schwarze PE, Camatini M. 'Differences in cytotoxicity versus pro-inflammatory potency of different PM fractions in human epithelial lung cells'. *Toxicology in Vitro*. 2010; 24(1):29–39. [PubMed: 19772914]
- Heinrich J, Pitz M, Bischof W, Krug N, Borm PJA. 'Endotoxin in fine (PM_{2.5}) and coarse (PM_{2.5-10}) particle mass of ambient aerosols. A temporo-spatial analysis'. *Atmospheric Environment*. 2003; 37(26):3659–3667.

- Hirayama T, Tamaki Y, Takakubo Y, Iwazaki K, Sasaki K, Ogino T, Goodman SB, Kontinen YT, Takagi M. 'Toll-Like Receptors and Their Adaptors are Regulated in Macrophages after Phagocytosis of Lipopolysaccharide-Coated Titanium Particles'. *Journal of Orthopaedic Research*. 2011; 29(7):984–992. [PubMed: 21308757]
- Host S, Larrieu S, Pascal L, Blanchard M, Declercq C, Fabre P, Jusot JF, Chardon B, Le Tertre A, Wagner V, Prouvost H, Lefranc A. 'Short-term associations between fine and coarse particles and hospital admissions for cardiorespiratory diseases in six French cities'. *Occupational and Environmental Medicine*. 2008; 65(8):544–551. [PubMed: 18056749]
- Jakab GJ, Hemenway DR. 'Concomitant Exposure to Carbon-Black Particulates Enhances Ozone-Induced Lung Inflammation and Suppression of Alveolar Macrophage Phagocytosis'. *Journal of Toxicology and Environmental Health*. 1994; 41(2):221–231. [PubMed: 8301700]
- Kim S, Jaques PA, Chang MC, Barone T, Xiong C, Friedlander SK, Sioutas C. 'Versatile aerosol concentration enrichment system (VACES) for simultaneous in vivo and in vitro evaluation of toxic effects of ultrafine, fine and coarse ambient particles - Part II: Field evaluation'. *Journal of Aerosol Science*. 2001a; 32(11):1299–1314.
- Kim S, Jaques PA, Chang MC, Froines JR, Sioutas C. 'Versatile aerosol concentration enrichment system (VACES) for simultaneous in vivo and in vitro evaluation of toxic effects of ultrafine, fine and coarse ambient particles - Part I: Development and laboratory characterization'. *Journal of Aerosol Science*. 2001b; 32(11):1281–1297.
- Lacher SE, Johnson C, Jessop F, Holian A, Migliaccio CT. 'Murine pulmonary inflammation model: a comparative study of anesthesia and instillation methods'. *Inhal Toxicol*. 2010; 22(1):77–83. [PubMed: 20017595]
- Mar TF, Larson TV, Stier RA, Claiborn C, Koenig JQ. 'An Analysis of the Association Between Respiratory Symptoms in Subjects with Asthma and Daily Air Pollution in Spokane, Washington'. *Inhalation Toxicology*. 2004; 16(13):809–815. [PubMed: 15513813]
- Migliaccio CT, Buford MC, Jessop F, Holian A. 'The IL-4Ralpha pathway in macrophages and its potential role in silica-induced pulmonary fibrosis'. *J Leukoc Biol*. 2008; 83(3):630–9. [PubMed: 18056481]
- Migliaccio CT, Hamilton RF Jr, Holian A. 'Increase in a distinct pulmonary macrophage subset possessing an antigen-presenting cell phenotype and in vitro APC activity following silica exposure'. *Toxicol Appl Pharmacol*. 2005; 205(2):168–76. [PubMed: 15893544]
- Migliaccio CT, Kobos E, King QO, Porter V, Jessop F, Ward T. 'Adverse effects of wood smoke PM_{2.5} exposure on macrophage functions'. *Inhalation Toxicology*. 2013; 25(2):67–76. [PubMed: 23363038]
- Monn C, Becker S. 'Cytotoxicity and induction of proinflammatory cytokines from human monocytes exposed to fine (PM_{2.5}) and coarse particles (PM_{10-2.5}) in outdoor and indoor air'. *Toxicology and Applied Pharmacology*. 1999; 155(3):245–252. [PubMed: 10079210]
- Mosser DM. 'The many faces of macrophage activation'. *Journal of leukocyte biology*. 2003; 73(2):209–212. [PubMed: 12554797]
- Ostro BD, Broadwin RACHEL, Lipsett MJ. 'Coarse and fine particles and daily mortality in the Coachella Valley, California: a follow-up study'. *Journal of Exposure Analysis & Environmental Epidemiology*. 2000; 10(5):412–419. [PubMed: 11051531]
- PapeLindstrom PA, Lydy MJ. 'Synergistic toxicity of atrazine and organophosphate insecticides contravenes the response addition mixture model'. *Environmental Toxicology and Chemistry*. 1997; 16(11):2415–2420.
- Pelucchi C, Negri E, Gallus S, Boffetta P, Tramacere I, La Vecchia C. 'Long-term particulate matter exposure and mortality: a review of European epidemiological studies'. *Bmc Public Health*. 2009; 9:453–460. [PubMed: 19995424]
- Pfau JC, Schneider JC, Archer AJ, Sentissi J, Leyva FJ, Cramton J. 'Environmental oxygen tension affects phenotype in cultured bone marrow-derived macrophages'. *American Journal of Physiology-Lung Cellular and Molecular Physiology*. 2004; 286(2):L354–L362. [PubMed: 14527932]
- Porter DW, Wu NQ, Hubbs AF, Mercer RR, Funk K, Meng FK, Li JT, Wolfarth MG, Battelli L, Friend S, Andrew M, Hamilton R, Sriram K, Yang F, Castranova V, Holian A. 'Differential Mouse

- Pulmonary Dose and Time Course Responses to Titanium Dioxide Nanospheres and Nanobelts'. *Toxicological Sciences*. 2013; 131(1):179–193. [PubMed: 22956629]
- Samet JM, Avila-Tang E, Boffetta P, Hannan LM, Olivo-Marston S, Thun MJ, Rudin CM. 'Lung cancer in never smokers: clinical epidemiology and environmental risk factors'. *Clinical cancer research : an official journal of the American Association for Cancer Research*. 2009; 15(18): 5626–45. [PubMed: 19755391]
- Schins RP, Lightbody JH, Borm PJ, Shi T, Donaldson K, Stone V. 'Inflammatory effects of coarse and fine particulate matter in relation to chemical and biological constituents'. *Toxicology and Applied Pharmacology*. 2004; 195(1):1–11. [PubMed: 14962500]
- Stoeger T, Reinhard C, Takenaka S, Schroepel A, Karg E, Ritter B, Heyder J, Schulz H. 'Instillation of six different ultrafine carbon particles indicates a surface area threshold dose for acute lung inflammation in mice'. *Environmental Health Perspectives*. 2006; 114(3):328–333. [PubMed: 16507453]
- Thorne PS. 'Inhalation toxicology models of endotoxin- and bioaerosol-induced inflammation'. *Toxicology*. 2000; 152(1-3):13–23. [PubMed: 11090935]
- Tong H, Cheng WY, Samet JM, Gilmour MI, Devlin RB. 'Differential cardiopulmonary effects of size-fractionated ambient particulate matter in mice'. *Cardiovascular toxicology*. 2010; 10(4):259–67. [PubMed: 20602262]
- Tschopp J. 'Induction of innate immune responses through Nalp3 inflammasome sensing of asbestos and silica'. *Cytokine*. 2008; 43(3):276–276.
- Tschopp J, Schroder K. 'NLRP3 inflammasome activation: the convergence of multiple signalling pathways on ROS production?'. *Nature Reviews Immunology*. 2010; 10(3):210–215.
- Underhill DM, Goodridge HS. 'Information processing during phagocytosis'. *Nature Reviews Immunology*. 2012; 12(7):492–502.
- Ward TJ, Lange T. 'The impact of wood smoke on ambient PM_{2.5} in northern Rocky Mountain valley communities'. *Environmental Pollution*. 2010; 158(3):723–729. [PubMed: 19897293]
- Ward TJ, Smith GC. 'The 2000/2001 Missoula Valley PM_{2.5} chemical mass balance study, including the 2000 wildfire season - seasonal source apportionment'. *Atmospheric Environment*. 2005; 39(4):709–717.
- Wegesser TC, Last JA. 'Mouse lung inflammation after instillation of particulate matter collected from a working dairy barn'. *Toxicology and Applied Pharmacology*. 2009; 236(3):348–357. [PubMed: 19272399]
- Weinmayr G, Romeo E, De Sario M, Weiland SK, Forastiere F. 'Short-Term Effects of PM₁₀ and NO₂ on Respiratory Health among Children with Asthma or Asthma-like Symptoms: A Systematic Review and Meta-Analysis'. *Environmental Health Perspectives*. 2010; 118(4):449–457. [PubMed: 20064785]
- Xu X, Sharma RK, Talbott EO, Zborowski JV, Rager J, Arena VC, Volz CD. 'PM₁₀ air pollution exposure during pregnancy and term low birth weight in Allegheny County, PA, 1994-2000'. *International archives of occupational and environmental health*. 2011; 84(3):251–7. [PubMed: 20496078]
- Yazdi AS, Guarda G, Riteau N, Drexler SK, Tardivel A, Couillin I, Tschopp J. 'Nanoparticles activate the NLR pyrin domain containing 3 (Nlrp3) inflammasome and cause pulmonary inflammation through release of IL-1 β and IL-1 α '. *Proceedings of the National Academy of Sciences*. 2010; 107(45):19449–19454.
- Zanobetti A, Redline S, Schwartz J, Rosen D, Patel S, O'Connor GT, Lebowitz M, Coull BA, Gold DR. 'Associations of PM₁₀ with sleep and sleep-disordered breathing in adults from seven U.S. urban areas'. *American Journal of Respiratory and Critical Care Medicine*. 2010; 182(6):819–25. [PubMed: 20508218]

Figure 1. Levels of cytokine production by mass of PM added per well (containing 100 000 BMdMs). The left and right graphs are IL-1 β and TNF- α levels, respectively, in response to PM and controls.

Figure 2. Cellular uptake of ambient PM₁₀. (A) 40X magnification of BMdMs containing PM₁₀, (B) Oil immersion 100X magnification showing fungi (spp. *Torula herbarum*) inside macrophage.

Figure 3.
PM Endotoxin content from winter and summer collected ambient particles.

Figure 4. Additivity values from different ratios of spring PM_{2.5} and PM_c mixtures. The left and right graphs are additivity values calculated from IL-1 β and TNF- α levels, respectively.

Table 1

Average and $\pm$ standard deviation of ambient PM concentrations ($\mu\text{g}/\text{m}^3$) in 2012, Missoula, MT.

	Winter Nov-Feb	Spring Mar-May	Summer June-Aug	Fall Sept-Oct
PM_C	9.3 $\pm$ 7.7	9.6 $\pm$ 7.7	13.8 $\pm$ 10.7	25.0 $\pm$ 30.4
PM_{2.5}	9.1 $\pm$ 4.9	5.9 $\pm$ 2.6	7.8 $\pm$ 6.4	18.1 $\pm$ 18.5
PM₁₀	18.4 $\pm$ 10.0	15.5 $\pm$ 9.8	21.6 $\pm$ 12.4	43.1 $\pm$ 37.0

Table 2

Average and $\pm$ standard deviation concentrations ($\mu\text{g}/\text{m}^3$) during winter and summer days when ambient particles were harvested.

	Winter	Spring	Summer
	Feb, 2012	May, 2012	July, 2011
PM_C	22.6 $\pm$ 9.2	7.4 $\pm$ 1.9	9.7 $\pm$ 4.4
PM_{2.5}	12.3 $\pm$ 3.0	5.3 $\pm$ 1.3	7.4 $\pm$ 2.1
PM₁₀	34.9 $\pm$ 8.5	12.7 $\pm$ 3.2	17.1 $\pm$ 6.5

Table 3

P-values comparing cytokine production of IL-1 β or TNF- α between two PM exposure types at 5,20, or 40 μ g.

	Winter PM _{2.5}	Winter PM _c	Summer PM _{2.5}	Summer PM _c
Winter PM _{2.5}	-----	*, 0.03, 0.01	*, 0.27, 0.16	*, 0.001, 0.003
Winter PM _c	<i>0.005, 0.0002, 0.0007</i>	-----	0.11, 0.04, 0.01	0.16, 0.06, 0.02
Summer PM _{2.5}	<i>0.009, 0.002, 0.17</i>	<i>0.008, 0.001, 0.001</i>	-----	0.10, 0.02, 0.01
Summer PM _c	<i>0.008, 0.008, 0.06</i>	<i>0.01, 0.01, 0.02</i>	<i>0.06, 0.09, 0.29</i>	-----

Each of the grouped (three) reported p-values from left to right are from 5, 20, and 40 μ g exposures, respectively. *Italicized* = TNF- α , normal type = IL-1 β .

* = cytokines not detected, cannot compare. Null hypothesis = Cytokines levels are the same between exposure treatments.