Differentiated thyroid tumors: surgical indications

R. LUCCHINI¹, M. MONACELLI¹, S. SANTOPRETE², R. TRIOLA¹, C. CONTI¹,
R. PECORIELLO¹, P. FAVORITI¹, M.S. DI PATRIZI¹, I. BARILLARO¹, A. BOCCOLINI¹,
S. AVENIA¹, M. D'AJELLO¹, A. SANGUINETTI¹, N. AVENIA¹

SUMMARY: Differentiated thyroid tumors: surgical indications.

R. LUCCHINI, M. MONACELLI, S. SANTOPRETE, R. TRIOLA, C. CONTI,
R. PECORIELLO, P. FAVORITI, M.S. DI PATRIZI, I. BARILLARO,
A. BOCCOLINI, S. AVENIA, M. D'AJELLO, A. SANGUINETTI, N. AVENIA

Thyroid gland tumors represent 1% of malignant tumors. In Italy

their incidence is in constant growth. The aggressiveness depends on the histological type. The relative non-aggressive grade of different forms of tumors is the basis for discussing the treatment of choice: total thyroidectomy vs lobectomy with or without lymphadenectomy of the sixth level in the absence of metastasis. Authors report about their experience, and they advocate, given the high percentage of multicentric forms, total thyroidectomy as treatment of choice.

KEY WORDS: Thyroid - Differentiated tumor - Surgery.

Introduction

Thyroid gland tumors represent 0.7 to 1% of malignant tumors, with an incidence estimated to be currently in Italy about 3600 new cases/year, in constant growth. 94% of these is represented by Differentiated Thyroid Cancer (DTC) i.e. papillary (80%), follicular (11%) and Hürthle cells (3%), 6% from anaplastic carcinoma. The aggressiveness is different depending on whether it is differentiated tumor, with which the majority of patients leads a life almost normal mean survival with 98% to 20 years, or of the variety that is anaplastic universally fatal within weeks of diagnosis (1). The relative non-aggressive grade of different forms means that even today we are discussing about what is the surgical treatment of choice for this type of tumor: total thyroidectomy *versus* lobectomy, with or without associated lymphectomy of the VI level even in the absence of lymph node metastases (2-6). In this review, we want to report our 7-

year experience on the surgical treatment of differentiated thyroid cancer.

Patients and methods

From January 2005 to December 2011, at the "AFOI (Area Funzionale Omogenea Interaziendale) of Endocrine Surgery of the Neck and Soft Tissues", in the Umbria region, Italy, 538 interventions for DTC were performed. Since 2005 we have seen a progressive increase in the number of cancers up to 2010 when we recorded a three times increase of cases in comparison to 2005. The increase is just partially justified from the technical improvement of diagnostic instruments. All the patients were subjected to total thyroidectomy; central compartment lymphadenectomy was performed for cancers > 2cm and for cancers < 2cm associated with lymphadenomegaly. Patients were discharged in the second to third postoperative day (patients outside the Region). Six patients (1%) had dysphonia and 30 patients hypocalcemia, in 15 of these the hypocalcemia has been definitive. Histologically 226 patients (42.7%) had multifocal distribution of tumor and 240 (44.6%) had lymph node metastases in the central compartment and 51 had monofocal tumors (Fig. 1).

Of the 226 patients with multifocal distribution of neoplastic disease, 133 (59%) were female and 93 (41%) males (Fig. 2)

In relation to the size, multifocality was found in 41 tumors with a diameter > 2 cm (18%), in 142 with diameter between 1 and 2 cm (63%) and in 43 with diameter < 1 cm (19%). In relation to the age, multifocality was found in 147 patients over 45 years (65%) and in 79 patients under 45 years (35%) (Fig. 3).

Of the patients with metastatic lymph nodes in central Compartment 174 were females (72.5%) and 66 males (27.5%) (Fig. 3).

Of these, 146 patients were older than 45 years (61%) and 94

University of Perugia, Terni, Italy

¹ Endocrine Surgical Unit

² Thoracic Surgical Unit

© Copyright 2013, CIC Edizioni Internazionali, Roma

Fig. 1 - DTC in our experience.

Fig. 2 - DTC plurifocal distribution for age and sex.

Fig. 3 - DTC + lymph node metastasis VI level.

under 45 years (39%); as regards the size, 50 patient (21%) had a tumor with diameter > 2 cm, 163 (68%) with diameter between 1 and 2 cm, and 26 (11%) with diameter <1 cm (Fig. 3).

In DTC with a diameter > 2 cm surgically treated, the positivity rate of the central compartment was 47%, as well as 47% was for cancer of diameter between 1 and 2 cm. The positivity rate for tumors <1 cm was 27%.

Discussion

Malignant tumors of the thyroid fit 11th place among all malignancies and the first of the endocrine system. The mortality, elevated to the forms anaplastic, is rather low for the differentiated. The rare deaths are recorded in elderly patients with aggressive forms poorly differentiated or locally (7, 8). In order to facilitate the choices of the appropriate treatments other prognostic variables are taken into considerations. These can be relating to patient (age, sex), tumor (size, multicentricity, histologic grade, histologic type, extrathyroidal invasion, lymph node metastasis, distant metastasis) or surgical procedure (complete and incomplete resection). These variables are then represented in the most common classification systems: AMES (Age, Metastasis, Extension, Size), AGES (Age, Grade, Extension, Size), MACE (Metastasis, Age, Completeness, Invasion, Size) and TNM, at the end of a proper identification of subjects at low and high risk. The role of the different prognostic factors in the definition of risk groups varies substantially between the authors (2-19). Shaha has catalogued all patients in low, medium and high risk with a mortality rate of 1%, respectively, 13%, 43% respectively (2). The risk can be discriminatory factor in choosing the surgical and post-surgical therapeutic strategy. The correct stratification of the risk can be made only on surgical specimen (20).

Molecular analysis of FNAB samples or surgical specimen provide useful information (21, 22). Genetic alterations (BRAF, ret / PTC, RAS, TRK for papillary carcinoma and RAS, PAX8-PPAR γ , PIK3CA, PTEN for follicular carcinoma) are meaningful indicators of the tumor aggressiveness (23-42). In micro carcinomas the percentage of metastatic spread to lymph nodes of the central compartment reaches up to 40% (3, 4). Percentage of recurrence of disease, in relation to age (> or <50 years), is not significantly different from tumors above and below 1 cm (5-7). If literature is concordant for the lymphadenectomy of level VI only in the presence of pathological lymph nodes, controversial is the opinion on total thyroidectomy versus lobectomy (5, 15). Everyone agrees on the total thyroidectomy in patients at high risk, in youth with lymphadenopathy, in bulky tumors and/or extracapsular disease and with cytological diagnosis of

poorly differentiated cancer (13, 14, 16, 20, 23, 25). For the new guidelines (20), total thyroidectomy is the most appropriate treatment for nodules > 1 cm with FNAB-positive for neoplasia in the presence of other nodules or contralateral lymph node metastases and/or distant metastases, with a history of neck irradiation or family history of thyroid cancer, for patients older than 45 years.

The lobectomy is sufficient for subcentimeter and unifocal cancer, low-risk cancer in the absence of pathological lymph nodes.

Rebecca I. Brown (23) consider total thyroidectomy the treatment of choice because prospective studies have showed increased and decreased disease-free period of relapse, even allowing for the treatment ablation with iodine 131 (21-22). We believe, from the literature given the high incidence of multicentric tumors (30-85%), and the percentage relatively low complications, according to some authors, (9, 6, 23) always make the total thyroidectomy (TT) because:

- in 40-70% of cases there are microscopic foci in the contralateral lobe;
- Total thyroidectomy removes the entire thyroid tissue allowing: the use of thyroglobulin recurrence as a cancer or distant metastasis marker; easier replacement therapy with thyroxine; effective ablative treatment with radioiodine;
- TT reduces the chance of anaplastic or poorly differentiated tumor in residual parenchyma;
- the correct assessment of medium or low risk is only postoperative, on the basis of tumor size and extracapsular infiltration (3, 4).

Conclusions

As unanimously recognized, the goal of an adequate surgical approach to DTC is to: remove the primary tumor and lymph nodes affected by the disease; to minimize the morbidity related to treatment; to facilitate the staging of the disease, and the treatment with 131 I (9); to allow an accurate surveillance in the long term; and reduce the risk of recurrent disease and metastatic spread. Then we consider TT appropriate for DTC, considering the possible multifocality and dissemination in lymph nodes of the central compartment also for tumors with a diameter <1 cm that the approach. However, surgery should consist in the execution of total thyroidectomy and central compartment lymphadenectomy when pathological. In this context, the formulation of guidelines and a multidisciplinary approach allows more uniform and shared approach to diagnostic work up and treatment and more meaningful comparison of the data.

References

- Linee guida per la gestione dei pazienti affetti da noduli tiroidei e carcinoma della tiroide, Ottobre 2008. I quaderni, n°13. A cura della Direzione Regionale Sanità e Servizi Sociali della regione Umbria.
- Shaha AR. Advances in the management of thyroid cancer. *Int J Surgery* 2005;3:213-220.
- Besic N. Extent of thyroidectomy and lymphadenectomy in 254 patients with papillary thyroid microcarcinoma: a single-institution experience. *Ann Surg Oncol* 2009;16(4):920-8.
- Salter KD, Andersen PE, Cohen JI, Schuff KG, Lester L, Shindo ML, Sauer D, Gross ND. Central nodal metastases in papillary thyroid carcinoma based on tumor histologic type and focality. *Arch Otolaryngol Head Neck Surg* 2010;136(7):692-6.
- Noguchi S, Yamashita H, Uchino S, Watanabe S. Papillary microcarcinoma. *World J Surg* 2008;32:747-753.
- Arora N. Papillary thyroid carcinoma and microcarcinoma: is there a need to distinguish the two? *Thyroid* 2009 May;19(5):473-7.
- Clark OH. Total thyroidectomy: the treatment of choice for patient with differentiated thyroid cancer. *Ann Surg* 1982;196:361-70.
- Tuttle RM, Tala H, Shah J, Leboeuf R, Ghossein R, Gonen M, Brokhin M, Omry G, Fagin JA, Shaha A. Estimating risk of recurrence in differentiated thyroid cancer after total thyroidectomy and radioactive iodine remnant ablation: using response to therapy variables to modify the initial risk estimates predicted by new American thyroid association staging system. *Thyroid* 2010;20:1341-50.
- Avenia N, Cristofani R, Puma F, Fonsi GB, Scalercio V, Urbani M, Ricci E, Servoli A. Il trattamento chirurgico del cancro della tiroide: analisi di 207 casi trattati. *Osp Ital Chir* 1999;5:451-6.
- Milan SA, Sosa JA, Roman SA. Chirurgia mini-invasiva per il cancro invasivo della tiroide. *Min Chir* 2010 Febbraio;65(1):39-43.
- Ducic Y, Young L, McIntyre J. Approccio "multi-target" nel trattamento del cancro della tiroide. *Min Chir* 2010 Febbraio;65(1):59-69.
- Zarebczan B, Chen H. Problematiche nella chirurgia del cancro della tiroide. *Min Chir* 2010;65(1):83-93.
- McHenry CR, Phitayakorn R. Follicular adenoma and carcinoma of the thyroid gland. *Oncologist* 2011;16(5):585-93. Epub 2011 Apr 11.
- Ahmadi N, Grewal, Davidson BJ. Patterns of Cervical Lymph Node Metastases in Primary and Recurrent Papillary Thyroid Cancer. *J Oncol* 2011;2011:735678. Epub 2011 Nov 17.
- Kuru B, Topgul K. The 2009 Revised American Thyroid Association Guidelines for Thyroid Cancer: The Extent of Surgery for Thyroid Carcinoma Less Than One and One Half Centimeters or Low-Risk Thyroid Carcinoma. *Thyroid* 2011 Dec 16.
- Milas Z., Shin J., Milas M. New guidelines for the management of thyroid nodules and differentiated thyroid cancer. *Minerva Endocrinologica* 2011 March;36(1):53-70.
- Falvo L, Catania A, D'Andrea V, Marzullo A, Giustiniani MC, De Antoni E. Prognostic importance of histologic vascular invasion in papillary thyroid carcinoma. *Ann Surg* 2005; 241:640-6.
- Falvo L, Catania A, Sorrenti S, D'Andrea V, Berni A, De Stefano M, De Antoni E. Prognostic significance of the age factor in the thyroid cancer: statistical analysis. *J Surg Oncol* 2004;88:217-22.
- Falvo L, D'Ercole C, Sorrenti S, D'Andrea V, Catania A, Berni A, Grilli P, De Antoni E. Papillary microcarcinoma of the thyroid gland: analysis of prognostic factors including histological subtype. *Eur J Surg Suppl* 2003 Jul;(588):28-32.
- González-González A, Mate-Valdezate A, Parra-Arroyo A, Tenías-Burillo JM. New guidelines for the management of thyroid nodules and differentiated thyroid cancer. *Minerva Endocrinol* 2011 Mar;36(1):7-12.
- Baldini E, Sorrenti S, Catania A, Guaitoli E, Prinzi N, Mocini R, Nardi F, D'Armiento E, Bianchini M, Favoriti P, Di Matteo FM, Ruggieri M, De Antoni E, Ulisse S. Diagnostic utility of thyroglobulin measurement in the fine needle aspirates from cervical lymph nodes: a case report. *G Chir* 2012;33:387-91.
- Baldini EP, Sorrenti S, Di Gioia C, De Vito C, Antonelli A, Gnesi L, Carbotta G, D'Armiento E, Miccoli P, De Antoni E, Ulisse S. Cervical lymph node metastases from thyroid cancer: does thyroglobulin and calcitonin measurement in fine needle aspirates improve the diagnostic value of cytology? *BMC Clin Pathol*;2013;13:7.
- Brown RL, de Souza JA, Cohen EE. Thyroid cancer: burden of illness and management of disease. *J Cancer* 2011 Apr 4;2:193-9.
- Moretti S, De Falco V, Tamburrino A, Barbi F, Tavano M, Avenia N, Santeusano F, Santoro M, Macchiarulo A, Puxeddu E. Insights into the molecular function of the inactivating mutations of B-Raf involving the DFG motif. *Biochim Biophys Acta* 2009 Nov;1793(11):1634-45. Epub 2009 Sep 6.
- Puxeddu E, Durante C, Avenia N, Filetti S, Russo D. Clinical implications of BRAF mutation in thyroid carcinoma. *Trends Endocrinol Metab* 2008 May-Jun;19(4):138-45. Epub 2008 Mar 11.
- Hay ID, Bergstralh EJ, Goellner JR, Ebersold JR, Grant CS. Predicting outcome in papillary thyroid carcinoma: development of a reliable prognostic scoring system in a cohort of 1779 patients surgically treated at one institution during 1940 through 1989. *Surgery* 1993;114(6):1050-7.
- Mazzaferri EL, Young RL. Papillary thyroid carcinoma: 10 year follow up report of the impact of therapy in 576 patients. *Am J Med* 1981;70(3):511-8.
- Cirocchi R, Trastulli S, Sanguinetti A, Cattorini L, Covarelli P, Giannotti D, Di Rocco G, Rondelli F, Barberini F, Boselli C, Santoro A, Gullà N, Redler A, Avenia N. Recurrent differentiated thyroid cancer: to cut or burn. *World J Surg Oncol* 2011 Aug 12;9:89.
- Moretti S, De Falco V, Tamburrino A, Barbi F, Tavano M, Avenia N, Santeusano F, Santoro M, Macchiarulo A, Puxeddu E. Insights into the molecular function of the inactivating mutations of B-Raf involving the DFG motif. *Biochim Biophys Acta* 2009 Nov;1793(11):1634-45.
- Rosato L, Miccoli P, Pinchera A, Lombardi G, Romano M, Avenia N, Bastagli A, Bellantone R, De Palma M, De Toma G, Gasparri G, Lampugnani R, Marini PL, Nasi PG, Pellizzo MR, Pezzullo L, Piccoli M, Testini M. Diagnostic, therapeutic and health-care management protocols in thyroid surgery. 2nd Consensus Conference (U.E.C. CLUB)]. *G Chir* 2009 Mar;30(3):73-86.
- Lucchini R, Puxeddu E, Calzolari F, Burzelli F, Monacelli M, D'Ajello F, Macaluso R, Giammartino C, Ragusa M, De Feo P, Cavaliere A, Avenia N. Recurrences of thyroid well differentiated cancer: ultrasonography-guided surgical treatment. *Minerva Chir* 2008 Aug 63(4):257-60.
- Puxeddu E, Durante C, Avenia N, Filetti S, Russo D. Clinical implications of BRAF mutation in thyroid carcinoma. *Trends Endocrinol Metab* 2008 May-Jun;19(4):138-45.

33. Puxeddu E, Susta F, Orvietani PL, Chiasserini D, Barbi F, Moretti S, Cavaliere A, Santeusano F, Avenia N, Binaglia L. Identification of differentially expressed proteins in papillary thyroid carcinomas with V600E mutation of BRAF. *Proteomics Clin Appl* 2007 Jul;1(7):672-80.
34. Durante C, Puxeddu E, Ferretti E, Morisi R, Moretti S, Bruno R, Barbi F, Avenia N, Scipioni A, Verrienti A, Tosi E, Cavaliere A, Gulino A, Filetti S, Russo D. BRAF mutations in papillary thyroid carcinomas inhibit genes involved in iodine metabolism. *J Clin Endocrinol Metab* 2007 Jul;92(7):2840-3. Epub 2007 May 8.
35. Fugazzola L, Puxeddu E, Avenia N, Romei C, Cirello V, Cavaliere A, Faviana P, Mannavola D, Moretti S, Rossi S, Sculli M, Bottici V, Beck-Peccoz P, Pacini F, Pinchera A, Santeusano F, Elisei R. Correlation between B-RAFV600E mutation and clinico-pathologic parameters in papillary thyroid carcinoma: data from a multicentric Italian study and review of the literature. *Endocr Relat Cancer* 2006 Jun;13(2):455-64.
36. Moretti S, Macchiarulo A, De Falco V, Avenia N, Barbi F, Carta C, Cavaliere A, Melillo RM, Passeri L, Santeusano F, Tartaglia M, Santoro M, Puxeddu E. Biochemical and molecular characterization of the novel BRAF(V599Ins) mutation detected in a classic papillary thyroid carcinoma. *Oncogene* 2006 Jul 13;25(30):4235-40.
37. Carta C, Moretti S, Passeri L, Barbi F, Avenia N, Cavaliere A, Monacelli M, Macchiarulo A, Santeusano F, Tartaglia M, Puxeddu E. Genotyping of an Italian papillary thyroid carcinoma cohort revealed high prevalence of BRAF mutations, absence of RAS mutations and allowed the detection of a new mutation of BRAF oncoprotein (BRAF(V599Ins)). *Clin Endocrinol (Oxf)* 2006 Jan;64(1):105-9.
38. Puxeddu E, Moretti S, Elisei R, Romei C, Pascucci R, Martinelli M, Marino C, Avenia N, Rossi ED, Fadda G, Cavaliere A, Ribacchi R, Falorni A, Pontecorvi A, Pacini F, Pinchera A, Santeusano F. BRAF(V599E) mutation is the leading genetic event in adult sporadic papillary thyroid carcinomas. *J Clin Endocrinol Metab* 2004 May;89(5):2414-20.
39. Rosato L, Avenia N, Bernante P, De Palma M, Gulino G, Nasi PG, Pelizzo MR, Pezzullo L. Complications of thyroid surgery: analysis of a multicentric study on 14,934 patients operated on in Italy over 5 years. *World J Surg* 2004 Mar;28(3):271-6.
40. Puxeddu E, Moretti S, Giannico A, Martinelli M, Marino C, Avenia N, Cristofani R, Farabi R, Reboldi G, Ribacchi R, Pontecorvi A, Santeusano F. Ret/PTC activation does not influence clinical and pathological features of adult papillary thyroid carcinomas. *Eur J Endocrinol* 2003 May;148(5):505-13.
41. Ulisse S, Baldini E, Sorrenti S, Barollo S, Prinzi N, Catania A, Nescia A, Gnessi L, Pelizzo MR, Mian C, De Vito C, Calvanese A, Palermo S, Persechino S, De Antoni E, D'Armiento M. In papillary thyroid carcinoma BRAFV600E is associated with increased expression of the urokinase plasminogen activator and its cognate receptor, but not with disease-free interval. *Clin Endocrinol* 2012;77:780-6.
42. Ulisse S, Baldini E, Sorrenti S, Barollo S, Gnessi L, Catania A, Pelizzo MR, Nardi F, Mian C, De Antoni E, D'Armiento M, Frati L. High expression of the urokinase plasminogen activator and its cognate receptor associates with advanced stages and reduced disease-free interval in papillary thyroid carcinoma. *J Clin Endocrinol Metab* 2011;96:504-8.