Published in final edited form as:

Neurochem Res. 2014 June ; 39(6): 1057–1067. doi:10.1007/s11064-013-1156-3.

Unexpected Properties of δ -Containing GABA_A Receptors in Response to Ligands Interacting with the α + β - Site

Pantea Mirheydari[#],

Department of Biochemistry and Molecular Biology, Center for Brain Research, Medical University Vienna, Spitalgasse 4, 1090 Vienna, Austria

Joachim Ramerstorfer[#],

Department of Biochemistry and Molecular Biology, Center for Brain Research, Medical University Vienna, Spitalgasse 4, 1090 Vienna, Austria

Zdravko Varagic,

Department of Biochemistry and Molecular Biology, Center for Brain Research, Medical University Vienna, Spitalgasse 4, 1090 Vienna, Austria

Petra Scholze,

Department of Pathobiology of the Nervous System, Center for Brain Research, Medical University Vienna, Spitalgasse 4, 1090 Vienna, Austria

Laurin Wimmer,

Institute of Applied Synthetic Chemistry, Vienna University of Technology, Getreidemarkt 6/163, 1060 Vienna, Austria

Marko M. Mihovilovic,

Institute of Applied Synthetic Chemistry, Vienna University of Technology, Getreidemarkt 6/163, 1060 Vienna, Austria

Werner Sieghart, and

Department of Biochemistry and Molecular Biology, Center for Brain Research, Medical University Vienna, Spitalgasse 4, 1090 Vienna, Austria

Margot Ernst

Department of Biochemistry and Molecular Biology, Center for Brain Research, Medical University Vienna, Spitalgasse 4, 1090 Vienna, Austria. Margot.Ernst@meduniwien.ac.at

[#] These authors contributed equally to this work.

Abstract

GABA_A receptors are the major inhibitory neurotransmitter receptors in the central nervous system and are the targets of many clinically important drugs, which modulate GABA induced chloride flux by interacting with separate and distinct allosteric binding sites. Recently, we described an allosteric modulation occurring upon binding of pyrazoloquinolinones to a novel

binding site at the extracellular $\alpha + \beta$ - interface. Here, we investigated the effect of 4-(8-methoxy-3-oxo-3,5-dihydro-2H-pyrazolo[4,3-c]quinolin-2-yl)benzonitrile (the pyrazoloquinolinone LAU 177) at several $\alpha\beta$, $\alpha\beta\gamma$ and $\alpha\beta\delta$ receptor subtypes. LAU 177 enhanced GABA-induced currents at all receptors investigated, and the extent of modulation depended on the type of α and β subunits present within the receptors. Whereas the presence of a $\gamma 2$ subunit within $\alpha\beta\gamma 2$ receptors did not dramatically change LAU 177 induced modulation of GABA currents compared to $\alpha\beta$ receptors, we observed an unexpected threefold increase in modulatory efficacy of this compound at $\alpha 1\beta 2,3\delta$ receptors. Steric hindrance experiments as well as inhibition by the functional $\alpha + \beta$ - site antagonist LAU 157 indicated that the effects of LAU 177 at all receptors investigated were mediated via the $\alpha + \beta$ - interface. The stronger enhancement of GABA-induced currents by LAU 177 at $\alpha 1\beta 3\delta$ receptors was not observed at $\alpha 4,6\beta 3\delta$ receptors. Other experiments indicated that this enhancement of modulatory efficacy at $\alpha 1\beta 3\delta$ receptors was not observed with another $\alpha + \beta$ - modulator, and that the efficacy of modulation by $\alpha + \beta$ - ligands is influenced by all subunits present in the receptor complex and by structural details of the respective ligand.

Keywords

GABA_A; CGS 9895; Pyrazoloquinolinones; $\alpha + \beta$ - Binding site; Positive modulators; Null modulators; δ Subunit; Extrasynaptic receptor

Introduction

GABA_A receptors are ligand-gated chloride channels composed of five subunits. The existence of 6 α , 3 β , 3 γ , δ , ϵ , π , θ , and 3 ρ subunits gives rise to a multiplicity of GABA_A receptor subtypes with distinct subunit composition and pharmacological and electrophysiological properties. Most of these receptors are composed however of one γ , two α and two β subunits. The δ , ϵ , and π subunits have been thought to replace the γ subunit in $\alpha\beta\gamma$ receptor subtypes [1]. However, the subunit stoichiometry and arrangement of receptors containing the δ or ϵ subunit have recently been questioned by demonstrating that several recombinant concatenated δ - or ϵ -containing receptors with different subunit arrangements can be formed [2-6]. The ρ subunits can either form homo-oligomers or hetero-oligomers with other ρ subunits [1].

GABA_A receptors are the site of action of a variety of clinically important drugs, such as benzodiazepines, barbiturates, neuroactive steroids, anesthetics, convulsants, and others [7]. All these drugs seem to allosterically interact with binding sites that are partially or completely distinct from each other. Modelling studies have identified a total of at least 16 solvent accessible spaces within GABA_A receptors that also could represent drug binding sites [8]. Thus, within the extracellular domain of GABA_A receptors composed of 2 α , 2 β , and 1 γ subunit, at least four binding sites seem to exist, each one located at a subunit interface. Each subunit per definition contains a plus (+) and a minus (−) side and the (+) side of one subunit forms an interface with the (−) side of the neighbouring subunit [9]. The two gamma-aminobutyric acid (GABA) binding sites are located at the two $\beta + \alpha$ - interfaces [10], the benzodiazepine binding site is located at the $\alpha + \gamma$ - interface [11] and a recently

identified binding site for pyrazoloquinolinones such as CGS 9895 has been located at the $\alpha + \beta$ - interface (Fig. 1a) [12]. It can be assumed that the remaining $\gamma + \beta$ - interface might also be the site of action of some drugs, but so far, direct evidence for this assumption is lacking. In addition to these extracellular binding sites at the subunit interfaces, another type of binding site has been identified within individual subunits of the extracellular domain of the GABA activated bacterial *Erwinia chrysanthemi* ligand-gated ion channel (ELIC) [13]. In the trans membrane (TM) domain, mutagenesis and photo labelling studies have suggested binding sites for volatile anesthetics, intravenous anesthetics, steroids, barbiturates, and ethanol [14-17]. These binding sites were partially assigned to subunit interfaces, or to the space inside the four helix bundle of each subunit of the GABA_A receptor [18-22]. The actual existence of binding sites at these positions was demonstrated by crystallization studies using homologous proteins [23, 24]. Recently, a novel binding site for the endocannabinoid 2-arachidonyl glycerol was reported and localized between the TM3 and TM4 helices of the $\beta 2$ subunit [25].

The $\alpha +$ side not only contributes to the $\alpha + \beta$ - interface but also to the $\alpha + \gamma$ - interface. It thus was no surprise that the pyrazoloquinolinone CGS 9895 is able to interact with both interfaces. This compound acts as a high affinity null modulator at the benzodiazepine binding site ($\alpha + \gamma$ - interface) and as a low potency positive allosteric modulator via the $\alpha + \beta$ - interface [12]. These observations are consistent with previous findings indicating that many of the drugs interacting with GABA_A receptors seem to do that via more than one binding site, as indicated by their different actions at different drug concentrations [7, 20].

In a subsequent study we identified 29 structural analogues of CGS 9895 that either behaved as positive allosteric modulators or null modulators via the $\alpha + \beta$ - interface of GABA_A receptors [26]. 16 of these compounds were then further investigated for their effects at GABA_A receptor subtypes composed of $\alpha 1,2,3,5\beta 3$ or $\alpha 1-6\beta 3\gamma 2$ subunits [27]. Results indicated that most of the compounds investigated exhibit comparable potency and efficacy for $\alpha\beta$ and $\alpha\beta\gamma$ receptors containing the same type of α or β subunit. Some small differences in the effects elicited in $\alpha\beta$ and $\alpha\beta\gamma$ receptors were explained by a possible allosteric interaction of the compounds bound to the benzodiazepine site and the $\alpha + \beta$ - site.

To possibly identify more receptor subtype-selective ligands in our compound library, in the present study we investigated some of the previously published compounds that so far have not been studied for their effects at various receptor subtypes [26, 27]. For that we started with LAU 177, one of the compounds exhibiting the highest efficacy at $\alpha 1\beta 3$ or $\alpha 1\beta 3\gamma 2$ receptors [26]. Recently, evidence accumulated that extra synaptic receptors such as $\alpha 1,4,6\beta\delta$ and $\alpha 5\beta\gamma$ receptors might have important functions in health and disease [28]. We therefore extended these studies by not only measuring the effects of this compound at receptors composed of $\alpha 1,2,3,5\beta 3$ or $\alpha 1-6\beta 3\gamma 2$ subunits, but also at those composed of $\alpha 1,4,6\beta 3\delta$ subunits. Surprisingly, LAU 177 exhibited pronounced differences in its efficacy between $\alpha 1\beta 3$, $\alpha 1\beta 3\gamma 2$, and $\alpha 1\beta 3\delta$ receptors. These and other results indicated that the presence of the $\gamma 2$ or δ subunit in GABA_A receptors somehow contribute to the efficacy of $\alpha + \beta$ - site ligands.

Methods

Two Electrode Voltage Clamp (TEV)

In vitro transcription of mRNA was based on the cDNA expression vectors encoding for GABA_A receptor subunits $\alpha 1$ –6, $\beta 1$ –3, $\gamma 2$ and δ (all from rat) [29]. After linearizing the cDNA vectors with appropriate restriction endonucleases, capped transcripts were produced using the *mMES-SAGE mMACHINE*® T7 transcription kit (Ambion, TX, USA). The capped transcripts were polyadenylated using yeast poly (A) polymerase (USB, OH, USA) and were diluted and stored in diethylpyrocarbonate-treated water at -70°C .

The methods for isolating, culturing, injecting, and defolliculating of oocytes were identical with those described by E. Sigel [30]. Mature female *Xenopus laevis* (Nasco, WI, USA) were anaesthetized in a bath of ice-cold 0.17 % Tricain (Ethyl-m-aminobenzoat, Sigma, MO, USA) before decapitation and removal of the frog's ovary. Stage 5–6 oocytes with the follicle cell layer around them were singled out of the ovary using a platinum wire loop. Oocytes were stored and incubated at 18°C in modified Barths' Medium [88 mM NaCl, 10 mM HEPES–NaOH (pH 7.4), 2.4 mM NaHCO₃, 1 mM KCl, 0.82 mM MgSO₄, 0.41 mM CaCl₂, 0.34 mM Ca(NO₃)₂] that was supplemented with 100 U/ml penicillin and 100 $\mu\text{g/ml}$ streptomycin. Oocytes with follicle cell layer still around them were injected with an aqueous solution of mRNA. A total of 2.5–4 ng of mRNA per oocyte was injected. Subunit ratio was 1:1:5 for $\alpha\times\beta 3\gamma 2$ receptors, 3:1:5 for $\alpha\times\beta 3\delta$ and 1:1 for $\alpha\times\beta 3$ receptors consisting of wild-type or mutated α subunits together with wild-type or mutated $\beta 3$ subunits. After injection of mRNA, oocytes were incubated for at least 24 h for $\alpha\beta$ and $\alpha\beta 6$ receptors and for at least 36 h for $\alpha\beta 2$ receptors before the enveloping follicle cell layers were removed. Collagenase-treatment (type IA, Sigma, MO, USA) and mechanical defolliculation of the oocytes was performed as described previously.

For electrophysiological recordings, oocytes were placed on a nylon-grid in a bath of Xenopus Ringer solution (XR, containing 90 mM NaCl, 5 mM HEPES–NaOH (pH 7.4), 1 mM MgCl₂, 1 mM KCl and 1 mM CaCl₂). For current measurements the oocytes were impaled with two microelectrodes (1–2 M Ω) which were filled with 2 M KCl. The oocytes were constantly washed by a flow of 6 ml/min XR that could be switched to XR containing GABA and/or drugs. Drugs were diluted into XR from DMSO-solutions resulting in a final concentration of 0.1 % DMSO perfusing the oocytes. Drugs were pre-applied for 30 s before the addition of GABA, which was then coapplied with the drugs until a peak response was observed. Between two applications, oocytes were washed in XR for up to 15 min to ensure full recovery from desensitization. Maximum currents measured in mRNA injected oocytes were in the published [26, 31] range for all wild type receptors. To test for modulation of GABA induced currents by compounds, a GABA concentration titrated to trigger 3–7 % of the respective maximum GABA-elicited current of the individual oocyte (=GABA EC₃) was applied to the cell together with various concentrations of compounds to be tested. All recordings were performed at room temperature at a holding potential of -60 mV using a Warner OC-725C two-electrode voltage clamp (Warner Instrument, Hamden, CT, USA) or a Dagan CA-1B Oocyte Clamp or a Dagan TEV-200A two-electrode voltage clamp (Dagan Corporation, Minneapolis, MN, USA). Data were digitized, recorded and measured using a

Digidata 1322A data acquisition system (Axon Instruments, Union City, CA, USA). Data were analyzed using GraphPad Prism. Data for GABA dependent dose-response curves were fitted to the equation $Y = \text{bottom} + (\text{top} - \text{bottom}) / (1 + 10^{(\text{LogEC}_{50} - X) * nH})$, where EC_{50} is the concentration of the compound that increases the amplitude of the GABA-evoked current by 50 %, and nH is the Hill coefficient. The bottom was restrained to 100 %, reflecting the GABA control current. Data are given as mean $\pm$ SEM (standard error of mean) from at least three oocytes of two or more oocyte batches. Statistical significance was determined by unpaired Student's t test and paired Student's t test for GABA concentration-response curves in the absence or presence of modulator at $\alpha 1\beta 3\delta$ receptors at a confidence interval of $P < 0.05$.

MTSEA-Biotin—Steric Hindrance

2 mM MTSEA-biotin (*N*-Biotinylaminoethyl methane-thiosulfonate) solution was freshly made in XR buffer containing the respective GABA- EC_3 concentration. Defolliculated oocytes were immediately immersed in the MTSEA-biotin solution for 3 min and washed with XR for 5 min. After the washing step, cells were used the same day for the electrophysiological recordings described above.

Materials

GABA_A Receptor Subunits and Point Mutations

cDNAs of rat GABA_A receptor subunits $\alpha 1$, $\alpha 4$, $\beta 1$, $\beta 2$, $\beta 3$, and $\gamma 2S$ were cloned as described [32]. cDNAs of the rat subunits $\alpha 2$, $\alpha 3$, and $\alpha 5$ were gifts from P. Malherbe, that of $\alpha 6$ was a gift of P. Seeburg, and that of δ was a gift of C. Czajkowski. The mutated construct $\alpha 1S204C$ was a gift from E. Sigel. For the generation of mutated $\beta 3$ subunit, this subunit was subcloned into the pCDM8 expression vector (Invitrogen, San Diego, CA, USA) as described previously [33]. Mutated subunits were constructed by PCR amplification using the wild-type subunit as a template. For this, PCR primers were used to construct point mutations within the subunits by the 'gene splicing by overlap extension' technique [34]. The PCR primers for $\beta 3Q64C$ contained *Xma*I and *Xho*I restriction sites, which were used to clone the $\beta 3$ fragments into pCI vector (Promega, Madison, WI, USA). The mutated subunits were confirmed by sequencing.

Compound Synthesis

Synthesis of LAU compounds was performed in analogy to previously outlined synthetic routes [35, 36].

Investigated Compounds

The following compounds were used: (LAU 177): 4-(8-methoxy-3-oxo-3,5-dihydro-2H-pyrazolo[4,3-c]quinolin-2-yl)benzonitrile. (LAU 157): 8-chloro-2-(4-nitrophenyl)-2H-pyrazolo[4,3-c]quinolin-3(5H)-one. (PZ-II-028): 8-chloro-2-(4-methoxyphenyl)-2H-pyrazolo[4,3-c]quinolin-3(5H)-one (was a gift from J. Cook). (DS2): 4-Chloro-*N*-[2-(2-thienyl)imidazo[1,2-*a*]pyridin-3-yl]benzamide (Tocris, Bristol, UK). (Tracazolate): 4-(butylamino)-1-ethyl-6-methyl-1H-pyrazolo[3,4-*b*]pyridine-5-carboxylate (Sigma Aldrich, AT).

Results

In a search for GABA_A receptor subtype-selective compounds we started to investigate the properties of the pyrazoloquinolinone LAU 177 (Fig. 2a) in more detail. This compound is a structural analogue of CGS 9895 [12], and as reported previously [26] and similar to CGS 9895, LAU 177 acts as a high affinity ligand at the benzodiazepine binding site of $\alpha 1\beta 3\gamma 2$ receptors (IC_{50} of [³H]Ro15-1788 binding of 0.75 ± 0.81 nM), and as a strong positive allosteric modulator ($EC_{50} = 1$ μ M, stimulation to 1063 ± 128 % of GABA EC_3 at 10 μ M concentration) at $\alpha 1\beta 3$ receptors. For the latter experiments $\alpha 1\beta 3$ receptors were used, to exclude effects mediated via the high affinity benzodiazepine binding site at the $\alpha + \gamma$ -interface (Fig. 1, Supplementary Table 1). To confirm that LAU 177 mediates its effect at $\alpha 1\beta 3$ receptors via the $\alpha 1 + \beta 3$ - interface, we here again employed the substituted cysteine accessibility method to introduce a steric hindrance into the $\alpha 1 + \beta 3$ - interface of $\alpha 1\beta 3$ receptors. The point mutations $\alpha 1S204C$ (loop C of the $\alpha 1+$ side) and $\beta 3Q64C$ (loop D of the $\beta 3$ - side) have been shown previously to not significantly change the potency or efficacy of GABA for enhancing GABA-induced currents at $\alpha 1\beta 3$ or $\alpha 1\beta 3\gamma 2$ receptors [12]. Recombinant $\alpha 1S204C\beta 3Q64C$ receptors were expressed in *Xenopus* oocytes and the effects of various concentrations of LAU 177 were investigated and compared with those at $\alpha 1\beta 3$ receptors. As shown in Fig. 2b, c, in agreement with previous results [26] LAU 177 enhanced GABA-induced currents at $\alpha 1\beta 3$ receptors in a concentration dependent way up to 1152 ± 145 % of GABA EC_3 . LAU 177 also enhanced GABA-induced currents in $\alpha 1S204C\beta 3Q64C$ receptors to a similar extent with a comparable potency. In the presence of MTSEA-biotin, however, current enhancement by LAU 177 was drastically reduced at the mutated receptor (Fig. 2b, c), indicating that LAU 177 exerted most of its action via the $\alpha 1 + \beta 3$ - binding site, as expected. In previous studies no change in potency and efficacy of a total of four different pyrazoloquinolinones has been observed when wild-type instead of mutated $\alpha 1\beta 3$ receptors were incubated with MTSEA-biotin [12, 26].

In other experiments, the effects of LAU 177 were compared at various $\alpha\beta$ and $\alpha\beta\gamma 2$ receptor subtypes (Fig. 3a, b). In agreement with previous results with other pyrazoloquinolinones [26] a similar concentration-dependent effect of LAU 177 at GABA EC_3 was obtained in $\alpha 1\beta 3$ and $\alpha 1\beta 3\gamma 2$ receptors, (maximum stimulation at $\alpha 1\beta 3\gamma 2$ receptors to about 1179 ± 143 % of GABA EC_3 at 10 μ M concentrations). The absence of strong LAU 177 effects at $\alpha 1\beta 3\gamma 2$ receptors at 10–100 nM concentrations, that are sufficient to saturate the benzodiazepine binding site of these receptors, suggested that this compound acts as a high affinity null modulator at the benzodiazepine binding site of GABA_A receptors, similar to CGS 9895 and other pyrazoloquinolinones [26, 27]. The similarity in the dose response curves between $\alpha 1\beta 3$ and $\alpha 1\beta 3\gamma 2$ receptors (Fig. 3a, b, Supplementary Tables 1 and 2) suggested that LAU 177 mediates most of its effect at $\alpha 1\beta 3\gamma 2$ receptors via the $\alpha 1 + \beta 3$ - interface.

To investigate a possible receptor subtype-selectivity, the effects of LAU 177 were also investigated at $\alpha 2\beta 3$, $\alpha 3\beta 3$, and $\alpha 5\beta 3$ receptors (Fig. 3a, Supplementary Table 1). LAU 177 was a comparably strong modulator of GABA EC_3 at $\alpha 2\beta 3$ and $\alpha 3\beta 3$ receptors, reaching a stimulation of up to 981 ± 20 and 913 ± 178 % of GABA EC_3 at 10 μ M concentrations. The effects of LAU 177 at $\alpha 5\beta 3$ receptors, however, were weaker and resulted in an

enhancement of the GABA-induced current to about 528 ± 103 % GABA $10 \mu\text{M}$. Whereas in $\alpha 2\beta 3\gamma 2$ and $\alpha 3\beta 3\gamma 2$ receptors the effects of LAU 177 were comparable to those of $\alpha 2\beta 3$ and $\alpha 3\beta 3$ receptors, the effects of this compound at $\alpha 5\beta 3\gamma 2$ was stronger than at $\alpha 5\beta 3$ receptors (Fig. 3b, Supplementary Tables 1 and 2). As with several other pyrazoloquinolinones [27], LAU 177 exhibited a stronger stimulation at $\alpha 6\beta 3\gamma 2$ receptors than at all other receptors investigated. The effects of LAU 177 at $\alpha 4\beta 3\gamma 2$ receptors, however, were comparable to those observed at all the other $\alpha \times \beta 3\gamma 2$ receptors (Fig. 3b, Supplementary Table 2).

So far, we did not investigate the effects of pyrazoloquinolinones at GABA_A receptors containing a δ instead of a $\gamma 2$ subunit. Such receptors are located exclusively extra- and peri-synaptically, exhibit a high sensitivity to GABA, show little desensitization and are believed to be one of the primary mediators of tonic inhibition [37]. So far, only $\alpha 1\beta \delta$ [38], $\alpha 4\beta \delta$ [39] or $\alpha 6\beta \delta$ receptors [40] have been more or less unequivocally identified in the mammalian brain. We therefore investigated only these receptor subtypes in the present study. To confirm the actual incorporation of the delta subunit into $\alpha \beta \delta$ receptors, GABA-induced currents at all *Xenopus* oocytes injected with mRNAs of α , β and δ subunits were tested by the addition of $10 \mu\text{M}$ of the imidazopyridine DS2 [41]. At this concentration DS2 dramatically enhanced GABA-induced currents at $\alpha \beta \delta$, but not at $\alpha \beta$ or $\alpha \beta \gamma 2$ receptors [41] (experiments not shown). Since the pyrazoloquinolinones mediate their effects via the $\alpha + \beta$ - interface, it was tacitly assumed that the effects of these compounds would be comparable whether a δ or a $\gamma 2$ subunit is present in the receptors. Surprisingly, however, LAU 177 exhibited a much stronger effect at $\alpha 1\beta 3\delta$ (stimulation of GABA EC_3 to 3285 ± 257 % at $10 \mu\text{M}$ concentration) than at $\alpha 1\beta 3\gamma 2$ receptors (stimulation of GABA EC_3 to 1179 ± 143 % at $10 \mu\text{M}$ concentration) (Fig. 3b, c, Supplementary Tables 2 and 3). Similarly, the effects of LAU 177 at $\alpha 4\beta 3\delta$ receptors (stimulation of GABA EC_3 to 1365 ± 117 % at $10 \mu\text{M}$ concentration) were stronger than at $\alpha 4\beta 3\gamma 2$ receptors (stimulation of GABA EC_3 to 981 ± 52 % at $10 \mu\text{M}$ concentration), whereas the effects of this compound at $\alpha 6\beta 3\gamma 2$ (stimulation of GABA EC_3 to 1622 ± 87 % at $10 \mu\text{M}$ concentration) and $\alpha 6\beta 3\delta$ (stimulation of GABA EC_3 to 1639 ± 244 % at $10 \mu\text{M}$ concentration) receptors were comparable (Supplementary Tables 2 and 3).

The strong potentiation of the GABA current by LAU 177 at $\alpha 1\beta 3\delta$ receptors was similar to the effects of other compounds observed at these receptors. Specifically, it has been demonstrated that neurosteroids, tracazolate, and DS2 are able to strongly enhance GABA-induced currents at δ -containing receptors compared to the much weaker effects at receptors not containing the δ -subunit. [31, 41] To further investigate this strong effect of LAU 177, a GABA concentration-effect curve was generated at $\alpha 1\beta 3\delta$ receptors in the absence or presence of LAU 177 (Fig. 4a, b). In agreement with previous results [31], GABA elicited currents at $\alpha 1\beta 3\delta$ receptors are very small. Whereas GABA was able to induce currents of $8 \mu\text{A}$ in $\alpha 1\beta 3$ or of $16 \mu\text{A}$ at $\alpha 1\beta 3\gamma 2$ receptors, the maximal GABA-induced effect at $\alpha 1\beta 3\delta$ receptors was only about $1 \mu\text{A}$. In the presence of LAU 177, however, the GABA-induced current was dramatically potentiated up to $13 \mu\text{A}$ at 1 mM GABA. This effect was similar to that of tracazolate or THDOC described previously [31], indicating that LAU 177 also is able to dramatically enhance the efficacy of GABA for opening the GABA_A receptor-associated chloride channel. Interestingly, however, in contrast to the effects observed with

tracazolate or THDOC [31], the potency of GABA for enhancing chloride currents was not significantly changed by LAU 177 (Fig. 4c).

As shown for $\alpha 1\beta 3\gamma 2$ receptors and CGS 9895 [12] or PZ-II-028 [27], the effects of LAU 177 at $\alpha 1\beta 3\delta$ receptors also strongly depended on the type of beta subunit present in the receptors (Fig. 5). LAU 177 exhibited similar efficacy but reduced potency at $\alpha 1\beta 2\delta$ receptors ($EC_{50} > 10.7 \mu M$) when compared with $\alpha 1\beta 3\delta$ receptors ($EC_{50} = 1.0 \mu M$). The efficacy of this compound at $\alpha 1\beta 1\delta$ receptors, however, was drastically reduced (Fig. 5, Supplementary Table 4) compared to $\alpha 1\beta 2,3\delta$ receptors.

Thus, the effects of LAU 177 on $GABA_A$ receptors depend on all subunits in the receptor complex—not only the subtype of α and β subunits, but also the presence of a γ or a δ subunit effect potency and efficacy to some extent. To investigate further whether the strong modulatory effects of LAU 177 at $\alpha 1\beta 3\delta$ receptors were also mediated via the pyrazoloquinolinone binding site at the $\alpha 1+$ $\beta 3-$ interface, steric hindrance experiments were again performed. Here, it was interesting to investigate whether the effect of LAU 177 could be inhibited via both the $\alpha 1+$ and the $\beta 3-$ side. As shown in Fig. 6a, LAU 177 exhibited a comparable effect at 100 nM and 1 μM concentrations at $\alpha 1\beta 3\delta$ or $\alpha 1\beta 3Q64C\delta$ receptors. This point mutation, however, dramatically enhanced the effect of LAU 177 in $\alpha 1\beta 3Q64C\delta$ receptors at 10 and 30 μM concentration as compared to $\alpha 1\beta 3\delta$ receptors. On incubation with MTSEA-biotin this effect of LAU 177 was drastically reduced at all LAU 177 concentrations investigated. Similarly, the point mutation $\alpha 1S204C$ dramatically enhanced the effect of LAU 177 at 10 and 30 μM concentration, but at 1 μM concentration the effect of LAU 177 was smaller in the mutated receptor as compared to the wild-type receptor (Fig. 6b). On incubation of $\alpha 1S204C\beta 3\delta$ receptors with MTSEA-biotin, the effects of LAU 177 were also dramatically reduced at 1, 10, and 100 μM concentrations. Together, these data indicate that the effects of LAU 177 can be reduced by steric hindrance introduced via the $\alpha 1+$ as well as via the $\beta 3-$ side of the interface, indicating that both sides contributed to the binding of LAU 177. These data indicated that most of the effects of LAU 177 at $\alpha 1\beta 3\delta$ receptors are mediated via an interaction of this compound with the $\alpha 1+$ $\beta 3-$ site of $GABA_A$ receptors.

An alternative way to demonstrate that the effects of LAU 177 were mediated via the $\alpha 1+$ $\beta 3-$ interface of $\alpha 1\beta 3\delta$ receptors is to use a compound acting as a null modulator (antagonist) of this binding site. LAU 157 is one of the five null modulators for $\alpha 1\beta 3$ receptors previously identified [26]. So far however, this compound was not investigated for its effect at other receptors subtypes. We thus investigated whether LAU 157 (Fig. 7a) behaves as a null modulator at $\alpha \beta \delta$ receptors. Results shown in Fig. 7b indicate that LAU 157 did not significantly modulate GABA EC_3 at $\alpha 1\beta 3$, $\alpha 1\beta 3\delta$, $\alpha 1\beta 2\delta$ receptors. To investigate whether this lack of effect was due to LAU 157 behaving as a null modulator or whether LAU 157 did not bind at all at the $\alpha+$ $\beta-$ interface of these receptors, experiments were performed investigating the inhibition of the effects of LAU 177 at $\alpha 1\beta 3\delta$, $\alpha 1\beta 2\delta$, or $\alpha 1\beta 3$ receptors by LAU 157. As shown in Fig. 7c, results indicated that LAU 157 was able to completely inhibit the effects of LAU 177 in all receptors investigated, suggesting that it is a null modulator at the $\alpha+$ $\beta-$ interface and was also able to inhibit the effects of LAU 177 at $\alpha 1\beta 3\delta$ receptors.

To investigate whether this strong modulation of $\alpha 1\beta 3\delta$ receptors by LAU 177 as compared to $\alpha 1\beta 3\gamma 2$ receptors was a general property of pyrazoloquinolinones, the effects of PZ-II-028 [26, 27], were also analyzed at $\gamma 2$ and δ containing receptors in analogous experiments. In contrast to LAU 177, the modulation of GABA-induced currents by PZ-II-028 at $\alpha 1\beta 3\delta$ receptors (stimulation to 1186 ± 89 % at $10 \mu\text{M}$ concentration) was only slightly enhanced compared to that at $\alpha 1\beta 3\gamma 2$ receptors (stimulation to 939 ± 64 at $10 \mu\text{M}$ concentration [27]). The modulation by this compound of GABA-induced currents at $\alpha 4\beta 3\gamma 2$ receptors (226 ± 14 % [27]) was also enhanced at $\alpha 4\beta 3\delta$ receptors (425 ± 26 %). But the strong stimulation of GABA-induced current at $\alpha 6\beta 3\gamma 2$ receptors by PZ-II-028 (stimulation to 1871 ± 28 % at $10 \mu\text{M}$ concentration, [27]) no longer was observed in $\alpha 6\beta 3\delta$ receptors (stimulation to 1152 ± 106 % at $10 \mu\text{M}$ concentration). The effects of this compound class are thus not only dependent on the molecular structure of the compound but also on the receptor composition (α , β subtypes, $\gamma 2$, or δ subunit). Further experiments will have to be performed to investigate the effects of PZ-II-028 and those of other pyrazoloquinolinones in $\alpha\beta$, $\alpha\beta\gamma 2$ and $\alpha\beta\delta$ receptors in more detail.

Tracazolate [31] and the imidazopyridine DS2 [41] also display strong enhancement of GABA currents in δ -containing receptors, and the efficacy is determined by all subunits present in the receptor complex. This analogy prompted us to investigate whether tracazolate or DS2 might exert their action via the extracellular $\alpha + \beta$ - binding site. Our experiments however indicated that increasing concentrations (up to $30 \mu\text{M}$) of the $\alpha + \beta$ - site antagonist LAU 157 did not inhibit the effects of $1 \mu\text{M}$ DS2 (stimulation to 400 % of GABA EC_3) or of $3 \mu\text{M}$ tracazolate (stimulation to 900 % of GABA EC_3) at $\alpha 1\beta 3\delta$ receptors (experiments not shown). This seems to indicate that neither tracazolate nor DS2 exert their actions via the pyrazoloquinolinone binding site at the $\alpha + \beta$ - interface.

Discussion

In this study, we investigated the effects of the pyrazoloquinolinone LAU 177 in more detail. The nM affinity of this compound for the benzodiazepine binding site at $\alpha 1\beta 3\gamma 2$ receptors, its lack of modulation of GABA-induced currents at this receptor in the 10–100 nM range, as well as the comparable concentration-effect curve at $\alpha 1\beta 3$ and $\alpha 1\beta 3\gamma 2$ receptors indicate that this compound, similar to other pyrazoloquinolinones [12, 26, 27], is a high affinity null modulator at the benzodiazepine binding site and a low potency high efficacy modulator via the extracellular $\alpha 1 + \beta 3$ - interface. This conclusion was supported by steric hindrance experiments at $\alpha 1\beta 3$ receptors as well as by the use of LAU 157, an $\alpha 1 + \beta 3$ - site antagonist, that was able to completely inhibit the action of LAU 177 at $\alpha 1\beta 3$ receptors.

Modulation by LAU 177 of GABA_A receptor subtypes composed of $\alpha 1, 2, 3, 5\beta 3$ or $\alpha 1-6\beta 3\gamma 2$ subunits, was slightly different in potency and efficacy, indicating that the modulation of this compound is dependent on the type of the α subunit present in the receptor. This again is consistent with LAU 177 eliciting modulation by binding to the $\alpha + \beta$ - interface. The extent of modulation of GABA currents was comparable in $\alpha 1-3\beta 3$ or $\alpha 1-3\beta 3\gamma 2$ receptors containing the same α subunit type. Only at $\alpha 5\beta 3\gamma 2$ receptors we observed a stronger modulation of LAU 177 than at $\alpha 5\beta 3$ receptors, As with several other

pyrazoloquinolinones [27], LAU 177 exhibited a distinctly stronger effect at receptors composed of $\alpha 6\beta 3\gamma 2$ subunits than at receptors containing other α subunit types together with $\beta 3$ and $\gamma 2$.

Interestingly, however, LAU 177 exhibited a much stronger modulation of GABA-induced currents at $\alpha 1\beta 3\delta$ receptors than at $\alpha 1\beta 3$ or $\alpha 1\beta 3\gamma 2$ receptors. Steric hindrance experiments indicated that the effects of this compound at $\alpha 1\beta 3\delta$ receptors were also mediated via the $\alpha + \beta -$ interface and this conclusion was supported by the use of the $\alpha + \beta -$ site antagonist LAU 157, that was able to completely inhibit the effects of LAU 177 at $\alpha 1\beta 3\delta$ receptors. This conclusion was further supported by the finding that the extra stimulation of LAU 177 at δ -containing receptors also depended on the type of α and β subunits. The α subunit dependence, however, was much stronger than in $\alpha\beta$ or $\alpha\beta\gamma 2$ receptors. Thus, whereas the effects of LAU 177 were strongly enhanced at $\alpha 1\beta 3\delta$ and $\alpha 4\beta 3\delta$ as compared to $\alpha 1\beta 3\gamma 2$ and $\alpha 4\beta 3\gamma 2$ receptors, there was no significant change of efficacy at $\alpha 6\beta 3\delta$ receptors as compared to $\alpha 6\beta 3\gamma 2$ receptors. In addition, LAU 177 strongly modulated GABA-induced currents at $\alpha 1\beta 3\delta$ and $\alpha 1\beta 2\delta$ receptors, with a lower potency at $\alpha 1\beta 2\delta$ receptors. In contrast, LAU 177 exhibited a quite low modulation at $\alpha 1\beta 1\delta$ receptors.

Taken together, all evidence indicates that LAU 177 exerts its modulatory effects in $\alpha 1\beta 3$, $\alpha 1\beta 3\gamma 2$ and $\alpha 1\beta 3\delta$ receptors by binding to the extracellular pocket at the $\alpha 1 + \beta 3 -$ interface. Nevertheless, the extent of modulation was much stronger at $\alpha 1\beta 3\delta$ receptors than at the other two receptors. The differences in the effects of this compound can thus only have been induced by the presence of the delta subunit and its contribution to state transitions or state stabilization.

The much stronger modulation by LAU 177 of $\alpha 1\beta 3\delta$ compared to $\alpha 1\beta 3\gamma 2$ receptors, however, was not observed with PZ-II-028, another pyrazoloquinolinone. Screening experiments indicated that this compound exhibited no or at most a weak, statistically marginal, increase in the modulation of $\alpha 1\beta 3\delta$ over $\alpha 1\beta 3\gamma 2$ receptors. In contrast, PZ-II-028 exhibited a large decrease in the modulation of $\alpha 6\beta 3\delta$ as compared to $\alpha 6\beta 3\gamma 2$ receptors, whereas LAU 177 exhibited no difference in the enhancement of GABA-induced currents at $\alpha 6\beta 3\delta$ and $\alpha 6\beta 3\gamma 2$ receptors. From a different point of view it could also be stated that the $\gamma 2$ subunit in this case could enhance the effects of PZ-II-028 as compared to δ -containing receptor subtypes. As demonstrated previously, the superstimulation of $\alpha 6\beta 3\gamma 2$ by PZ-II-028 was additionally dependent on the $\beta 3$ subunit and was absent in $\beta 2$ containing receptors [27]. Since it is assumed that $\alpha\beta$ receptors are composed of 2 α and 3 β subunits [33, 42, 43], a replacement of a $\beta 3$ by a $\gamma 2$ or a δ subunit in $\alpha 1,6\beta 2,3$, $\alpha 1,6\beta 2,3\gamma 2$, or $\alpha 1,6\beta 2,3\delta$ receptors changes the efficacy of compounds differentially, depending on the structure of the pyrazoloquinolinone as well as on the types of α and β subunits present in the receptors. The actual effects of the pyrazoloquinolinones thus clearly depend on all subunits present in the receptors, although the effects of these compounds are mediated via the $\alpha + \beta -$ interface. The mechanisms of these changes in efficacy currently are not known.

In agreement with previous results [31] we demonstrated that GABA only weakly activated $\alpha 1\beta 3\delta$ receptors, indicating that GABA is a partial agonist at these receptors. The addition

of 10 μ M LAU 177 dramatically (about 12 times) enhanced the conductance of $\alpha 1\beta 3\delta$ receptors induced by maximal GABA concentrations. A similar effect has been demonstrated previously using tracazolate or THDOC [31], or DS2 [41]. In contrast to these compounds that also were able to enhance the potency of GABA consistently, LAU 177 only enhanced the efficacy of GABA but not its potency, suggesting a difference in the actions of LAU 177, tracazolate, THDOC, or DS2. This conclusion was supported by our finding that the $\alpha + \beta -$ site antagonist LAU 157 was unable to block the effects of DS2 or tracazolate (experiments not shown). From these observations we conclude that LAU 177 has a unique profile of action in $\alpha\beta$, $\alpha\beta\gamma$ and $\alpha\beta\delta$ receptors that results from its binding to the pyrazoloquinolinone binding site at the extracellular $\alpha + \beta -$ interface. The efficacy of LAU 177 is determined by ligand specific features, by the binding site forming subunits, and by the nature of the third subunit in the complex. In addition, the site of action and mechanism by which the third subunit and LAU 177 act together is distinct from that of DS2 and tracazolate. Whereas the site of action of DS2 and tracazolate is not known, LAU 177 is acting via the extracellular $\alpha + \beta -$ interface.

Investigations with recombinant concatenated δ -containing receptors have indicated that several types of δ receptors can be formed, in which the δ subunit can assume different positions and replace either the $\gamma 2$ or one of the β subunits of $\alpha 1\beta\gamma 2$ receptors [3, 4]. Each one of these receptors displayed a specific set of properties, and there are also receptors that obviously cannot be activated by GABA in the absence of the steroid THDOC, suggesting that these receptors are essentially silent in the absence of the steroid [3, 4]. If these different receptors also can be formed from non-concatenated subunits, it can be assumed that a mixture of δ -receptors with different subunit arrangements might be formed on injection of *Xenopus* oocytes with GABA_A receptor subunit mRNAs and that the low GABA activation of this receptor mixture could have been caused by a low percentage of receptors that can be directly activated by GABA. Depending on the type of a subunit, the composition of the receptor mixture could have changed possibly explaining the different allosteric modulation of δ -containing receptor subtypes by LAU 177. Further experiments would have to be performed with concatenated receptors to investigate this possibility.

In summary, LAU 177 is a modulator of GABA_A receptors which can uncover extrasynaptic $\alpha 1\beta 2,3\delta$ receptors that due to their low GABA-induced currents are functionally more or less silent in the absence of modulatory agents such as neurosteroids [4, 31]. In contrast to previously described modulators exhibiting similar effects, such as tracazolate or DS2, LAU 177 interacts with a known binding site, the extracellular $\alpha + \beta -$ site, making it more suitable for a rational design of drugs exhibiting such an action profile. The existence of the functional $\alpha + \beta -$ site antagonist LAU 157 not only is useful for in vivo studies as it can terminate the effect of LAU 177 or similar compounds, but also can help to identify compounds from different structural classes interacting with this binding site. Thus, the path forward is to use LAU 177 and related pyrazoloquinolinones as a pharmacophore template to develop more compounds acting as superstimulators on $\alpha 1\beta 2,3\delta$ receptors and to further improve their pharmacological profiles.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support by the graduate school program MolTag (Austrian Science Fund FWF, grant nr. W1232) to L.W. and P. M. is gratefully acknowledged. The mutated construct $\alpha 1S204C$ was a gift from E. Sigel, Department of Pharmacology, University of Bern.

References

1. Olsen RW, Sieghart W. International union of pharmacology. LXX. Subtypes of gamma-aminobutyric acid (A) receptors: classification on the basis of subunit composition, pharmacology, and function. Update. *Pharmacol Rev.* 2008; 60(3):243–260. [PubMed: 18790874]
2. Barrera NP, et al. Atomic force microscopy reveals the stoichiometry and subunit arrangement of the $\alpha 4\beta 3\delta$ GABA(A) receptor. *Mol Pharmacol.* 2008; 73(3):960–967. [PubMed: 18079275]
3. Baur R, Kaur KH, Sigel E. Diversity of structure and function of $\alpha 1\alpha 6\beta 3\delta$ GABA_A receptors: comparison with $\alpha 1\beta 3\delta$ and $\alpha 6\beta 3\delta$ receptors. *J Biol Chem.* 2010; 285(23):17398–17405. [PubMed: 20382738]
4. Kaur KH, Baur R, Sigel E. Unanticipated structural and functional properties of delta-subunit-containing GABA_A receptors. *J Biol Chem.* 2009; 284(12):7889–7896. [PubMed: 19141615]
5. Shu HJ, et al. Characteristics of concatemeric GABA(A) receptors containing $\alpha 4/6$ subunits expressed in *Xenopus* oocytes. *Br J Pharmacol.* 2012; 165(7):2228–2243. [PubMed: 21950777]
6. Bollan KA, et al. The promiscuous role of the epsilon subunit in GABA_A receptor biogenesis. *Mol Cell Neurosci.* 2008; 37(3):610–621. [PubMed: 18206389]
7. Sieghart W. Structure and pharmacology of gamma-aminobutyric acid A receptor subtypes. *Pharmacol Rev.* 1995; 47(2):181–234. [PubMed: 7568326]
8. Ernst M, et al. Comparative models of GABA_A receptor extracellular and transmembrane domains: important insights in pharmacology and function. *Mol Pharmacol.* 2005; 68(5):1291–1300. [PubMed: 16103045]
9. Ernst M, et al. Comparative modelling of GABA(A) receptors: limits, insights, future developments. *Neuroscience.* 2003; 119(4):933–943. [PubMed: 12831854]
10. Smith GB, Olsen RW. Functional domains of GABA_A receptors. *Trends Pharmacol Sci.* 1995; 16(5):162–168. [PubMed: 7624971]
11. Sigel E. Mapping of the benzodiazepine recognition site on GABA(A) receptors. *Curr Top Med Chem.* 2002; 2(8):833–839. [PubMed: 12171574]
12. Ramerstorfer J, et al. The GABA_A receptor $\alpha + \beta$ interface: a novel target for subtype selective drugs. *J Neurosci.* 2011; 31(3):870–877. [PubMed: 21248110]
13. Spurny R, et al. Pentameric ligand-gated ion channel ELIC is activated by GABA and modulated by benzodiazepines. *Proc Natl Acad Sci U S A.* 2012; 109(44):E3028–E3034. [PubMed: 23035248]
14. Belelli D, et al. The interaction of the general anesthetic etomidate with the gamma-aminobutyric acid type A receptor is influenced by a single amino acid. *Proc Natl Acad Sci U S A.* 1997; 94(20):11031–11036. [PubMed: 9380754]
15. Jenkins A, et al. Evidence for a common binding cavity for three general anesthetics within the GABA_A receptor. *J Neurosci.* 2001; 21(6):RC136. [PubMed: 11245705]
16. Mihic SJ, et al. Sites of alcohol and volatile anaesthetic action on GABA(A) and glycine receptors. *Nature.* 1997; 389(6649):385–389. [PubMed: 9311780]
17. Yamakura T, et al. Anesthetics and ion channels: molecular models and sites of action. *Annu Rev Pharmacol Toxicol.* 2001; 41:23–51. [PubMed: 11264449]

18. Li GD, et al. Identification of a GABA_A receptor anesthetic binding site at subunit interfaces by photolabeling with an etomidate analog. *J Neurosci.* 2006; 26(45):11599–11605. [PubMed: 17093081]
19. Chiara DC, et al. Mapping general anesthetic binding site(s) in human alpha1 beta3 gamma-aminobutyric acid type A receptors with [(3)H]TDBzl-etomidate, a photoreactive etomidate analogue. *Biochemistry.* 2012; 51(4):836–847. [PubMed: 22243422]
20. Hosie AM, et al. Endogenous neurosteroids regulate GABA_A receptors through two discrete transmembrane sites. *Nature.* 2006; 444(7118):486–489. [PubMed: 17108970]
21. Hosie AM, et al. Conserved site for neurosteroid modulation of GABA(A) receptors. *Neuropharmacology.* 2009; 56(1):149–154. [PubMed: 18762201]
22. Chiara DC, et al. Specificity of intersubunit general anesthetic-binding sites in the transmembrane domain of the human $\alpha 1\beta 3\gamma 2$ gamma-aminobutyric acid type A (GABA_A) receptor. *J Biol Chem.* 2013; 288(27):19343–19357. [PubMed: 23677991]
23. Nury H, et al. X-ray structures of general anaesthetics bound to a pentameric ligand-gated ion channel. *Nature.* 2011; 469(7330):428–431. [PubMed: 21248852]
24. Spurny R, et al. Multisite binding of a general anesthetic to the prokaryotic pentameric *Erwinia chrysanthemi* ligand-gated ion channel (ELIC). *J Biol Chem.* 2013; 288(12):8355–8364. [PubMed: 23364792]
25. Baur R, et al. Molecular analysis of the site for 2-arachidonylglycerol (2-AG) on the $\beta 2$ subunit of GABA_A receptors. *J Neurochem.* 2013; 126(1):29–36. [PubMed: 23600744]
26. Varagic Z, et al. Identification of novel positive allosteric modulators and null modulators at the GABA_A receptor alpha+ beta–interface. *Br J Pharmacol.* 2013; 169(2):371–383. [PubMed: 23472852]
27. Varagic Z, et al. Subtype selectivity of alpha+ beta– site ligands of GABA_A receptors: identification of the first highly specific positive modulators at $\alpha 6\beta 2/3\gamma 2$ receptors. *Br J Pharmacol.* 2013; 169(2):384–399. [PubMed: 23472935]
28. Brickley SG, Mody I. Extrasynaptic GABA(A) receptors: their function in the CNS and implications for disease. *Neuron.* 2012; 73(1):23–34. [PubMed: 22243744]
29. Ramerstorfer J, et al. The point mutation gamma 2F77I changes the potency and efficacy of benzodiazepine site ligands in different GABA_A receptor subtypes. *Eur J Pharmacol.* 2010; 636(1–3):18–27. [PubMed: 20303942]
30. Sigel E, et al. The effect of subunit composition of rat brain GABA_A receptors on channel function. *Neuron.* 1990; 5(5):703–711. [PubMed: 1699569]
31. Zheleznova N, Sedelnikova A, Weiss DS. $\alpha 1\beta 2\delta$, a silent GABA_A receptor: recruitment by tracazolate and neurosteroids. *Br J Pharmacol.* 2008; 153(5):1062–1071. [PubMed: 18204487]
32. Ebert V, Scholze P, Sieghart W. Extensive heterogeneity of recombinant gamma-aminobutyric acid A receptors expressed in $\alpha 4\beta 3\gamma 2$ -transfected human embryonic kidney 293 cells. *Neuropharmacology.* 1996; 35(9-10):1323–1330. [PubMed: 9014148]
33. Tretter V, et al. Stoichiometry and assembly of a recombinant GABA_A receptor subtype. *J Neurosci.* 1997; 17(8):2728–2737. [PubMed: 9092594]
34. Horton RM, et al. Gene splicing by overlap extension. *Methods Enzymol.* 1993; 217:270–279. [PubMed: 8474334]
35. Fryer RI, et al. Structure-activity relationship studies at the benzodiazepine receptor (BZR): a comparison of the substituent effects of pyrazoloquinolinone analogs. *J Med Chem.* 1993; 36(11):1669–1673. [PubMed: 8388472]
36. Hoerlein G, et al. Heterocyclen durch Anellierung an 4-pyridinole, II thieno[3,2-c]pyridin-3-ole. *Liebigs Annalen Chem.* 1979; 3:387–391.
37. Farrant M, Nusser Z. Variations on an inhibitory theme: phasic and tonic activation of GABA(A) receptors. *Nat Rev Neurosci.* 2005; 6(3):215–229. [PubMed: 15738957]
38. Glykys J, et al. A new naturally occurring GABA(A) receptor subunit partnership with high sensitivity to ethanol. *Nat Neurosci.* 2007; 10(1):40–48. [PubMed: 17159992]
39. Sun C, Sieghart W, Kapur J. Distribution of $\alpha 1$, $\alpha 4$, $\gamma 2$, and delta subunits of GABA_A receptors in hippocampal granule cells. *Brain Res.* 2004; 1029(2):207–216. [PubMed: 15542076]

40. Nusser Z, Sieghart W, Somogyi P. Segregation of different GABA_A receptors to synaptic and extrasynaptic membranes of cerebellar granule cells. *J Neurosci.* 1998; 18(5):1693–1703. [PubMed: 9464994]
41. Jensen ML, et al. A study of subunit selectivity, mechanism and site of action of the delta selective compound 2 (DS2) at human recombinant and rodent native GABA(A) receptors. *Br J Pharmacol.* 2013; 168(5):1118–1132. [PubMed: 23061935]
42. Baumann SW, Baur R, Sigel E. Subunit arrangement of gamma-aminobutyric acid type A receptors. *J Biol Chem.* 2001; 276(39):36275–36280. [PubMed: 11466317]
43. Farrar SJ, et al. Stoichiometry of a ligand-gated ion channel determined by fluorescence energy transfer. *J Biol Chem.* 1999; 274(15):10100–10104. [PubMed: 10187791]

Fig. 1. Top view onto the extracellular domain of GABA_A receptors composed of αβγ or αβ subunits.

Each subunit has a plus (+) and a minus (-) side assigned. Binding sites for GABA are located at the interfaces formed by a “-” side of an alpha subunit and a “+” side of a beta subunit. **a** αβγ receptors composed of 2α, 2β and one γ subunit. The binding site for benzodiazepine site ligands (Bz) is located at the interface formed by the “+” side of an α and the “-” side of the γ subunit. The CGS 9895 binding site is located at the interface formed by the “+” side of an α and the “-” side of a β subunit. **b** αβ receptors formed of 2 α and 3 β subunits exhibit two CGS 9895 binding sites

Fig. 2. Steric hindrance at the α + β - interface drastically reduces the effects of LAU 177 at α 1 β 3 receptors.

a Structure of LAU 177. **b** Representative current traces of GABA EC₃ in the absence or presence of 10 μ M LAU 177 at α 1 β 3 or α 1S204C β 3Q64C receptors without or with steric hindrance (MB treated). **c** Concentration—response effects of LAU 177 at α 1 β 3 ($n = 3$) or α 1S204C β 3Q64C receptors ($n = 4$) in the absence or presence of MTSEA-biotin. MTSEA-biotin significantly reduced the effect of 10 μ M LAU 177 at α 1S204C β 3Q64C receptors ($p < 0.005$; unpaired Student's t test; $n = 4$). Data are mean values $\pm$ SEM

Fig. 3. Effects of LAU 177 at different GABA_A receptor subtypes.

a Concentration-response curves of LAU 177 at α1β3, α2β3, α3β3 and α5β3 receptors (n = 3). **b** Concentration-response curves of LAU 177 at α1-6β3γ2 receptors (n = 3–7). **c** Concentration-response curves of LAU 177 at α1β3δ, α4β3δ and α6β3δ receptors (n = 4–7). Data are mean values ± SEM

Fig. 4. LAU 177 strongly enhances GABA evoked currents at $\alpha 1\beta 36$ receptors without changing GABA-potency.

a Representative traces of a low GABA concentration (1 μ M) and a high GABA concentration (1 mM) in absence or presence of 10 μ M LAU 177. **b** Concentration dependent currents of GABA (*filled square*, $n = 4$) and GABA plus 10 μ M LAU 177 (*open square*, $n = 4$). **c** Effects of GABA and GABA plus 10 μ M LAU 177 are normalized to maximum evoked current. A concentration of 10 μ M LAU 177 showed no significant change of GABA EC₅₀ ($p > 0.05$; paired student's t test; $n = 4$). Data are mean values $\pm$ SEM

Fig. 5. β -subunit dependent effects of LAU 177 at $\alpha 1\beta 8$ receptors.

The effects of LAU 177 at $\alpha 1\beta 38$ ($n = 13$) and $\alpha 1\beta 28$ ($n = 9$) receptors are comparable (stimulation to 3285 ± 257 % at $\alpha 1\beta 38$ and to 2843 ± 396 % at $\alpha 1\beta 28$ at $10 \mu\text{M}$ concentrations) although LAU 177 exhibits a reduced potency at $\alpha 1\beta 28$ receptors. However, the effects of LAU 177 at $\alpha 1\beta 18$ receptors ($n = 9$) are drastically reduced. Data are mean values $\pm$ SEM

Fig. 6. Steric hindrance at $\alpha 1\beta 38$ receptors via the β - or α + side.

a Introduction of point mutation Q64C at the “-” side of the β subunit strongly enhances the effects of LAU 177 (to 6941 ± 1268 % at $10 \mu\text{M}$; $n = 3$) compared to $\alpha 1\beta 38$ wild-type receptors (3372 ± 455 % at $10 \mu\text{M}$; $n = 5$). Incubation with MTSEA biotin dramatically reduces this effect to 1754 ± 568 % (at $10 \mu\text{M}$; $n = 3$) ($p < 0.05$; unpaired Student’s t test). **b** Introduction of the point mutation S204C at the “+” side of the α subunit strongly enhances the effects of LAU 177 (to 5278 ± 1008 % at $10 \mu\text{M}$; $n = 3$) as compared to $\alpha 1\beta 38$ receptors (3327 ± 455 % at $10 \mu\text{M}$; $n = 5$). Incubation with MTSEA biotin dramatically reduces this effect to 1884 ± 311 % (at $10 \mu\text{M}$; $n = 4$) ($p < 0.05$; unpaired Student’s t test). Data are mean values $\pm$ SEM

Fig. 7. LAU 157 inhibits effects of LAU 177.

a Structure of LAU 157. **b** Concentration-dependent effect of LAU 157 at $\alpha 1\beta 3\delta$ ($n = 3$), $\alpha 1\beta 2\delta$ ($n = 3$) and $\alpha 1\beta 3$ ($n = 3$). **c** Increasing concentrations of LAU 157 dose dependently inhibit the effect of 1 μ M LAU 177 at $\alpha 1\beta 3\delta$ ($n = 4$), $\alpha 1\beta 2\delta$ ($n = 3$) and $\alpha 1\beta 3$ ($n = 4$), reaching complete inhibition. Data are mean values $\pm$ SEM