Published in final edited form as:

J Comb Chem. 2010 September 13; 12(5): 664–669. doi:10.1021/cc100068a.

A Novel and Efficient One-Step Parallel Synthesis of Dibenzopyranones *via* Suzuki-Miyaura Cross Coupling

Kodumuru Vishnumurthy* and Alexandros Makriyannis

Center for Drug Discovery, 360 Huntington Avenue, 116 Mugar Hall, Boston, Massachusetts 02115-5000, U.S.A.

Abstract

Microwave promoted novel and efficient one-step parallel synthesis of dibenzopyranones and heterocyclic analogues from bromo arylcarboxylates and *o*-hydroxyarylboronic acids *via* Suzuki-Miyaura cross coupling reaction is described. Spontaneous lactonization gave dibenzopyranones and heterocyclic analogues bearing electron donating and withdrawing groups on both aromatic rings in good to excellent yields.

Keywords

Suzuki-Miyaura cross coupling; microwave based synthesis; benzopyranones

Introduction

Dibenzopyranone is a privileged scaffold in many natural products¹ such as alternariol, graphis lactones, autumnariol, autumnariniol, altenuisol and biologically active compounds.² Such lactones have been used as intermediates in the synthesis of several pharmaceutically interesting compounds including progesterone, androgen, glucocorticoid modulators,³ and endothelial cell proliferation inhibitors.⁴ Furthermore, dibenzopyranones occur naturally in many food sources including citrus fruits, herbs, and vegetables.⁵ There are several methods available for the synthesis of dibenzopyranones. The most popular involves Suzuki-Miyaura cross coupling followed by Lewis acid⁶ or metal⁷ mediated lactonization. More recently the Diels-Alder cycloaddition of 4-cyanocoumarins,⁸ *tert*-butyllithium-mediated cyclization of bromobenzylfluorophenyl ethers,⁹ and ruthenium-catalysed cyclotrimerization of aryl diynes¹⁰ have been reported. In 2002, Abbott Laboratories reported a practicable and scalable synthesis of glucocorticoid receptor A-224817.0 through a Negishi cross coupling.¹¹ However, none of these methods are suitable for the parallel synthesis approach to generating large numbers of analogues for drug discovery research. The disadvantages include low overall yields, long reaction times, low temperature conditions, multi-step sequences, and the need to purify intermediates. Although the previous method of coupling of *o*-methoxyphenylboronic acids with bromo arylcarboxylates **1** followed by borontribromide⁶ or hydroiodic acid mediated lactonization¹² has potential for parallel synthesis, but suffers from the following drawbacks: a) it is two step procedure requiring minimum the purification of two intermediates, b) alkoxy and alkoxyester substituents undergo demethylation in the presence of borontribromide,¹³ and c) BBr₃-mediated

* To whom correspondence should be addressed. Tel.: 617-373-7620; Fax: 617-373-7493; kvishnumurthy@hotmail.com; v.kodumuru@neu.edu.

Supporting Information Available. General experimental details, yields, and compound characterization data (¹H NMR, IR and MS analysis) of all final products. This information is available free of charge via the internet at <http://pubs.acs.org>.

lactonization⁶ requires the reaction to be performed at -78 °C, while HI-mediated lactonization requires to be conducted at 110 °C.¹²

Recently, we reported¹³ that dibenzopyranone analogues act as CB2 agonists and that they are being investigated for the treatment of neuropathic pain, inflammation, and Alzheimer's diseases. As part of an ongoing lead optimization project, we required a facile route to prepare a library of dibenzopyranones and heterocyclic analogues **3**. We have, therefore, investigated a straightforward route for dibenzopyranones **3** via tandem Suzuki coupling of bromo arylcarboxylates **1** with *o*-hydroxyarylboronic acids **2** followed by lactonization (Scheme 1).

Also, we utilized microwave assisted organic synthesis, which has attracted increasing attention because of its potential to accelerate drug discovery research.¹⁵ Reaction times of many organic transformations are often dramatically reduced from hours to minutes or seconds. Moreover, there have been several reports of microwave-assisted syntheses of combinatorial libraries.¹⁶ We now report a novel and efficient one step parallel synthesis of a library of dibenzopyranones and heterocyclic analogues **3** via Suzuki-Miyaura cross coupling of bromo arylcarboxylates **1** with *o*-hydroxyarylboronic acids **2**.

Results and Discussion

To develop an efficient methodology for the synthesis of dibenzopyranones and heterocyclic analogues **3**, using bromo arylcarboxylates **1** and *o*-hydroxyarylboronic acids **2**, various catalysts, ligands, bases and solvents were first screened for the Suzuki-Miyaura cross coupling reaction (Table 1). At the outset of the studies, commercially available methyl 2-bromo-5-methoxybenzoate **1{1}**, methyl 2-bromo-4-fluorobenzoate **1{2}**, and *o*-hydroxyphenylboronic acid **2{1}** were selected as representative reactants (Table 1). We initially examined the reaction conditions reported for chloro(di-2-norbornylphosphino)(2'-dimethylamino-1,1'-biphenyl-2-yl)palladium (II), an air-stable catalyst that is highly active for C-C and C-N coupling reactions with aryl chlorides.¹⁷ When a solution of **1{1}** or **1{2}** (1 equiv), **2{1}** (1.2 equiv), potassium fluoride (5 equiv), and 5 mol% of the above catalyst in dioxane was heated in a microwave reactor at 130 °C for 15 min, the desired product i.e., 8-methoxy-6H-benzo[c]chromen-6-one **3{1}** or 9-fluoro-6H-benzo[c]chromen-6-one **3{2}**, was obtained in 54 and 48% isolated yields, respectively (entry 1, Table 1). Coupling under Buchwald's conditions using highly active palladium catalyst, i.e.,¹⁸ potassium fluoride, Pd(OAc)₂ and (*o*-biphenyl)P(*t*-Bu)₂ in THF at room temperature, afforded **3{1}** and **3{2}** in 45% and 47% yields, respectively (entry 2, Table 1). To improve the yields for the desired products **3**, we screened several catalysts such as Pd(PCy₃)₂Cl₂, PdCl₂(PPh₃)₂, Pd(PPh₃)₄, 1,1'-bis(diphenylphosphino) ferrocene dichloropalladium (II); bases such as NaOAc, Cs₂CO₃, CsF, KF; ligand (*o*-biphenyl)P(*t*-Bu)₂ with different equivalents and combinations of reagents and microwave irradiation temperatures in THF, NMP, DME or dioxane. The results of formation of products **3{1}** and **3{2}** are shown in Table 1. From the screening results, the best conditions for coupling of **1{1}** or **1{2}** with **2{1}** were found to be the use of Pd(PPh₃)₄ as a catalyst in the presence of Cs₂CO₃ in DME at 125 °C under microwave irradiation for 15 min (Table 1, entry 6).

Utilising the above optimized conditions, Suzuki-Miyaura cross coupling of various bromo arylcarboxylates **1** (Figure 1) with *o*-hydroxyarylboronic acids **2** (Figure 2) was conducted and the results are summarized in Table 2. The coupling of *o*-hydroxyarylboronic acids **2** with bromo arylcarboxylates **1** bearing a range of electron donating and electron withdrawing groups gave corresponding benzopyranones **3** in good to excellent yields (entries 1-19, Table 2). In addition, bromothiophene carboxylates (entry 20, 21), bromopyridine carboxylates (entries 22 and 23), and fused bromoaryl carboxylates (entries

24 and 25) also reacted well to give the corresponding products (entries 20-25, Table 2). Clearly, the tandem coupling followed by lactonization is a facile protocol to generate a library of compounds **3**.

Conclusion

A simple, rapid and versatile one-step synthetic route to dibenzopyranones **3** and their heterocyclic analogues has been developed. The tandem C-C bond formation *via* Suzuki-Miyaura cross coupling followed by lactonization affords the desired dibenzopyranones **3** in a manner suitable for parallel synthesis. The protocol is complementary to the existing syntheses for dibenzopyranones in terms of number of synthetic steps, ease of accessibility of starting materials, and suitability for combinatorial format.

Experimental Section

General

Unless otherwise noted, reagents and solvents were used as received from commercial suppliers. All palladium catalysts were purchased from Strem chemicals, Inc., and boronic acids were purchased from Combi-Blocks Inc., unless otherwise mentioned. ¹H NMR spectra were recorded on a Varian Inova 500 MHz and 400 MHz spectrometers; the chemical shifts are reported in parts per million (δ), and the coupling constants are reported in hertz. IR spectra were obtained on a Bruker ALPHA ATR spectrometer. The high resolution mass spectral (HRMS) data (EI) were obtained on a Micromass 70-VSE instrument. The final compounds were purified with WATERS preparative LC/MS autopurification system, unless otherwise mentioned. The microwave reactions were conducted in a Emrys optimizer from Personal Chemistry, and generally 10 reactions were performed sequentially for parallel library synthesis.

Methyl 2-bromo-5-(trifluoromethyl)benzoate (**1{7}**)

To a solution of 2-bromo-5-(trifluoromethyl) benzoic acid (5.00 g, 18.58 mmol) in methanol (20 mL) was added thionyl chloride (1.0 mL, 13.7 mmol) and the resulting mixture was refluxed for 8h. After cooling to room temperature, the solvent was concentrated to dryness under reduced pressure. The residue was dissolved in dichloromethane (40 mL), washed with saturated aqueous sodium bicarbonate solution (50 mL) followed by water (100 mL). The organic layer was dried over anhydrous Na₂SO₄, and the filtrate was concentrated to dryness under reduced pressure to give **1{7}** (4.98 g, 95% yield). ¹H-NMR (CDCl₃, 400 MHz) δ 8.07 (s, 1H), 7.81 (d, *J* = 8.0 Hz, 1H), 7.59-7.56 (m, 1H), 3.97 (s, 3H).

General procedure for the preparation of 2-hydroxyarylboronic acids (**1{27}** -**1{32}**)

To a solution of corresponding methoxy boronicacids (1 mmol) in anhydrous dichloromethane (15 mL), was added boron tribromide (1.2 mmol) at -78°C under nitrogen atmosphere. The resulting solution was stirred for 15 minutes at -78°C. The cooling bath was removed and the reaction mixture was continued to stir for 30 minutes at room temperature. The reaction was quenched with anhydrous methanol (1 mL) at -78°C, concentrated *in vacuo* to dryness. The desired corresponding *o*-hydroxy boronic acids were confirmed by LC/MS and used as it is without further purification.

Illustrative example: Preparation of 8-methoxy-6H-benzo[c]chromen-6-one (**3{1}**)

19 A solution of methyl 3-methoxybenzoate (**1{1}**, 0.5 mmol), 2-hydroxyarylboronic acid (**2{1}**, 0.65 mmol, 1.3 equiv), cesium carbonate (2 mmol, 4 equiv) in a mixture of dimethoxyethane (5 mL) and water (0.75 mL) was degassed with argon for 5 minutes. Tetrakis(triphenylphosphine)palladium (0) (10 mol%) was added, the reaction tube was

sealed. The reaction was heated in a microwave (Emrys optimizer from Personal Chemistry) at 125 °C (power 300 W) for 15 minutes. After cooling at room temperature, the reaction mixture was diluted with water (30 mL), extracted with dichloromethane (2 × 15 mL). The combined organic layers were passed through a phase separator and concentrated *in vacuo*. The residue was purified with WATERS preparative LC/MS autopurification system to afford the title compound in 98% yield. ¹H-NMR (CDCl₃, 500 MHz) δ 8.05 (d, *J* = 8.5 Hz, 1H), 7.99 (d, *J* = 8.0 Hz, 1H), 7.82 (d, *J* = 3.0 Hz, 1H), 7.45-7.31 (m, 4H), 3.95 (s, 3H); IR (cm⁻¹) 1707.9, 1610.1, 1479.8, 1276.1, 1024.4, 824.1, 742.1; LC/MS 226.98 (M+1); HRMS (EI): *m/z* calcd for C₁₄H₁₀O₃ 226.0630, found 226.0635.

9-Fluoro-6H-benzo[c]chromen-6-one (3{2})

Yield 96%, white solid, ¹H-NMR (CDCl₃, 500 MHz) δ 8.44-8.41 (m, 1H), 7.95 (d, *J* = 8.0 Hz, 1H), 7.74-7.72 (m, 1H), 7.54-7.50 (m, 1H), 7.38-7.34 (m, 2H), 7.29-7.25 (m, 1H); IR (cm⁻¹) 1731.6, 1611.2, 1420.8, 1300.8, 1187.5, 877.3, 742.1; LC/MS 215.0 (M+1); HRMS (EI): *m/z* calcd for C₁₃H₇FO₂ 214.0430, found 214.0422.

8-Fluoro-6H-benzo[c]chromen-6-one (3{3})

Yield 88%, white solid, ¹H-NMR (CDCl₃, 500 MHz) δ 8.14-8.12 (m, 1H), 8.06-8.03 (m, 1H), 8.02-7.99 (m, 1H), 7.57-7.53 (m, 1H), 7.50-7.47 (m, 1H), 7.38-7.34 (m, 2H); LC/MS 214.9 (M+1); HRMS (EI): *m/z* calcd for C₁₃H₇FO₂ 214.0430, found 214.0437.

8-Methyl-6H-benzo[c]chromen-6-one (3{4})

^{6a} Yield 88%, white solid, ¹H-NMR (CDCl₃, 500 MHz) δ 8.18 (s, 1H), 8.02-7.98 (m, 2H), 7.63-7.61 (m, 1H), 7.46-7.43 (m, 1H), 7.35-7.30 (m, 2H), 2.49 (s, 3H); LC/MS 211.0 (M+1); HRMS (EI): *m/z* calcd for C₁₄H₁₀O₂ 210.0681, found 210.0682.

8-(Trifluoromethoxy)-6H-benzo[c]chromen-6-one (3{5})

Yield 80%, white solid, ¹H-NMR (CDCl₃, 500 MHz) δ 8.25 (s, 1H), 8.19 (d, *J* = 8.5 Hz, 1H), 8.04 (d, *J* = 8.0, 1H), 7.69-7.66 (m, 1H), 7.54-7.51 (m, 1H), 7.41-7.36 (m, 2H); LC/MS 280.9 (M+1); HRMS (EI): *m/z* calcd for C₁₄H₇F₃O₂ 280.0347, found 280.0388.

8-Nitro-6H-benzo[c]chromen-6-one (3{6})

^{6a} Yield 88%, white solid, ¹H-NMR (CDCl₃, 500 MHz) δ 9.25 (d, *J* = 2.5 Hz, 1H), 8.09 (d, *J* = 9.0, 2.5 Hz, 1H), 8.31 (d, *J* = 8.5 Hz, 1H), 8.14-8.12 (m, 1H), 7.65-7.62 (m, 1H), 7.46-7.42 (m, 2H); LC/MS 241.9 (M+1); HRMS (EI): *m/z* calcd for C₁₃H₇NO₄ 241.0375, found 241.0358.

8-(Trifluoromethyl)-6H-benzo[c]chromen-6-one (3{7})

Yield 83%, white solid, ¹H-NMR (CDCl₃, 500 MHz) δ 8.70 (s, 1H), 8.27 (d, *J* = 8.0 Hz, 1H), 8.11 (dd, *J* = 8.0, 1.5 Hz, 1H), 8.05 (dd, *J* = 8.5, 2.0 Hz, 1H), 7.60-7.56 (m, 1H), 7.43-7.39 (m, 2H); LC/MS 264.9 (M+1); HRMS (EI): *m/z* calcd for C₁₄H₇F₃O₂ 264.0398, found 264.0407.

Methyl 6-oxo-6H-benzo[c]chromene-8-carboxylate (3{8})

Yield 80%, white solid, ¹H-NMR (CDCl₃, 500 MHz) δ 9.08 (s, 1H), 8.47 (d, *J* = 8.5 Hz, 1H), 8.23 (d, *J* = 8.5 Hz, 1H), 8.13-8.11 (m, 1H), 7.57-7.55 (m, 1H), 7.42-7.39 (m, 2H), 3.99 (s, 3H); LC/MS 254.9 (M+1); HRMS (EI): *m/z* calcd for C₁₅H₁₀O₄ 254.0579, found 254.0570.

6H-Benzo[c]chromen-6-one (3{9})

Yield 85%, white solid, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 8.42 (d, J = 8.0 Hz, 1H), 8.14 (d, J = 7.5 Hz, 1H), 8.08 (d, J = 8.0 Hz, 1H), 7.86-7.83 (m, 1H), 7.61-7.58 (m, 1H), 7.51-7.48 (m, 1H), 7.39-7.33 (m, 2H); IR (cm^{-1}) 1725.5, 1604.9, 1431.6, 1302.1, 1262.2, 1076.2, 743.9; LC/MS 196.9 (M+1); HRMS (EI): m/z calcd for $\text{C}_{13}\text{H}_8\text{O}_2$ 196.0529, found 196.0524.

Methyl 6-oxo-6H-benzo[c]chromene-9-carboxylate (3{10})

Yield 90%, white solid, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 8.79 (s, 1H), 8.47 (d, J = 8.5 Hz, 1H), 8.20-8.15 (m, 2H), 7.53-7.51 (m, 1H), 7.40-7.37 (m, 2H), 4.03 (s, 3H); LC/MS 254.9 (M+1); HRMS (EI): m/z calcd for $\text{C}_{15}\text{H}_{10}\text{O}_4$ 254.0579, found 254.0571.

9-Chloro-6H-benzo[c]chromen-6-one (3{11})

Yield 92%, white solid, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 8.35 (d, J = 8.5 Hz, 1H), 8.64 (d, J = 2.0 Hz, 1H), 8.02-8.00 (m, 1H), 7.56-7.51 (m, 2H), 7.40-7.35 (m, 2H); LC/MS 230.9, 232.9 (M+1); HRMS (EI): m/z calcd for $\text{C}_{13}\text{H}_7\text{ClO}_2$ 230.0140, found 230.0134.

2-Chloro-8-methoxy-6H-benzo[c]chromen-6-one (3{12})

Yield 89%, white solid, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 7.97 (d, J = 8.5 Hz, 1H), 7.93 (d, J = 2.5 Hz, 1H), 7.80 (d, J = 2.0 Hz, 1H), 7.41 (dd, J = 8.5, 2.5 Hz, 1H), 7.37 (dd, J = 8.5, 2.5 Hz, 1H), 7.30 (d, J = 8.5 Hz, 1H); LC/MS 260.9, 263.0 (M+1); HRMS (EI): m/z calcd for $\text{C}_{14}\text{H}_9\text{ClO}_3$ 260.0240, found 260.0225.

9-Methyl-6H-benzo[c]chromen-6-one (3{13})

^{6a} Yield 92%, white solid, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 8.27 (d, J = 8.0 Hz, 1H), 8.05 (d, J = 1.0 Hz, 1H), 7.89 (s, 1H), 7.48-7.45 (m, 1H), 7.39-7.31 (m, 2H), 2.56 (s, 3H); LC/MS 211.0 (M+1); HRMS (EI): m/z calcd for $\text{C}_{14}\text{H}_{10}\text{O}_2$ 210.0680, found 210.0674.

8, 9-Dimethoxy-6H-benzo[c]chromen-6-one (3{14})

Yield 84%, white solid, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 7.96 (d, J = 8.00 Hz, 1H), 7.75 (s, 1H), 7.61-7.28 (m, 2H), 7.37-7.31 (m, 2H); LC/MS 257.0 (M+1); HRMS (EI): m/z calcd for $\text{C}_{15}\text{H}_{12}\text{O}_4$ 256.0735, found 256.0729.

7-Methyl-6H-benzo[c]chromen-6-one (3{15})

Yield 92%, white solid, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 8.04-7.98 (m, 2H), 7.67-7.64 (m, 1H), 7.46-7.43 (m, 1H), 7.38-7.37 (m, 1H), 7.33-7.28 (m, 2H), 2.87 (s, 3H); IR (cm^{-1}) 1716.2, 1461.1, 1242.9, 1206.4, 1049.0, 785.1, 751.5; LC/MS 211.0 (M+1); HRMS (EI): m/z calcd for $\text{C}_{14}\text{H}_{10}\text{O}_2$ 210.0680, found 210.0675.

10-Methyl-6H-benzo[c]chromen-6-one (3{16})

Yield 86%, white solid, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 8.39-8.30 (m, 2H), 7.66-7.64 (m, 1H), 7.49-7.46 (m, 2H), 7.42-7.39 (m, 1H), 7.35-7.32 (m, 1H), 2.91 (s, 3H); LC/MS 211.0 (M+1); HRMS (EI): m/z calcd for $\text{C}_{14}\text{H}_{10}\text{O}_2$ 210.0680, found 210.0682.

7-Chloro-6H-benzo[c]chromen-6-one (3{17})

The residue was purified with Biotage SP1 flash autopurification system to afford the white solid in 68% yield, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 8.52 (d, J = 3.0 Hz, 1H), 7.82 (dd, J = 8.0, 1.5 Hz, 1H), 7.79-7.78 (m, 1H), 7.43-7.40 (m, 1H), 7.35-7.33 (m, 1H), 7.30-7.26 (m, 1H); LC/MS 230.9, 232.9 (M+1); HRMS (EI): m/z calcd for $\text{C}_{13}\text{H}_7\text{ClO}_2$ 230.0134, found 230.0126.

8,9-Difluoro-6H-benzo[c]chromen-6-one (3{18})

Yield 86%, white solid, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 8.21-8.18 (m, 1H), 7.92-7.86 (m, 2H), 7.55-7.51 (m, 1H), 7.40-7.36 (m, 2H); LC/MS 232.9 (M+1); HRMS (EI): m/z calcd for $\text{C}_{13}\text{H}_6\text{F}_2\text{O}_2$ 232.0335, found 232.0342.

2,9-Difluoro-6H-benzo[c]chromen-6-one (3{19})

Yield 94%, white solid, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 8.47-8.44 (m, 1H), 7.67-7.62 (m, 2H), 7.39-7.31 (m, 2H), 7.27-7.23 (m, 1H); LC/MS 232.9 (M+1); HRMS (EI): m/z calcd for $\text{C}_{13}\text{H}_6\text{F}_2\text{O}_2$ 232.0335, found 232.0343.

4H-Thieno[3,4-c]chromen-4-one (3{20})

Yield 85%, white solid, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 8.52 (d, $J = 3.0$ Hz, 1H), 7.82 (dd, $J = 8.0, 1.5$ Hz, 1H), 7.79-7.78 (m, 1H), 7.43-7.40 (m, 1H), 7.35-7.33 (m, 1H), 7.30-7.26 (m, 1H); LC/MS 202.9 (M+1); HRMS (EI): m/z calcd for $\text{C}_{11}\text{H}_6\text{O}_2\text{S}$ 202.0088, found 202.0083.

4H-Thieno[2,3-c]chromen-4-one (3{21})

Yield 84%, white solid, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 7.94 (d, $J = 5.0$ Hz, 1H), 7.86 (d, $J = 8.0$ Hz, 1H), 7.67 (d, $J = 5.0$ Hz, 1H), 7.53-7.45 (m, 2H), 7.38-7.34 (m, 1H); LC/MS 202.9 (M+1); HRMS (EI): m/z calcd for $\text{C}_{11}\text{H}_6\text{O}_2\text{S}$ 202.0088, found 202.0093.

5H-Chromeno[4,3-b]pyridin-5-one (3{22})

Yield 95%, white solid, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 9.03-9.02 (m, 1H), 8.63-8.57 (m, 2H), 7.60-7.57 (m, 1H), 7.54-7.51 (m, 1H), 7.42-7.38 (m, 2H); LC/MS 197.9 (M+1); HRMS (EI): m/z calcd for $\text{C}_{12}\text{H}_7\text{NO}_2$ 197.0476, found 197.0482.

5H-Chromeno[4,3-c]pyridin-5-one (3{23})

Yield 83%, white solid, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 9.58 (s, 1H), 8.88-8.87 (m, 1H), 8.21-8.16 (m, 2H), 7.58-7.55 (m, 1H), 7.44-7.41 (m, 2H); LC/MS 198.0 (M+1); HRMS (EI): m/z calcd for $\text{C}_{12}\text{H}_7\text{NO}_2$ 197.0476, found 197.0469.

6H-Naphtho[2,1-c]chromen-6-one (3{24})

Yield 87%, white solid, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 8.87 (d, $J = 7.5$ Hz, 1H), 8.52 (d, $J = 8.0$ Hz, 1H), 8.33 (d, $J = 8.5$ Hz, 1H), 8.01-7.94 (m, 2H), 7.75-7.68 (m, 2H), 7.56-7.49 (m, 2H), 7.42-7.39 (m, 1H); LC/MS 246.9 (M+1); HRMS (EI): m/z calcd for $\text{C}_{17}\text{H}_{10}\text{O}_2$ 246.0680, found 246.0689.

6H-[1]Benzothieno[2,3-c]chromen-6-one (3{25})

Yield 94%, white solid, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 8.65-8.62 (m, 1H), 8.49 (d, $J = 8.0$ Hz, 1H), 8.03-8.01 (m, 1H), 7.65-7.61 (m, 2H), 7.58-7.52 (m, 2H), 7.47-7.44 (m, 1H); LC/MS 252.9 (M+1); HRMS (EI): m/z calcd for $\text{C}_{15}\text{H}_8\text{O}_2\text{S}$ 252.0245, found 252.0241.

Methyl 3-fluoro-6-oxo-6H-benzo[c]chromene-8-carboxylate (3{26})

Yield 88%, white solid, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 9.05 (s, 1H), 8.46 (dd, $J = 7.0, 2.0$ Hz, 1H), 8.13-8.12 (m, 2H), 7.14-7.12 (m, 2H), 3.99 (s, 3H); LC/MS 273.2 (M+1); HRMS (EI): m/z calcd for $\text{C}_{15}\text{H}_9\text{FO}_4$ 272.0430, found 272.0428.

2-Methyl-8-(trifluoromethyl)-6H-benzo[c]chromen-6-one (3{27})

The residue was purified with Biotage SP1 flash autopurification system to afford the white solid in 69% yield, $^1\text{H-NMR}$ (CDCl_3 , 500 MHz) δ 8.67 (s, 1H), 8.24-8.23 (d, $J = 8.0$ Hz,

1H), 8.03-8.02 (d, $J = 2.0$, 1H), 7.87 (s, 1H), 7.35-7.27 (m, 2H), 2.49 (s, 3H); LC/MS 279.9 (M+1); HRMS (EI): m/z calcd for $C_{15}H_9F_3O_2$ 278.0554, found 278.0545.

4-Chloro-9-fluoro-6H-benzo[c]chromen-6-one (3{28})

Yield 71%, white solid, 1H -NMR ($CDCl_3$, 500 MHz) δ 8.47-8.44 (m, 1H), 7.88 (dd, $J = 7.0$, 1.0 Hz, 1H), 7.74 (dd, $J = 7.5$, 2.5 Hz, 1H), 7.59 (dd, $J = 6.0$, 1.5 Hz, 1H), 7.34-7.28 (m, 2H); LC/MS 249.8 (M+1); HRMS (EI): m/z calcd for $C_{13}H_6ClFO_2$ 248.0040, found 248.0029.

2-Methoxy-8-(trifluoromethyl)-6H-benzo[c]chromen-6-one (3{29})

Yield 85%, white solid, 1H -NMR ($CDCl_3$, 500 MHz) δ 8.69 (s, 1H), 8.19 (d, $J = 8.5$ Hz, 1H), 8.04 (d, $J = 2.0$ Hz, 1H), 7.51 (s, 1H), 7.34 (m, 1H), 7.14-7.12 (m, 1H); LC/MS 295.1 (M+1); HRMS (EI): m/z calcd for $C_{15}H_9F_3O_3$ 294.0503, found 294.0512.

3-Chloro-8-methyl-6H-benzo[c]chromen-6-one (3{30})

Yield 92%, white solid, 1H -NMR ($CDCl_3$, 500 MHz) δ 8.19 (s, 1H), 7.97-7.94 (m, 2H), 7.65-7.64 (m, 1H), 7.36-7.28 (m, 2H), 2.50 (s, 3H); LC/MS 245.6, 247.2 (M+1); HRMS (EI): m/z calcd for $C_{14}H_9ClO_2$ 244.0291, found 244.0306.

9-Chloro-1,2-difluoro-6H-benzo[c]chromen-6-one (3{31})

Yield 70%, white solid, 1H -NMR ($CDCl_3$, 500 MHz) δ 8.51-8.50 (m, 1H), 8.40 (d, $J = 8.0$ Hz, 1H), 7.64 (dd, 7.0, 2.0 Hz, 1H), 7.37-7.33 (m, 1H), 7.19-7.17 (m, 1H); LC/MS 267.8, 269.9 (M+1); HRMS (EI): m/z calcd for $C_{13}H_5ClF_2O_2$ 266.9946, found 265.9955.

2-Hydroxy-8-(trifluoromethyl)-6H-benzo[c]chromen-6-one (3{32})

The residue was purified with Biotage SP1 flash autopurification system to afford the white solid in 72% yield, 1H -NMR (CD_3OD , 500 MHz) δ 9.33-9.30 (m, 1H), 9.24 (s, 1H), 9.05-9.03 (m, 1H), 8.48-8.47 (m, 1H), 8.13-8.11 (m, 1H), 7.90-7.87 (m, 1H); LC/MS 281.1 (M+1); HRMS (EI): m/z calcd for $C_{14}H_7F_3O_3$ 280.0428, found 280.0431.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research work was supported by Grants DA007215 and DA003801 from the National Institute of Health and National Institute of Drug Abuse. This paper is dedicated to Prof. John R. Scheffer, University of British Columbia on the occasion of his 70th birthday. One of the authors (KV) would like to thank Prof. J. Narasimha Moorthy, Indian Institute of Technology, Kanpur for helpful suggestions.

Reference and Notes

- (a) Koch K, Podlech J, Pfeiffer E, Metzler M. J Org Chem. 2005; 70:3275–3276. [PubMed: 15822993] (b) Abe H, Nishioka K, Takeda S, Arai M, Takeuchi Y, Harayama T. Tetrahedron Lett. 2005; 46:3197–3200. (c) Sidwell WTL, Fritz H, Tamm C. Helv Chim Acta. 1971; 54:207–215. (d) Raistrick H, Stilings CE, Thomas R. Biochemistry. 1953; 55:421–433. (e) Pero RW, Harvan D. Tetrahedron Lett. 1973; 12:945–948.
- (a) Garino C, Bihel F, Pietrancosta N, Laras Y, Quelever G, Woo I, Klein P, Bain J, Boucher JL, Kraus JL. Bioorg Med Chem Lett. 2005; 15:135–138. [PubMed: 15582426] (b) Sun W, Cama LD, Birzin ET, Warrier S, Locco L, Mosley R, Hammond ML, Rohrer SP. Bioorg Med Chem Lett. 2006; 16:1468–1472. [PubMed: 16412638]

3. (a) Edwards JP, West SJ, Marschke KB, Mais DE, Gottardis MM, Jones TK. *J Med Chem.* 1998; 41:303–310. [PubMed: 9464361] (b) Hamann LG, Higuchi RI, Zhi L, Edwards JP, Wang XN, Marschke KB, Kong JW, Farmer LJ, Jones TK. *J Med Chem.* 1998; 41:623–639. [PubMed: 9484511] (c) Coghlan MJ, Kym PR, Elmore SW, Wang AX, Luly JR, Wilcox D, Stashko M, Lin CW, Miner J, Tyree C, Nakane M, Jacobson P, Lane BC. *J Med Chem.* 2001; 44:2879–2885. [PubMed: 11520196]
4. Schmidt JM, Tremblay GB, Page M, Mercure J, Feher M, Dunn-Dufault R, Peter MG, Redden PR. *J Med Chem.* 2003; 46:1289–1292. [PubMed: 12672229]
5. Myrray, R.; Mendez, J.; Brown, S. *The Natural Coumarins: Occurrence, Chemistry and Biochemistry.* John Wiley & Sons; New York: 1982. p. 97-111.
6. (a) Zhou QJ, Worm K, Dolle RE. *J Org Chem.* 2004; 69:5147–5149. [PubMed: 15255755] (b) Kemperman GJ, Ter Horst B, Van de Goor D, Roeters T, Bergwerff J, van der Eem R, Basten J. *Eur J Org Chem.* 2006; 14:3169–3174. (c) Hussain I, Nguyen VTH, Yawer TT, Fiscer C, Reinke H, Langer P. *J Org Chem.* 2007; 72:6255–6258. [PubMed: 17604399] (d) Alo BI, Patil PA, Sharp MJ, Siddiqui MA, Snieckus V. *J Org Chem.* 1991; 56:3763–3768.
7. Thasana N, Worayuthakarn R, Kradanrat P, Hohn E, Young L, Ruchirawat S. *J Org Chem.* 2007; 72:9379–9382. [PubMed: 17975927]
8. Jung ME, Allen DA. *Org Lett.* 2009; 11:757–760. [PubMed: 19138152]
9. Sanz R, Fernandez Y, Castroviejo MP, Perez A, Fananas FJ. *Eur J Org Chem.* 2007; 1:62–69.
10. Teske JA, Deiters A. *Org Lett.* 2008; 10:2195–2198. [PubMed: 18452299]
11. Ku YY, Grieme T, Raju P, Morton HE, Rozema M, King SA. *J Org Chem.* 2003; 68:3238–3240. [PubMed: 12688796]
12. Jilani JA. *Chem, Pap.* 2007; 61:410–412.
13. Khanolkar AD, Lu D, Ibrahim M, Duclos R Jr, Thakur GA, Malan TP Jr, Porreca F, Veerappan V, Tian X, George C, Parrish DA, Papahatjis DP, Makriyannis A. *J Med Chem.* 2007; 50:6493–6500. [PubMed: 18038967]
14. McOmie JFW, Watts ML, West DE. *Tetrahedron.* 1968; 24:2289–2292.
15. (a) Dallinger D, Kappe O. *Chem Rev.* 2007; 107:2563–2591. [PubMed: 17451275] (b) Lidstrom P, Tierney J, Wathey B, Westman J. *Tetrahedron.* 2001; 57:9225–9283.
16. (a) Kennedy JP, Williams L, Bridges DN, Weaver D, Lindsley CW. *J Comb Chem.* 2008; 10:345–354. [PubMed: 18220367] (b) Lew A, Krutzik PO, Hart ME, Chamberlin R. *J Comb Chem.* 2002; 4:95–105. [PubMed: 11886281]
17. Liu B, Moffett KK, Joseph RW, Dorsey BD. *Tetrahedron Lett.* 2005; 46:1779–1782.
18. Wolfe JP, Singer RA, Yang BH, Buchwald SL. *J Am Chem Soc.* 1999; 121:9550–9561.
19. Zhang W, Pugh G. *Tetrahedron Lett.* 2001; 42:5613–5615.

Figure 1.
Diversity of bromo arylcarboxylates **1**

Figure 2.
Diversity of *o*-hydroxyarylboronic acids 2

Scheme 1. Suzuki-Miyaura cross coupling of bromo arylcarboxylates 1 and *o*-hydroxyarylboronic acids 2

Table 1

Optimization of Suzuki-Miyaura cross coupling reaction conditions.

<p> $1\{1\}$ X = H Y = OCH₃ $1\{2\}$ X = F Y = H </p>			
Entry	Conditions	Yield ^a (%)	
		3{1}	3{2}
1	Chloro(di-2-norbornylphosphino)(2'-dimethylamino-1,1'-biphenyl-2-yl)palladium (II) (5 mol %), KF (5 equiv), dioxane, 130 °C, 15 min	54	48
2	Pd(OAc) ₂ (1 mol %), (<i>o</i> -biphenyl)P(t-Bu) ₂ (2 mol %), KF (3 equiv), THF, rt, 24 h	45	47
3	PdCl ₂ (PPh ₃) ₂ (10 mol %), Cs ₂ CO ₃ (4 equiv), DME, H ₂ O, 130 °C, 15 min	38	41
4	1,1'-bis(diphenylphosphino) ferrocene dichloropalladium (II) (10 mol %), NaOAc (5 equiv), DME, H ₂ O, 150 °C, 15 min	31	29
5	Pd(PCy ₃) ₂ Cl ₂ (5 mol %), CsF (2 equiv), NMP, 130 °C, 15 min	31	32
6	Pd(PPh ₃) ₄ (10 mol %), Cs ₂ CO ₃ (4 equiv), DME, H ₂ O, 125 °C, 15 min	98	96

^a Isolated product after purification by silica gel column chromatography.

Table 2Synthesis of a library of benzopyranone and heterocyclic analogues **3**.

<p>Ar = ary R¹ and R² = EDG/EWG</p>				
Entry	1	2	Product	Yield ^a (%)
1	1{1}	2{1}	3{1}	98 ^{b,19}
2	1{2}	2{1}	3{2}	96
3	1{3}	2{1}	3{3}	88
4	1{4}	2{1}	3{4}	88 ^{b,6a}
5	1{5}	2{1}	3{5}	80
6	1{6}	2{1}	3{6}	88 ^{b,6a}
7	1{7}	2{1}	3{7}	83
8	1{8}	2{1}	3{8}	80

<p>Ar = aryl R¹ and R² = EDG/EWG</p>				
Entry	1	2	Product	Yield ^a (%)
9	1{9}	2{1}	 3{9}	85
10	1{10}	2{1}	 3{10}	90
11	1{11}	2{1}	 3{11}	92
12	1{12}	2{2}	 3{12}	89
13	1{13}	2{1}	 3{13}	92 ^{b,6a}
14	1{14}	2{1}	 3{14}	84
15	1{15}	2{1}	 3{15}	92
16	1{16}	2{1}	 3{16}	86

<p>Ar = ary R¹ and R² = EDG/EWG</p>				
Entry	1	2	Product	Yield ^a (%)
17	1{17}	2{1}	3{17}	68
18	1{18}	2{1}	3{18}	86
19	1{2}	2{3}	3{19}	94
20	1{19}	2{1}	3{20}	85
21	1{20}	2{1}	3{21}	84
22	1{21}	2{1}	3{22}	95
23	1{22}	2{1}	3{23}	83
24	1{23}	2{1}	3{24}	87

<p>Ar = aryl R¹ and R² = EDG/EWG</p>				
Entry	1	2	Product	Yield ^a (%)
25	1{24}	2{1}	3{25}	94
26	1{8}	2{4}	3{26}	88
27	1{7}	2{5}	3{27}	69
28	1{2}	2{6}	3{28}	71
29	1{7}	2{7}	3{29}	85
30	1{4}	2{9}	3{30}	92
31	1{11}	2{8}	3{31}	70
32	1{7}	2{10}	3{32}	72

^a Isolated product after purification by silica gel column chromatography.

^b Known compounds, see references 6a and 19.