Received:
26 November 2012

Revised:
26 April 2013

Accepted:
7 May 2013

doi: 10.1259/bjr.20120614

Cite this article as:

Obadina ET, Torrealba JM, Kanne JP. Acute pulmonary injury: high-resolution CT and histopathological spectrum. *Br J Radiol* 2013;86: 20120614.

PICTORIAL REVIEW

Acute pulmonary injury: high-resolution CT and histopathological spectrum

¹E T OBADINA, MD, ²J M TORREALBA, MD and ¹J P KANNE, MD, FCCP

¹Department of Radiology, University of Wisconsin School of Medicine and Public Health, Madison, WI, USA

²Department of Pathology, University of Texas Southwestern Medical Center, Dallas, TX, USA

Address correspondence to: Dr Jeffrey P. Kanne

E-mail: kanne@wisc.edu

ABSTRACT

Acute lung injury usually causes hypoxaemic respiratory failure and acute respiratory distress syndrome (ARDS). Although diffuse alveolar damage is the hallmark of ARDS, other histopathological patterns of injury, such as acute and fibrinoid organising pneumonia, can be associated with acute respiratory failure. Acute eosinophilic pneumonia can also cause acute hypoxaemic respiratory failure and mimic ARDS. This pictorial essay reviews the high-resolution CT findings of acute lung injury and the correlative histopathological findings.

Patients with acute pulmonary injury (API) typically present with dyspnoea, tachypnoea and hypoxaemia and develop acute hypoxaemic respiratory failure, most commonly manifesting as acute respiratory distress syndrome (ARDS) [1]. ARDS is defined as acute hypoxaemia with a ratio of partial pressure of arterial oxygen to the fraction of inspired oxygen ($\text{PaO}_2:\text{FIO}_2$) ≤ 200 mmHg, bilateral lung opacities with a radiographic appearance consistent with pulmonary oedema and no clinical evidence of cardiac failure [2]. Acute lung injury (ALI) is less severe than ARDS and has similar diagnostic criteria, with the exception of $\text{PaO}_2:\text{FIO}_2 < 300$ mmHg [2].

Most patients with clinical ARDS and ALI have histological evidence of diffuse alveolar damage (DAD). Other histological patterns associated with ALI and ARDS include acute eosinophilic pneumonia (AEP) and acute fibrinous and organising pneumonia (AFOP). Organising pneumonia can develop in the setting of ALI but will not be discussed here. Patients with API invariably have abnormal chest radiographs, given the widespread histopathological abnormalities. High-resolution CT (HRCT) is typically performed early in the disease course to aid in establishing a diagnosis of API. Once a diagnosis of API is established, most patients are followed up with chest radiography, as needed. HRCT may be used in selected cases when problems, such as superimposed infection and its own respective complications, arise. In this pictorial essay, we illustrate HRCT findings associated with API and correlate the histopathological features (Table 1).

DIFFUSE ALVEOLAR DAMAGE

DAD is the most common histological manifestation of ARDS and ALI. Patients typically present with severe hypoxaemia and require mechanical ventilation [1]. Numerous causes of DAD have been described, including infections, inhalants, drugs, shock, sepsis and radiation [3]. The exact cause of DAD may not be apparent from imaging alone, as the imaging findings of DAD are similar regardless of the cause. However, integration of clinical and earlier imaging findings may suggest the precipitating event, such as aspiration or infection. Although the precise mechanism of injury is unclear, capillary endothelial and alveolar epithelial damage result in exudation of fluid and products of cellular breakdown. With time, pneumocyte hyperplasia and fibroblast proliferation ensue [1].

The histological appearance of DAD is divided into three phases: acute or exudative phase, organising or proliferative phase and chronic or fibrotic phase. The acute phase (Figure 1) occurs during the first week after the initial insult. The main histological findings seen during this phase are capillary congestion, interstitial and intra-alveolar oedema associated with fibrin and formation of hyaline membranes, which are most prominent at 3–7 days [3,4]. Thrombi may develop as a result of localised alterations in the coagulation pathway. The organising phase (Figure 2) occurs after 1–2 weeks and is characterised by fibroblast proliferation admixed with scattered mononuclear inflammatory cells, resulting in the formation of organised granulation tissue [4,5]. Type-2 pneumocyte

Table 1. Radiological and pathological findings of acute pulmonary injury

Pattern of injury	HRCT findings	Distribution	Histopathology
Diffuse alveolar damage (DAD acute)	Consolidation, ground-glass opacity, crazy paving	Diffuse with posterior and basal predominance	Capillary congestion, interstitial and intra-alveolar oedema associated with fibrin, hyaline membranes
DAD (organising and fibrotic)	Development of traction bronchiectasis and reticulation	Anterior predominance	Fibroblast, scattered mononuclear inflammatory cells, organised granulation tissue, Type-2 pneumocyte hyperplasia, interstitial fibrosis (late)
Acute fibrinous and organising pneumonia	Consolidation and ground-glass opacity	Peripheral and peribronchial with basal predominance	Fresh intra-alveolar fibrin ("fibrin balls"), mild interstitial widening, lymphocytic infiltrates, organising fibroblastic tissue, hyaline membranes absent
Acute eosinophilic pneumonia	Consolidation, ground-glass opacity, thickening of interlobular septa and bronchovascular bundles, small pleural effusions, absent cardiomegaly	Random or peripheral	Intra-alveolar fibrin, macrophages, hyaline membranes, numerous eosinophils (in interstitium, blood vessel walls), eosinophilic microabscesses

HRCT, high-resolution CT.

Figure 1. Diffuse alveolar damage (acute phase). The photomicrograph shows diffuse hyaline membranes (arrows) lining the alveolar walls along with the erythrocytes (arrowheads) in air spaces. An interstitial mononuclear inflammatory infiltrate is present. Haematoxylin and eosin 400×.

hyperplasia also occurs and may show considerable cytological atypia [4]. Oedema and hyaline membranes are not prominent during this phase [4]. The fibrotic phase is characterised by features of interstitial fibrosis, including alveolar septal thickening from collagen deposition [3].

Initially, HRCT shows heterogeneous foci of consolidation and ground-glass opacity with a gravitationally dependent gradient, with more consolidation in the posterobasal portions of the lungs. A diffuse "crazy-paving" pattern can also be seen [5] (Figures 3 and 4). Air bronchograms and small pleural effusions are common. With organisation and fibrosis, reticulation and traction bronchiectasis may develop (Figure 5), often having an

Figure 2. Diffuse alveolar damage (organising phase). The photomicrograph shows hyaline membranes (arrows) incorporated in the alveolar septa with fibroblast proliferation and septal widening. Interstitial inflammatory infiltrates may persist at this stage. Haematoxylin and eosin 200×.

Figure 3. A 54-year-old female with daptomycin-induced diffuse alveolar damage (DAD). Transverse (a) and coronal (b) high-resolution CT images at presentation show peripheral and basal predominant foci of consolidation with halos of ground-glass opacity (arrows). Over the course of a week, the patient developed acute respiratory distress syndrome. Early DAD can have an appearance similar to organising pneumonia, as in this case, but patients with DAD usually deteriorate rapidly.

anterior predominance. Complications include pneumothorax, pneumomediastinum, interstitial emphysema and formation of subpleural bullae or cysts [5].

ACUTE FIBRINOUS AND ORGANISING PNEUMONIA

AFOP is a recently described histological pattern associated with ALI. Most patients present with severe respiratory failure similar to ARDS. However, a small subset of patients may have a milder course [1]. AFOP can be idiopathic or related to a wide range of insults similar to ARDS.

The dominant histopathological finding in AFOP is fresh intra-alveolar fibrin ("fibrin balls") (Figure 6) [1,6]. In contrast to

Figure 5. An 80-year-old female with acute respiratory distress syndrome following surgery. (a) The high-resolution CT (HRCT) image shows patchy consolidation and ground-glass opacity in the lower lobes with mild septal thickening (arrow-heads). Small pleural effusions are present. (b) The HRCT image taken 3 months later shows interstitial fibrosis characterised by reticulation, traction bronchiectasis (arrows) and ground-glass opacity.

Figure 4. A 71-year-old male with acute respiratory distress syndrome caused by sepsis. The high-resolution CT image shows bilateral consolidation predominantly affecting the dependent areas of the lungs and ground-glass opacity and septal thickening anteriorly. Small pleural effusions (arrow-heads) are present.

Figure 6. Acute fibrinous and organising pneumonia. The photomicrograph shows predominantly intra-alveolar fibrin aggregates ("fibrin balls"). Associated mild interstitial mononuclear infiltrate is also present. Haematoxylin and eosin 200 $\times$.

DAD, hyaline membranes are absent. Other associated histological findings include mild interstitial widening, lymphocytic infiltrates and organising fibroblastic tissue.

HRCT findings of AFOP are not well described but include peripheral and peribronchial foci of consolidation and ground-glass opacity, similar to organising pneumonia, but greater in extent with superimposed features more typical of DAD, such as more extensive ground-glass opacity and consolidation and a gravitationally dependent gradient (Figures 7 and 8, [6]).

ACUTE EOSINOPHILIC PNEUMONIA

AEP is characterised by the acute onset of fever, hypoxaemia and severe respiratory distress, in the absence of infection [7]. AEP may be idiopathic or may result from toxic inhalation, drug reaction and cigarette smoking [1,8]. Peripheral blood eosinophilia is often absent in contrast to chronic eosinophilic pneumonia [7]. Eosinophilia on bronchoalveolar lavage (>25% eosinophils) or lung biopsy is the key. In contrast to other forms

Figure 7. A 57-year-old male with daptomycin-induced acute fibrinoid and organising pneumonia. (a) High-resolution CT (HRCT) image at presentation shows bilateral central peribronchial ground-glass opacity (arrows) with mild septal thickening. (b) The HRCT image 17 days later shows extensive peribronchial, subpleural (arrows) and perilobular consolidation and ground-glass opacity, similar to but more extensive than organising pneumonia. (c) The HRCT image 10 weeks after presentation shows residual bands of perilobular consolidation (arrows) and mild bronchial dilation (arrowheads).

Figure 8. A 58-year-old female with rheumatoid arthritis and acute fibrinoid and organising pneumonia: transverse (a) and coronal (b). The high-resolution CT images show patchy consolidation and ground-glass opacity in a random distribution.

Figure 9. Acute eosinophilic pneumonia. The photomicrograph shows interstitial widening accompanied by mixed infiltrates of lymphocytes, macrophages and eosinophils. Focal alveolar fibroblastic proliferation is also present (arrow). Haematoxylin and eosin 200 $\times$.

of API, patients with AEP usually have a dramatic response to corticosteroids, with rapid resolution of clinical signs and symptoms and radiographic abnormalities [7,8].

The histopathological findings of AEP include intra-alveolar fibrin, macrophages, hyaline membranes and numerous eosinophils (Figure 9, [1]). Eosinophils may be present in the interstitium and blood vessel walls and may form eosinophilic microabscesses [1].

The HRCT findings of AEP include bilateral consolidation and ground-glass opacity in a random or peripheral distribution. Smooth interlobular septal thickening, bronchovascular bundle thickening and small pleural effusions are frequently present

Figure 10. An 18-year-old female with acute eosinophilic pneumonia resulting from new-onset cigarette smoking. The high-resolution CT image shows diffuse septal thickening (arrowheads) and multiple peripheral foci of lung consolidation (arrows).

and, in the absence of cardiomegaly, are helpful clues during diagnosis (Figure 10, [8]).

CONCLUSION

Although the HRCT findings of consolidation and ground-glass opacity in general and in the setting of API in particular are often non-specific, certain features may suggest the underlying pathological process. Distribution of abnormalities is important in narrowing the differential diagnosis with peripheral lung opacities more common in eosinophilic pneumonia and peribronchovascular opacities in AFOP. Knowledge of the distinguishing HRCT findings and correlation with the clinical scenarios in which these injuries occur allow the radiologist to add value to the clinical evaluation and management of these critically ill patients.

REFERENCES

1. Beasley MB. The pathologist's approach to acute lung injury. *Arch Pathol Lab Med* 2010; 134:719–27. doi: 10.1043/1543-2165-134.5.719.
2. Bernard GR, Artigas A, Brigham KL, Carlet J, Falke K, Hudson L, et al. The American-European Consensus Conference on ARDS. Definition, mechanisms, relevant outcomes, and clinical trial coordination. *Am J Respir Crit Care Med* 1994;149:818–24.
3. Castro CY. ARDS and diffuse alveolar damage: a pathologist's perspective. *Semin Thorac Cardiovasc Surg* 2006;18:13–19. doi: 10.1053/j.semtcvs.2006.02.001.
4. Sarmiento X, Guardiola JJ, Almirall J, Mesalles E, Mate JL, Soler M, et al. Discrepancy between clinical criteria for acute respiratory distress syndrome secondary to community acquired pneumonia with autopsy findings of diffuse alveolar damage. *Respir Med* 2011;105:1170–5.
5. Ketani LH, Godwin JD. A new view of pulmonary oedema and acute respiratory distress syndrome. *J Thorac Imaging* 1998;13:147–71.
6. Beasley MB, Franks TJ, Galvin JR, Gochuico B, Travis WD. Acute fibrinous and organizing pneumonia: a histological pattern of lung injury and possible variant of diffuse alveolar damage. *Arch Pathol Lab Med* 2002;126:1064–70. doi: 10.1043/0003-9985(2002)126<1064:AFAOP>2.0.CO;2.
7. Tazelaar HD, Linz LJ, Colby TV, Myers JL, Limper AH. Acute eosinophilic pneumonia: histopathologic findings in nine patients. *Am J Respir Crit Care Med* 1997;155:296–302. doi: 10.1164/ajrccm.155.1.9001328.
8. Daimon T, Johkoh T, Sumikawa H, Honda O, Fujimoto K, Koga T, et al. Acute eosinophilic pneumonia: thin-section CT findings in 29 patients. *Eur J Radiol* 2008;65:462–7. doi: 10.1016/j.ejrad.2007.04.012.