Published in final edited form as:

Hum Relat. 2012 June ; 65(6): 705–728. doi:10.1177/0018726711435250.

Workforce Disengagement Stressors and Retiree Alcohol Misuse: The Mediating Effects of Sleep Problems and the Moderating Effects of Gender

Elena Belogolovsky,

Technion-Israel Institute of Technology, Israel

Peter Bamberger, and

Tel Aviv University, Israel & Cornell University, USA

Samuel Bacharach

Cornell University, USA

Abstract

We generate and test a moderated mediation model of the effects of two retirement-related stressors (namely, financial and marital) on the severity of alcohol misuse among retirees. We posit that in addition to using alcohol to cope with stressors in retirement, alcohol may also be used to self-medicate the secondary, sleep-related effects of such stressors, and that gender serves as a key boundary condition, moderating the impact of such stressors on sleep-related problems, and of sleep-related problems on alcohol misuse. Using longitudinal data collected from a sample of 292 retirees, our findings generally support this model, suggesting that both stressors are associated with the severity of alcohol misuse among male retirees. Moreover, our findings demonstrate that -- for male retirees -- the effect of both stressors on the severity of alcohol misuse is to a large extent secondary to the stressors themselves, mediated by the sleep-related problems they may generate.

Keywords

alcohol misuse; sleep-related problems; stressors; retirement

Although workforce disengagement, be it in the form of unemployment, underemployment or retirement, can have significant mental health implications (Lin & Lueng, 2010), few studies have examined the impact of workforce disengagement on alcohol misuse, with those that have generating largely inconsistent findings (Dooley et al., 1998; NIAAA, 2006; Brennan, Schutte & Moos, 2010). For example, although cross-sectional and aggregate longitudinal studies find community unemployment rates to be associated with increased alcohol sales (Weeks & Drengacz, 1982) and treatment program admissions (Brenner, 1975), results of longitudinal studies at the individual level are more mixed with some studies finding unemployment to significantly increase the risk of an alcohol disorder diagnosis (Dooley et al., 1992), and others finding either null (Hammer, 1992) or highly contingent effects (Lahelma et al., 1995). Similarly, although several studies have identified a link between retirement and older adults' drinking behavior (Perriera & Sloan, 2001), others suggest that there is little or no difference between retirees' and non-retirees' patterns

of alcohol consumption (Ekerdt et al., 1989; Midanik et al., 1995), and still others find retirement to be associated with *reduced* alcohol consumption (Rosenkoetter et al., 2001).

In light of these mixed findings, several scholars have recently proposed that rather than trying to identify a direct link between workforce disengagement and drinking, research should focus on individuals' appraisal of their disengagement experience as a determinant of their mental health (Lin & Lueng, 2010) in general, and drinking behavior (Bacharach et al., 2007; 2008; Brennan et al., 2010) in particular. Such a perspective is grounded on findings which suggest that most of the adverse mental health consequences associated with such disengagement stem from a handful of disengagement-related stressors, with economic hardship and marital conflict being among those most commonly cited (Kessler et al., 1988; Price et al., 2002; McKee-Ryan et al., 2005). Although a recent study demonstrated the impact of such stressors on the general mental health among the unemployed (Lin & Lueng, 2010), to date, we have little understanding of the implications of these same stressors on more specific mental health outcomes such as alcohol misuse, particularly among those disengaging from the workforce on a more permanent basis, namely retirees. The absence of such research is particularly disturbing in that alcohol misuse is a serious and growing health problem among older adults, affecting nearly the same proportion of adults aged 60+ as those between the ages of 18 and 45 (approximately 20% of those in that cohort in the U.S.A., 10% –17% of those in that cohort in the U.K., and over 25% of males in that cohort in Italy -- Blow, 1998; Barnes et al., 2010; Hallgren, Hogberg & Andreasson, 2010), and resulting in direct hospital costs for this population in the USA of over \$60 billion/year (Schonfeld et al 1995; Dar, 2006). Accordingly, a primary objective of the current research is to enhance our understanding of the extent to which financial and marital stressors in retirement may be linked to the onset or exacerbation of alcohol misuse among retirees.

Additionally, we seek to understand *how* such stressors may affect alcohol misuse among older adults. Research examining the stress-alcohol relationship has largely been based on the tension reduction hypothesis which posits that alcohol misuse may serve as a means by which to mitigate the strain generated by exposure to stressors (Conger, 1956; Frone, 1999; 2008). This perspective suggests a direct effect of financial and marital stressors on alcohol misuse. However, from a resource ecology perspective (Hobfoll, 1988), there may be reason to believe that at least some of the effect of such stressors on alcohol misuse may be more indirect, with individuals turning to alcohol to medicate the physiological consequences of stress-related resource drain such as sleep-related problems. While subtle, this difference in how financial and marital stressors influence alcohol use may have significant implications with respect to both prevention and treatment. Accordingly, a second objective of the current study is to assess the degree to which sleep-related problems mediate the association between financial and marital stressors on retiree alcohol misuse.

Finally, while research suggests that males are at greater risk of alcohol misuse than females (NIAAA, 1998; Su et al., 1997), to date research on the consequences of disengagement-related stressors on mental health has yet to assess whether these consequences are the same for men and women. Accordingly, a third objective of the current study is to examine the degree to which gender may serve as a moderator of the relationships noted above.

In sum, in the current study we examine how two key stressors in retirement may be linked to the severity of alcohol misuse, how sleep-related problems may mediate this association, and how gender may moderate this pattern of relationships. More specifically, we propose a moderated-mediation model according to which financial and marital stressors serve as primary antecedents of shifts in retiree alcohol misuse, but whose effects are mediated by the impact such events may have on sleep-related problems and moderated by gender (see Figure 1). Defining retirement as “the exit from an organizational position or career path of

considerable duration, taken by individuals after middle age, and taken with the intention of reduced psychological commitment to work thereafter” (Feldman, 1994: 287), we test this model on the basis of a three-wave prospective design, regressing the pre- to 10-year post-retirement change in older employees’ alcohol misuse on stressors experienced 5 – 6 years after retiring, and sleep-related problems 1 – 2 years after that. Our findings contribute to both the workforce disengagement and alcohol literatures by offering insight into the psycho-physiological dynamics governing how and for whom stressors experienced in retirement may affect well-being in general, and alcohol misuse in particular.

Direct Effects of Stressors in Retirement on Retiree Alcohol Misuse

While for many, if not most individuals, retirement represents relief from the stresses and strains associated with work, for others, it may be associated with the precipitation or exacerbation of situational stressors (Atchley, 1989). Two stressors have received particular attention in the retirement literature namely those associated with financial/economic concerns (Beehr, 1986; Wang, 2007; Wang et al., 2008) and marital/family conflict (Kim & Feldman, 2000; Shultz et al., 1998; Szinovacz, 2003). Research suggests that for some, retirement may be associated with financial stressors as income from retirement sources (e.g., pension, social security) fails to match income previously earned on the job (Shultz, 1985; Keith, 1993). Indeed, studies suggest that discontinued sources of income, liquidated assets, and servicing debt on a fixed income could lead to increased mood disturbances even if these economic strains are anticipated (Dew & Yorgeson, 2010). Similarly, retirement may precipitate or exacerbate marital stressors (Kim & Moen, 2001) as couples adjust to new roles at home (Richardson & Kilty, 1991; Szinovacz, 1996) or to reduced income levels (Dew & Yorgeson, 2010). Such stressors may be particularly severe to the extent that couples’ retirement is asynchronous (i.e., one partner retires before the other) and/or to the extent that retirement generates a sense of status dissonance relative to traditional gendered roles at home (Moen et al., 2001). According to the tension reduction theory (Conger, 1956; Frone 2008), exposure to stressors induces alcohol misuse as individuals attempt to mitigate experienced tension and strain. The mechanisms by which substance use may reduce tension or strain include reduction of self-awareness (Hull, 1981), or redirection of one’s focus away from stress-producing cues toward more immediate cues (Steele & Josephs, 1988). Similarly, conservation of resources theory (Hobfoll, 1989) suggests that such stressors may result in the draining of psychological resources as individuals cope or attempt to cope with the loss or threatened loss of key personal resources such as energies (e.g., financial resources) and supportive relationships (e.g., marriage), and that, as a result, individuals may opt to use alcohol in an attempt to stem the depletion of psychological resources. More specifically, while retirement itself may be framed as resulting in the loss or potential loss of key resources such as money or support, secondary resource loss may occur as energy and social resources are utilized to stem the risk of primary resource loss. COR theory posits that as resources become depleted, individuals tend to adopt less efficient or maladaptive loss-control strategies and that this may result in the emergence of loss-spirals -- a pattern of increasingly more rapid depletion of the resources needed in order to be able to offset stress, improve one’s condition and reduce future stress vulnerability (Hobfoll et al., 2003: 633). Alcohol misuse may represent one maladaptive approach by which individuals attempt to address resource drain or depletion (Bacharach et al., 2008; Muys, 2007). Accordingly, consistent with the notion of tension-reduction, COR theory suggests that as resources become depleted, there may be an increased tendency to reject more resource-costly, problem-focused strategies of coping in favor of coping strategies viewed as resource-conserving (at least in the short-run) such as those involving physical or psychological withdrawal (Hobfoll, 2002) like alcohol misuse. Accordingly, we posit: *Hypothesis 1: The higher the intensity of financial (H1a) and marital (H1b) stressors in retirement, the greater the increase in the severity of alcohol misuse from pre-retirement levels.*

The Mediating Role of Sleep-Related Problems

Although alcohol may be used as a maladaptive means by which to mitigate strain or conserve resources in the face of stressors experienced in retirement (Frone, 2008; Hobfoll et al., 2003), another possibility is that alcohol misuse is *secondary* to such problems; used as a means by which to self-medicate the problems potentially generated by stressor-related resource depletion. Given that sleep disorders are common manifestations of resource drain and depletion (Soderstrom et al., 2004; Vela Bueno et al., 2008), as well as common health problems among older adults (Kalimo et al., 2000; Schubert et al., 2002), we propose that the sleep-related problems potentially generated by financial and marital stressors may explain much of the impact of such stressors on alcohol misuse.

Research indicates that aside from being linked to a number of demographic and hereditary factors, sleep-related problems may also be precipitated by resource drain and serve as a manifestation of resource depletion (Grossi et al., 2003; Melamed et al., 2006, Soderstrom et al., 2004). Indeed, in several studies, exhausted subjects consistently reported more sleep complaints, shorter sleep duration and frequent napping, and poor sleep quality than did vital subjects (Armon, 2009; Nicolson & van Diest, 2000; van Diest, 1990). Likely underlying this association between resource drain and sleep-related problems is the tendency of individuals to ruminate about their inability to fall asleep (Lundh & Broman, 2000; Middelkoop et al., 1996) and/or become hypervigilant while trying to fall asleep (Akerstedt et al., 2002; Arata et al., 2000). Such behaviors may be self-reinforcing and result in the disturbance of the hypothalamic-pituitary-adrenal (HPA) axis (Melamed et al., 2006) and the precipitation or exacerbation of sleep-disturbance cycles (Hall et al., 2008; Roth & Roehrs, 2003).

Research also suggests that alcohol is commonly used as a means of self-medication for those experiencing sleep problems with the motivation being the belief that alcohol will stem rumination, help them relax and thus fall asleep (Johnson, 1997; Roth, 2005). Indeed, more than 30% of individuals with persistent insomnia report using alcohol in such a manner (Ancoli-Israel Roth, 1999). Studies also suggest that the self-medication of sleep problems on the basis of alcohol misuse is particularly pervasive among older populations (Johnson, 1997; Roth, 2005).

Thus, beyond using alcohol as a maladaptive means of resource conservation in the face of stressors experienced in retirement, the linkages noted above suggest that alcohol may also be used as a means by which to self-medicate the effects of resource depletion (i.e., exhaustion) generated by these stressful experiences. Accordingly, we posit that at least part of the impact that stressors in retirement may have on alcohol misuse may be explained by sleep-related problems, or in other words: *Hypothesis 2: The impact of financial (H2a) and marital (H2b) stressors experienced in retirement on the severity of alcohol misuse is mediated by sleep-related problems.*

The Moderating Role of Gender

Gender likely serves as a critical conditioning factor for both the stressors in retirement → sleep-related problems and the sleep-related problems → retiree alcohol misuse relationships. Regarding the former, a higher prevalence of sleep-related problems among females, and particularly older females, has consistently been found across numerous epidemiological studies since the 1970s (Bixler, et al., 1979; LeBlanc et al., 2009; Li et al., 2002; Quan et al., 2005). More importantly for our model, there is evidence that environmental stressors may take a greater toll on women's sleep than on men's (Kushnir & Melamed, 1992). A number of factors have been proposed as underlying such an amplified

effect of stressors on sleep (Li et al., 2002). For example, several researchers argue that a higher prevalence of psychiatric morbidities (e.g., anxiety) among women may increase the likelihood (relative to men) of their manifesting sleep-related problems in response to heightened stress (Lindberg et al., 1997; Middelkoop et al., 1996). Biological differences in sex steroids have also been postulated to account for women's heightened sleep-related sensitivity to environmental stressors with several researchers proposing that hormone-related changes in the post-menopausal state may increase the sensitivity of female's sleep cycles to such stressors (Landis & Moe, 2004; Lee-Chiong, 2006; Manber & Armitage, 1999). Thus, we propose: *Hypothesis 3: Gender moderates the first stage of the sleep-mediated impact of financial (H3a) and marital (H3b) stressors in retirement on the severity of retiree alcohol misuse such that the impact of these stressors on sleep-related problems is stronger for female than male retirees.*

The notion that gender conditions the association between sleep-related problems and the severity of alcohol misuse (the second stage of the sleep-mediated impact of stressors in retirement on alcohol misuse severity) is consistent with research indicating that women and men react differently to sleep loss (Blatter et al., 2006). We posit that sleep-related problems will be more strongly associated with alcohol misuse for male than female retirees for two reasons. First, women tend to cope with sleep related problems by using tranquilizers/sleeping pills, while men may be more likely to use alcohol as a means by which to self-medicate sleep-related problems and the distress that such problems may generate (Lex, 1991). Second, a number of studies indicate that while the pharmacological effects of alcohol may be stress-reducing for both men and women, the cognitive impact is different. For example, several studies indicate that simply the belief that alcohol has been consumed may actually *increase* distress among women (Abrams & Wilson, 1979; Sutker et al., 1982), while other research indicates that women expect less relaxation after drinking than do men (Rohsenow, 1983). Cooper et al.'s (1992) findings – that the association between life event stress and alcohol misuse is significant and robust for men but not for women – lends further support to these assertions, as does a recent study of over 20,000 Finish adults finding men experiencing sleep problems to be nearly five times more likely to engage in periodic heavy drinking than women experiencing similar sleep-related problems (Hublin et al., 2007). Accordingly we posit: *Hypothesis 4: Gender moderates the second stage of the sleep-mediated impact of stressors in retirement on the severity of retiree alcohol misuse such that the impact of sleep-related problems on alcohol misuse severity is stronger for male than female retirees.*

METHOD

Sample and Procedure

We test our hypotheses using data from a longitudinal study of older workers who, at the time of initial data collection (T_0 in 2000), were within 6 months of being eligible to retire on the basis of full retirement benefits. Of the 2812 individuals in the target sample (identified through the membership files of nine American unions representing transportation, manufacturing and construction workers), 1279 agreed to participate (i.e., 46% response rate), with all data collected on the basis of telephone interviewing. Approximately half of the 1279 participants chose to retire upon becoming eligible for these benefits (i.e., within a year of the initial interview).

Given our focus on retirees in the current study, we analyzed data collected from the subset of those remaining in the study six years after initial contact and who had opted to fully retire (i.e., completely withdraw from the workforce). Accordingly, T_1 data were collected in late 2006, with T_2 data collected approximately 18 months later and T_3 data collected 2 years after T_2 . We opted for a two-year interval between T_2 and T_3 based on the research of

Weissman et al. (1997) and Crum et al. (2004) which suggests that while the effect of sleep-related problems on alcohol misuse may already be statistically capturable within a year of the onset of sleep problems, longer follow up periods likely reduce the risk of confounds and increase the robustness of the effect. Of the 927 participants remaining in the study at T₁ (i.e., a drop-out rate of 39.5 percent from the 1279 initially agreeing to participate at T₀ in 2000, the bulk of which resulted from outdated contact information), 868 participated in the T₃ survey. At T₁, the number of respondents from each employment sector was as follows: 660 respondents were members of 3 unions representing transport workers, 149 respondents were members of 2 unions representing manufacturing workers, and 118 respondents were members of 4 construction workers' unions. In order to be included in the study, the 868 remaining participants at T₃ had to have fully retired by T₁ (n = 673 or 73% of the T₁ sample). Additionally, because hospitalization would likely have a direct impact on drinking, they could not have been hospitalized during the past year (n=577). Finally, we retained only those for whom we had complete data along all study variables at all 3 points of data collection (n=292 or 43% of those who retired by T₁). T-test analysis indicates that those dropping out of or excluded from the sample between times T₁ and T₃ were not significantly different along any of the study variables from those remaining. Moreover, there was no significant change in the overall composition of the sample from T₁ to T₃, with the latter composed of 221 male respondents and 71 females (i.e., 76% male) having a mean age of 67.2 years (SD=4.3). 234 of the T₃ study participants were Caucasian (80.14%).

Measures

Dependent variable—*Severity of Alcohol Misuse* was assessed on the basis of the Drinking Problems Index (DPI), an instrument specially designed to assess the severity of drinking problems among older adults (Finney, Moos, & Brennan, 1991). We did so both prior to retirement eligibility (i.e., T₀) and subsequent to retirement at T₁ and T₃. Based on their analysis of the comparative validity of alternative measures of problem drinking (e.g., the CAGE instrument), Bamberger, Sonnenstuhl and Vashdi (2006: 132) find the DPI to offer high levels of both sensitivity and specificity, and conclude that, “the DPI offers significant advantages over the CAGE when the target population includes large numbers of older adults employed or soon to retire.”

For the DPI, respondents were read the following statement: “People often report that a variety of things happen as a result of their drinking.” Then, they were asked to respond to 17 problem-based items and indicate how often such problems were experienced in the last 12 months with response categories ranging from “never” (1) to “often” (5). A sample item is: “How often have you had a family member worry or complain about your drinking?” In order for the DPI scale (calculated by summing all the responses to the 17 items) to have a minimum response of zero (indicating NO drinking problems), per Finney et al. (1991), we transformed all the items from a 1 to 5 scale to a 0 to 4 scale. Additionally, as recommended by Finney et al. (1991:398), in order to decrease the influence of two “liberal” items (i.e., “got a buzz/high” and “drink to forget worries”), responses to these two items were dichotomized (responses 0 to 2 were recoded as 0, and responses 3 and 4 were recoded to equal 1 prior to their inclusion in the summed, scale score). As such, the potential scoring range for the DPI was 0 to 62 (top score of 4 for 15 items plus top score of two for two items). Despite the count nature of this variable, inter-item reliability is high (alpha for the DPI = .76 at T₀ and .85 at T₃).

Independent variables—*Financial Stressors* (T₁) were assessed on the basis of 5 items from Ilfeld's (1976) financial stressors measure. Participants were asked to report on a five-point scale (1=never, 5=often), how often in the past year it happened that they did not have enough money to afford the kind of food they or their family should have, the kind of

medical care they or their family should have, etc. Cronbach alpha was .78. *Marital Stressors* (T_1) were measured using two items from Spanier's (1976) Dyadic Adjustment Scale (DAS). The items were drawn from two DAS two subscales, namely satisfaction (e.g., 'How often do you and your mate "get on each other's nerves"?'') and conflict (e.g., 'How often do you and your partner quarrel?'). The items were rated on a 5-point scale (0=never, 4=several times a day). Cronbach alpha was .78.

Mediator variable—*Sleep-related Problems* were measured at T_2 using the Pittsburgh Sleep Quality Index (PSQI; Buysse et al., 1989). This self-administered questionnaire assesses quality of sleep during the previous month and contains 19 self-rated questions covering 7 components including subjective sleep quality, sleep duration and disturbance, and daytime dysfunction. Each component is scored from 0–3, yielding a global score between 0–21 (higher scores = greater sleep-related problems).

Moderator variable—*Gender* was coded such that 0=male and 1=female.

Control variables—Because post-retirement drinking may be associated with a variety of demographic, health, and situational factors, we controlled for the following variables at T_1 : *Age* (in years, ranging from 53–74); *Marital Status* (1=married, else=0); *Minority* (0=Caucasian; 1=other); and *Education* -- measured on the basis of a 6-level categorical variable with up to some high school=1, high school degree or equivalent=2, some college=3, associates degree or equivalent=4; bachelors degree or equivalent=5, advanced (MA/MSc or PhD) degree=6. Additionally, we controlled for *Years Since Retirement* at T_3 , and *Recent Physical Health* at T_1 -- measured on the basis of 6 items adapted from Moos and Moos (1994) by asking participants report whether they experienced certain symptoms in the past month (e.g., trouble breathing, pain in back/spine). Finally, *Social desirability* at T_1 was included as a control variable because it may in some cases create spurious relationships, whereas in others, it might suppresses relationships that might otherwise be significant. Using the self-deceptive enhancement component of Paulhus' (1991) Balanced Inventory of Desirable Responding instrument, social desirability was measured on the basis of a 5-point response format (1 = not at all true, 5 = very true), with a higher value indicating greater self-deceptive enhancement and consequently an increased risk of a social desirability bias. As Paulhus recommends, we summed the responses for the 19 items. Finally, in order to capture the impact of the hypothesized variables on the change in individual drinking behavior from pre-retirement levels, we controlled for the baseline level of alcohol misuse (i.e., DPI at T_0).

Analytical procedure

Our dependent variable (severity of alcohol misuse) has two important characteristics that influence the choice of the appropriate analytical method. First, it is non-negative. Second, it exhibits a pattern of over-dispersion (i.e., a high frequency of abstinence or moderate consumption with no problems relative to other levels of problem drinking). In such cases, where over-dispersion may occur as a result of excess zeros, a zero-inflated count model can be used. Zero-inflated count models assume that the process generating the excess zeros is qualitatively different from the process that generates the non-zeros (Greene, 2008; Tu, 2006). With our data exhibiting evidence of excess zeros as well as over-dispersion due to unobserved heterogeneity, we opted to test our hypotheses on the basis of a Zero-inflated Poisson (ZIP) model (Cheung, 2002; Lambert, 1992). ZIP models correct for these types of distributions by assuming a Poisson distribution and incorporating a correction for over-dispersion (Hernandez-Avila et al., 2006; Zorn, 1996).

Following the approach recommended by Edwards and Lambert (2007) and Muller, Judd and Yzerbyt (2005) we tested our hypotheses in four steps. First, we tested the main effect model suggested by Hypotheses 1a and 1b. We then tested the simple mediation model suggested by Hypotheses 2a and 2b. Finally, we incorporated into this same model gender as a moderator variable in order to test for moderated-mediation (posited in Hypotheses 3a, 3b and 4, respectively). Prior to the analyses, the continuous variables (i.e., marital and financial stressors and sleep-related problems) were mean-centered as recommended by Aiken and West (1991) and simple slopes for the effects of sleep-related problems (M) on alcohol misuse at T₃ (Y) were evaluated on the basis of Aiken and West (1991). We assess the significance of the indirect effects on the basis of bootstrapped confidence intervals (Edwards & Lambert, 2007; Preacher, Rucker & Hayes., 2007). Additionally, we assess the relative predictive utility of the alternative models that we test on the basis of differences in the pseudo R^2 and $-2\log$ likelihood associated with these comparative models.

Finally and as noted above, restrictions on the nature of the causal influences between our variables of interest were included in these models. More specifically, Rogosa (1980) elaborated on the problems of inferring causality from a longitudinal panel data in which the relations between two variables are examined without controlling for the appropriate baseline variables. It has been argued that controlling for the dependent variable at baseline (Y₀ or in this case, alcohol misuse prior to retirement) when examining the effect of X (i.e., stressors,) on Y₃ (i.e., alcohol misuse at T₃) may not only allow one to rule out the rival hypothesis that alcohol misuse causes stressors, but also greatly reduces the threat of spuriousness (Allison, 1990: 94). Accordingly, in the current study we applied the regressor variable method, where Y₃ is regressed on both Y₀ and X, in order to control for alcohol misuse at baseline (i.e., before retirement) (see Allison, 1990).

RESULTS

Descriptive Statistics

Means, standard deviations, and correlations among the variables are displayed in Table 1. Based on these results, it appears that the mean alcohol misuse (i.e., DPI) score remained stable from the assessment at baseline (i.e., before retirement) to the assessment at Time 3 (M=.56, SD=1.53; M=.57, SD=1.96, T₀ and T₃ respectively). Notable are the significant positive correlations between sleep-related problems at T₂ and the severity of alcohol misuse at T₃ ($r=.21$, $p<.001$). Moreover, the findings show a significant positive correlation between the severity of alcohol misuse at T₁ and the severity of alcohol misuse at T₃ ($r=.52$, $p<.001$).

Hypotheses Testing

To provide a baseline against which to compare the predictive utility of our more specified models, we first tested a control model capturing the effects of pre-retirement alcohol misuse (i.e., T₀). This “Control model” (Model 1; Table 2) shows a significant positive effect of alcohol misuse at T₀ on alcohol misuse at T₃ (estimate =.15, $p<.001$).

Tests of simple mediation—According to Baron and Kenny (1986), four conditions are necessary to establish mediation: (a) the independent and dependent variables must be significantly related; (b) the independent and mediating variables must be significantly related; (c) the mediator and dependent variable must be significantly related; and (d) the relationship between the independent variable and dependent variable should be non-significant or weaker when the mediator is added.

The regression results for testing mediation are reported in Table 2. First, the results in Column 2 of Table 2 show that, after taking the control variables into account, financial (but not marital) stressors at T_1 were positively related to alcohol misuse at T_3 (estimate = 1.49, $p < .001$), supporting Hypothesis 1a. Second, the results also indicate that beyond the effects of alcohol misuse at T_1 (estimate = .30, $p < .001$), financial and marital stressors at T_1 (estimate = 1.77, $p < .001$; estimate = .47, $p < .05$, respectively) were positively related to sleep-related problems at T_2 (column 3, Table 2). Finally, sleep-related problems at T_2 were positively but non-significantly (estimate = .06, ns) related to the severity of alcohol misuse at T_3 (column 4, Table 2), thus violating a key condition for mediation (Baron & Kenny, 1986) and suggesting that in contrast to Hypotheses 2a–2b, sleep-related problems at T_2 fail to mediate the impact of financial and marital stressors on the severity of alcohol misuse at T_3 . However, given that we also posited that both stages of the proposed mediation are moderated by gender (Hypotheses 3a and 3b), we next investigate the possibility of moderated mediation.

Tests of moderated mediation—The results of the tests for the moderated-mediation implied by Hypotheses 3a, 3b, and 4 are shown in Model 4 of Table 2. This model tests: (a) whether gender (W), as specified by Hypotheses 3a and 3b, amplifies the strength of the first stage of mediation [i.e., the relationship between stressors (X_1 and X_2), and sleep-related problems (M)], and (b) whether, as specified by Hypothesis 4, it moderates the second stage of mediation [i.e., the relationship between sleep-related problems (M) and the severity of alcohol misuse (Y)].

As is evident from the left-hand column of Model 4 (“Moderated-Mediation Model”), the impact of financial (but not marital) stressors (X_1 ; estimate = 5.97, $p < .001$) on sleep-related problems is significantly moderated by gender. As recommended by Aiken and West (1991), we used simple slopes analysis to assess the degree to which these effects were consistent with those hypothesized. As shown in Figure 2, while the slope of financial stressors at T_1 on sleep-related problems at T_2 was positive and significant for both men and women, consistent with Hypothesis 3a, it was amplified for females (estimate = 8.48 vs. 1.79 for women and men, respectively, $p < .001$ for both). Accordingly, with respect to the moderation of the first stage of the hypothesized mediation, our data suggest support only for Hypothesis 3a and not for 3b.

Full support for moderated mediation was found with respect to the second stage of mediation, i.e., the relationship between sleep-related problems (M) and alcohol misuse (Y). Consistent with Hypothesis 4, the results indicate that the interaction of sleep-related problems at T_2 (M) * gender (W) (estimate = $-.16$, $p < .01$) is significant in predicting the severity of alcohol misuse at T_3 (see the right-most column of Table 2). The results of the simple slopes analysis for this model (shown in Figure 3) indicate that – consistent with Hypothesis 4 -- while the slope of sleep-related problems at T_2 on alcohol misuse at T_3 for women was non-significant (estimate = .10, ns), for men, the slope was positive and significant (estimate = .19, $p < .001$).

To further assess the degree to which these sleep-mediated effects are moderated by gender, we tested the significance of the indirect effects on the basis of the bootstrapped confidence intervals. The bootstrap analysis indicated that while the indirect effects of both financial and marital stressors as specified in model 4 of Table 2 were significant for men (effects = .28 [90% CI = 0.01 to 0.73]) and .10 [90% C.I. = 0.01 to 0.32] respectively), they were not significant for women (the confidence interval included a zero)¹. We assessed the relative

¹We used 90% confidence intervals to correspond to one-tailed, .05 hypothesis tests, which according to Preacher, Zyphur and Zhang (2011:217) is justified in mediation research.

size of the indirect effect for male retirees by comparing it to the size of the total effect of financial and marital stressors as specified in Model 4. Respectively, these total effects were .59 and .15, suggesting that while the indirect effects via sleep problems explained approximately half of the effect of total financial stressors on male retirees' alcohol misuse, marital stressors explained approximately 2/3 of the effect of marital stressors on male retirees' alcohol misuse.

Further evidence in support of a significant moderated mediation effect is provided on the basis of: (a) the significant $\Delta-2$ Log Likelihood, and (b) a significant increase in R^2 . In terms of the former, $\Delta-2$ Log Likelihood (relative to Model 2) = 10.50 ($p < .05$) indicating that the inclusion of the interaction of sleep problems*gender significantly contributes to the model's explanatory potential. In terms of the latter, since the alcohol misuse models were assessed on the basis of ZIP regressions, effect sizes were estimated based on Cox and Snell's (1989) generalized R^2 . Although such likelihood-based R^2 estimates are typically lower than those obtained in linear regression, they still provide an indication of the relative explanatory strength (Hosmer & Lemeshow, 2000) of a model. Accordingly, our results indicate that the interaction model ("Model 4" of Table 2) ($R^2 = .32$) added .02 ($p < .05$) to the generalized R^2 of the total effect model ("Model 2" of Table 2), indicating a significant increase in explanatory strength.

Finally, in order to confirm the causal ordering assumed by our model, we tested the reversed specification, namely, that alcohol misuse severity affects financial and marital stressors via sleep-related problems. The results indicate that after controlling for the same variables included in the models presented above, alcohol misuse at T_1 was positively related to sleep-related problems at T_2 (estimate = .32, $p < .001$). However, it is important to note that the model tested above, by incorporating the effect of alcohol misuse severity at T_1 on sleep-related problems at T_2 , already took such a reciprocal relationship into account and demonstrated that financial stressors have an effect on subsequent sleep-related problems above and beyond the effects of alcohol misuse severity at T_1 . Moreover, alcohol misuse at T_1 was not significantly related to either financial or marital stressors at T_3 (estimate = .00 and .15, respectively, ns for both), and sleep-related problems at T_2 were not significantly related to the financial or marital stressors at T_3 (estimate = -.00 and .02, respectively, ns for both). Furthermore, neither of the two stages of the sleep-mediated, alcohol misuse $\rightarrow$ marital and financial stressors relationship was significantly moderated by gender. These findings suggest little if any support for the reverse causal pathway (i.e., that alcohol misuse, via sleep-related problems, generates marital or financial stress).

Taken as a whole, these results indicate substantial support for Hypotheses 1a and 3a, and support for Hypothesis 4, suggesting that (a) gender moderates the first stage of the sleep-mediated impact of financial stressors on the severity of retiree alcohol misuse such that the impact of these stressors on sleep-related problems is even more pronounced for females, and (b) gender moderates the second stage of the sleep-mediated impact of financial stressors on the severity of retiree alcohol misuse, such that the impact of sleep-related problems on alcohol misuse is amplified for men.

DISCUSSION

The current study aimed at addressing three main gaps in research on the alcohol-related consequences of retirement as a form of workforce withdrawal, namely: (1) How stressors in retirement (as opposed to retirement per se) may affect an important adjustment-related behavior (i.e., alcohol misuse), (2) how the impact of these stressors on the severity of retiree alcohol misuse may be secondary to the stressors themselves, operating instead via the sleep-related problems precipitated or exacerbated by such stressors, and (3) how gender

may serve as a moderator of these relationships. While findings from previous research regarding the link between workforce withdrawal and alcohol misuse are equivocal, our results indicate that -- consistent with both prior research on unemployment (Lin & Leung, 2010) and resource-based treatments of retirement (Wang & Shultz, 2010) ---- two key stressors (financial and marital) experienced in retirement are associated with the precipitation or exacerbation of alcohol misuse among male retirees. Moreover, our findings demonstrate that -- at least for male retirees -- the impact of such stressors experienced in retirement on alcohol misuse is, at least in part, secondary to the stressors themselves, mediated by the sleep-related problems such stressors may generate. That is, while we found no evidence that sleep-related problems mediate the impact of financial and marital stressors in retirement on female retirees' alcohol misuse, our findings suggest that male retirees may use alcohol as a mode of self-medication (Roth, 2005; Rumble & Morgan, 1992) in order to cope with a critical, *secondary* effect of such stressors, namely sleep-related problems. This finding suggests that while male retirees may, indeed, turn to alcohol to cope with financial and marital stressors, female retirees are unlikely to do so, perhaps relying on other coping behaviors (e.g., talking to friends or relatives, or addressing the problem source), viewing these other mechanisms as offering similar, if not greater, relief with fewer negative side effects.

The findings with respect to financial and marital stressors experienced in retirement are largely consistent with our hypotheses. We found evidence of a robust, albeit distal effect of both stressors on the change in alcohol misuse from pre-retirement levels. And while sleep-related problems were not found to mediate the effect of such stressors on alcohol misuse in general, a robust mediation effect *was* found when gender was taken into consideration. More specifically, the second stage of the mediation was found to be significant among male retirees only. Interestingly, although we also posited gender moderation in the first stage of the mediation (i.e., between both stressors and sleep-related problems), gender was found to amplify the impact only of financial stressors experienced in retirement on sleep-related problems. For marital stressors experienced in retirement, the effect on sleep-related problems was no greater for female retirees than male retirees. While the literature on sleep disorders among older adults suggests that stressors may be a more robust risk factor for women than men (Middelkoop et al., 1996), our findings suggest that marital stressors experienced in retirement may be of equal salience with respect to sleep-related problems for both female and male retirees, and that feeling content in one's marriage may be beneficial for both male and female retirees' sleep patterns.

Our findings are generally consistent with related findings in the geriatrics, sleep and alcohol literatures which suggest that stressors experienced by retirees may play a significant role in precipitating or exacerbating sleep-related problems (Middelkoop et al., 1996), and that alcohol-based self-medication is a common means by which particularly male retirees may attempt to cope with such problems (Roth, 2005). No less importantly, these findings are also consistent with a number of contemporary retirement adjustment perspectives (Wang & Shultz, 2010) and particularly those taking a more ecological approach (e.g., Wang, Henkens, and van Solinge, 2011). Consistent with such an approach, our findings suggest that the mental health consequences of disruptions in the maintenance of life patterns in retirement (experienced as financial or marital stressors) stem in no small part to the drain such disruptions may place on retirees' psychological resources and the potentially maladaptive ways in which some retirees attempt to address such resource outflows.

However, beyond merely providing additional evidence in support of such perspectives on retirement adjustment, our findings offer an important theoretical contribution by offering insight into the psycho-physiological mechanisms by which such stressors experienced in

retirement affect retirees' well-being. More specifically, consistent with COR theory (Hobfoll et al., 2003), they suggest that the drain on resources caused by such stressors experienced in retirement may manifest itself physiologically in the form of sleep-related problems, and that behaviors indicative of maladjustment such as alcohol misuse may increase in severity as individuals adopt maladaptive strategies for coping with such problems. We view this as an important step forward in understanding the relatively widespread misuse of alcohol among older adults noted in the introduction. Contrary to the tension reduction hypotheses and the idea that older adults use alcohol as a means by which to address the more dysphoric aspects of retirement (Perreira & Sloan, 2001), our results indicate that, at least for male retirees, any increase in the severity of alcohol misuse may be just as likely to be secondary to such a sense of dysphoria, having to do with the individual's attempt to self-medicate the sleep-related problems generated by such stressors in retirement.

Limitations

The findings of our study should be considered in light of its limitations, which may also offer important research opportunities. First, while our study was designed to track the change in alcohol misuse from before retirement, it was beyond the scope of our study to model the impact of change in stressors in retirement. Future research might apply growth modeling to examine how shifts in alcohol misuse at multiple points in time in retirement may map on to such stressor-based changes. Second, our rather strict inclusion criteria (i.e., full retirement by T₁; complete data along all study variables at all 3 points of data collection; not hospitalized during the past year), limited the size of our sample size to 292 individuals. While our sample bias checks suggest that these strict criteria did not limit the external validity of our findings, they may have limited statistical power. Studies aiming to replicate or extend our results should therefore consider increasing the size of the baseline sample. Finally, given the limitations imposed by relying on participants' subjective assessment of sleep-related problems, future research may benefit from the use of more physiological measures (e.g., electroencephalography).

Practical Implications

Despite these limitations, our findings suggest two important implications for policy and practice. First, they highlight the need to recognize financial and marital stressors in retirement as potential risk factors for male retiree well-being, and for organizations to assist workers nearing retirement eligibility to engage in the kind of retirement planning activities that might reduce such risks. Second, our findings underscore the key role potentially played by sleep-related problems as a mediator of the effects of financial stressors in retirement on male retirees' physical and emotional well-being. Accordingly, they suggest the need to go beyond simply helping older adults adjust to workforce disengagement so as to mitigate the stressors associated with it. Rather they suggest the need to increase the awareness of retirees' primary care physicians to the stressors often faced in workforce disengagement and their sleep-related implications so that these professionals might be able to offer more effective and less risky means by which retirees might address or manage sleep-related problems. They also suggest the value of extending employee assistance services to retirees as a means by which to facilitate early intervention and thereby reduce the risk that such problems in retirement may lead to the exacerbation of alcohol misuse and/or the precipitation of alcohol dependence.

Acknowledgments

This research was funded by a grant from the National Institute on Alcohol Abuse and Alcoholism # 5 R01 AA011976.

References

- Abrams DB, Wilson GT. Effects of alcohol on social anxiety in women: Cognitive versus physiological arousal. *Journal of Abnormal Psychology*. 1979; 88:161–173. [PubMed: 447899]
- Aiken, LS.; West, SG. Multiple regression: Testing and interpreting interactions. Newbury Park, CA: Sage Publications; 1991.
- Åkerstedt T, Knutsson A, Westerholm P, Theorell T, Alfredsson L, Kecklund G. Sleep disturbances, work stress and work hours: A cross-sectional study. *Journal of Psychosomatic Research*. 2002; 53(3):741–748. [PubMed: 12217447]
- Allison PD. Change scores as dependent variables in regression analysis. *Sociological Methodology*. 1990; 20:93–114.
- Ancoli-Israel S, Roth T. Characteristics of insomnia in the United States: Results of the 1991 national sleep foundation survey. I. Sleep. 1999; 22(Suppl 2):S347–S353. [PubMed: 10394606]
- Arata CM, Picou JS, Johnson GD, McNally TS. Coping with technological disaster: An application of the conservation of resources model to the Exxonvaldez oil spill. *Journal of Traumatic Stress*. 2000; 13(1):23–39. [PubMed: 10761172]
- Armon G. Do burnout and insomnia predict each other's levels of change over time independently of the job demand control–support (JDC–S) model? *Stress and Health*. 2009; 25:333–342.
- Atchley RC. A continuity theory of normal aging. *The Gerontologist*. 1989; 29(2):183–190. [PubMed: 2519525]
- Atchley, RC. Continuity and adaptation in aging: Creating positive experiences. 1st ed.. The Johns Hopkins University Press; 1999.
- Bacharach SB, Bamberger PA, Doveh E. Firefighters, critical incidents, and drinking to cope: The adequacy of unit-level performance resources as a source of vulnerability and protection. *Journal of Applied Psychology*. 2008; 93:155–169. [PubMed: 18211142]
- Bacharach S, Bamberger PA, Doveh E, Cohen A. Retirement, social support and drinking behavior: A cohort analysis of males with a baseline history of problem drinking. *Journal of Drug Issues*. 2007; 37:717–736. [PubMed: 18612374]
- Bacharach S, Bamberger PA, Sonnenstuhl WJ, Vashdi DR. Retirement and drug abuse: The conditioning role of age and retirement trajectory. *Addictive Behaviors*. 2008; 33(12):1610–1614. [PubMed: 18771858]
- Bamberger PA, Sonnenstuhl WJ, Vashdi D. Screening older, blue-collar workers for drinking problems: An assessment of the efficacy of the drinking problems index. *Journal of Occupational Health Psychology*. 2006; 11(1):119–134. [PubMed: 16551179]
- Barnes AJ, Moore AA, Xu H, Ang A, Tallen L, Mirkin M, Ettner S. Prevalence and correlates of at-risk drinking among older adults: the Project SHARE study. *Journal of General Internal Medicine*. 2010; 25:840–846. [PubMed: 20396975]
- Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: Conceptual, strategic, and statistical considerations. *Journal of Personality and Social Psychology*. 1986; 51(6):1173–1182. [PubMed: 3806354]
- Beehr TA. The process of retirement: A review and recommendations for future investigation. *Personnel Psychology*. 1986; 39:31–55.
- Benshoff JJ, Harrawood LK. Substance abuse and the elderly: Unique issues and concerns. *Journal of Rehabilitation*. 2003; 69:43–48.
- Bixler EO, Kales A, Soldatos CR, Kales JD, Healey S. Prevalence of sleep disorders in the Los Angeles metropolitan area. *American Journal of Psychiatry*. 1979; 136:1257–1262. [PubMed: 314756]
- Blatter K, Graw P, Münch M, Knoblauch V, Wirz-Justice A, Cajochen C. Gender and age differences in psychomotor vigilance performance under differential sleep pressure conditions. *Behavioural Brain Research*. 2006; 168:312–317. [PubMed: 16386807]
- Blow, F. Substance abuse among older adults. treatment improvement protocol (TIP) series no. 26. Rockville, MD: U.S.: Department of Health and Human Services, Public Health Service, Substance Abuse and Mental Health Services Administration, Center for Substance Abuse Treatment; 1998.

- Blow, FC.; Barry, KL.; Fuller, BE. Analysis of the national health and nutrition examination survey (NHANES): Longitudinal analysis of drinking over the life span. In: Korper, SP.; Council, CL., editors. Substance use by older adults: Estimates of future impact on treatment system. Rockville, MD: Substance Abuse and Mental Health Services Administration, Office of Applied Studies; 2002.
- Brennan PL, Schutte KK, Moos RH. Retired status and older adults' 10-year drinking trajectories. *Journal of Studies on Alcohol and Drugs*. 2010; 2010; 71(2):165–168. [PubMed: 20230712]
- Brenner M. Trends in alcohol consumption and associated illness: Some effects of economic changes. *American Journal of Public Health*. 1975; 65:1270–1291.
- Buysse DJ, Reynolds CF 3rd, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh sleep quality index: A new instrument for psychiatric practice and research. *Psychiatry Research*. 1989; 28(2):193–213. [PubMed: 2748771]
- Cheung YB. Zero-inflated models for regression analysis of count data: A study of growth and development. *Statistics in Medicine*. 2002; 21(10):1461–1469. [PubMed: 12185896]
- Conger JJ. Alcoholism: Theory, problem and challenge. II. Reinforcement theory and the dynamics of alcoholism. *Journal of Studies on Alcohol*. 1956; 13:296–305.
- Cooper ML, Russell M, Skinner JB, Frone MR, Mudar P. Stress and alcohol use: Moderating effects of gender, coping, and alcohol expectancies. *Journal of Abnormal Psychology*. 1992; 101(1):139–152. [PubMed: 1537960]
- Cox, DR.; Snell, EJ. Analysis of binary data. London: Chapman and Hall; 1989.
- Crum RM, Storr CL, Chan Y, et al. Sleep disturbance and risk for alcohol-related problems. *American Journal of Psychiatry*. 2004; 161(7):1197–1203. [PubMed: 15229051]
- Dar K. Alcohol use disorders in elderly people: fact or fiction? *Advances in Psychiatric Treatment*. 2006; 12:173–181.
- Davey A, Szinovacz ME. Dimensions of marital quality and retirement. *Journal of Family Issues*. 2004; 25:431–464.
- Dew J, Yorgason J. Economic pressure and marital conflict in retirement-aged couples. *Journal of Family Issues*. 2010; 31(2):164–188.
- Dohrenwend BP, Dohrenwend BS. Sex differences in. psychiatric disorders. *American Journal of Sociology*. 1976; 81:1447–1454.
- Dooley D, Prause J. Underemployment and alcohol misuse in the national Longitudinal Survey of Youth. *Journal of Studies on Alcohol*. 1998; 59:669–680. [PubMed: 9811088]
- Dooley D, Catalano R, Hough R. Unemployment and alcohol disorder in 1910 and 1990: Drift versus social causation. *Journal Occupational and Organizational Psychology*. 1992; 65:277–290.
- Edwards JR, Lambert LS. Methods for integrating moderation and mediation: A general analytical framework using moderated path analysis. *Psychological Methods*. 2007; 12(1):1–22. [PubMed: 17402809]
- Ekerdt DJ, De Labry LO, Glynn RJ, Davis RW. Change in drinking behaviors with retirement: Findings from the normative aging study. *Journal of Studies on Alcohol*. 1989; 50:347–353. [PubMed: 2787876]
- Feldman DC. The decision to retire early: A review and conceptualization. *The Academy of Management Review*. 1994; 19(2):285–311.
- Finney JW, Moos RH, Brennan PL. The drinking problems index: A measure to assess alcohol-related problems among older adults. *Journal of Substance Abuse*. 1991; 3(4):395–404. [PubMed: 1821293]
- Frone MR. Work stress and alcohol use. *Alcohol Research and Health*. 1999; 23:284–291.
- Frone, MR. Employee alcohol and illicit drug use: Scope, causes, and organizational consequences. In: Cooper, CL.; Barling, J., editors. *The Sage Handbook of Organizational Behavior* (Vol. 1: Micro approaches). Thousand Oaks, CA: Sage; 2008. p. 519–540.
- Greene, WN. *Econometric analysis*. Upper Saddle River, NJ: Pearson/Prentice Hall; 2008.
- Grossi G, Perski A, Evengard B, Blomkvist V, Orth-Gomer K. Physiological correlates of burnout among women. *Journal of Psychosomatic Research*. 2003; 55:309–316. [PubMed: 14507541]

- Hall M, Buysse DJ, Nofzinger EA, Reynolds CF 3rd, Thompson W, Mazumdar S, Monk TH. Financial strain is a significant correlate of sleep continuity disturbances in late-life. *Biological Psychology*. 2008; 77(2):217–222. [PubMed: 18055094]
- Hallgren, M.; Hogberg, P.; Andreasson, S. Alcohol Consumption among Elderly European Union Citizens. Stockholm: Swedish National Institute of Public Health; 2010. (Downloaded on Oct. 23, 2011 from <http://www.abuel.org/docs/25alcoholelderly.pdf>)
- Hammer T. Unemployment and use of drugs and alcohol among young people: a longitudinal study in the general population. *British Journal of Addiction*. 1992; 87:1571–1581. [PubMed: 1458036]
- Hernandez-Avila CA, Song C, Kuo L, Tennen H, Armeli S, Kranzler HR. Targeted versus daily naltrexone: Secondary analysis of effects on average daily drinking. *Alcoholism, Clinical and Experimental Research*. 2006; 30(5):860–865.
- Hobfoll SE. Social and psychological resources and adaptation. *Review of General Psychology*. 2002; 6:307–324.
- Hobfoll SE, Johnson RJ, Ennis N, Jackson AP. Resource loss, resource gain, and emotional outcomes among inner city women. *Journal of Personality and Social Psychology*. 2003; 84(3):632–643. [PubMed: 12635922]
- Hobfoll, SE. The ecology of stress. Washington, DC: Hemisphere; 1988.
- Hobfoll SE. Conservation of resources: A new attempt at conceptualizing stress. *American Psychologist*. 1989; 44(3):513–524. [PubMed: 2648906]
- Horwitz AV, White HR. Gender role orientations and styles of pathology among adolescents. *Journal of Health and Social Behavior*. 1987; 28:158–170. [PubMed: 3611702]
- Hosmer, DW.; Lemeshow, S. Applied logistic regression. New York: Wiley; 2000.
- Hublin C, Partinen M, Koskenvuo M, Kaprio J. Sleep and mortality: A population-based 22-year follow-up study. *Sleep*. 2007; 30(10):1245–1253. [PubMed: 17969458]
- Hull JG. A self-awareness model of the causes and effects of alcohol consumption. *Journal of Abnormal Psychology*. 1981; 90:586–600. [PubMed: 7320328]
- Ilfeld FW. Further validation of a psychiatric symptom index in a normal population. *Psychological Reports*. 1976; 39:1215–1228.
- Johnson JE. Insomnia, alcohol, and over-the-counter drug use in old-old urban women. *Journal of Community Health Nursing*. 1997; 14(3):181–188. [PubMed: 9282538]
- Kalimo, R.; Tenkanen, L.; Härmä, M.; Poppius, E.; Heinsalmi, P. Job stress and sleep disorders: Findings from the helsinki heart study. John Wiley & Sons, Ltd; 2000.
- Keith VM. Gender, financial strain, and psychological distress among older adults. *Research on Aging*. 1993; 15:123–147.
- Kessler RC, Turner JB, House JS. Effects of unemployment on health in a community survey: Main, modifying, and mediating effects. *Journal of Social Issues*. 1988; 44(4):69–85.
- Kim S, Feldman DC. Working in retirement: The antecedents of bridge employment and its consequences for quality of life in retirement. *Academy of Management Journal*. 2000; 43:1195–1210.
- Kushnir T, Melamed S. The Gulf War and its impact on burnout and well-being of working civilians. *Psychological Medicine*. 1992; 22:987–995. [PubMed: 1488493]
- Lahelma E, Kangas R, Manderbacka K. Drinking and unemployment: Contrasting patterns among men and women. *Drug and Alcohol Dependence*. 1995; 37:71–82. [PubMed: 7882876]
- Lambert D. Zero-inflated poisson regression, with an application to defects in manufacturing. *Technometrics*. 1992; 34(1):1–14.
- Landis CA, Moe KE. Sleep and menopause. *The Nursing Clinics of North America*. 2004; 39(1):97–115. [PubMed: 15062730]
- LeBlanc M, Merette C, Savard J, Ivers H, Baillargeon L, Morin CM. Incidence and risk factors of insomnia in a population-based sample. *Sleep*. 2009; 32(8):1027–1037. [PubMed: 19725254]
- Lee-Chiong, TL. Sleep: a comprehensive handbook. J. Wiley & Sons; 2006.
- Levin, SM.; Kruger, J. Substance abuse among older adults: A guide for social service providers. Rockville, MD: Substance abuse and Mental Health Services Administration; 2000.

- Li RH, Wing YK, Ho SC, Fong SY. Gender differences in insomnia—a study in the Hong Kong Chinese population. *Journal of Psychosomatic Research*. 2002; 53:601–609. [PubMed: 12127178]
- Lindberg E, Janson C, Gislason T, Bjornsson E, Hetta J, Boman G. Sleep disturbances in a young adult population: can gender differences be explained by differences in psychological status? *Sleep*. 1997; 20:381–387. [PubMed: 9302720]
- Lundh LG, Broman JE. Insomnia as an interaction between sleep-interfering and sleep-interpreting processes. *Journal of Psychosomatic Research*. 2000; 49:1–12. [PubMed: 11053596]
- Manber R, Armitage R. Sex, steroids, and sleep: a review. *Sleep*. 1999; 22(5):540–555. [PubMed: 10450590]
- McKee-Ryan FM, Song Z, Wanberg CR, Kinicki AJ. Psychological and physical well-being during unemployment: A meta-analytic study. *Journal of Applied Psychology*. 2005; 90(1):53–76. [PubMed: 15641890]
- Melamed S, Shirom A, Toker S, Berliner S, Shapira I. Burnout and risk of cardiovascular disease: evidence, possible causal paths, and promising research directions. *Psychological Bulletin*. 2006; 132(3):327–353. [PubMed: 16719565]
- Midanik LT, Soghikian K, Ransom LJ, Tekawa IS. The effect of retirement on mental health and health behaviors: The Kaiser Permanente Retirement Study. *Journals of Gerontology Series B: Psychological Sciences and Social Sciences*. 1995; 50:S59–S61.
- Middelkoop HA, Smilde-van den Doel DA, Neven AK, Kamphuisen HA, Springer CP. Subjective sleep characteristics of 1,485 males and females aged 50–93: Effects of sex and age factors related to self-evaluated quality of sleep. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*. 1996; 51(3):M108–M115.
- Moen P, Kim JE, Hofmeister H. Couples' Work/Retirement transitions, gender, and marital quality. *Social Psychology Quarterly*. 2001; 64(1):55–71.
- Moos, R.; Moos, B. *Life Stressors and Social Resources Inventory: Adult Form manual*. Odessa, FL: Psychological Assessment Resources; 1994.
- Muys, M. Characteristics of life in exile: vulnerability factors for substance use. In: Fountain, J.; Korf, DJ., editors. *Drugs in Society: European Perspectives*. Oxford: U.K. Radcliffe Publishing; 2007. p. 62-75.
- National Institute on Alcohol Abuse and Alcoholism. *Drinking in the United States: Main Findings From the 1992 National Longitudinal Alcohol Epidemiologic Survey (NLAES)*. U.S. Alcohol Epidemiologic Data Reference Manual. 1st ed.. Vol. 6. Bethesda, MD: The Institute; 1998.
- National Institute on Alcohol Abuse and Alcoholism. Unpublished data from the 2001–2002 National Epidemiologic Survey on Alcohol and Related Conditions (NESARC), a nationwide survey of 43,093 U.S. adults aged 18 or older. 2004
- National Institute on Alcohol Abuse and Alcoholism. Report to the Extramural Advisory Board, August 16–17, 2006, Division of Epidemiology and Prevention Research Strategic Planning Document, Bethesda, MD: National Institute on Alcohol Abuse and Alcoholism. 2006
- Nicolson NA, van Diest R. Salivary cortisol patterns in vital exhaustion. *Journal of Psychosomatic Research*. 2000; 49:335–342. [PubMed: 11164057]
- Paulhus, DL. Measurement and control of response bias. In: Robinson, JP.; Shaver, PR.; Wrightsman, LS., editors. *Measures of personality and social psychological attitudes*. New York: Academic Press; 1991. p. 17-59.
- Pedersen SL, McCarthy DM. Person-environment transactions in youth drinking and driving. *Psychology of Addictive Behaviors: Journal of the Society of Psychologists in Addictive Behaviors*. 2008; 22(3):340–348. [PubMed: 18778127]
- Perreira KM, Sloan FA. Life events and alcohol consumption among mature adults: A longitudinal analysis. *Journal of Studies on Alcohol*. 2001; 62(4):501–508. [PubMed: 11513228]
- Preacher KJ, Zyphur MJ, Zhang Z. A general multilevel SEM framework for assessing multilevel mediation. *Psychological Methods*. 2011; 15:209–233. [PubMed: 20822249]
- Preacher KJ, Rucker DD, Hayes AF. Addressing moderated mediation hypotheses: Theory, methods and prescriptions. *Multivariate Behavioral Research*. 2007; 42:185–227.

- Price RH, Choi JN, Vinokur AD. Links in the chain of adversity following job loss: How financial strain and loss of personal control lead to depression, impaired functioning, and poor health. *Journal of Occupational Health Psychology*. 2002; 7(4):302–312. [PubMed: 12396064]
- Quan SF, Katz R, Olson J, Bonekat W, Enright PL, Young T, Newman A. Factors associated with incidence and persistence of symptoms of disturbed sleep in an elderly cohort: The cardiovascular health study. *The American Journal of the Medical Sciences*. 2005; 329(4):163–172. [PubMed: 15832098]
- Richardson V, Kilty KM. Adjustment to retirement: Continuity vs. discontinuity. *International Journal of Aging & Human Development*. 1991; 33(2):151–169. [PubMed: 1955209]
- Rogosa D. A critique of cross-lagged correlation. *Psychological Bulletin*. 1980; 88(2):245–258.
- Rohsenow DJ. Drinking habits and expectancies about alcohol's effects for self versus others. *Journal of Consulting and Clinical Psychology*. 1983; 51:752–756. [PubMed: 6630690]
- Rosenkoetter MM, Garris JM, Engdahl RA. Postretirement use of time: Implications for preretirement planning and postretirement management. *Activities, Adaptation & Aging*. 2001; 25(3–4):1–18.
- Roth T. Prevalence, associated risks, and treatment patterns of insomnia. *The Journal of Clinical Psychiatry*. 2005; 66(Suppl 9):10–13. quiz 42–3. [PubMed: 16336036]
- Roth T, Roehrs T. Insomnia: Epidemiology, characteristics, and consequences. *Clinical Cornerstone*. 2003; 5(3):5–15. [PubMed: 14626537]
- Rumble R, Morgan K. Hypnotics, sleep, and mortality in elderly people. *Journal of the American Geriatrics Society*. 1992; 40(8):787–791. [PubMed: 1634722]
- Schonfeld L, Dupree LW, Rohrer GE. Age-specific differences between younger and older alcohol abusers. *Journal of Clinical Geropsychology*. 1995; 1:219–227.
- Schubert CR, Cruickshanks KJ, Dalton DS, Klein BE, Klein R, Nondahl DM. Prevalence of sleep problems and quality of life in an older population. *Sleep*. 2002; 25(8):889–893. [PubMed: 12489896]
- Shultz, JH. *The economics of aging*. Belmont, CA: Wadsworth; 1985.
- Shultz KS, Morton KR, Weckerle JR. The influence of push and pull factors on voluntary and involuntary early retirees' retirement decision and adjustment. *Journal of Vocational Behavior*. 1998; 53:45–57.
- Söderström M, Ekstedt M, Akerstedt T, Nilsson J, Axelsson BA. Sleep and sleepiness in young individuals with high burnout scores. *Sleep*. 2004; 27:1369–1377. [PubMed: 15586790]
- Spanier GB. Measuring dyadic adjustment: New scales for assessing the quality of marriage and similar dyads. *Journal of Marriage and Family*. 1976; 38(1):15–28.
- Steele CM, Josephs RA. Drinking your troubles away II: An attention-allocation model of alcohol's effect on psychological stress. *Journal of Abnormal Psychology*. 1988; 97:196–205. [PubMed: 3385073]
- Su, SS.; Larison, C.; Ghadially, R., et al. *Substance Use Among Women in the United States. SAMHSA Analytic Series A-3*. Rockville, MD: Substance Abuse and Mental Health Services Administration; 1997.
- Sulander, T. *Functional ability and health behaviours trends and associations among elderly people, 1985–2003*. Doctoral Dissertation. Department of Epidemiology and Health Promotion & #x00026; Department of Social Policy, National Public Health Institute & University of Helsinki; 2005.
- Sutker PB, Allain AN, Brantley P, Randall C. Acute alcohol intoxication, negative affect, and autonomic arousal in women and men. *Addictive Behaviors*. 1982; 7:17–25. [PubMed: 7200716]
- Szinovacz M. Couples Employment/Retirement patterns and perceptions of marital quality. *Research on Aging*. 1996; 18(2):243–268.
- Szinovacz, ME. Contexts and pathways: Retirement as institution, process, and experience. In: Adams &, GA.; Adams Beehr, TA., editors. *Retirement: Reasons, processes, and results*. New York: Springer; 2003. p. 6–52.
- Tu, W. Zero-inflated data. In: El Shaarawi, AH.; Piegorsch, WW., editors. *Encyclopedia of Environmetrics*. Chichester: John Wiley & Sons; 2006.

- vanDiest R. Subjective sleep characteristics as coronary risk factors, their association with Type A behavior and vital exhaustion. *Journal of Psychosomatic Research*. 1990; 34:415–426. [PubMed: 2376842]
- Vela-Bueno A, Moreno-Jiménez B, Rodríguez-Muñoz A, Olavarrieta-Bernardino S, Fernández-Mendoza J, José De la Cruz-Troca J, Bixler EO, Vgontzas AN. Insomnia and sleep quality among primary care physicians with low and high burnout levels. *Journal of Psychosomatic Research*. 2008; 64(4):435–442. [PubMed: 18374744]
- Wang M, Shultz KS. Employee retirement: A review and recommendations for future investigation. *Journal of Management*. 2010; 36:172–206.
- Wang M. Profiling retirees in the retirement transition and adjustment process: Examining the longitudinal change patterns of retirees' psychological well-being. *Journal of Applied Psychology*. 2007; 92:455–474. [PubMed: 17371091]
- Wang M, Zhan Y, Liu S, Shultz K. Antecedents of bridge employment: A longitudinal investigation. *Journal of Applied Psychology*. 2008; 93:818–830. [PubMed: 18642986]
- Weeks EC, Drengacz S. The Non-economic Impact of Community Economic Shock. *Journal of Health and Human Resources Administration*. 1982; 4:303–318.
- Weissman MM, Greenwald S, Niño-Murcia G, Dement WC. The morbidity of insomnia uncomplicated by psychiatric disorders. *General Hospital Psychiatry*. 1997; 19:245–250. [PubMed: 9327253]
- Zorn, CJW. Evaluating zero-inflated and hurdle poisson specifications. *Midwest Political Science Association*; 1996.

Figure 1.
Conceptual Model

Figure 2.
Moderation of the Effect of (centered) Financial Stressors in Retirement at T_1 (X) on Sleep-Related Problems T_2 (M)

Figure 3.
Moderation of the Effect of (centered) Sleep-Related Problems T_2 (M) on Severity of Alcohol Misuse at T_3 (Y).

Table 1

Means, Standard Deviations and Intercorrelations of the Variables (n=292)

	Mean	SD	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)
(1) Alcohol Misuse at T ₃	.57	1.96													
(2) Alcohol Misuse at pre-retirement (i.e., T ₀)	.56	1.53	.52 ^c												
(3) Alcohol Misuse at T ₁	.62	1.74	.65 ^c	.67 ^c											
(4) Health related symptoms at T ₁	.19	.19	.06	.20 ^c	.11										
(5) Age at T ₁	67.21	4.36	-.14 ^c	-.10	-.10	-.03									
(6) Minority (0=No, 1= Yes)	.20	.40	.09	-.03	.03	-.11	-.07								
(7) Education	2.86	1.14	.04	-.02	.03	.000	-.33 ^c	-.12 ^a							
(8) Marital Status (0=other, 1=married)	.94	.24	.002	-.02	-.02	-.04	.06	-.05	-.12 ^a						
(9) Years since retirement at T ₃	5.87	1.09	.01	.01	-.04	-.03	.33 ^c	-.18 ^b	-.26	.10					
(10) Social Desirability at T ₁	54.31	7.50	-.09	-.08	-.10	-.27 ^c	.04	.04	-.05	.03	.12 ^a				
(11) Financial Stressors at T ₁	1.11	.29	.07	.04	.004	.22 ^c	.01	-.06	-.05	-.01	.04	-.12 ^a			
(12) Marital Stressors at T ₁	.97	.70	.11	.16 ^b	.19 ^a	.21 ^c	-.07	-.09	.03	.02	-.08	-.21 ^c	.13 ^a		
(13) Sleep-Related Problems at T ₂	5.94	2.58	.21 ^c	.27 ^c	.26 ^a	.26 ^c	-.13	-.002	.04	-.04	-.05	-.11	.25 ^c	.23 ^c	
(14) Gender (0=male, 1= female)	.24	.43	.03	.04	.11	.13 ^a	-.43 ^c	-.02	.53 ^c	-.09	-.36 ^c	-.08	-.07	.08	.12 ^a

^aP 0.05;^bP 0.01;^cP 0.001

Results of Moderated Mediation Analysis (n=292)

Dep. Variable	Control Model (Model 1)		Total Effect Model (Model 2)		Mediation Model (Model 3)				Moderated Mediation Model (Model 4)			
	Alcohol Misuse at T ₃		Alcohol Misuse at T ₃ X->Y		Sleep-Related Problems X->M		Alcohol Misuse at T ₃ M->Y		Sleep-Related Problems X->M		Alcohol Misuse at T ₃ M->Y	
Indep. Variable	Est.	SE	Est.	SE	Est.	SE	Est.	SE	Est.	SE	Est.	SE
Intercept	4.06 ^b	1.54	1.13	1.75	6.68 ^a	2.85	1.20	1.78	7.19 ^b	2.84	3.80 ^a	1.97
Heath related symptoms at T ₁	-.01	.49	.03	.50	2.39 ^b	.80	-.01	.50	2.32 ^b	.79	.02	.52
Age at T ₁	-.05 ^b	.02	-.01	.02	-.06	.04	-.01	.02	-.03	.04	-.02	.02
Minority (0= Caucasians, 1= other)	1.02 ^c	.22	.56 ^a	.27	.17	.36	.63 ^a	.27	.25	.36	.30	.29
Education	.11	.09	.04	.09	.01	.13	.06	.09	-.02	.15	.17	.11
Marital Status (0=other, 1=married)	-.47	.39	-.05	.45	-.31	.58	.00	.45	-.06	.58	.22	.44
Years since retirement	.16 ^a	.08	.05	.09	.01	.14	.08	.09	.02	.14	-.06	.10
Social Desirability at T ₁	-.03 ^b	.01	-.04 ^c	.01	.00	.02	-.05 ^c	.01	.004	.02	-.05 ^c	.01
Alcohol Misuse at pre-retirement (i.e., T ₀)	.15 ^c	.03	.17 ^c	.03	---	---	.14 ^c	.03	---	---	.14 ^c	.03
Alcohol Misuse at T ₁	---	---	---	---	.30 ^c	.08	---	---	.26 ^c	.08	---	---
Financial Stressors at T ₁ (X ₁)			1.49 ^c	.37	1.77 ^c	.49	1.08 ^a	.48	1.36 ^b	.51	.31	.54
Marital Stressors at T ₁ (X ₂)			.04	.16	.47 ^a	.21	.01	.16	.48 ^a	.23	.05	.15
Gender (0= male, 1= female) (W)									.60	.42	-.32	.28
Financial Stressors * Gender (X ₁ *W)									5.97 ^c	1.83		
Marital Stressors * Gender (X ₂ *W)									.09	.50		
Sleep-Related Problems at T ₂ (M)							.06	.04			.21 ^c	.06
Sleep Problems * Gender (M*W)											-.16 ^b	.07
R-Square	.26		.30		.19		.30		.22		.32	
Δ R-Square			.04 ^a		---		0		---		.02 ^a	
			(rel. to Model 1)				(rel. to Model 2)				(rel. to Model 2)	
2 log likelihood	499.97		484.62		---		482.57		---		474.12	

Dep. Variable	Control Model (Model 1)		Total Effect Model (Model 2)		Mediation Model (Model 3)		Moderated Mediation Model (Model 4)	
	Alcohol Misuse at T ₃		Alcohol Misuse at T ₃ X->Y		Sleep-Related Problems X->M		Alcohol Misuse at T ₃ M->Y	
	Est.	SE	Est.	SE	Est.	SE	Est.	SE
$\Delta -2 \log \text{likelihood}$			15.35 ^c (rel. to Model 1)	---	2.05 (rel. to Model 2)	---	10.50 ^d (rel. to Model 2)	

Note: Except for Model 3 and Model 4 “Sleep-Related Problem X->M” (linear regression), all analyses were run using ZIP regression.

^aP 0.05;

^bP 0.01;

^cP 0.001