Published in final edited form as:

Res Social Adm Pharm. 2014 ; 10(2): 378–386. doi:10.1016/j.sapharm.2013.07.001.

Relationships between health literacy and heart failure knowledge, self-efficacy, and self-care adherence

Aleda M. H. Chen, Pharm.D., Ph.D. [Assistant Professor of Pharmacy Practice],
Cedarville University School of Pharmacy, Cedarville, Ohio

Karen S. Yehle, Ph.D., M.S., R.N., F.A.H.A. [Associate Professor of Nursing],
Purdue University School of Nursing & Center on Aging and the Life Course, West Lafayette, Indiana

Nancy M. Albert, Ph.D., CCNS, CCRN, NE-BC, FAHA, FCCM [Senior Director],
Cleveland Clinic Nursing Research & Innovation and CNS, Cleveland, Ohio

Kenneth F. Ferraro, Ph.D. [Distinguished Professor],
Purdue University Department of Sociology & Center on Aging and the Life Course, West Lafayette, Indiana

Holly L. Mason, Ph.D. [Associate Dean & Professor],
Purdue University College of Pharmacy, West Lafayette, Indiana

Matthew M. Murawski, Ph.D., R.Ph. [Associate Professor], and
Purdue University College of Pharmacy, West Lafayette, Indiana

Kimberly S. Plake, Ph.D. [Associate Professor of Pharmacy Practice]
Purdue University College of Pharmacy & Center on Aging and the Life Course, West Lafayette, Indiana

Abstract

Background—It has been argued that only 12 percent of adults have the necessary health literacy to manage their health care effectively, which can lead to difficulties in self-care activities, such as medication adherence. Prior research suggests that health literacy may influence knowledge, self-efficacy and self-care, but this has not been fully examined.

Objective—To test a model to explain the relationships between health literacy, heart failure knowledge, self-efficacy, and self-care.

Methods—Prior to receiving clinic-based education, newly referred patients to 3 heart failure clinics completed assessments of health literacy, heart failure knowledge, self-efficacy, self-care, and demographics. Structural equation modeling was completed to examine the strength of the inter-variable relationships.

© 2013 Elsevier Inc. All rights reserved.

Corresponding Author: Aleda M. H. Chen, Cedarville University, 251 N. Main St., Cedarville, OH 45314, Tel: 937-766-7454, Fax: 937-766-7410, amchen@cedarville.edu.

Portions of this project were completed while Aleda Chen was a graduate research assistant at Purdue, University College of Pharmacy, West Lafayette, Indiana

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Results—Of 81 participants recruited, data from 63 patients were complete. Health literacy was independently associated with knowledge ($p<0.001$). Health literacy was not related to self-care. Self-efficacy was independently-associated with self-care adherence ($p=0.016$). No other relationships were statistically significant. The model had good fit (comparative fit index=1.000) and explained 33.6% of the variance in knowledge and 27.6% in self-care.

Conclusions—Health literacy influences knowledge about heart failure but not self-care adherence. Instead, self-efficacy influenced self-care adherence. Future research should incorporate additional factors that may better model the relationships between health literacy, knowledge, self-efficacy, and self-care.

Keywords

heart failure; health literacy; self-care; self-efficacy; heart failure knowledge

INTRODUCTION

Nearly 6 million Americans have heart failure, a chronic, progressive condition that accounts for significant morbidity and mortality.¹ Heart failure incidence is predicted to increase by 25 percent in the next two decades and may lead to a dramatic increase in healthcare costs.¹ Costly hospitalizations and heart failure exacerbations can be reduced with self-care adherence.^{2,3} According to Orem's Theory of Self-Care, self-care is a regulatory function, whereby individuals are given and assume the functions and responsibility of care for themselves, and when individuals are not willing or able to perform these functions, there are self-care deficits.⁴ Patients engage in self-care maintenance—tasks to prevent symptoms, such as adhering to sodium restrictions—and self-care management—activities to respond to symptoms—to prevent these deficits, i.e., improve or maintain their functioning.^{2,3} The bulk of the empirical evidence, however, indicates that most patients do not adhere well to self-care recommendations, such as adhering to their medications and reducing sodium intake.² Addressing potential barriers to self-care behavior may help patients achieve better outcomes.⁵

Patients with heart failure typically gain disease-specific knowledge and then apply the knowledge to specific heart-failure situations,⁶ as successful self-care utilizes both the skill and knowledge of individuals.^{4,6} Many individuals with heart failure lack knowledge regarding their self-care, such as behaviors that maintain stability, what symptoms require monitoring, and what to do when symptoms occur.⁶ Patient challenges increase when there are barriers to gaining knowledge, such as low health literacy (difficulty understanding health information⁷), which is associated with less disease knowledge.^{8–13} Lack of disease-specific knowledge also may affect confidence, or self-efficacy, regarding the ability to adhere to complex self-care regimens. Self-care confidence is derived from the concept of self-efficacy from Bandura's Social Cognitive Theory.^{2,14} The level of self-efficacy an individual possesses influences adherence to goals and responses to challenges. If individuals are not confident in their decisions, appropriate self-care may not occur.^{14,15} The role of health literacy in this process is unclear.^{16,17}

The objective of this study was to test a model examining the relationships between years of formal education, health literacy, heart failure knowledge, self-efficacy, and self-care.

METHODS

This study used a cross-sectional, correlational design and survey methods. Institutional Review Board (IRB) approval was obtained from Purdue University as well as each heart failure clinic.

Model Development

To explain the proposed relationships between health literacy, heart failure knowledge, self-efficacy, and self-care, a model was proposed based on Orem's Theory of Self-Care, Bandura's Social Cognitive Theory, and a review of the literature (Figure 1, Model 1).^{4,14} The goal of developing this model was to better characterize patients who presented for initial appointments and received individual education about heart failure and self-care in outpatient clinics.

The amount of formal education individuals have completed affects literacy, and general literacy levels are the foundation for and are associated with health literacy.^{7,18,19} Patient educational attainment, i.e., amount of formal education, is associated with health literacy.^{7,17–19} Health literacy, in turn, may impact patients' self-care decision-making, ability to gain knowledge regarding their condition during traditional clinic-based education, and their confidence in making self-care decisions. If patients have not gained enough knowledge, they may be unable to perform or adhere to self-care activities. Additionally, lack of knowledge may undermine patient self-efficacy, and without sufficient self-efficacy, individuals may be less likely to change or start a new health behavior.¹⁴

The hypothesized model is displayed in Figure 1 as Model 1, but three alternative specifications derived out of the literature, Models 2–4 in Figure 1, also were tested. The hypothesized model specified that (1) formal education would be associated with health literacy and directly effect heart failure knowledge; (2) health literacy would directly affect heart failure knowledge and self-efficacy; (3) health literacy would indirectly affect self-efficacy through heart failure knowledge; and (4) health literacy would indirectly affect self-care through heart failure knowledge and self-efficacy.

Alternative specifications were derived out of the literature suggesting health literacy may not be directly associated with self-care.¹⁷ There also was some question as to whether years of formal education (i.e., educational attainment) was directly related to heart failure knowledge or influenced knowledge through health literacy, which led to alternative pathways.

Participants and Procedures

Participants were recruited by researchers or clinic nurses who did not provide direct patient care from 2009 to 2011 at 3 heart failure clinics: Cleveland Clinic heart failure clinic in the Heart and Vascular Institute (Cleveland, OH), Indiana University Health-Bloomington Hospital HEARTTEAM Cardiopulmonary Rehabilitation and Congestive Heart Failure Center (Bloomington, IN), and Community Health Network Indiana Heart Hospital Healthy Hearts Center (Indianapolis, IN). Patients were invited to participate if they were a new referral to the heart failure clinic, at least 18 years of age, could read and speak English, and had no cognitive impairment (as deemed by clinical judgment). Patients were excluded if they resided in a skilled nursing facility or received home healthcare services. After consent was obtained, study instruments were completed in written format by participants prior to receiving traditional clinic-based education at the time of their first office appointment. Study researchers or clinic nurses who did not provide direct patient care administered the instruments and timed the completion of the health literacy instrument.

Measures

Health literacy was measured using the Short-Form Test of Functional Health Literacy (S-TOFHLA), a valid and reliable measure with scores ranging from 0–36.²⁰ The S-TOFHLA contains 36 reading comprehension items, based on examples of commonly-used materials in the healthcare system, and must be completed within a 7-minute time-frame. There are

three scoring ranges: inadequate (0–16 points), marginal (17–22 points), and adequate (23–36 points). The S-TOFHLA is a reliable and valid measure of health literacy: Cronbach's alpha is 0.98, suggesting a strong internal consistency across measures, while correlation with other established measures of health literacy (Test of Functional Health Literacy in Adults (TOFHLA, $r=0.91$) and the Rapid Estimation of Adult Literacy in Medicine (REALM, $r=0.80$)) suggests the S-TOFHLA's criterion validity was adequate.²⁰

The Heart Failure Knowledge Questionnaire (HFKQ) was used to measure patients' knowledge of heart failure related to pathology, symptoms, medications, and self-management, a reliable measure with scores from 0–15.⁶ The HFKQ consists of 14 close-ended items and 1 open-ended answer. No cutoffs were established to measure adequate knowledge, but scores range from 0 (lack of knowledge) to 15 (knowledgeable). Reliability of the HFKQ was established in recently-discharged patients with heart failure (Cronbach's alpha of 0.62).⁶

The Self-Care Heart Failure Index v.6 (SCHFI), a valid and reliable 22-item instrument, was used to evaluate patient's self-care maintenance and management adherence as well as self-efficacy in performing self-care through 3 subscales.^{3,21} Each item is rated on a four-point response scale by the participant. There are three subscales: maintenance, management, and confidence (self-efficacy). Scores on each subscale are standardized to 100 points, and scores can range from 0–100. To score Subscale B (self-care management), patients must have experienced an exacerbation of heart failure within the prior 3 months. The instrument authors recommend that a score of 70 can be used as the threshold for adequate self-care adherence on individual subscales.^{3,21} The SCHFI appears to have a high degree of internal consistency reliability (maintenance: $\alpha=0.553$, management: $\alpha=0.597$)^{3,21}; additional testing (confidence/self-efficacy: $\alpha=0.827$, combined maintenance/management: $\alpha=0.798$).²²

Demographic information also was obtained. The following patient demographics were obtained: gender, age, marital status, co-habitation, presence of someone in whom to confide, quality of support, ethnicity/race, years of education, highest educational degree obtained, employment status, income, smoking history, alcohol use, exercise recommendation, time spent exercising, height, weight, insurance, place of residence (i.e., at home, retirement community, assisted living, or other), and number of prescription medications.

Data Analysis

Statistical analyses were conducted using SAS v. 9.2 (SAS Institute, Inc., Cary, North Carolina) with an *a priori* level of 0.05 for statistical significance. Descriptive statistics were performed. A power analysis was performed to determine the sample size needed to achieve a power of 0.8 with an alpha of 0.05, a sample size of at least 57 participants was needed for correlational analyses. Pearson correlations were used to measure associations between educational attainment, health literacy, knowledge, self-efficacy, and self-care.

In performing structural equation modeling (SEM), there are many methods for calculating appropriate sample size. Some suggest that 5–20 observations per parameter estimated or at least 200 observations (whichever is greater) are desirable.²³ Not all studies, particularly where there is no incentive for participation can achieve a sample size of 200. If larger sample sizes are not obtainable, some researchers have suggested that 4 observations per parameter provide stable estimates. It also is recommended that models be simplified as much as possible and use reliable measures.²⁴ With 11 parameters (i.e., paths) in the most complex model and 5 observations per parameter, a minimum of 55 participants with complete data were needed.

A total of five variables and their relationships were tested: years of formal education (as measured by the demographic questionnaire), health literacy level (S-TOFHLA scores), knowledge (HFKQ scores), self-efficacy (SCHFI confidence subscale), and self-care. Since self-care is a process where patients perform behaviors that maintain stability (maintenance) and respond to symptoms (self-care management),²¹ self-care maintenance and self-care management were combined into a latent variable (self-care), which reduced model complexity. Participant S-TOFHLA sum scores were used, rather than category, to reduce the complexity of the structural equation model. Other researchers have utilized the S-TOFHLA as a continuous variable rather than a categorical variable in association and regression analyses and structural equation modeling to understand relationships between variables.^{25–28}

Model fit was assessed using maximum likelihood estimation, with conservative cut-offs for several fit statistics, including accountability for smaller sample size: a Chi-square statistic with a p-value greater than 0.05 (indicates observed covariance matrix is similar to model-predicted covariance), a Root Mean Square Error of Approximation (RMSEA) less than 0.05, a Goodness of Fit Index (GFI) greater than 0.95, a Normed Fit Index (NFI) greater than 0.95, and a Comparative Fit Index (CFI) greater than 0.95.^{23,29–31} If more than one model met all of these criteria, then the most parsimonious model was chosen as the best-fitting model.²³

RESULTS

Participant Characteristics and Associations

A total of 81 participants provided baseline data; however, after removing participants with incomplete data, the analyses were limited to 63 participants (see Table 1). Patients were removed for the following reasons: (1) patient did not have an exacerbation of heart failure within the past three months and, therefore, were ineligible to complete the section of the SCHFI regarding self-care management or (2) patient did not complete an item. Compared to the 81 participants who enrolled in the study, the 63 participants used for analyses were not significantly different ($p>0.05$, data not shown). Participants, on average, were older, white, achieved at least a high school education, and were prescribed 10 prescription medications on a regular basis. Most participants had adequate health literacy (scores ≥ 23 on the S-TOHFLA) but were not adherent in self-care (score < 70). Self-efficacy among participants also was not adequate, and participants answered less than 55% of heart failure knowledge questions correctly (see Table 3). Health literacy was positively associated with years of formal education ($p=0.001$) and heart failure knowledge ($p<0.001$). Years of formal education were positively associated with knowledge ($p=0.001$). Self-efficacy, self-care maintenance, and self-care management were not associated with health literacy, years of formal education, and heart failure knowledge ($p>0.05$; see Table 3).

Structure Equation Model Comparisons

Examining the criteria for model fit revealed that Model 1 had the best fit (see Table 4). All four models met criteria for good fit, but only Models 1 and 2 met all of the pre-specified fit criteria. Model 1 was chosen over Model 2 as it was more parsimonious. The highest percentage of the variance in knowledge (33.6%) and self-care (27.6%) were explained by Model 1. No model explained much of the variance in self-efficacy (see Figure 1). There was an independent effect of health literacy on knowledge. Health literacy was neither directly nor indirectly related to self-efficacy or self-care. Self-efficacy independently affected self-care. Knowledge was not directly related to self-efficacy.

DISCUSSION

In this study, the importance of health literacy on patients' understanding of basic knowledge about heart failure was revealed and underscores the importance of educational efforts in the clinical setting. There were independent effects for health literacy on knowledge and for self-efficacy on self-care but no indirect effects for health literacy on self-care or self-efficacy as hypothesized. Although there was a significant bivariate relationship between years of formal education and heart failure knowledge, in the structural equation model, health literacy was the primary influence on knowledge about heart failure. The implication is that patients with low health literacy may not understand the value of heart failure self-care behaviors. Further, patients also may believe the information they already have about heart failure self-care adherence is accurate, even when it may not be formed from evidence-based scientific information. Thus, actions taken also may not be based on current evidence.

Consistent with other studies, positive associations were found between health literacy and patient knowledge in heart failure,^{16,32} and this relationship also has been observed for other diseases and chronic conditions.^{8–10,12,13} Although some investigators (with similar sample sizes) have found an association between health literacy and self-efficacy^{16,17} and between health literacy and self-care in cross-sectional studies,¹⁷ there were no associations between these variables in this study. Experience with performing self-care and managing symptoms may improve self-efficacy over longer periods of time, as other investigators primarily examined patients who were not newly-diagnosed.^{16,17} When patients experience success in performing self-care, their self-efficacy may improve by seeing their actions produce positive results. The continuous cycle of self-efficacy and self-care may explain why there were no statistically significant associations between health literacy, self-care, and self-efficacy in this sample.

A model in which health literacy was assumed to have direct effects on knowledge, indirect effects on self-efficacy through knowledge, and indirect effects on self-care through knowledge and self-efficacy was found to be a good fit for the data. Macabasco and colleagues evaluated the relationship between these same factors and health-related quality of life and, similarly, found that health literacy had a direct effect on knowledge and self-efficacy had a direct effect on self-care. However, researchers also found the effect of health literacy was mediated by knowledge and self-efficacy, in contrast to this study.³² There are potential reasons for differences in findings between studies: use of different measures and patient recruitment. Despite differences, both studies revealed the critical role of adequate health literacy in heart failure knowledge. Furthermore, the results of these studies emphasized the importance of patient self-efficacy on performance of self-care.

Since this model explained 33.6% percent of the variance in knowledge and 27.6% of the variance in self-care, it is likely that there were other important factors that would explain relationships between knowledge, self-efficacy, and self-care. Motivation to perform self-care or values patients have for specific self-care behaviors may be essential components that were not included in this model. Patients must value and be motivated and willing to change behaviors, as changes can be challenging to incorporate into daily life.² Future research should include patient factors not studied here or in other research to improve the model of health literacy and self-care in heart failure.

Limitations

Findings may be limited due to higher health literacy of this sample. Sites for this project were chosen in an attempt to obtain more diversity in health literacy levels, and while 20 participants (31.7%) with inadequate or marginal health literacy were recruited, there were

more participants with adequate health literacy than marginal or inadequate health literacy. Since the estimates of low health literacy among patients with heart failure are between 17.5–41%,^{18,19,33,34} the distribution of health literacy in this study appears to be representative of the general heart failure population. Also, new referrals to heart failure clinics may not equal a new diagnosis of heart failure. Patients may have had heart failure for some time and could have been treated by a primary care physician or other healthcare provider before referral to the heart failure clinic. Finally, this sample also may be more educated about heart failure, but the levels of heart failure knowledge, self-efficacy, and self-care scores at the beginning of study were not at desired levels (see Table 2).

Given that this study was cross-sectional in nature and examined the relationships between these variables in newly referred patients, the influence of health literacy on knowledge, self-efficacy, and self-care over time should be assessed as relationships may change with time and within the context of traditional clinic-based education. Other limitations in this study include the naturalistic setting, use of self-report measures, and small recruitment from one site (Community Health Network), as well as the absence of data on patient heart failure classification or prior education about heart failure. Utilizing a naturalistic setting could result in unknown confounding factors and ultimately bias results, but this setting also has higher external validity. Moreover, the use of self-report measures may introduce bias, although the risk of this was minimized by utilizing previously-validated measures. The sample size for this study was adequate to test the structural equation model examining the relationships between health literacy, knowledge, self-efficacy, and self-care, but there was not sufficient sample size to add additional demographic parameters to the model that could further explain relationships with health literacy as demonstrated in other modeling research.³⁵ There also were some participants excluded due to incomplete data, which could have altered the results.

CONCLUSION

Although health literacy influences patient knowledge, health literacy and knowledge do not fully explain why patients perform self-care. Instead, self-efficacy was found to be independently-related to self-care. The models tested clarified some relationships between health literacy and self-care, but relationships between health literacy, knowledge, self-efficacy, and self-care appear to be complex and merit further study. Future research should examine additional factors that may influence heart failure self-care, such as motivation to perform self-care.

Acknowledgments

This work was supported by a seed grant from the Purdue University Regenstrief Center for Healthcare Engineering, the Clifford Kinley Trust (Purdue University), the

American Association of Heart Failure Nurses Bernard Saperstein Grant, and the Delta Omicron Chapter of Sigma Theta Tau International. Support for Aleda Chen while a graduate student was provided by the National Institute on Aging (T32AG025671) and the Purdue University Center on Aging and the Life Course as well as from the American Foundation for Pharmaceutical Excellence. The authors would like to thank Susie Carter, RN, BC, FAACVPR, AACC, Manager of Cardiopulmonary Rehab at the Advanced Heart Care Center, Indiana University Health Bloomington Hospital, and Jennifer Forney BSN, RN, Ellen Slifcak BA, RN and Susan Krajewski BSN, RN, MPA, Cleveland Clinic for their assistance and support of this project.

References

1. Go AS, Mozaffarian D, Roger VL, et al. Heart disease and stroke statistics—2013 update. *Circulation*. 2013; 127:e6–220. [PubMed: 23239837]

2. Riegel B, Lee CS, Dickson VV, Carlson B. An update on the self-care of heart failure index. *J Cardiovasc Nurs*. 2009; 24:485–497. [PubMed: 19786884]
3. Riegel B, Moser DK, Anker SD, et al. State of the science. Promoting self-care in persons with heart failure. A scientific statement from the American Heart Association. *Circulation*. 2009; 120:1141–1163. [PubMed: 19720935]
4. Orem, D. Self-care and health-promotion: Understanding self-care. In: McLaughlin, K.; Taylor, SG., editors. *Self-care theory in nursing: selected papers of Dorothea Orem*. New York, NY: Springer Publishing Co, Inc; 2003.
5. Evangelista LS, Rasmusson KD, Laramie AS, et al. Health literacy and the patient with heart failure - Implications for patient care and research: a consensus statement of the Heart Failure Society of America. *J Cardiac Fail*. 2010; 16:9–16.
6. Artinian NT, Magnan M, Sloan M, Lange MP. Self-care behaviors among patients with heart failure. *Heart Lung*. 2002; 31:161–172. [PubMed: 12011807]
7. Institute of Medicine. *Health literacy: a prescription to end confusion*. National Academy of Sciences; 2004.
8. Gazmararian JA, Williams MV, Peel J, Baker DW. Health literacy and knowledge of chronic disease. *Patient Educ Couns*. 2003; 51:267–275. [PubMed: 14630383]
9. Kalichman SC, Benotsch E, Suarez T, Catz S, Miller J, Rompa D. Health literacy and health-related knowledge among persons living with HIV/AIDS. *Am J Prev Med*. 2000; 18:325–331. [PubMed: 10788736]
10. Miller D, Brownlee C, McCoy T, Pignone M. The effect of health literacy on knowledge and receipt of colorectal cancer screening: a survey study. *BMC Fam Pract*. 2007; 8:16. [PubMed: 17394668]
11. Paasche-Orlow MK, Riekert KA, Bilderback A, et al. Tailored education may reduce health literacy disparities in asthma self-management. *Am J Respir Crit Care Med*. 2005; 172:980–986. [PubMed: 16081544]
12. Pandit AU, Tang JW, Bailey SC, et al. Education, literacy, and health: mediating effects on hypertension knowledge and control. *Patient Educ Couns*. 2009; 75:381–385. [PubMed: 19442477]
13. Peterson NB, Dwyer KA, Mulvaney SA, Dietrich MA, Rothman RL. The influence of health literacy on colorectal cancer screening knowledge, beliefs and behavior. *J Natl Med Assoc*. 2007; 99:1105–1112. [PubMed: 17987913]
14. Bandura A. Self efficacy: toward a unifying theory of behavioral change. *Psychol Rev*. 1977; 84:191–215. [PubMed: 847061]
15. Yehle KS, Plake KS. Self-efficacy and educational interventions in heart failure: a review of the literature. *J Cardiovasc Nurs*. 2010; 25:175–188. [PubMed: 20386241]
16. Dennison CR, McEntee ML, Samuel L, et al. Adequate health literacy is associated with higher heart failure knowledge and self-care confidence in hospitalized patients. *J Cardiovasc Nurs*. 2011; 26:359–367. [PubMed: 21099698]
17. Chen AMH, Yehle KS, Plake KS, Murawski MM, Mason HL. Health literacy and self-care of patients with heart failure. *J Cardiovasc Nurs*. 2011; 26:446–451. [PubMed: 21263340]
18. Peterson PN, Shetterly SM, Clarke CL, et al. Health literacy and outcomes among patients with heart failure. *J Amer Med Assoc*. 2011; 305:1695–1701.
19. Morrow D, Wu J, Murray MD, et al. Correlates of health literacy in patients with chronic heart failure. *Gerontologist*. 2006; 46:669–676. [PubMed: 17050758]
20. Baker DW, Williams MV, Parker RM, Gazmararian JA, Nurss J. Development of a brief test to measure functional health literacy. *Patient Educ Couns*. 1999; 38:33–42. [PubMed: 14528569]
21. Riegel B, Carlson B, Moser DK, Sebern M, Hicks FD, Roland V. Psychometric testing of the self-care of heart failure index. *J Cardiac Fail*. 2004; 10:350–360.
22. Yehle KS, Sands LP, Rhynders PA, Newton GD. The effect of shared medical visits on knowledge and self-care in patients with heart failure: a pilot study. *Heart Lung*. 2009; 38(1):25–33. [PubMed: 19150528]
23. Kline, RB., editor. *Principles and practice of structural equation modeling*. 3. New York: The Guilford Press; 2011.

24. Holbert RL, Stephenson MT. Structural equation modeling in the communication sciences, 1995–2000. *Health Communication Research*. 2002; 28:531–551.
25. Federman AD, Sano M, Wolf MS, Siu AL, Halm EA. Health literacy and cognitive performance in older adults. *J Am Geriatr Soc*. 2009; 57:1475–1480. [PubMed: 19515101]
26. Sudore RL, Landefeld CS, Williams BA, Barnes DE, Lindquist K, Schillinger D. Use of a modified informed consent process among vulnerable patients. *J Gen Intern Med*. 2006; 21:867–873. [PubMed: 16881949]
27. Sarkar U, Fisher L, Schillinger D. Is self-efficacy associated with diabetes self-management across race/ethnicity and health literacy? *Diabetes Care*. 2006; 29:823–829. [PubMed: 16567822]
28. Schillinger D, Barton LR, Karter AJ, Wang F, Adler N. Does literacy mediate the relationship between education and health outcomes? A study of a low-income population with diabetes. *Public Health Reports*. 2006; 121:245–254. [PubMed: 16640146]
29. Hooper D, Coughlan J, Mullen MR. Structural equation modeling: guidelines for determining model fit. *The Electronic Journal of Business Research Methods*. 2008; 6:53–60.
30. Steiger JH. Point estimation, hypothesis testing, and interval estimation using the RMSEA: some comments and a reply to Hayduk and Glaser. *Struct Equ Modeling*. 2000; 7:149–162.
31. Schreiber JB. Core reporting practices in structural equation modeling. *Res Social Adm Pharm*. 2008; 4:83–97. [PubMed: 18555963]
32. Macabasco-O’Connell A, DeWalt D, Brouckson KA, et al. Relationship between literacy, knowledge, self-care behaviors, and heart failure-related quality of life among patients with heart failure. *J Gen Intern Med*. 2011; 26:979–986. [PubMed: 21369770]
33. Murray MD, Young J, Hoke S, et al. Pharmacist intervention to improve medication adherence in heart failure: a randomized trial. *Arch Intern Med*. 2007; 146:714–725.
34. DeWalt DA, Corr KE, Kosner MC, et al. A heart failure self-management program for patients of all literacy levels: a randomized, controlled trial. *BMC Health Serv Res*. 2006; 6:30–40. [PubMed: 16533388]
35. Schmitt MR, Miller MJ, Harrison DL, et al. Communicating non-steroidal anti-inflammatory drug risks: verbal counseling, written medicine information, and patients’ risk awareness. *Patient Educ Couns*. 2011; 83:391–397. [PubMed: 21129892]

Figure 1.
Structural Equation Models Tested

Table 1**Participant Characteristics (N=63)**

Age, mean (SD), y	62.1	(13.7)
Years of Education, mean (SD), y	13.7	(2.9)
Prescription Medications, mean (SD)	10.23	(5.5)
Recruitment Site, N(%)		
Bloomington Hospital	25	(39.7)
Community Health Network	5	(7.9)
Cleveland Clinic	33	(52.4)
Male, N(%)	33	(52.4)
Married, N(%)	39	(61.9)
Married	34	(66.7)
Ethnicity/Race, N(%)		
Black/African American	7	(11.1)
White/Caucasian	54	(85.7)
Hispanic/Latino	1	(1.6)
American Indian/Alaskan Native	1	(1.6)
Employment Status, N(%)		
Full-Time Employed	20	(31.7)
Sick Leave/Disability	10	(15.9)
Unemployed or Retired	33	(52.4)
Perceived Financial Status, N(%)		
More than Enough to Make Ends Meet	25	(39.7)
Enough to Make Ends Meet	29	(46.0)
Not Enough to Make ends Meet	9	(14.3)
Health Literacy Category,^a N(%)		
Inadequate (Range: 0–16)	10	(15.9)
Marginal (Range: 17–22)	10	(15.9)
Adequate (Range: 23–36)	43	(68.3)

^a As measured by the Short-Form Test of Functional Health Literacy in Adults (S-TOFHLA)

Table 2

Health Literacy, Knowledge, Self-Efficacy, and Self-Care Scores (N=61)

Measure	Mean $\pm$ SD	Possible Range	Meaning
Health Literacy ^a	27.4 $\pm$ 9.3	0–36	Adequate health literacy
Self-Care Maintenance ^b	67.6 $\pm$ 17.8	0–100	Not adequate adherence
Self-Care Management ^b	64.7 $\pm$ 21.6	0–100	Not adequate adherence
Self-Efficacy ^b	67.3 $\pm$ 19.7	0–100	Not adequate
Heart Failure (HF) Knowledge, Overall ^c	8.1 $\pm$ 2.6	0–15	54% correct

HF Knowledge, Individual Items Correct Answer ^c	N	%
Definition of heart failure	43	69.4
Inappropriate weight gain	21	33.9
Mechanism of ACE Inhibitors	17	27.4
Side effects of ACE Inhibitors	15	24.2
Mechanism of digoxin	14	22.6
Side effects of digoxin	24	38.7
HF exacerbation symptom	46	74.2
Mechanism of diuretics	52	83.9
Side effects of diuretics	9	14.5
Appropriate alcohol use	41	66.1
Definition of advanced directive	39	62.9
Sodium in a food label	48	77.4
Food item with lowest sodium	56	90.3
Proper heart failure self-care	23	37.1
Reasons for rehospitalization	37	59.7

^a As measured by the Short-Form Test of Functional Health Literacy in Adults (S-TOFHLA)^b As measured by the Self-Care of Heart Failure Index v.6 (SCHFI®)^c As measured by the Heart Failure Knowledge Questionnaire (HFKQ)

Table 3
Correlations of Health Literacy and Years of Formal Education on Outcome Variables

	Years of Formal Education Pearson Corr. (Sig.)	Heart Failure Knowledge Pearson Corr. (Sig.)	Self-Efficacy for Self-Care Pearson Corr. (Sig.)	Self-Care Maintenance Adherence Pearson Corr. (Sig.)	Self-Care Management Adherence Pearson Corr. (Sig.)
Health Literacy Score	0.418 (p=0.001)	0.548 (p<0.001)	0.201 (p=0.114)	0.116 (p=0.366)	0.233 (p=0.066)
Years of Formal Education	----	0.402 (p=0.001)	0.186 (p=0.145)	0.239 (p=0.060)	0.176 (p=0.169)
Heart Failure Knowledge		----	0.123 (p=0.335)	0.182 (p=0.153)	0.226 (p=0.075)
Self-Efficacy			----	0.306 (p=0.015)	0.334 (p=0.007)
Self-Care Maintenance Adherence				----	0.285 (p=0.023)

Table 4

Comparison of Structural Equation Models for Maximum Likelihood Estimation

	Model 1	Model 2	Model 3	Model 4
χ^2	3.0466	2.9076	6.3392	6.2001
DF	4	3	5	4
$\text{Pr} > \chi^2$	0.5501*	0.4061*	0.2746*	0.1847*
$\Delta \text{ in } \chi^2$	-	-0.1390	+3.2926	+3.1535
RMSEA	0.0000*	0.0000*	0.0657	0.0942
GFI	0.9840*	0.9846*	0.9681*	0.9862*
NFI	0.9511*	0.9534*	0.8983	0.9005
CFI	1.0000*	1.0000*	0.9717*	0.9535*

Key: DF = Degrees of Freedom, Pr = Probability, RMSEA = Root mean square error of approximation, GFI = Goodness of Fit Index, NFI = Normed fit index, CFI = Comparative fix index

* Met conservative cut-off for fit statistic