Themed Section: Midkine

REVIEW

Midkine and cytoplasmic maturation of mammalian oocytes in the context of ovarian follicle physiology

Shuntaro Ikeda* and Masayasu Yamada

*Laboratory of Reproductive Biology, Graduate School of Agriculture, Kyoto University, Kyoto Japan***Correspondence**

Shuntaro Ikeda and Masayasu Yamada, Laboratory of Reproductive Biology, Graduate School of Agriculture, Kyoto University, Kyoto 606-8502, Japan. E-mail: ikedash@kais.kyoto-u.ac.jp; masayasu@kais.kyoto-u.ac.jp

*Present address: Laboratory of Animal Physiology and Functional Anatomy, Graduate School of Agriculture, Kyoto University, 606-8502 Kyoto, Japan.

Keywords

midkine; oocyte maturation; *in vitro* maturation; cytoplasmic maturation; developmental competence; granulosa cell; cumulus cell; follicle; ovary

Received

6 June 2013

Revised

12 July 2013

Accepted

21 July 2013

Midkine (MK) was originally characterized as a member of a distinct family of neurotrophic factors functioning in the CNS. However, it was later discovered that MK is abundantly expressed in ovarian follicles. Since then, the physiological roles of this molecule in the ovary have been steadily investigated. During the *in vitro* maturation (IVM) of oocytes MK was shown to promote the cytoplasmic maturation of oocytes, as indicated by post-fertilization development. This effect of MK could be mediated via its pro-survival (anti-apoptotic) effects on the cumulus-granulosa cells that surround oocytes. The oocyte competence-promoting effects of MK are discussed in the context of the recently discovered involvement of MK in the full maturation of ovarian follicles. MK was at the frontline of a new paradigm for neurotrophic factors as oocytotrophic factors. MK may promote the developmental competence of oocytes via common signalling molecules with the other neurotrophic factor(s). Alternatively or concomitantly, MK may also interact with various transmembrane molecules on cumulus-granulosa cells, which are important for ovarian follicle growth, dominance and differentiation, and act as a unique pro-survival factor in ovarian follicles, such that MK promotes oocyte competence. MK, along with other ovarian neurotrophic factors, may contribute to the optimization of the IVM system.

LINKED ARTICLES

This article is part of a themed section on Midkine. To view the other articles in this section visit <http://dx.doi.org/10.1111/bph.2014.171.issue-4>

Abbreviations

ALK, anaplastic lymphoma tyrosine kinase; ART, assisted reproductive technology; BDNF, brain-derived neurotrophic factor; CEO, cumulus-enclosed oocyte; DO, denuded oocyte; DKO, double deficient; GV, germinal vesicle; GDNF, glial cell-derived neurotrophic factor; HSPGs, heparan sulphate proteoglycans; IVC, *in vitro* culture; IVF, *in vitro* fertilization; IVM, *in vitro* maturation; LM-PCR, ligation-mediated PCR; LRP1, low density lipoprotein receptor-related protein 1; LH, luteinizing hormone; MII, metaphase of the second meiotic division; MK, midkine; NRG-1, neuregulin 1; Pro-1, prophase of the first meiotic division; PTN, pleiotrophin; PTPR, receptor tyrosine phosphatase; Trk, tropomyosin-related kinase

Introduction

Midkine (MK) was originally identified as the product of a gene, the expression of which increased in the early stage of retinoic acid-induced differentiation of murine embryonal

carcinoma cells, and it was shown to be abundant in the mid-gestation stages (days 8 to 11) of murine embryonic development (Kadomatsu *et al.*, 1988). Along with pleiotrophin (PTN), another member of a distinct family of growth factors, MK is known to exhibit neurotrophic properties that

promote various neuronal functions such as neurite outgrowth (Rauvala, 1989; Muramatsu *et al.*, 1993), survival (Owada *et al.*, 1999a,b; Jin *et al.*, 2009) and migration (Maeda and Noda, 1998; Maeda *et al.*, 1999). However, MK is also expressed in adult ovaries and the involvement of MK in ovarian physiology has been gradually investigated. As described below, MK is exclusively expressed in the granulosa cells of ovarian follicles (Karino *et al.*, 1995; Gomez *et al.*, 2003; Muramatsu *et al.*, 2006; Hirota *et al.*, 2007). Granulosa cells are somatic cells that form multiple layers inside the basal lamina of follicles and enclose oocytes. In well-developed antral follicles, the complex of granulosa cell layers and an oocyte forms a cumulus oophorus, a mound-like structure projecting into the antrum, and granulosa cells in the cumulus oophorus are often called cumulus cells (Figure 1A). We previously demonstrated that MK promotes the cytoplasmic maturation of oocytes, as indicated by their ability to develop to the blastocyst stage, and it has been suggested that this action of MK is due to its pro-survival effects on cumulus-granulosa cells (Ikeda *et al.*, 2000a,b; 2006). These findings preceded those of other studies, where it was shown that various neurotrophic factors have the ability to act as oocyte competence-promoting factors (De Sousa *et al.*, 2004; Mao *et al.*, 2012; Linher-Melville and Li, 2013) and MK was demonstrated to be involved in the full maturation of ovarian follicles (Muramatsu *et al.*, 2006). In this review, we describe the effects of MK on the cytoplasmic maturation of oocytes and associated processes in terms of ovarian follicle physiology. The molecular target nomenclature used in this review conforms to that in *British Journal of Pharmacology's* Concise guide to PHARMACOLOGY (Alexander *et al.*, 2013).

MK expression in mammalian ovarian follicles

Various tissues express MK in adults (Muramatsu *et al.*, 1993; for the mouse expressed sequence tag profile, see <http://www.ncbi.nlm.nih.gov/UniGene/ESTProfileViewer.cgi?uglist=Mm.906>), and, among them, the ovary is one of the representative tissues that expresses MK (Ohshima *et al.*, 1994; Karino *et al.*, 1995; Minegishi *et al.*, 1996; Hirota *et al.*, 2007). MK mRNA is exclusively expressed in the granulosa cells of ovarian follicles across species (Karino *et al.*, 1995; Gomez *et al.*, 2003; Muramatsu *et al.*, 2006; Hirota *et al.*, 2007). Furthermore, consistent with the expression of its mRNA in follicular cells, follicular fluid contains high concentrations of MK (Ohshima *et al.*, 1994; Hirota *et al.*, 2007). These findings indicate that MK is abundantly expressed in ovarian follicles and provide an insight into the possible involvement of MK in oocyte maturation, as the ovarian follicle is the microenvironment for oocyte maturation in mammals (Hennet and Combelles, 2012).

Oocyte maturation

Fully grown oocytes in the ovaries of all animals are meiotically arrested at the prophase of the first meiotic division (Pro-I). Oocytes at this stage contain a large nucleus called the germinal vesicle (GV). In most mammals, oocytes are released from the Pro-I arrest in response to a pre-ovulatory surge of pituitary-derived luteinizing hormones (LH) and then resume meiosis. The resumption of meiosis goes through dissolution of the GV nuclear envelope (GV breakdown), chromatin condensation, the assembly of the meiosis I spindles, chromosome migration and asymmetric cell division to reduce the chromosome number through the extrusion of the first polar body, and oocytes enter second arrest at the metaphase of the second meiotic division (MII). Fertilization triggers the release of oocytes from the MII arrest, and they complete meiosis and proceed to mitosis (Masui and Clarke, 1979; Masui, 2001). Oocyte maturation is defined as the re-initiation and completion of the first meiotic division, the subsequent progression to MII and the completion of nuclear and cytoplasmic processes that are essential for fertilization and early embryo

Figure 1

(A) Schematic diagram of an antral follicle. (B) Bovine CEOs recovered from ovaries. Bar represents 300 μ m.

development (Kupker *et al.*, 1998). The meiotic process in oocyte maturation from Pro-I to MII arrest is often referred to as 'nuclear maturation' and this term postulates that oocyte maturation entails 'cytoplasmic maturation', all accompanying events that are necessary to support subsequent embryonic development, which are recognized to include organellar modifications, post-transcriptional control of polyadenylation and protein accumulation (Hennet and Combettes, 2012; Li and Albertini, 2013). In contrast to nuclear maturation, which is a fairly well-defined process, cytoplasmic maturation has not been clearly elucidated either in molecular and morphological terms. However, its existence is evident because (i) oocytes that complete nuclear maturation do not necessarily carry out the subsequent developmental process and (ii) the ability of nuclear-matured oocytes to complete post-maturation developmental processes (fertilization, completion of meiosis, pronuclear formation and pre-implantation development and further) is affected by the environmental conditions present during nuclear maturation (Leibfried-Rutledge *et al.*, 1987; van de Leemput *et al.*, 1999; Rizados *et al.*, 2002; Lonergan, 2007; Hashimoto, 2009; Leidenfrost *et al.*, 2011).

Oocyte maturation in most mammals occurs in the ovarian antral follicles preceding ovulation and is a prerequisite for subsequent fertilization. Mammalian oocytes in ovarian antral follicles are enclosed in multiple layers of cumulus-granulosa cells, and oocyte maturation occurs under communication with these ovarian somatic cells (Li and Albertini, 2013).

***In vitro* maturation (IVM)**

IVM is a culture method of fully grown oocytes that allows them to undergo oocyte maturation *in vitro*. IVM has been experimentally established in various species (Edwards, 1965) and is used practically for the *in vitro* production of cattle embryos, when it is combined with the subsequent *in vitro* fertilization (IVF) and *in vitro* culture (IVC) of embryos (a set of three culture procedures is often referred to as IVM-F-C; Lonergan, 2007; Hashimoto, 2009). Furthermore, IVM has received increasing attention as an attractive alternative to gonadotrophin stimulation for conventional human IVF (Jurema and Nogueira, 2006; Smits *et al.*, 2011; Nogueira *et al.*, 2012; Chian *et al.*, 2013).

MK as an oocyte competence-promoting factor

The blastocyst stage is the general end point of the culture in the bovine IVM-F-C system because blastocysts can be non-surgically transferred to the uterus of live animals and are expected to implant and result in pregnancy. Although a high proportion (>70%) of cultured oocytes undergo nuclear maturation, fertilization and post-fertilization early cleavage in the current bovine IVM-F-C programme, they do not necessarily develop to the blastocyst stage (Leidenfrost *et al.*, 2011). The culture conditions of IVM significantly affect the developmental competence of oocytes after IVF (Lonergan, 2007; Hashimoto, 2009). Furthermore, it is well known that

post-fertilization development of IVM-derived oocytes to the blastocyst stage is generally less successful than that of oocytes matured *in vivo* (Leibfried-Rutledge *et al.*, 1987; van de Leemput *et al.*, 1999; Rizados *et al.*, 2002). These phenomena have provided evidence for the 'cytoplasmic maturation' described above. In other words, the ability of oocytes to develop to the blastocyst stage is a practical criterion for the cytoplasmic maturation of oocytes. The reduced cytoplasmic maturation observed in IVM-derived oocytes relative to *in vivo* matured oocytes is often attributed to the lack of physiological factors in IVM that regulate the acquisition of developmental competence of oocytes. *In vivo*, cumulus-enclosed oocytes (CEOs; Figure 1B) undergo maturation in the follicular fluid of antral follicles. Therefore, the effects of adding follicular fluid or its components to IVM medium have been extensively examined in an attempt to improve the IVM system, such that it confers optimal developmental competence to oocytes (Gordon, 2003). We focused on a heparin-binding growth factor, MK in the search for oocyte competence-promoting factors in bovine follicular fluid. MK was chosen because when we partitioned pooled bovine follicular fluid into heparin-binding and non-binding fractions by affinity chromatography and examined the effects of each fraction on the IVM of bovine CEOs, the heparin-binding fraction markedly enhanced the ability of CEOs to develop to the blastocyst stage after IVF (Ikeda *et al.*, 1999).

In studies that examined the effects of MK (Ikeda *et al.*, 2000a,b), bovine CEOs at the GV stage obtained from small ovarian follicles (2–6 mm in diameter) were cultured in a serum-free IVM medium for 24 h. CEOs that were presumed to be mature were then subjected to IVF with capacitated bovine spermatozoa. Following IVF, presumptive zygotes were freed from cumulus cells and transferred into IVC medium, and then cultured until day 8 (IVF = day 0). When purified mouse or bovine recombinant MK was added to the IVM medium at physiological concentrations, the percentage of oocytes reaching the blastocyst stage by day 8 after IVF was significantly higher in the MK-treated groups than in the untreated group but the proportion of oocytes reaching the MII stage and the cleavage stage after IVF were not affected. These results indicate that MK has the ability to promote the cytoplasmic maturation of bovine CEOs without affecting nuclear maturation, and suggest that MK in follicular fluid is a physiological regulator that enhances the developmental competence of bovine CEOs.

MK is a heparin-binding protein that is thought to interact with a heparin-like carbohydrate chain in heparan sulphate proteoglycans (HSPGs) on the target cell surface. Heparin, a highly sulphated glycosaminoglycan and putative antagonist of the heparan sulphate moiety of cell-linked HSPGs, was shown to inhibit MK action on neurite outgrowth (Kaneda *et al.*, 1996a,b). Heparitinase treatment, which digests heparan sulphate, also diminished the effects of MK on neurite cells (Kaneda *et al.*, 1996b). The promoting effects of MK during IVM on the developmental competence of bovine CEOs were also strongly inhibited by the pre-incubation of MK with heparin (Ikeda *et al.*, 2000a) or by treatment of CEOs with heparitinase (Ikeda *et al.*, 2000b), which suggests that the heparin-like moiety of the heparan sulphate chain on the cell surface of CEOs is required for MK to exert its oocyte competence-promoting effects.

In addition, supplementation of the IVM medium with MK also enhanced the cytoplasmic maturation of mouse CEOs with the same mode of action as that observed in bovine CEOs (Yamada *et al.*, 2012).

MK action on oocytes via cumulus-granulosa cells

Cumulus cells are actively involved in the development of CEOs including Pro-I arrest of oocytes until the pre-ovulatory LH surge, mediating the maturation-inducing signal of the LH surge to the oocytes, and support the cytoplasmic maturation of oocytes during the maturation period (Tanghe *et al.*, 2002). As oocytes are cultured as CEOs in IVM, we attempted to clarify whether the presence of cumulus (or granulosa) cells is required for the promoting effects of MK on developmental competence after IVF. When oocytes mechanically freed from cumulus cells (denuded oocytes, DOs) were matured in IVM medium with or without MK in the presence or absence of isolated cumulus or granulosa cells and were subjected to IVF, the promoting effects of MK on the developmental competence of DOs to the blastocyst stage after IVF were exerted only in the presence of these cells (Ikeda *et al.*, 2000a; 2006). In addition, we prepared the conditioned media of granulosa cells cultured with or without MK (CMMK+ or CMMK-, respectively) and found that the addition of CMMK+ into the IVM medium of DOs significantly enhanced post-fertilization blastocyst development relative to the no additive control and CMMK- supplemented groups (Ikeda *et al.*, 2006). These results collectively suggest that the developmental competence-promoting effect of MK during IVM is mediated by cumulus-granulosa cells rather than by a direct action on oocytes.

Pro-survival effects of MK on cumulus-granulosa cells

Many reports have demonstrated the importance of the presence of cumulus-granulosa cells during an IVM culture in terms of the developmental competence of oocytes (Zhang *et al.*, 1995; Kim *et al.*, 1997; Hashimoto *et al.*, 1998; Luciano *et al.*, 2005). As described above, the primary site of action of MK was shown to be cumulus cells rather than oocytes for its developmental competence-promoting effects. This finding prompted us to explore the action of MK on cumulus cells. Cumulus-cells in CEOs spontaneously undergo apoptosis during IVM (Ikeda *et al.*, 2003; Yuan *et al.*, 2005). However, MK exhibits anti-apoptotic effects on certain types of cells, including neuronal (Owada *et al.*, 1999a,b) and tumour (Qi *et al.*, 2000; Ohuchida *et al.*, 2004) cells. Therefore, we hypothesized that MK enhanced the viability of cumulus cells in CEOs during IVM through its anti-apoptotic effects. Using ligation-mediated PCR, which can detect apoptotic internucleosomal fragmentation in genomic DNA with high sensitivity, we examined the effects of MK on the apoptosis of cumulus cells in bovine CEOs during the IVM period. Fragmentation was scarcely detected at the start of IVM, but increased as the IVM culture progressed. The degree of fragmentation was significantly

lower in MK-treated CEOs than in untreated CEOs. This apoptosis-suppressing effect of MK on cumulus cells was further confirmed *in situ* by the conventional TUNEL assay (Ikeda *et al.*, 2006). Hirota *et al.* (2007), using the 5-bromo-2'-deoxyuridine (BrdU) incorporation assay, also found that MK stimulated the proliferation of cultured luteinized granulosa cells. As cumulus cells have an important role in IVM and there is a negative correlation between apoptosis in cumulus cells and the developmental competence of oocytes, we proposed a model in which the higher the extent of apoptosis in cumulus cells during the IVM period the greater the impairment of developmental competence of cultured oocytes (Ikeda *et al.*, 2003; 2006). Although a negative correlation between cumulus apoptosis and oocyte competence was not observed when oocyte growth-based classification was applied for CEOs before IVM (Anguita *et al.*, 2007; Janowski *et al.*, 2012), many other findings are supportive of apoptosis in cumulus cells as a predictive parameter for oocyte competence after oocyte maturation (Alisch *et al.*, 2003; Corn *et al.*, 2005; Leroy *et al.*, 2005; Yuan *et al.*, 2005).

When we proposed the negative relationship between the developmental competence of oocytes and apoptosis of surrounding cumulus cells during IVM, we inferred that the degree of apoptosis in cumulus cells of *in vivo* matured bovine CEOs, which are considered to be highly competent, was low (Ikeda *et al.*, 2003; 2006). However, at that time, this had not been demonstrated. Salhab *et al.* (2013) recently evaluated the degree of cumulus apoptosis in *in vivo* matured bovine CEOs using TUNEL and found that the degree of apoptosis was significantly lower than that of the IVM counterparts, and was similar to that of CEOs before IVM. This finding further supports the possibility that a lower cumulus apoptosis is associated with oocyte developmental competence.

Taken together these results indicate that the survival of cumulus cells during the IVM period may have a causal relationship with the developmental competence of concomitantly cultured oocytes, and MK may act as a pro-survival factor of the cumulus cells.

Knockout model of MK family in relation to follicular physiology

Oocyte maturation and its ovulation in live mammals are closely related to the survival of antral follicles to attain the final pre-ovulatory stage, which is escape from the degeneration of follicles called atresia, the fate of the massive majority of follicles (McGee and Hsueh, 2000). The involvement of the MK family in ovarian follicular survival including ovulation has been clearly demonstrated in the double-deficient mouse model of MK and PTN (Muramatsu *et al.*, 2006). Although double-deficient (DKO) male mice were fertile in this study, most DKO females (19/24, 79%) were infertile. In DKO females, the number of mature follicles, which have a single, fused antrum, and that of ovulated oocytes upon superovulation were significantly lower than those in the wild-type female.

Apoptosis is a fundamental mechanism involved in follicular atresia (Hsueh *et al.*, 1994; Tilly, 1996); thus, from results revealed in *in vitro* studies, it is suggested that MK plays a role

in follicular survival by direct pro-survival actions on follicular granulosa cells (Ikeda *et al.*, 2006; Hirota *et al.*, 2007). Alternatively, the angiogenic properties of MK and PTN (Muramatsu, 2002; Zhang and Deuel, 1999) may also be involved in follicular survival, because angiogenesis is crucial for ovarian function including follicular growth, antrum formation, ovulation and follicle-luteal transition (Robinson *et al.*, 2009). However, the authors of the DKO study (Muramatsu *et al.*, 2006) noted that the density and size of the capillaries in the ovary were not significantly different between wild-type and DKO mice and suggested that functional deficits in the ovaries of DKO mice were not due to abnormalities in angiogenesis. Therefore, MK, as well as PTN, may be able to promote follicular survival via direct and local actions on follicular cells, which is separate from angiogenesis.

Neurotrophic factors are oocytotrophic?

After the discovery of MK as an oocyte competence-promoting factor, other neurotrophic factors, that is, brain-derived neurotrophic factor (BDNF; Kawamura *et al.*, 2005; Martins da Silva *et al.*, 2005), glial cell-derived neurotrophic factor (GDNF; Linher *et al.*, 2007) and neuregulin 1 (NRG-1; Noma *et al.*, 2011; Mao *et al.*, 2012) were successively

reported to have analogous oocyte competence-promoting effects with some variations (Table 1). These neurotrophic factors, similar to MK, are expressed not only in the CNS but also in peripheral tissues including the ovary: BDNF and GDNF were also reviewed by De Sousa *et al.* (2004) and/or Linher-Melville and Li (2013) and the ovarian expression of NRG-1 was documented in the work by Noma *et al.* (2011). The originally identified and/or well-established physiological activity of these factors is promoting the survival of various classes of neurons (Barde *et al.*, 1982; Lin *et al.*, 1993; Bermingham-McDonogh *et al.*, 1996; Gerecke *et al.*, 2004) similar to MK (Muramatsu and Muramatsu, 1991; Michikawa *et al.*, 1993; Satoh *et al.*, 1993). This suggests that neurotrophic and oocytotrophic signals may have features in common.

In particular, BDNF is included in the neurotrophin family, which is distinct from the MK family, and the signal transduction system of this family is well characterized: the biological effects of neurotrophins are mediated through the activation of one or more of the three members of the tropomyosin-related kinase (Trk) family (TrkA, TrkB and TrkC) and all neurotrophins activate the p75 neurotrophin receptor, a member of the TNF receptor superfamily (Skaper, 2012). Among these receptors, BDNF binds to and activates TrkB or p75 (Klein *et al.*, 1991). The major signal transduction pathways activated by TrkB are Ras-MAPK, PI3K-PKB and

Table 1

Ovarian neurotrophic factors that affect cytoplasmic maturation of CEOs during IVM

Factors	Receptor reported to be expressed in CEOs	Nuclear maturation (reported species)	Cytoplasmic maturation (reported species)	Direct action on oocytes in terms of nuclear (N) or cytoplasmic (C) maturation	Pro-survival effects on cumulus or granulosa cells <i>in vitro</i>
MK	LRP1 ^[1-3] , Notch2 ^[4,5] , $\alpha 6\beta 1$ integrin ^[6,7] , HSPGs ^[8,9]	-(cow) ^[18,19]	$\uparrow$ (mouse) ^[24] $\uparrow$ (cow) ^[18,19,25]	No (C) ^[18,25]	Yes ^[2,25]
BDNF	TrkB ^[10-14] , p75 ^[10-13]	$\uparrow$ (mouse) ^[10,14] $\uparrow$ (cow) ^[20] $\uparrow$ (pig) ^[12] $\uparrow$ (human) ^[21] -(mouse) ^[22] -(cow) ^[11] -(human) ^[13]	$\uparrow$ (mouse) ^[10, 22] $\uparrow$ (cow) ^[11] $\uparrow$ (pig) ^[12] -(cow) ^[20a] $\downarrow$ (human) ^[13]	Yes (N) ^[14,21] Yes (C) ^[11]	Yes ^[26] No ^[27]
GDNF	GFR α 1/Ret ^[15,16]	$\uparrow$ (mouse) ^[15] $\uparrow$ (pig) ^[16] $\uparrow$ (human) ^[21]	$\uparrow$ (pig) ^[16] -(mouse) ^[15]	Yes (N) ^[15,21]	?
NRG-1	ErbB2-3 ^[17]	$\uparrow$ (pig) ^[23] $\downarrow$ (mouse) ^[17]	$\uparrow$ (pig) ^[23] -(mouse) ^[17b]	?	Yes ^[17]

$\uparrow$, positive effect; -, no effect; $\downarrow$, negative effect; ?, no reference.

1, Azhar *et al.*, 2006; 2, Hirota *et al.*, 2007; 3, Ireland *et al.*, 2004; 4, Johnson *et al.*, 2001; 5, Zhang *et al.*, 2011; 6, Fujiwara *et al.*, 1997; 7, Honda *et al.*, 1995; 8, Princivale *et al.*, 2001; 9, Watson *et al.*, 2012; 10, Kawamura *et al.*, 2005; 11, Martins da Silva *et al.*, 2005; 12, Lee *et al.*, 2007; 13, Anderson *et al.*, 2010; 14, Seifer *et al.*, 2002; 15, Kawamura *et al.*, 2008; 16, Linher *et al.*, 2007; 17, Noma *et al.*, 2011; 18, Ikeda *et al.*, 2000a; 19, Ikeda *et al.*, 2000b; 20, Hong *et al.*, 2009; 21, Zhao *et al.*, 2011; 22, Zhang *et al.*, 2010a; 23, Mao *et al.*, 2012; 24, Yamada *et al.*, 2012; 25, Ikeda *et al.*, 2006; 26, Yi *et al.*, 2008; 27, Zhang *et al.*, 2010b.

^aPositive effect in the presence of metformin.

^bPositive effect in the presence of amphiregulin.

GFR α 1, GDNF family receptor α 1.

Figure 2

Schematic model of the actions of MK that result in the promotion of oocyte developmental competence. In this model, MK, upon binding to its receptor complex on cumulus-granulosa cells, activates the MAPK and PI3K pathways, which are common signalling pathways activated by BDNF or NRG-1, the other oocytetrophic factors in ovarian follicles (black arrows). Each component of the MK receptor complexes on granulosa cells functions as a crucial regulator of follicle growth, dominance and differentiation. LRP1 and $\alpha 6\beta 1$ integrin regulate steroidogenesis. Notch2 enhances granulosa cell proliferation. HSPGs contribute to oocyte-cumulus communications mediated by oocyte-secreted paracrine factors in maturing CEOs. MK activates these multifunctions of granulosa (cumulus) cells towards the oocyte full maturation and ovulation, and consequently promotes the developmental competence of oocytes (white arrows). The model does not mean that the MAPK and PI3K pathways are exclusive of the phenomena linked by white arrows.

PLC γ -PKC pathways (Thiele *et al.*, 2009; Skaper, 2012). NRG-1, a member of the EGF family, binds to and activates the receptor tyrosine kinases ErbB2-4 and downstream signal transduction includes the MAPK and PI3K pathways (Citri and Yarden, 2006; Yarden and Sliwkowski, 2001).

With regard to MK, its candidate receptors include receptor tyrosine phosphatase (PTPR) type Z1 (Maeda *et al.*, 1999), anaplastic lymphoma tyrosine kinase (ALK; Stoica *et al.*, 2002; Kuo *et al.*, 2007), low density lipoprotein receptor-related protein 1 (LRP1; Muramatsu *et al.*, 2000), Notch2 (Huang *et al.*, 2008), $\alpha 4\beta 1$ or $\alpha 6\beta 1$ integrins (Muramatsu *et al.*, 2004) and HSPGs (Kojima *et al.*, 1996; Nakanishi *et al.*, 1997; Kurosawa *et al.*, 2001). Some of these components are thought to form complexes depending on the cellular context mediating MK signalling (Kadomatsu *et al.*, 2013). The signal transduction of MK includes the MAPK and PI3K pathways (Owada *et al.*, 1999a; Qi *et al.*, 2001; Sandra *et al.*, 2004), which cross-talk with each other (Aksamitiene *et al.*, 2012) and are in common with the signal transduction system of BDNF and NRG-1. Therefore, MK may enhance the developmental competence of oocytes via these signalling cascades. In addition, the incomplete suppression of fertility in the MK-PTN DKO female mice discussed in the previous section might reflect the redundancy in the activation of the signalling by various factors including BDNF and NRG-1. Furthermore, it is noteworthy that the conditional disruption of MAPK (ERK1/2) in mouse ovarian granulosa cells resulted in ovulation failure (Fan *et al.*, 2009), which is analogous to the phenotype of the DKO females. This finding indicates that the MAPK pathway is critical at least for the ovulation and supports the involvement of MK in the pathway. However, in contrast to the case of DKO mice, the disruption

of MAPK did not suppress the formation of mature follicles. Therefore, the effect of MK in the ovary cannot be explained solely by the activation of the MAPK pathway.

Among the candidates for the MK receptor listed above, LRP1 (Ireland *et al.*, 2004; Azhar *et al.*, 2006; Hirota *et al.*, 2007), Notch2 (Johnson *et al.*, 2001; Zhang *et al.*, 2011), $\alpha 6\beta 1$ integrin (Honda *et al.*, 1995; Fujiwara *et al.*, 1997) and HSPGs (Princiville *et al.*, 2001; Watson *et al.*, 2012) were all shown to be expressed in ovarian granulosa cells. LRP1 and $\alpha 6\beta 1$ integrin have been implicated in granulosa cell steroidogenesis, which is important for ovarian follicle growth, dominance and differentiation (Fujiwara *et al.*, 1997; Ireland *et al.*, 2004; Azhar *et al.*, 2006). Notch2 regulates granulosa cell apoptosis and proliferation and, hence, is involved in follicle development (Zhang *et al.*, 2011). HSPGs are thought to play an essential role in oocyte-cumulus cell communication and mediate oocyte-secreted factor signalling on cumulus cells in the periovulatory follicle (Watson *et al.*, 2012). MK has close relationships to these players. Therefore, MK may exert multifunctions as a unique pro-survival factor in ovarian follicles, which consequently promotes oocyte competence. The model of MK function proposed in this section is depicted in Figure 2. However, although MK could exert its effects through any of the receptors mentioned above, MK also affects the function of granulosa cells (Ikeda *et al.*, 2006; Hirota *et al.*, 2007) despite the absence of representative receptor components, that is, PTPRZ1 and ALK on this target (Shiota *et al.*, 2003; Ren *et al.*, 2012). Hence, follicular granulosa cells may remain at the frontier in the search for the unique signal transductions of MK. A comprehensive understanding of MK signalling in granulosa cells is required to establish MK as a key player in ovarian physiology.

Concluding remarks

In this review, we have provided an overview of the roles for MK in the cytoplasmic maturation of oocytes, with recent topics concerning MK and other ovarian neurotrophic factors. MK was shown to promote the cytoplasmic maturation of oocytes during IVM of oocytes and these effects are thought to be mediated via its pro-survival (anti-apoptotic) effects on cumulus-granulosa cells surrounding oocytes. MK can be seen as a pro-survival factor of granulosa cells that is involved in follicular development including oocyte full maturation.

Practical applications of IVM for GV stage oocytes are currently limited to the production of domestic animals (mainly cattle). IVM applications in human assisted reproductive technology (ART) is developing with the expectation of a more patient-friendly alternative to ovarian stimulation and fertility preservation in young women with cancer in combination with the cryopreservation of ovarian tissue (Smitz *et al.*, 2011; Nogueira *et al.*, 2012; Chian *et al.*, 2013). MK along with other ovarian neurotrophic factors may contribute to the optimization of the IVM system not only in livestock production, but also in human infertility treatment in ART.

Conflict of interest

The authors declare no conflict of interest.

References

- Aksamitiene E, Kiyatkin A, Kholodenko BN (2012). Cross-talk between mitogenic Ras/MAPK and survival PI3K/Akt pathways: a fine balance. *Biochem Soc Trans* 40: 139–146.
- Alexander SPH, Benson HE, Faccenda E, Pawson AJ, Sharman JL, Catterall WA, Spedding M, Peters JA, Harmar AJ and CGTP Collaborators (2013). The Concise Guide to PHARMACOLOGY 2013/14: Overview. *Br J Pharmacol* 170: 1449–1867.
- Alish A, Ruping K, Koster F, Schopper B, Baum M, Finas D *et al.* (2003). Cumulus cell apoptosis as a predictor for oocyte quality in artificial reproduction technique. *Zentralbl Gynakol* 125: 452–457.
- Anderson RA, Bayne RA, Gardner J, De Sousa PA (2010). Brain-derived neurotrophic factor is a regulator of human oocyte maturation and early embryo development. *Fertil Steril* 93: 1394–1406.
- Anguita B, Vandaele L, Mateusen B, Maes D, Van Soom A (2007). Developmental competence of bovine oocytes is not related to apoptosis incidence in oocytes, cumulus cells and blastocysts. *Theriogenology* 67: 537–549.
- Azhar S, Medicherla S, Shen WJ, Fujioka Y, Fong LG, Reaven E *et al.* (2006). LDL and cAMP cooperate to regulate the functional expression of the LRP in rat ovarian granulosa cells. *J Lipid Res* 47: 2538–2550.
- Barde YA, Edgar D, Thoenen H (1982). Purification of a new neurotrophic factor from mammalian brain. *EMBO J* 1: 549–553.
- Birmingham-McDonogh O, McCabe KL, Reh TA (1996). Effects of GGF/neuregulins on neuronal survival and neurite outgrowth correlate with erbB2/neu expression in developing rat retina. *Development* 122: 1427–1438.
- Chian RC, Uzelac PS, Nargund G (2013). In vitro maturation of human immature oocytes for fertility preservation. *Fertil Steril* 99: 1173–1181.
- Citri A, Yarden Y (2006). EGF-ERBB signalling: towards the systems level. *Nat Rev Mol Cell Biol* 7: 505–516.
- Corn CM, Hauser-Kronberger C, Moser M, Tews G, Ebner T (2005). Predictive value of cumulus cell apoptosis with regard to blastocyst development of corresponding gametes. *Fertil Steril* 84: 627–633.
- De Sousa PA, Da Silva SJ, Anderson RA (2004). Neurotrophin signaling in oocyte survival and developmental competence: a paradigm for cellular totipotency. *Cloning Stem Cells* 6: 375–385.
- Edwards RG (1965). Maturation in vitro of mouse, sheep, cow, pig, rhesus monkey and human ovarian oocytes. *Nature* 208: 349–351.
- Fan HY, Liu Z, Shimada M, Sterneck E, Johnson PF, Hedrick SM *et al.* (2009). MAPK3/1 (ERK1/2) in ovarian granulosa cells are essential for female fertility. *Science* 324: 938–941.
- Fujiwara H, Honda T, Ueda M, Nakamura K, Yamada S, Maeda M *et al.* (1997). Laminin suppresses progesterone production by human luteinizing granulosa cells via interaction with integrin alpha 6 beta 1. *J Clin Endocrinol Metab* 82: 2122–2128.
- Gerecke KM, Wyss JM, Carroll SL (2004). Neuregulin-1beta induces neurite extension and arborization in cultured hippocampal neurons. *Mol Cell Neurosci* 27: 379–393.
- Gomez E, Royo LJ, Duque P, Carneiro G, Hidalgo C, Goyache F *et al.* (2003). 9-cis-retinoic acid during in vitro maturation improves development of the bovine oocyte and increases midkine but not IGF-I expression in cumulus-granulosa cells. *Mol Reprod Dev* 66: 247–255.
- Gordon I (2003). Maturing the oocyte. In: Gordon I (ed.). *Laboratory Production of Cattle Embryos*, 2nd edn. CAB International: Wallingford, pp. 112–157.
- Hashimoto S (2009). Application of in vitro maturation to assisted reproductive technology. *J Reprod Dev* 55: 1–10.
- Hashimoto S, Saeki K, Nagao Y, Minami N, Yamada M, Utsumi K (1998). Effects of cumulus cell density during in vitro maturation of the developmental competence of bovine oocytes. *Theriogenology* 49: 1451–1463.
- Hennet ML, Combelles CM (2012). The antral follicle: a microenvironment for oocyte differentiation. *Int J Dev Biol* 56: 819–831.
- Hirota Y, Osuga Y, Nose E, Koga K, Yoshino O, Hirata T *et al.* (2007). The presence of midkine and its possible implication in human ovarian follicles. *Am J Reprod Immunol* 58: 367–373.
- Honda T, Fujiwara H, Ueda M, Maeda M, Mori T (1995). Integrin alpha 6 is a differentiation antigen of human granulosa cells. *J Clin Endocrinol Metab* 80: 2899–2905.
- Hong SG, Jang G, Oh HJ, Koo OJ, Park JE, Park HJ *et al.* (2009). The effects of brain-derived neurotrophic factor and metformin on in vitro developmental competence of bovine oocytes. *Zygote* 17: 187–193.
- Hsueh AJ, Billig H, Tsafiri A (1994). Ovarian follicle atresia: a hormonally controlled apoptotic process. *Endocr Rev* 15: 707–724.
- Huang Y, Hoque MO, Wu F, Trink B, Sidransky D, Ratovitski EA (2008). Midkine induces epithelial-mesenchymal transition through Notch2/Jak2-Stat3 signaling in human keratinocytes. *Cell Cycle* 7: 1613–1622.

- Ikeda S, Azuma T, Hashimoto S, Yamada M (1999). In vitro maturation of bovine oocytes with fractions of bovine follicular fluid separated by heparin affinity chromatography. *J Reprod Dev* 45: 397–404.
- Ikeda S, Ichihara-Tanaka K, Azuma T, Muramatsu T, Yamada M (2000a). Effects of midkine during in vitro maturation of bovine oocytes on subsequent developmental competence. *Biol Reprod* 63: 1067–1074.
- Ikeda S, Nishikimi A, Ichihara-Tanaka K, Muramatsu T, Yamada M (2000b). cDNA cloning of bovine midkine and production of the recombinant protein, which affects in vitro maturation of bovine oocytes. *Mol Reprod Dev* 57: 99–107.
- Ikeda S, Imai H, Yamada M (2003). Apoptosis in cumulus cells during in vitro maturation of bovine cumulus-enclosed oocytes. *Reproduction* 125: 369–376.
- Ikeda S, Saeki K, Imai H, Yamada M (2006). Abilities of cumulus and granulosa cells to enhance the developmental competence of bovine oocytes during in vitro maturation period are promoted by midkine; a possible implication of its apoptosis suppressing effects. *Reproduction* 132: 549–557.
- Ireland JL, Jimenez-Krassel F, Winn ME, Burns DS, Ireland JJ (2004). Evidence for autocrine or paracrine roles of alpha2-macroglobulin in regulation of estradiol production by granulosa cells and development of dominant follicles. *Endocrinology* 145: 2784–2794.
- Janowski D, Salilew-Wondim D, Torner H, Tesfaye D, Ghanem N, Tomek W *et al.* (2012). Incidence of apoptosis and transcript abundance in bovine follicular cells is associated with the quality of the enclosed oocyte. *Theriogenology* 78: 656–669. e651–655.
- Jin L, Jianghai C, Juan L, Hao K (2009). Pleiotrophin and peripheral nerve injury. *Neurosurg Rev* 32: 387–393.
- Johnson J, Espinoza T, McGaughey RW, Rawls A, Wilson-Rawls J (2001). Notch pathway genes are expressed in mammalian ovarian follicles. *Mech Dev* 109: 355–361.
- Jurema MW, Nogueira D (2006). In vitro maturation of human oocytes for assisted reproduction. *Fertil Steril* 86: 1277–1291.
- Kadomatsu K, Tomomura M, Muramatsu T (1988). cDNA cloning and sequencing of a new gene intensely expressed in early differentiation stages of embryonal carcinoma cells and in mid-gestation period of mouse embryogenesis. *Biochem Biophys Res Commun* 151: 1312–1318.
- Kadomatsu K, Kishida S, Tsubota S (2013). The heparin-binding growth factor midkine: the biological activities and candidate receptors. *J Biochem* 153: 511–521.
- Kaneda N, Talukder AH, Ishihara M, Hara S, Yoshida K, Muramatsu T (1996a). Structural characteristics of heparin-like domain required for interaction of midkine with embryonic neurons. *Biochem Biophys Res Commun* 220: 108–112.
- Kaneda N, Talukder AH, Nishiyama H, Koizumi S, Muramatsu T (1996b). Midkine, a heparin-binding growth/differentiation factor, exhibits nerve cell adhesion and guidance activity for neurite outgrowth in vitro. *J Biochem* 119: 1150–1156.
- Karino S, Minegishi T, Ohyama Y, Tano M, Nakamura K, Miyamoto K *et al.* (1995). Regulation and localization of midkine in rat ovary. *FEBS Lett* 362: 147–150.
- Kawamura K, Kawamura N, Mulders SM, Sollewijn Gelpke MD, Hsueh AJ (2005). Ovarian brain-derived neurotrophic factor (BDNF) promotes the development of oocytes into preimplantation embryos. *Proc Natl Acad Sci U S A* 102: 9206–9211.
- Kawamura K, Ye Y, Kawamura N, Jing L, Groenen P, Gelpke MS *et al.* (2008). Completion of meiosis I of preovulatory oocytes and facilitation of preimplantation embryo development by glial cell line-derived neurotrophic factor. *Dev Biol* 315: 189–202.
- Kim KS, Minami N, Yamada M, Utsumi K (1997). Follicular cells affect the fertilizability and developmental competency of bovine oocytes in vitro. *Reprod Fertil Dev* 9: 763–766.
- Klein R, Nanduri V, Jing SA, Lamballe F, Tapley P, Bryant S *et al.* (1991). The trkB tyrosine protein kinase is a receptor for brain-derived neurotrophic factor and neurotrophin-3. *Cell* 66: 395–403.
- Kojima T, Katsumi A, Yamazaki T, Muramatsu T, Nagasaka T, Ohsumi K *et al.* (1996). Human ryudocan from endothelium-like cells binds basic fibroblast growth factor, midkine, and tissue factor pathway inhibitor. *J Biol Chem* 271: 5914–5920.
- Kuo AH, Stoica GE, Riegel AT, Wellstein A (2007). Recruitment of insulin receptor substrate-1 and activation of NF-kappaB essential for midkine growth signaling through anaplastic lymphoma kinase. *Oncogene* 26: 859–869.
- Kupker W, Diedrich K, Edwards RG (1998). Principles of mammalian fertilization. *Hum Reprod* 13 (Suppl 1): 20–32.
- Kurosawa N, Chen GY, Kadomatsu K, Ikematsu S, Sakuma S, Muramatsu T (2001). Glypican-2 binds to midkine: the role of glypican-2 in neuronal cell adhesion and neurite outgrowth. *Glycoconj J* 18: 499–507.
- Lee E, Jeong YI, Park SM, Lee JY, Kim JH, Park SW *et al.* (2007). Beneficial effects of brain-derived neurotrophic factor on in vitro maturation of porcine oocytes. *Reproduction* 134: 405–414.
- van de Leemput EE, Vos PL, Zeinstra EC, Bevers MM, van der Weijden GC, Dieleman SJ (1999). Improved in vitro embryo development using in vivo matured oocytes from heifers superovulated with a controlled preovulatory LH surge. *Theriogenology* 52: 335–349.
- Leibfried-Rutledge ML, Critser ES, Eyestone WH, Northey DL, First NL (1987). Development potential of bovine oocytes matured in vitro or in vivo. *Biol Reprod* 36: 376–383.
- Leidenfrost S, Boelhauve M, Reichenbach M, Gungor T, Reichenbach HD, Sinowatz F *et al.* (2011). Cell arrest and cell death in mammalian preimplantation development: lessons from the bovine model. *Plos ONE* 6: e22121.
- Leroy JL, Vanholder T, Mateusen B, Christophe A, Opsomer G, de Kruif A *et al.* (2005). Non-esterified fatty acids in follicular fluid of dairy cows and their effect on developmental capacity of bovine oocytes in vitro. *Reproduction* 130: 485–495.
- Li R, Albertini DF (2013). The road to maturation: somatic cell interaction and self-organization of the mammalian oocyte. *Nat Rev Mol Cell Biol* 14: 141–152.
- Lin LF, Doherty DH, Lile JD, Bektesh S, Collins F (1993). GDNF: a glial cell line-derived neurotrophic factor for midbrain dopaminergic neurons. *Science* 260: 1130–1132.
- Linher K, Wu D, Li J (2007). Glial cell line-derived neurotrophic factor: an intraovarian factor that enhances oocyte developmental competence in vitro. *Endocrinology* 148: 4292–4301.
- Linher-Melville K, Li J (2013). The roles of glial cell line-derived neurotrophic factor, brain-derived neurotrophic factor and nerve growth factor during the final stage of folliculogenesis: a focus on oocyte maturation. *Reproduction* 145: R43–R54.
- Loneragan P (2007). State-of-the-art embryo technologies in cattle. *Soc Reprod Fertil Suppl* 64: 315–325.
- Luciano AM, Lodde V, Beretta MS, Colleoni S, Lauria A, Modena S (2005). Developmental capability of denuded bovine oocyte in a

co-culture system with intact cumulus-oocyte complexes: role of cumulus cells, cyclic adenosine 3',5'-monophosphate, and glutathione. *Mol Reprod Dev* 71: 389–397.

McGee EA, Hsueh AJ (2000). Initial and cyclic recruitment of ovarian follicles. *Endocr Rev* 21: 200–214.

Maeda N, Noda M (1998). Involvement of receptor-like protein tyrosine phosphatase zeta/RPTPbeta and its ligand pleiotrophin/heparin-binding growth-associated molecule (HB-GAM) in neuronal migration. *J Cell Biol* 142: 203–216.

Maeda N, Ichihara-Tanaka K, Kimura T, Kadomatsu K, Muramatsu T, Noda M (1999). A receptor-like protein-tyrosine phosphatase PTPzeta/RPTPbeta binds a heparin-binding growth factor midkine. Involvement of arginine 78 of midkine in the high affinity binding to PTPzeta. *J Biol Chem* 274: 12474–12479.

Mao J, Whitworth KM, Spate LD, Walters EM, Zhao J, Prather RS (2012). Regulation of oocyte mitochondrial DNA copy number by follicular fluid, EGF, and neuregulin 1 during in vitro maturation affects embryo development in pigs. *Theriogenology* 78: 887–897.

Martins da Silva SJ, Gardner JO, Taylor JE, Springbett A, De Sousa PA, Anderson RA (2005). Brain-derived neurotrophic factor promotes bovine oocyte cytoplasmic competence for embryo development. *Reproduction* 129: 423–434.

Masui Y (2001). From oocyte maturation to the in vitro cell cycle: the history of discoveries of maturation-promoting factor (MPF) and cytotrophic factor (CSF). *Differentiation* 69: 1–17.

Masui Y, Clarke HJ (1979). Oocyte maturation. *Int Rev Cytol* 57: 185–282.

Michikawa M, Kikuchi S, Muramatsu H, Muramatsu T, Kim SU (1993). Retinoic acid responsive gene product, midkine, has neurotrophic functions for mouse spinal cord and dorsal root ganglion neurons in culture. *J Neurosci Res* 35: 530–539.

Minegishi T, Karino S, Tano M, Ibuki Y, Miyamoto K (1996). Regulation of midkine messenger ribonucleic acid levels in cultured rat granulosa cells. *Biochem Biophys Res Commun* 229: 799–805.

Muramatsu H, Muramatsu T (1991). Purification of recombinant midkine and examination of its biological activities: functional comparison of new heparin binding factors. *Biochem Biophys Res Commun* 177: 652–658.

Muramatsu H, Shirahama H, Yonezawa S, Maruta H, Muramatsu T (1993). Midkine, a retinoic acid-inducible growth/differentiation factor: immunochemical evidence for the function and distribution. *Dev Biol* 159: 392–402.

Muramatsu H, Zou K, Sakaguchi N, Ikematsu S, Sakuma S, Muramatsu T (2000). LDL receptor-related protein as a component of the midkine receptor. *Biochem Biophys Res Commun* 270: 936–941.

Muramatsu H, Zou P, Suzuki H, Oda Y, Chen GY, Sakaguchi N *et al.* (2004). $\alpha 4 \beta 1$ - and $\alpha 6 \beta 1$ -integrins are functional receptors for midkine, a heparin-binding growth factor. *J Cell Sci* 117 (Pt 22): 5405–5415.

Muramatsu H, Zou P, Kurosawa N, Ichihara-Tanaka K, Maruyama K, Inoh K *et al.* (2006). Female infertility in mice deficient in midkine and pleiotrophin, which form a distinct family of growth factors. *Genes Cells* 11: 1405–1417.

Muramatsu T (2002). Midkine and pleiotrophin: two related proteins involved in development, survival, inflammation and tumorigenesis. *J Biochem* 132: 359–371.

Nakanishi T, Kadomatsu K, Okamoto T, Ichihara-Tanaka K, Kojima T, Saito H *et al.* (1997). Expression of syndecan-1 and -3 during

embryogenesis of the central nervous system in relation to binding with midkine. *J Biochem* 121: 197–205.

Nogueira D, Sadeu JC, Montagut J (2012). In vitro oocyte maturation: current status. *Semin Reprod Med* 30: 199–213.

Noma N, Kawashima I, Fan HY, Fujita Y, Kawai T, Tomoda Y *et al.* (2011). LH-induced neuregulin 1 (NRG1) type III transcripts control granulosa cell differentiation and oocyte maturation. *Mol Endocrinol* 25: 104–116.

Ohuchida T, Okamoto K, Akahane K, Higure A, Todoroki H, Abe Y *et al.* (2004). Midkine protects hepatocellular carcinoma cells against TRAIL-mediated apoptosis through down-regulation of caspase-3 activity. *Cancer* 100: 2430–2436.

Ohya Y, Miyamoto K, Minamino N, Matsuo H (1994). Isolation and identification of midkine and pleiotrophin in bovine follicular fluid. *Mol Cell Endocrinol* 105: 203–208.

Owada K, Sanjo N, Kobayashi T, Mizusawa H, Muramatsu H, Muramatsu T *et al.* (1999a). Midkine inhibits caspase-dependent apoptosis via the activation of mitogen-activated protein kinase and phosphatidylinositol 3-kinase in cultured neurons. *J Neurochem* 73: 2084–2092.

Owada K, Sanjo N, Kobayashi T, Kamata T, Mizusawa H, Muramatsu H *et al.* (1999b). Midkine inhibits apoptosis via extracellular signal regulated kinase (ERK) activation in PC12 cells. *J Med Dent Sci* 46: 45–51.

Princiville M, Hasan S, Hosseini G, de Agostini AI (2001). Anticoagulant heparan sulfate proteoglycans expression in the rat ovary peaks in preovulatory granulosa cells. *Glycobiology* 11: 183–194.

Qi M, Ikematsu S, Ichihara-Tanaka K, Sakuma S, Muramatsu T, Kadomatsu K (2000). Midkine rescues Wilms' tumor cells from cisplatin-induced apoptosis: regulation of Bcl-2 expression by midkine. *J Biochem* 127: 269–277.

Qi M, Ikematsu S, Maeda N, Ichihara-Tanaka K, Sakuma S, Noda M *et al.* (2001). Haptotactic migration induced by midkine. Involvement of protein-tyrosine phosphatase zeta. Mitogen-activated protein kinase, and phosphatidylinositol 3-kinase. *J Biol Chem* 276: 15868–15875.

Rauvala H (1989). An 18-kd heparin-binding protein of developing brain that is distinct from fibroblast growth factors. *EMBO J* 8: 2933–2941.

Ren H, Tan ZP, Zhu X, Crosby K, Haack H, Ren JM *et al.* (2012). Identification of anaplastic lymphoma kinase as a potential therapeutic target in ovarian cancer. *Cancer Res* 72: 3312–3323.

Rizos D, Ward F, Duffy P, Boland MP, Lonergan P (2002). Consequences of bovine oocyte maturation, fertilization or early embryo development in vitro versus in vivo: implications for blastocyst yield and blastocyst quality. *Mol Reprod Dev* 61: 234–248.

Robinson RS, Woad KJ, Hammond AJ, Laird M, Hunter MG, Mann GE (2009). Angiogenesis and vascular function in the ovary. *Reproduction* 138: 869–881.

Salhab M, Dhorne-Pollet S, Auclair S, Guyader-Joly C, Brisard D, Dalbès-Tran R *et al.* (2013). In vitro maturation of oocytes alters gene expression and signaling pathways in bovine cumulus cells. *Mol Reprod Dev* 80: 166–182.

Sandra F, Harada H, Nakamura N, Ohishi M (2004). Midkine induced growth of ameloblastoma through MAPK and Akt pathways. *Oral Oncol* 40: 274–280.

Satoh J, Muramatsu H, Moretto G, Muramatsu T, Chang HJ, Kim ST *et al.* (1993). Midkine that promotes survival of fetal human neurons is produced by fetal human astrocytes in culture. *Brain Res Dev Brain Res* 75: 201–205.

- Seifer DB, Feng B, Shelden RM, Chen S, Dreyfus CF (2002). Brain-derived neurotrophic factor: a novel human ovarian follicular protein. *J Clin Endocrinol Metab* 87: 655–659.
- Shiota M, Tanihiro T, Nakagawa Y, Aoki N, Ishida N, Miyazaki K *et al.* (2003). Protein tyrosine phosphatase PTP20 induces actin cytoskeleton reorganization by dephosphorylating p190 RhoGAP in rat ovarian granulosa cells stimulated with follicle-stimulating hormone. *Mol Endocrinol* 17: 534–549.
- Skaper SD (2012). The neurotrophin family of neurotrophic factors: an overview. *Methods Mol Biol* 846: 1–12.
- Smits JE, Thompson JG, Gilchrist RB (2011). The promise of in vitro maturation in assisted reproduction and fertility preservation. *Semin Reprod Med* 29: 24–37.
- Stoica GE, Kuo A, Powers C, Bowden ET, Sale EB, Riegel AT *et al.* (2002). Midkine binds to anaplastic lymphoma kinase (ALK) and acts as a growth factor for different cell types. *J Biol Chem* 277: 35990–35998.
- Tanghe S, Van Soom A, Nauwynck H, Coryn M, de Kruif A (2002). Minireview: functions of the cumulus oophorus during oocyte maturation, ovulation, and fertilization. *Mol Reprod Dev* 61: 414–424.
- Thiele CJ, Li Z, McKee AE (2009). On Trk – the TrkB signal transduction pathway is an increasingly important target in cancer biology. *Clin Cancer Res* 15: 5962–5967.
- Tilly JL (1996). The molecular basis of ovarian cell death during germ cell attrition, follicular atresia, and luteolysis. *Front Biosci* 1: d1–11.
- Watson LN, Mottershead DG, Dunning KR, Robker RL, Gilchrist RB, Russell DL (2012). Heparan sulfate proteoglycans regulate responses to oocyte paracrine signals in ovarian follicle morphogenesis. *Endocrinology* 153: 4544–4555.
- Yamada M, Isaji Y, Ikeda S (2012). Midkine, a factor promoting cytoplasmic maturation of oocytes. In: Ergüven M, Muramatsu T, Bilir A (eds). *Midkine: From Embryogenesis to Pathogenesis and Therapy*. Springer: Heidelberg, pp. 183–193.
- Yarden Y, Sliwkowski MX (2001). Untangling the ErbB signalling network. *Nat Rev Mol Cell Biol* 2: 127–137.
- Yi KL, Zhou X, Shi DS, Chen HH, Qin QL, Chen Y *et al.* (2008). The mRNA expression of brain-derived neurotrophic factor in oocytes and embryos and its effects on the development of early embryos in cattle. *Animal* 2: 1786–1794.
- Yuan YQ, Van Soom A, Leroy JL, Dewulf J, Van Zeveren A, de Kruif A *et al.* (2005). Apoptosis in cumulus cells, but not in oocytes, may influence bovine embryonic developmental competence. *Theriogenology* 63: 2147–2163.
- Zhang CP, Yang JL, Zhang J, Li L, Huang L, Ji SY *et al.* (2011). Notch signaling is involved in ovarian follicle development by regulating granulosa cell proliferation. *Endocrinology* 152: 2437–2447.
- Zhang L, Jiang S, Wozniak PJ, Yang X, Godke RA (1995). Cumulus cell function during bovine oocyte maturation, fertilization, and embryo development in vitro. *Mol Reprod Dev* 40: 338–344.
- Zhang L, Li J, Su P, Xiong C (2010a). The role of brain-derived neurotrophic factor in mouse oocyte maturation in vitro. *J Huazhong Univ Sci Technolog Med Sci* 30: 781–785.
- Zhang L, Liang Y, Liu Y, Xiong CL (2010b). The role of brain-derived neurotrophic factor in mouse oocyte maturation in vitro involves activation of protein kinase B. *Theriogenology* 73: 1096–1103.
- Zhang N, Deuel TF (1999). Pleiotrophin and midkine, a family of mitogenic and angiogenic heparin-binding growth and differentiation factors. *Curr Opin Hematol* 6: 44–50.
- Zhao P, Qiao J, Huang S, Zhang Y, Liu S, Yan LY *et al.* (2011). Gonadotrophin-induced paracrine regulation of human oocyte maturation by BDNF and GDNF secreted by granulosa cells. *Hum Reprod* 26: 695–702.