Published in final edited form as:

Psychosom Med. 2009 June ; 71(5): 541–548. doi:10.1097/PSY.0b013e31819e7526.

Socioeconomic Status, Antioxidant Micronutrients, and Correlates of Oxidative Damage: The Coronary Artery Risk Development in Young Adults (CARDIA) Study

Denise Janicki-Deverts, Ph.D.,

Department of Psychology, Carnegie Mellon University, Pittsburgh, PA 15213; Phone: (412) 268-3295; Fax: (412) 268-3294

Sheldon Cohen, Ph.D.,

Department of Psychology, Carnegie Mellon University, Pittsburgh, PA 15213

Karen A. Matthews, Ph.D.,

Department of Psychiatry, University of Pittsburgh, Pittsburgh, PA 15213

Myron D. Gross, Ph.D., and

Department of Laboratory Medicine and Pathology, School of Medicine, University of Minnesota, Minneapolis, MN 55455

David R. Jacobs Jr., Ph.D.

Division of Epidemiology and Community Health, School of Public Health, University of Minnesota, Minneapolis, MN 55455, also affiliated with the Department of Nutrition, University of Oslo, Oslo, Norway

Abstract

Objective—Risk for chronic disease increases with decreasing socioeconomic status (SES). One pathway by which low SES might influence disease risk is by promoting oxidative stress. The aim of the present study is to examine whether SES (education, occupation, income), is associated both cross-sectionally and prospectively with circulating concentrations of (a) two correlates of oxidative damage, F₂-isoprostanes (F₂-IsoPs) and gamma-glutamyltransferase (GGT); and (b) antioxidant nutrients (ascorbic acid and carotenoids). We also examine whether the proposed associations are mediated by smoking, alcohol consumption and depression.

Methods—Data from 1278 participants in the Coronary Artery Risk Development in Young Adults (CARDIA) study were used to examine the association of SES with oxidation correlates and antioxidant nutrients. Education, occupation, health behaviors, and body mass index (BMI) were assessed during Years 0, 10, and 15 of the study; income and depression at Years 10 and 15. F₂-IsoPs were measured at Year 15, GGT at Years 0 and 10, carotenoids at Years 0 and 15, and ascorbic acid at Years 10 and 15.

Results—Cross-sectionally, oxidation correlates decreased and antioxidant nutrients increased with increasing SES, estimated in several ways, independent of age, sex, race, and BMI. Prospectively, lower Year 0 education and occupation predicted greater increases in GGT and greater decreases in carotenoids over 10–15 years. Prospective associations of Year 0 SES with Year 15 carotenoids were independent of Year 15 SES. Smoking, drinking, and depression symptoms partially mediated these effects.

Conclusions—Circulating oxidation correlates increase and antioxidant nutrients decrease with decreasing SES, both cross-sectionally and prospectively.

Keywords

socioeconomic status; oxidative stress; antioxidants; prospective; CARDIA

Chronic disease risk increases steadily with decreasing socioeconomic status (SES) (1). Lower SES might increase disease risk through promotion of behaviors, environments, and psychological states that contribute to oxidative stress, a steady state imbalance favoring pro- to anti-oxidants. Pro-oxidants are molecules with a single unpaired electron (free radicals) that have the potential to damage healthy cells while antioxidants are chemical compounds that bind to free radicals, preventing cell damage. When the balance is shifted in the direction of pro-oxidants, an unfavorable and potentially damaging systemic environment is created that surpasses the body's detoxification capacity and in turn contributes to organic pathology (2).

Increasing prevalence of certain health risk behaviors with decreasing SES may contribute to higher levels of oxidative stress among persons of lower relative to higher status. Rates of smoking (3) and alcoholism (4) both have been found to increase with decreasing SES. In turn, smoking has been associated with higher levels of oxidative stress (5) and chronic alcohol consumption with increased free radical concentrations (6, 7). Low SES also could influence oxidative stress through increased exposure to environmental toxins (hazardous wastes, pollution) at home and at work (8).

Finally, psychosocial factors associated with lower SES, such as depressive symptomatology (9), could also mediate the association of decreasing SES with increasing vulnerability to oxidative stress. Depression may influence oxidative stress directly via activation of the physiological stress axes. Metabolism of catecholamines, for example, results in production of free-radical intermediates which irreversibly bind to and damage cell constituents (10). It is possible that increases in catecholamines in response to stimulation of the sympatho-adrenal-medullary axis by depression may contribute to this source of pro-oxidant activity.

An important measure approximating oxidative stress *in vivo* is circulating F₂-isoprostane concentration (F₂-IsoP) (11). F₂-IsoPs are oxidatively-damaged molecules that result from free radical peroxidation of arachidonic acid. One feature of F₂-IsoPs that has contributed to their frequent use marking oxidative stress is that their concentrations depend largely on production rather than metabolism and excretion, thus suggesting they accurately reflect *in vivo* levels of oxidative damage (12). Empirical support for using F₂-IsoPs as oxidative stress markers derives from comparative animal research wherein F₂-IsoPs have been found to increase following experimental oxidant injury (13) and decrease following administration of antioxidants (14).

Gamma-glutamyltransferase (GGT) is an ectoplasmic enzyme produced by many cell types. Traditionally, elevated serum GGT has been used as a clinical marker of excessive alcohol consumption. However, GGT concentrations also have been found to correlate with levels of biologic exposure to other known pro-oxidant-producing toxins (15, 16). Thus, GGT may be an indirect marker of risk for future oxidative damage (17). An earlier report from CARDIA showed that serum GGT at recruitment and 10 years later correlated with circulating markers of inflammation and oxidative damage at 15 years post-recruitment (18).

Antioxidants can be categorized as endogenous antioxidant enzymes and non-enzymatic antioxidant molecules. Non-enzymatic antioxidants include the innate compound glutathione as well as exogenous antioxidant vitamins obtained through the diet, such as α -tocopherol, ascorbic acid, and beta-carotene. Circulating antioxidant levels relate inversely to markers of

oxidative stress and inflammation, both cross-sectionally (19) and prospectively (20). Serum ascorbic acid and carotenoid levels also have been found to correlate inversely with serum GGT (21, 22).

Several cross-sectional studies have found circulating antioxidant nutrient levels to vary directly with SES. Higher occupational class has been associated with higher ascorbic acid (23, 24) and beta-carotene (25) concentrations, and higher education with higher ascorbic acid (24) and lycopene (a carotenoid) (26) concentrations. Aside from the association of lower GGT with higher education at CARDIA baseline (18), no published studies have examined whether SES is related to circulating F₂-IsoPs or GGT.

The present study has two primary aims. First, we examine whether SES is associated *cross-sectionally* with concurrent concentrations of (a) two correlates of oxidative damage (F₂-IsoPs and GGT), and (b) two dietary nutrients with known antioxidant properties (carotenoids and ascorbic acid). We predict that GGT and F₂-IsoPs (hereafter referred to as *oxidation correlates*) will *increase* and antioxidant nutrients will *decrease* with decreasing SES. Second, we examine whether SES during early adulthood predicts *change* in two of these biomarkers (GGT and carotenoids) over time. Specifically, we expect lower SES to predict greater increases in GGT and greater decreases in carotenoids over 10–15 years.

Methods

Subjects

In 1985–1986, 5115 black and white adults (46% male), aged 18–30 years, were recruited into CARDIA at four sites: Birmingham, AL; Chicago, IL; Minneapolis, MN; and Oakland, CA. Recruitment was stratified on age (18–24 years, 25–30 years), race (black, white), sex, and education (< 12 years, ≥ 12 years). Participants were recruited by telephone in three of the localities and by telephone and door-to-door contact in the fourth. Overall response rates ranged from 51–82% (see (27, 28)). Participants were examined at study entry (Year 0) and Years 2, 5, 7, 10, 15, and 20. The current analyses focus on Year 0, and the years that oxidative stress markers were assessed (10 and 15). To obtain a single sample for all analyses, participant data were excluded from the present analyses if any of the following information was missing (see *Measures* below): Year 0, 10, or 15 education or occupational class; Year 10 or 15 household income; Year 0 or 10 GGT; Year 15 F₂-IsoPs; Year 0 or 15 carotenoids; Year 10 or 15 ascorbic acid. We also excluded data from persons who reported being students at Year 0, as these individuals were in the process of achieving their “educational attainment” and likely were not yet established in the workforce. The resulting sample after exclusions were made was $n = 1278$. Relative to the original 5115 baseline CARDIA participants, the present sample was older (mean age at recruitment = 26.06 vs. 24.85), less likely to be female (51.2% vs. 54.5%), and less likely to be black (43.1% vs. 51.6%). Alternative analyses that maximized sample sizes in individual analyses (different sample for each analysis) result in identical conclusions despite often substantially greater sample size. Site institutional review committee approval and informed consent were obtained. This manuscript has been approved by the CARDIA steering committee.

Measures

The Young Adult Longitudinal Trends in Antioxidant (YALTA) ancillary study has assayed various circulating antioxidants and related biochemicals at several times throughout the CARDIA study.

F₂-IsoPs

Plasma F₂-IsoPs were measured by gas chromatography-mass spectrometry of blood samples collected at Year 15. This method provides a measure of total free F₂-IsoPs, can detect plasma concentrations of 5–500 pg/ml, and has high specificity and sensitivity (2). The assay is reproducible with an intraindividual variation of <13% and an interassay coefficient of variation (CV) of <10%. Samples were stable for at least 8 years when stored at –80°C (mean and SD: 56.7±5.7 pg/ml in 2000–2002 (n=135) and 58.1±7.0 pg/mL in 2008 (n=9), with similar values throughout). F₂-IsoPs were measured at the Molecular Epidemiology and Biomarker Research Laboratory, University of Minnesota.

GGT

Blood for GGT measurement was collected at Years 0 and 10, and frozen at –70° C within 90 minutes of blood drawing. Frozen samples were stored for a few days at the CARDIA clinic center until being shipped to the laboratory for analysis or to the Solomon Park storage facility for long-term storage at –70°C. Year 0 serum GGT was measured with a SMAC 12 continuous-flow analyzer (Technicon Instruments Corp.) at American Bioscience Laboratories (now Smith-Kline Beecham), and Year 10 GGT was measured colorimetrically by a nitroanilide methodology at Linco Research, Inc. As GGT values were not compatible across the two measurement methods, 103 samples frozen for 16 years were reanalyzed. Year 0 values were recalibrated using the following formula (see (21)): recalibrated Year 0 GGT = (1.9004 × original Year 0 GGT) + 2.7618. GGT concentrations from both years were natural log-transformed to reduce skew.

Antioxidant nutrients

Carotenoids—Blood for measurement of serum carotenoids was collected at Years 0 and 15. Serum carotenoid concentrations (α-carotene, β-carotene, β-cryptoxanthin, zeaxanthin-plus-lutein, and lycopene) were assayed by high performance liquid chromatography (HPLC) and stored at –70°C. For analytic purposes, a *sum of carotenoids* score (hereafter referred to as *carotenoids*) was computed by summing the z-scores for concentrations of α- and β-carotene, β-cryptoxanthin, and zeaxanthin-plus-lutein. Lycopene was excluded from the summed score because previous CARDIA reports showed that associations of lycopene with other variables differed substantially from those involving the other four carotenoids (20, 29–31).

Ascorbic acid—Plasma ascorbic acid was measured using HPLC from blood collected at Years 10 and 15. Specimens were collected in vials containing metaphosphoric acid and frozen for up to one year at –70°C.

All antioxidants were measured at the Molecular Epidemiology and Biomarker Research Laboratory.

SES

Education—Education was measured by highest grade of regular school completed. At each follow-up exam, participants were asked to select one code from 01 to 20, with 01 indicating the first year of elementary school and 20 indicating 4 years of graduate education.

Occupational class—Occupation data were collected at each follow-up exam. We used the Stevens and Cho Socioeconomic Index (SEI) (32), as our measure of occupational class. SEI scores are “predicted prestige ratings” assigned to census occupation codes. Scores are derived from a regression of the estimated proportion of people rating the prestige of an

occupation as “good” or “excellent” based on (a) the proportion of people in that occupation with at least some college education and (b) the proportion whose previous year’s personal income was \$10,000 in 1980 U.S. dollars (32). For the present study, SEI scores were applied to census codes associated with participants’ current occupations as per the sociodemographic interviews conducted at each exam.

Income—Household income data were collected beginning at the third follow-up exam (Year 5), by which time it was assumed that all participants had moved out of their parents’ homes. Income was divided into eight categories, ranging from <\$5,000 to \$75,000. For analytic purposes, income was recoded to the midpoint of each of the eight categories (\$87,500 for the highest category). Recoded income data were adjusted for household size by dividing by the square-root of the number of persons living in the home (33), and for inflation by dividing by the consumer purchasing index (CPI) relative to 1990 (Year 5). Resulting adjusted income scores were natural log-transformed to reduce skew.

Smoking, alcohol consumption, and BMI

Measurement of BMI (weight in kg/[height in m]²) and self-reported health practices were obtained during each CARDIA visit. Self-report measures included smoker status, drinker status, smoking rate (number of cigarettes usually smoked per day), and alcohol consumption (number of drinks usually consumed per week; one drink = one 12-oz glass, bottle or can of beer, one 5-oz glass of wine, one 1.5-oz shot of liquor). Smoker and drinker statuses at each exam were represented by a 3-level categorical variable (0 = never, 1 = former, 2 = current). For purposes of analysis, never and former smokers and drinkers were assigned scores of “0” for smoking rate and alcohol consumption, respectively.

Depression Symptoms

Symptoms of depression were measured at Year 15 using the Center for Epidemiologic Studies Depression (CES-D) Scale (34). Scores were natural-log transformed to approximate a normal distribution.

Statistical analyses

Partial correlations were used to examine cross-sectional associations of education, occupational class, and household income with concurrent levels of oxidation correlates and antioxidant nutrients. All analyses controlled for a set of standard covariates: age, sex, race, and concurrent BMI.

Multiple regression was used to conduct lag analyses examining prospective associations of Year 0 SES with Year 10 GGT and Year 15 carotenoids. All lag models controlled for the standard covariates, as well as for GGT or carotenoids measured concurrently with the predictor (Year 0), thus examining the *change* in the dependent variable since its baseline. To determine whether prospective associations between SES and the outcome variables were independent of cross-sectional SES effects, a second set of prospective analyses was conducted that included control for SES measured concurrently with the outcomes.

Results

Sample characteristics

Sample characteristics are displayed in Table 1. As expected, all three SES indicators increased from Year 0 to Year 15. GGT concentrations increased from Year 0 to Year 10, as did carotenoid concentrations from Year 0 to Year 15. Ascorbic acid levels increased slightly between Years 10 and 15.

Intercorrelations of independent and dependent variables

Independent variables—Correlations among SES indicators measured during a given exam year ranged from $r = .39$ – $.64$. Between years, education was most stable ($r = .85$), followed by income ($r = .73$) and occupation ($r = .50$ – $.68$). All correlations were significant at $p < .001$.

Dependent variables—At Years 0 and 10, respectively, higher GGT concentrations were associated with lower carotenoids ($r = -.14$) and ascorbic acid ($r = -.18$). At Year 15, higher F₂-IsoPs also were associated with lower carotenoids ($r = -.36$) and ascorbic acid ($r = -.18$). GGT concentrations were correlated across Years 0 and 10 ($r = .65$), carotenoids across Years 0 and 15 ($r = .60$), and ascorbic acid across Years 10 and 15 ($r = .34$). All correlations were significant at $p < .001$.

Cross-sectional analyses

SES and oxidation correlates—Cross-sectional analyses were conducted for each year that oxidation correlates were collected. Education and income showed the strongest associations with GGT, with higher levels of each being associated with lower concurrent GGT (partial $r = -.08$ to $-.11$, $p < .003$). Occupation also demonstrated associations in the predicted direction, but achieved statistical significance only at Year 10 (Year 0: partial $r = -.05$, $p < .10$; Year 10: partial $r = -.07$, $p < .02$).

Findings for Year 15 F₂-IsoPs were similar to those reported for Year 0 and 10 GGT in that education and especially income showed stronger cross-sectional associations with F₂-IsoPs than did occupation (education: $r = -.08$, $p < .004$; income: $r = -.21$, $p < .001$; occupation: $r = -.04$, $p = .12$).

SES and antioxidant nutrients—As expected, higher education, occupation and income each were associated with higher concurrent carotenoid concentrations at Years 0 and 15 (partial $r = .13$ – $.28$, $p < .001$) and higher ascorbic acid concentrations at Years 10 and 15 (partial $r = .09$ – $.20$, $p < .002$).

Prospective analyses

With lag analyses we examined the prospective associations of Year 0 education and occupation with concentrations of (a) the oxidation correlate GGT at Year 10 and (b) carotenoids at Year 15. Models included control for age, sex, race, Year 0 GGT or carotenoids, and BMI concurrent with the outcome. Because income was not measured at Year 0, we could not examine 10- and 15-year prospective associations with income as the predictor.

GGT—Higher Year 0 education was associated with lower GGT levels at Year 10 ($b = -.07$, $SE = .007$, $p < .01$, partial $R^2 = .004$). When Year 10 education was included as a covariate, the prospective effect of Year 0 education was reduced to nonsignificance ($p = .14$). Higher Year 0 occupation also was associated with lower GGT at Year 10 ($b = -.06$, $SE = .001$, $p < .05$, partial $R^2 = .003$). Inclusion of Year 10 occupation in the model also resulted in a loss of statistical significance for Year 0 occupation, but a marginal trend remained apparent ($p < .08$). Neither Year 10 education nor Year 10 occupation emerged as independent predictors when entered simultaneously with the analogous Year 0 SES indicator ($p > .60$).

Carotenoids—As expected, higher Year 0 education predicted higher carotenoid concentrations at Year 15 ($\beta = .12$, $SE = .006$, $p < .001$, partial $R^2 = .011$). When Year 15

education was included as a covariate, the association of Year 0 education with Year 15 carotenoids was unaffected. Higher Year 0 occupation also predicted higher Year 15 carotenoids ($\beta = .13$, $SE = .001$, $p < .001$, partial $R^2 = .013$); and there was no change in effect size when additional control was added for Year 15 occupation. Neither Year 15 education nor Year 15 occupational class emerged as an independent predictor when entered simultaneously with the analogous Year 0 SES indicator ($p > .13$).

Mediation of lagged association of Year 0 SES with Year 15 carotenoids—

Smoking and alcohol consumption each have been associated with lower circulating antioxidant concentrations (35, 36). Antioxidant concentrations also have been found to decrease with increasing symptoms of depression (36). Thus, we examined whether smoking, alcohol consumption, and depression symptoms may have mediated the associations of Year 0 education and occupation with Year 15 carotenoids.

With univariate analyses, we examined whether each of the potential mediators was associated with Year 15 carotenoid concentrations. Results of univariate analyses indicated that smoker status and daily cigarette consumption at Years 0 and 15 each were related to Year 15 carotenoids ($r = -.27$ to $-.28$, $p < .001$). Drinker status at Year 15 was positively related to concurrent carotenoid levels ($r = .09$, $p < .01$); however, Year 0 drinker status was unrelated to Year 15 carotenoids ($p > .66$). Weekly alcohol consumption at Years 0 and 15 were negatively related to Year 15 carotenoids ($r = -.11$, $p < .001$), as were Year 15 depression symptoms ($r = -.19$, $p < .001$).

To determine whether levels of each of the potential mediators increased with decreasing SES, we examined univariate associations of SES indicators with each of the mediators. Lower Year 0 education and occupational class each were associated with greater likelihood of being a smoker and greater cigarette consumption at Years 0 and 15 ($p < .001$). Likewise, both Year 0 SES variables were associated with drinker status at Years 0 and 15 ($p < .001$), such that lower SES was associated with a greater likelihood of being a consumer of alcohol.

In separate models that included the standard covariates and Year 15 SES, we examined the reduction in association of Year 0 SES with Year 15 carotenoids when each of the potential mediators was included in the model. When examining smoking and drinking as mediators, we included measures from Year 0 and Year 15, simultaneously.

As indicated by Table 2, smoking explained roughly a quarter of the association of Year 0 education with Year 15 carotenoids, and depression symptoms explained about 12% of the association. Explanatory effects of drinking were virtually non-existent. To determine the degree of overlap between potential mediators in accounting for the association of Year 0 education with Year 15 carotenoids, we entered smoking, drinking and depression symptoms simultaneously into the model. As shown in the last row of Table 2, the combination of all three mediators accounted for nearly one-third of the association between Year 0 education and Year 15 carotenoids. Because this proportion of the association was less than the roughly 40% that would have been accounted for had each of the three mediators made independent contributions to the association, we can assume some overlap in effects.

As indicated by Table 3, smoking explained approximately 20% of the association of Year 0 occupation with Year 15 carotenoids; the proportion explained by drinking was much smaller (2%). Year 15 depression symptoms did not contribute to the association of Year 0 occupation with Year 15 carotenoids. When examined in combination, the mediating variables explained almost one-quarter of the association of Year 0 occupation with Year 15

carotenoids, suggesting that smoking and, to a lesser extent, drinking each accounted for a unique proportion of the association.

SES by oxidative stress vulnerability—In exploratory analyses, we examined whether SES might have a greater influence on future GGT levels among persons who already were vulnerable to oxidative stress. Specifically, we asked the question of whether SES effects on future risk for oxidative damage might be stronger among individuals with reduced antioxidant defense. Thus, we examined the interaction of Year 0 education and occupation, respectively, with Year 0 carotenoid levels in the prediction of Year 10 GGT. These models included the standard covariates, Year 0 GGT, Year 0 carotenoids, Year 0 SES and the interaction of Year 0 carotenoids and Year 0 SES. Results showed no moderating effect of Year 0 carotenoids on the association of either education or occupation with Year 10 GGT ($p > .80$).

Discussion

Lower SES, estimated in several ways, was associated cross-sectionally with higher levels of oxidation correlates and lower levels of antioxidant nutrients. Lower Year 0 education and occupation also predicted greater increases in GGT and decreases in carotenoids over 10–15 years. All associations were independent of age, sex, race, and BMI; prospective associations were independent of baseline measures of the outcomes.

Although there is some variability in the size of the associations of education, occupation, and income with each of the outcomes, all three indicators seem to play a role. That correlations of SES with concurrent carotenoids concentrations were larger than correlations of SES with ascorbic acid, GGT or F₂-IsoPs might be explained by differences in how each outcome was measured. Carotenoid concentrations were estimated by aggregating across four separate carotenoid measures (α -carotene, β -carotene, β -cryptoxanthin, zeaxanthin-plus-lutein), whereas each of the other outcome variables was estimated using only a single measurement. Therefore, the carotenoid measure may be a more reliable and more sensitive indicator of the underlying construct it is intended to represent (e.g., total antioxidant capacity) than are the ascorbic acid, GGT or F₂-IsoP measures. An alternative explanation may be that the differing contributions of SES to the variance in each of the outcomes are indicative of a real difference in the association of SES with antioxidant capacity and risk for oxidative damage.

In addition to the cross-sectional analyses, we conducted lag analyses examining the prospective association of Year 0 SES with Year 10 GGT and Year 15 carotenoids, respectively. Results suggest that Year 0 education and occupation each predict changes in GGT and carotenoids across 10–15 years, independent of Year 0 biomarker levels. Year 0 education and occupation, respectively, remained independent predictors of Year 15 carotenoids when controlling for Year 15 SES, whereas associations with Year 10 GGT lost significance when controlling for Year 10 SES.

We propose three possible models to explain these findings: (1) Year 0 SES influences GGT and carotenoid concentrations measured 10–15 years later, regardless of subsequent SES; (2) The *stability* of the association of SES with GGT and carotenoids over time is the factor driving the association of Year 0 SES with Year 10 and 15 biomarker levels; and (3) Year 10 and 15 SES may be the most important determinant of concurrent GGT and carotenoids, respectively; thus, the primary reason Year 0 SES correlates with later outcomes is that Year 0 SES is a strong predictor of future SES. That control for Year 15 SES did not eliminate the association of Year 0 SES with Year 15 carotenoids reduces the plausibility of the third model and thus suggests that earlier exposure was key. By comparison, associations of Year

0 SES with Year 10 GGT lost significance when Year 10 SES was controlled. These findings, however, also fail to support the hypothesis that this association is primarily attributable to concurrent SES because neither Year 10 SES indicator emerged as an independent correlate of Year 10 GGT.

We examined the possibility that smoking, alcohol consumption, and depression symptoms might account for some of the prospective association of Year 0 education and occupation with Year 15 carotenoid concentrations. Neither singly nor in combination did the three proposed mediators explain the entire association (though the association of Year 0 education with carotenoids lost statistical significance). However, simultaneous inclusion of the three proposed mediators substantially reduced the variance in Year 15 carotenoids accounted for by Year 0 SES, with much of that reduction being attributable to the effects of smoking, with smaller reductions attributable to drinking and depression symptoms.

The results of these analyses suggest that smoking, drinking, and depression symptoms may comprise three potential pathways by which SES influences future antioxidant nutrient concentrations. Smoking may relate directly to reductions in antioxidants as exposure of human plasma *in vitro* to gaseous phase tobacco smoke has been found to result in an overall degradation in lipophilic antioxidant nutrients including all four of the carotenoids examined here (37). Both smoking and excessive alcohol consumption might indirectly influence carotenoid concentrations by promoting the generation of free radicals (5, 7). Because circulating carotenoids are known scavengers of free radicals (38), it is possible that excess generation of these pro-oxidant species by damaging properties of tobacco smoke and metabolism of ethanol may deplete available carotenoid stores. Depression has been associated increased production of pro-inflammatory cytokines (39), which in turn lead to production of pro-oxidant species (40). Thus depression symptoms, as well, may indirectly influence carotenoid concentrations via promotion of oxidative stress.

Because SES remained an independent predictor of carotenoid concentrations after controlling for previous carotenoid levels, and concurrent smoking, drinking, and depression symptoms, one might infer that additional factors not examined here may further explain the association. For example, exposure to environmental tobacco smoke has been associated with lower circulating antioxidant concentrations (41, 42). Thus, insofar as lower SES persons, regardless of their own smoking status, are more likely to share living or workspace with smokers, they may be at increased risk of oxidative stress. Poor dietary habits, especially low fruit and vegetable intake, can directly influence circulating antioxidant nutrient concentrations. The Year 0 cross-sectional association of higher fruit and vegetable intake with increased carotenoid levels has previously been demonstrated in CARDIA (30). However, when included in the multivariate lag analyses reported here, Year 0 fruits and vegetables were unrelated to Year 15 carotenoids, and thus, did not reduce the prospective association of Year 0 SES with Year 15 carotenoids (data not shown). It is likely that dietary habits at Year 15 would have demonstrated a stronger association with Year 15 carotenoids. These data, however, are not available.

Though independent in association, SES explained a rather small amount of the variance in biomarker concentrations, both cross-sectionally and prospectively. In regard to cross-sectional analyses, partial R^2 ranged from .002 to .044 for oxidative correlates and from .008 to .078 for antioxidant nutrients. By comparison, the more conservative prospective analyses—that controlled for the cross-sectional associations with Year 15 carotenoids, yielded smaller effects (partial R^2 = .003 to .008). These latter values are much smaller than those associated with the contributions of smoking at Years 0 and 15 to Year 15 carotenoids (univariate analyses: r^2 = .07–.08; analyses including standard controls and SES: partial R^2 = .01–.02), which are comparable to that which previously have been reported (35). Despite

the relative smallness of effect when compared with that associated with smoking, the effects of SES on carotenoid levels may still be of substantial importance when considered in terms of public health. Even modest differences in levels of a given risk factor can make a large difference in population attributable risk, as opposed to relative risk. Accordingly, were the entire population to shift downward in SES by even a small amount, the impact on antioxidant capacity and ultimately oxidative damage, could be considerable.

There are limitations to the present study. First, identical circulating antioxidants were not measured at all years. Likewise, F₂-IsoPs were measured only at Year 15. Thus, we were unable to examine prospective associations of Year 0 SES with ascorbic acid or F₂-IsoPs. Second, levels for each oxidative correlate and anti-oxidant marker were based on single measures at the relevant follow-ups. Because of variability in actual levels and in the assays, this measurement strategy could result in underestimates of the real effect sizes. Third, because Year 0 income data were not available, we could not examine 10- and 15-year prospective associations of income with oxidation correlates and antioxidant nutrients.

In conclusion, lower SES was associated both cross-sectionally and prospectively with higher oxidation correlate concentrations, and lower antioxidant nutrient levels. Prospective analyses controlling for concurrent SES suggest that education and occupational class during early adulthood may affect antioxidant levels measured 15 years later independently of current status. Impressively, the effects reported in this article were consistent across cross-sectional (at different exam years) and prospective analyses. It is not clear whether the obtained effect sizes are clinically significant. However, the very existence of the consistent associations reported here, regardless of their size, invite further examination of a possible link between SES and oxidative stress. Refinements in the conceptualization and measurement of both SES and oxidative stress may enable the detection of stronger associations between socioeconomic factors and risk for oxidative damage.

Acknowledgments

Work on this manuscript was supported by contracts University of Alabama at Birmingham, Coordinating Center, N01-HC-95095 University of Alabama at Birmingham, Field Center, N01-HC-48047 University of Minnesota, Field Center, N01-HC-48048 Northwestern University, Field Center, N01-HC-4049 Kaiser Foundation Research Institute, N01-HC-48050 University of California, Irvine, Echocardiography Reading Center, N01-HC-45134 Harbor-UCLA research Education Institute, and Computed Tomography Reading Center, N01-HC-05187 from the National Heart, Lung and Blood Institute and by the MacArthur Research Network on SES and Health through grants from the John D. and Catherine T. MacArthur Foundation. Preparation of the manuscript also was facilitated by the Pittsburgh Mind-Body Center (HL65111 & HL65112).

Abbreviations

BMI	body mass index
CARDIA	Coronary Artery Risk Development in Young Adults study
GGT	gamma-glutamyltransferase
F₂-IsoPs	F ₂ -isoprostanes
SEI	Stevens & Cho Socioeconomic Index
SES	socioeconomic status

References

1. Adler NE, Boyce T, Chesney MA. Socioeconomic status and health: The challenge of the gradient. *Am Psychol.* 1994; 49:15–24. [PubMed: 8122813]

2. Sies, H. Oxidative stress: Oxidants and antioxidants. New York, NY: Academic Press; 1991.
3. Barbeau EM, Krieger N, Soobader MJ. Working class matters: Socioeconomic disadvantage, race/ethnicity, gender, and smoking in NHIS 2000. *Am J Public Health*. 2004; 94:269–278. [PubMed: 14759942]
4. Hemmingsson T, Lundberg I, Diderichsen F, Allebeck P. Explanations of social class differences in alcoholism among young men. *Soc Sci Med*. 1998; 47:1399–1405. [PubMed: 9823036]
5. Reilly M, Delanty N, Lawson JA, FitzGerald GA. Modulation of oxidant stress in vivo in chronic cigarette smokers. *Circulation*. 1996; 94:19–25. [PubMed: 8964113]
6. Husain K. Vascular endothelial oxidative stress in alcohol-induced hypertension. *Cell Mol Biol*. 2007; 53:70–77. [PubMed: 17519114]
7. Zima T, Fialova L, Mestek O, Janebova M, Crkovska J, Malbohan I, Stipek S, Mikulikova L, Popov P. Oxidative stress, metabolism of ethanol and alcohol-related diseases. *Journal of Biomedical Science*. 2001; 8:59–70. [PubMed: 11173977]
8. Evans GW, Kantrowitz E. Socioeconomic status and health: the potential role of environmental risk exposure. *Annu Rev Public Health*. 2002; 23:303–331. [PubMed: 11910065]
9. Kessler RC, McGonagle KA, Zhao S, Nelson CB, Hughes M, Eshleman S, Wittchen HU, Kendler KS. Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States. Results from the National Comorbidity Survey. *Arch Gen Psychiatry*. 1994; 5:8–19. [PubMed: 8279933]
10. Bindoli A, Rigobello MP, Deeble DJ. Biochemical and toxicological properties of the oxidation products of catecholamines. *Free Radic Biol Med*. 1992; 13:391–405. [PubMed: 1398218]
11. Roberts LJ, Morrow JD. Measurement of F2-isoprostanes as an index of oxidative stress in vivo. *Free Radic Biol Med*. 2000; 28:505–513. [PubMed: 10719231]
12. Morrow JD. Quantification of isoprostanes as indices of oxidant stress and the risk of atherosclerosis in humans. *Arterioscler Thromb Vasc Biol*. 2005; 25:279–286. [PubMed: 15591226]
13. Morrow JD, Hill KE, Burk RF, Nammour TM, Badr KF, Roberts LJ. A series of prostaglandin F2-like compounds are produced in vivo in humans by a non-cyclooxygenase, free radical-catalyzed mechanism. *Pro Natl Acad Sci USA*. 1990; 87:9383–9387.
14. MacDonald-Wicks LK, Garg ML. Vitamin E supplementation in the mitigation of carbon tetrachloride induced oxidative stress in rats. *J Nutr Biochem*. 2003; 14:211–218. [PubMed: 12770645]
15. Hu SW, Cheng TJ, ChangChien GP, Chan CC. Association between dioxins/furans exposures and incinerator workers' hepatic function and blood lipids. *J Occup Environ Med*. 2003; 45:601–608. [PubMed: 12802213]
16. Lee DH, Jacobs DRJ. Association between serum concentrations of persistent organic pollutants and gamma glutamyltransferase: results from the National Health and Examination Survey 1999–2002. *Clin Chem*. 2006; 52:1825–1827. [PubMed: 16940464]
17. Lee DH, Blomhoff R, Jacobs DRJ. Is serum gamma glutamyltransferase a marker of oxidative stress? *Free Radic Res*. 2004; 38:535–539. [PubMed: 15346644]
18. Lee DH, Jacobs DRJ, Gross M, Kiefe CI, Roseman J, Lewis CE, Steffes M. [gamma]-Glutamyltransferase is a predictor of incident diabetes and hypertension: The CARDIA Study. *Clin Chem*. 2003; 49:1358–1366. [PubMed: 12881453]
19. van Herpen-Broekmans WM, Klopping-Ketelaars IA, Bots ML, Kluft C, Princen H, Hendriks HF, Tijburg LB, van Poppel G, Kardinaal AF. Serum carotenoids and vitamins in relation to markers of endothelial function and inflammation. *Eur J Epidemiol*. 2004; 19:915–921. [PubMed: 15575349]
20. Hozawa A, Jacobs DRJ, Steffes MW, Gross MD, Steffen LM, Lee DH. Relationships of circulating carotenoid concentrations with several markers of inflammation, oxidative stress, and endothelial dysfunction: The Coronary Artery Risk Development in Young Adults (CARDIA)/Young Adult Longitudinal Trends in Antioxidants (YALTA) study. *Clin Chem*. 2007; 53:447–455. [PubMed: 17234732]

21. Lee DH, Gross MD, Jacobs DRJ. Association of serum carotenoids and tocopherols with gamma-glutamyltransferase: The Cardiovascular Risk Development in Young Adults (CARDIA) Study. *Clin Chem*. 2004; 50:582–588. [PubMed: 14726472]
22. Lim JS, Yang JH, Chun BY, Kam S, Jacobs DRJ, Lee D. Is serum gamma-glutamyltransferase inversely associated with serum antioxidants as a marker of oxidative stress? *Free Radic Biol Med*. 2004; 37:1018–1023. [PubMed: 15336318]
23. Wrieden WL, Hannah MK, Bolton-Smith C, Tavendale R, Morrison C, Tunstall-Pedoe H. Plasma vitamin C and food choice in the third Glasgow MONICA population survey. *J Epidemiol Community Health*. 2000; 54:355–360. [PubMed: 10814656]
24. Shohaimi S, Bingham S, Welch A, Luben R, Day N, Wareham N, Khaw KT. Occupational social class, educational level and area deprivation independently predict plasma ascorbic acid concentration: A cross-sectional population based study in the Norfolk cohort of the European Prospective Investigation into Cancer (EPIC-Norfolk). *Eur J Clin Nutr*. 2004; 58:1432–1435. [PubMed: 15054419]
25. Kristenson M, Kucinskiene Z, Bergdahl B, Orth-Gomer K. Risk factors for coronary heart disease in different socioeconomic groups of Lithuania and Sweden--the LiVicordia Study. *Scand J Public Health*. 2001; 29:140–150. [PubMed: 11484867]
26. Ganji V, Kafai MR. Population determinants of serum lycopene concentrations in the United States: Data from the Third National Health and Nutrition Examination Survey, 1988–1994. *J Nutr*. 2005; 135:567–572. [PubMed: 15735095]
27. Cutter GR, Burke GL, Dyer AR, Friedman GD, Hilner JE, Hughes GH, Hulley SB, Jacobs DRJ, Liu K, Monolio TA, Oberman A, Perkins LL, Savage PJ, Serwitz JR, Sidney S, Wagenknecht LE. Cardiovascular risk factors in young adults: The CARDIA baseline monograph. *Control Clin Trials*. 1991; 12:1s–77s. [PubMed: 1851696]
28. Friedman GD, Cutter GR, Donahue RP, Hughes GH, Hulley SB, Jacobs DRJ, Liu K, Savage PJ. CARDIA: Study design, recruitment, and some characteristics of the examined subjects. *J Clin Epidemiol*. 1988; 41:1105–1116. [PubMed: 3204420]
29. Andersen LF, Jacobs DRJ, Gross MD, Schreiner P, Williams OD, Lee DH. Longitudinal associations between body mass index and serum carotenoids: The CARDIA study. *Br J Nutr*. 2006; 92:358–365. [PubMed: 16469154]
30. Hozawa A, Jacobs DRJ, Steffes MW, Gross MD, Steffen LM, Lee DH. Associations of serum carotenoid concentrations with the development of diabetes and with insulin concentration: interaction with smoking: The Coronary Artery Risk Development in Young Adults (CARDIA) Study. *Am J Epidemiol*. 2006; 163:929–937. [PubMed: 16597706]
31. Ohira T, Hozawa A, Iribarren C, Daviglus ML, Matthews KA, Gross MD, Jacobs DRJ. Longitudinal association of serum carotenoids and tocopherols with hostility: The CARDIA study. *Am J Epidemiol*. 2007 Epub ahead of print: October 10.
32. Stevens G, Cho JH. Socioeconomic indices and the new 1980 census occupational classification scheme. *Soc Sci Res*. 1985; 14:142–168.
33. Buhmann B, Rainwater L, Schmaus G, Smeeding TM. Equivalence scales, well-being, inequality, and poverty: Sensitivity estimates across ten countries using the Luxembourg Income Study (LIS) database. *Rev Income Wealth*. 1988; 34:115–142.
34. Radloff LS. The CES-D Scale: A self-report depression scale for research in the general population. *Appl Psychol Meas*. 1977; 1:385–401.
35. Rust P, Lehner P, Elmadfa I. Relationship between dietary intake, antioxidant status and smoking habits in female Austrian smokers. *Eur J Nutr*. 2001; 40:78–83. [PubMed: 11518203]
36. Tsuboi H, Tatsumi A, Yamamoto K, Kobayashi F, Shimoi K, Kinae N. Possible connections among job stress, depressive symptoms, lipid modulation and antioxidants. *J Affect Disord*. 2006; 91:63–70. [PubMed: 16430969]
37. Handelman GJ, Packer L, Cross CE. Destruction of tocopherols, carotenoids, and retinol in human plasma by cigarette smoke. *J Am Diet Assoc*. 1996; 96:693–702. [PubMed: 8675913]
38. El-Agamey A, Lowe GM, McGarvey DJ, Mortensen A, Phillip DM, Truscott TG, Young AJ. Carotenoid radical chemistry and antioxidant/pro-oxidant properties. *Arch Biochem Biophys*. 2004; 430:37–48. [PubMed: 15325910]

39. Maes, M. The immune pathophysiology of major depression. In: Honig, A.; Praag, HMv, editors. *Depression: Neurobiological, psychopathological and therapeutic advances*. London: John Wiley; 1997. p. 197-215.
40. Maziere C, Auclair M, Maziere JC. Tumor necrosis factor enhances low density lipoprotein oxidative modification by monocytes and endothelial cells. *FEBS Lett*. 1994; 338:43–46. [PubMed: 8307154]
41. Farchi S, Forastiere F, Pistelli R, Baldacci S, Simoni M, Perucci C, Viegi G. Exposure to environmental tobacco smoke is associated with lower plasma beta-carotene levels among nonsmoking women married to a smoker. *Cancer Epidemiol Biomarkers Prev*. 2001; 10:907–909. [PubMed: 11489760]
42. Dietrich M, Block G, Norkus EP, Hudes M, Traber MG, Cross CE, LP. Smoking and exposure to environmental tobacco smoke decrease some plasma antioxidants and increase gamma-tocopherol in vivo after adjustment for dietary antioxidant intakes. *Am J Clin Nutr*. 2003; 77:160–166. [PubMed: 12499336]

Table 1Sample characteristics^a.

	Year 0	Year 10	Year 15
% smokers			
Current	27.78 (355)	24.80 (317)	22.07 (282)
Former	14.24 (182)	19.72 (252)	22.85 (292)
Never	57.98 (741)	55.48 (709)	55.09 (704)
Median daily cigarette consumption	12.0 (5.0–20.0)	10.0 (5.0–20.0)	10.0 (5.0–20.0)
% consumers of alcohol			
Current	87.95 (1124)	80.36 (1027)	79.97 (1022)
Former	5.48 (70)	15.88 (203)	16.35 (209)
Never	6.57 (84)	3.76 (48)	3.68 (47)
Median weekly alcohol consumption	3.0 (0–8.0)	3.0 (0–8.0)	3.0 (0–8.0)
Median BMI (kg/m ²)	23.54 (21.48–26.30)	26.11 (23.32–30.00)	27.33 (24.16–31.78)
Median depression symptoms	NA	8.0 (0–56)	7.0 (0–54)
Median education (years)	14.0 (12.0–16.0)	15.0 (13.0–16.0)	15.0 (13.0–16.0)
Median occupational class	31.05 (22.58–54.42)	36.84 (23.88–52.99)	46.27 (25.23–54.48)
Median income (thousands \$US) ^b	NA	21.81 (12.61–36.32)	27.34 (15.80–46.87)
Median GGT (μg/ml)	8.00 (5.00–12.00)	18.00 (12.00–28.00)	NA
Median F ₂ -IsoPs (pg/ml)	NA	NA	50.60 (38.34–69.52)
Median carotenoids (μg/dl)	40.95 (29.31–56.47)	NA	55.74 (38.65–78.60)
Median ascorbic acid (mg/L)	NA	8.43 (5.86–10.68)	8.55 (5.90–11.12)

^aN = 1278; Values displayed as percent (*n*) or median (interquartile range)^bAdjusted for CPI relative to 1990 and number of persons living in the household.

Table 2

Change in association of Year 0 education with Year 15 carotenoids when potential mediators are included in the model.

	Association with Year 15 Carotenoids				
	<i>B</i>	<i>SEB</i>	β	$\Delta\beta$	Partial R^2
Year 0 education Standard covariates	.027	.010	.116**	-	.003
Year 0 education Standard covariates Year 0 & 15 drinker status and alcohol consumption	.028	.010	.117***	<.10%	.004
Year 0 education Standard covariates Year 0 & 15 smoker status and cigarette consumption	.020	.010	.085*	-27%	.002
Year 0 education Standard covariates Year 15 depression symptoms	.024	.010	.102*	-12%	.003
Year 0 education Standard covariates Year 0 & 15 drinker status and alcohol consumption Year 0 & 15 smoker status and cigarette consumption Year 15 depression symptoms	.019	.010	.079 [†]	-32%	.002

[†] $p < .06$.

* $p < .05$.

** $p < .01$.

*** $p < .001$.

Standard covariates: age, sex, race, Year 15 BMI, Year 0 carotenoids, Year 15 education

$\Delta\beta$ = change in β in comparison to the base model containing Year 0 education and the standard covariates

Table 3

Change in association of Year 0 occupation with Year 15 carotenoids when potential mediators are included in the model.

	Association with Year 15 Carotenoids			
	<i>B</i>	<i>SEB</i>	β	$\Delta\beta$ Partial <i>R</i> ²
Year 0 occupation Standard covariates	.003	.001	.108***	- .008
Year 0 occupation Standard covariates Year 0 & 15 drinker status and alcohol consumption	.003	.001	.106***	-2% .007
Year 0 occupation Standard covariates Year 0 & 15 smoker status and cigarette consumption	.002	.001	.085**	-21% .005
Year 0 occupation Standard covariates Year 15 depression symptoms	.003	.001	.107***	<.10% .008
Year 0 occupation Standard covariates Year 0 & 15 drinker status and alcohol consumption Year 0 & 15 smoker status and cigarette consumption Year 15 depression symptoms	.002	.001	.082**	-24% .004

[†] $p < .06$.

* $p < .05$.

** $p < .01$.

*** $p < .001$.

Standard covariates: age, sex, race, Year 15 BMI, Year 0 carotenoids, Year 15 occupational class