You Still Need More than CPAP for OSA Patients to Lose Weight

Response to Mysliwiec et al. Weight gain with CPAP: a complication of treatment?

J Clin Sleep Med 2014;10:347.

Stuart F. Quan, M.D., F.A.A.S.M.^{1,2}; Rohit Budhiraja, M.D., F.A.A.S.M.²; Denise P. Clarke, B.Sc.¹; James L. Goodwin, Ph.D.²; Daniel J. Gottlieb, M.D., M.P.H., F.A.A.S.M.^{1,6}; Deborah A. Nichols, M.S.³; Richard D. Simon, M.D., F.A.A.S.M.⁴; Terry W. Smith, B.S.²; James K. Walsh, Ph.D., F.A.A.S.M.⁵; Clete A. Kushida, M.D., F.A.A.S.M.³; Barbara Phillips, M.D., M.S.P.H., F.A.A.S.M.⁷

¹Division of Sleep Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA;

²Arizona Respiratory Center, University of Arizona, Tucson, AZ; ³Stanford, University, Stanford, CA; ⁴Providence St. Mary Medical Center, Walla Walla, WA; ⁵St. Luke's Hospital, Chesterfield, MO; ⁶VA Boston Healthcare System, Boston, MA;

⁷Division of Pulmonary, Critical Care and Sleep Medicine, University of Kentucky College of Medicine, Lexington, KY

In their comment regarding our recent paper demonstrating that persons with obstructive sleep apnea (OSA) gain weight after being treated with continuous positive airway pressure (CPAP) and the accompanying editorial,^{1,2} Mysliwiec and colleagues suggest that the observed increase in weight was related to an increase in lean body mass (LBM) rather than adipose tissue.³ Consequently, they contend that weight gain with CPAP may actually represent a beneficial effect.³ Inasmuch as we did not perform body composition studies on the participants in our study, we cannot confirm or refute whether an increase in LBM, adipose tissue, or both occurred.

Although we agree that CPAP can be associated with an increase in LBM in some individuals,^{4,5} it needs to be emphasized that the vast majority of persons with OSA are overweight if not frankly obese.⁶ Most, but not all studies report that visceral fat does not decrease with CPAP use.^{4,5,7-9} Moreover, in the study by Muntzer et al.,⁴ there was a trend for an increase in visceral fat with CPAP, although it did not reach statistical significance. However, waist circumference⁴ and subcutaneous fat⁵ increased. Yet, whether weight gain induced by CPAP in persons with OSA is reflected by increased LBM or adipose tissue is not as important as the failure to observe any weight reduction. A recent meta-analysis indicates that even metabolically healthy obese persons are at increased risk for cardiovascular disease and death, suggesting a "healthy obesity" phenotype does not exist.¹⁰ Weight reduction is recommended in all persons with OSA who are overweight and can result in remission or reduction in severity. Therefore, our findings still support the conclusion that CPAP alone will not lead to weight reduction, and that clinicians need to recommend or prescribe additional interventions to facilitate weight loss in their patients with OSA.

CITATION

Quan SF; Budhiraja R; Clarke DP; Goodwin JL; Gottlieb DJ; Nichols DA; Simon RD; Smith TW; Walsh JK; Kushida CA; Phillips B. You still

need more than CPAP for OSA patients to lose weight. *J Clin Sleep Med* 2014;10(3):349.

REFERENCES

1. Quan SF, Budhiraja R, Clarke DP, et al. Impact of treatment with continuous positive airway pressure (CPAP) on weight in obstructive sleep apnea. *J Clin Sleep Med* 2013;9:989-93.
2. Phillips B, Dhaon N. Weigh the options before starting CPAP. *J Clin Sleep Med* 2013;9:995-6.
3. Mysliwiec V, O'Reilly B, Roth BJ. Weight gain with CPAP: a complication of treatment? *J Clin Sleep Med* 2014;10:347.
4. Munzer T, Hegglin A, Stannek T, et al. Effects of long-term continuous positive airway pressure on body composition and IGF1. *Eur J Endocrinol* 2010;162:695-704.
5. Hoyos CM, Killick R, Yee BJ, Phillips CL, Grunstein RR, Liu PY. *Thorax* 2012;67:1081-9.
6. Young T, Peppard PE, Taheri S. Excess weight and sleep-disordered breathing. *J Appl Physiol* 2005;99:1592-9.
7. Sivam S, Phillips CL, Trenell MI, et al. Effects of 8 weeks of continuous positive airway pressure on abdominal adiposity in obstructive sleep apnoea. *Eur Respir J* 2012;40:913-8.
8. Trenell MI, Ward JA, Yee BJ, et al. Influence of constant positive airway pressure therapy on lipid storage, muscle metabolism and insulin action in obese patients with severe obstructive sleep apnoea syndrome. *Diabetes Obes Metab* 2007;9:679-87.
9. Kritikou K, Basta M, Tappouni R, et al. Sleep apnoea and visceral adiposity in middle-aged male and female subjects. *Eur Respir J* 2013;41:601-9.
10. Kramer CK, Zinman B, Retnakaran R. Are metabolically healthy overweight and obesity benign conditions? a systematic review and meta-analysis. *Ann Intern Med* 2013;159:758-69.

SUBMISSION & CORRESPONDENCE INFORMATION

Submitted for publication December, 2013

Accepted for publication January, 2014

Address correspondence to: Stuart F. Quan, M.D., Division of Sleep Medicine, Harvard Medical School, 401 Park Dr., 2nd Floor East, Boston, MA 02215; Tel: (617) 998-8842; Fax: (617) 998-8823; Email: Stuart.Quan@hms.harvard.edu

DISCLOSURE STATEMENT

Dr. Quan is the Editor-in-Chief of the *Journal of Clinical Sleep Medicine*. The other authors have indicated no financial conflicts of interest.