Published in final edited form as:

Antiviral Res. 2013 October ; 100(1): 14–19. doi:10.1016/j.antiviral.2013.07.002.

The antimicrobial agent C31G is effective for therapy for HSV-1 ocular keratitis in the rabbit eye model

James M. Hill^{1,2,3,4}, Ethan M. Stern^{1,5}, Partha S. Bhattacharjee^{1,6}, Daniel Malamud⁷, Christian Clement¹, Paulo Rodriguez^{2,8}, Walter J. Lukiw^{1,3,9}, Augusto C. Ochoa^{8,10}, Timothy P. Foster^{2,8}, Cruz Velasco¹¹, and Harris E. McFerrin Jr.⁶

James M. Hill: jhill@lsuhsc.edu; Ethan M. Stern: ester1@lsuhsc.edu; Partha S. Bhattacharjee: pbhattach@xula.edu; Daniel Malamud: daniel.malamud@nyu.edu; Christian Clement: cclm1@lsuhsc.edu; Paulo Rodriguez: prodri1@lsuhsc.edu; Walter J. Lukiw: wlukiw@lsuhsc.edu; Augusto C. Ochoa: aochoa@lsuhsc.edu; Timothy P. Foster: tfoste@lsuhsc.edu; Cruz Velasco: cvelas@lsuhsc.edu; Harris E. McFerrin: hmcferri@xula.edu

¹Louisiana State University Health Sciences Center (LSUHSC) Department of Ophthalmology, 533 Bolivar Street New Orleans, LA 70112

²Department of Microbiology, LSUHSC

³Neuroscience Center, LSUHSC, 2020 Gravier Street, 8th Floor, New Orleans, LA 70112

⁴Department of Pharmacology, LSUHSC, 1901 Perdido Street, New Orleans, LA 70112

⁵Tulane University, 6823 St. Charles Avenue, New Orleans, LA 70118

⁶Department of Biology, Xavier University of Louisiana, 1 Drexel Drive, New Orleans, LA 70125

⁷Department of Basic Sciences, NYU College of Dentistry, 345 East 24th St, Room 904-S, New York, NY 10010

⁸Stanley S. Scott Cancer Center, LSUHSC, 533 Bolivar Street, Room 459, New Orleans, LA 70112

⁹Department of Neurology, LSUHSC, 1542 Tulane Avenue, New Orleans, LA 70112

¹⁰Department of Pediatrics, LSUHSC, 200 Henry Clay Avenue, New Orleans, LA 70118

¹¹School of Public Health, LSUHSC, 2020 Gravier St, New Orleans, LA 70112

1. Summary

The amphoteric C31G solution contains equimolar alkyl dimethylglycine and alkyl dimethyl amine oxide buffered with citric acid. C31G acts as a broad spectrum antiviral and an antibacterial. No previous *in vivo* studies have been done to test C31G in an animal model of HSV-1 ocular keratitis. We assessed the anti-herpetic activity of C31G in the rabbit eye model using three treatment groups: (1) 1% trifluorothymidine (TFT); (2) 0.25% C31G plus 0.5% hydroxypropyl methylcellulose (HPMC); and (3) 0.5% HPMC. Scarified rabbit corneas were inoculated with the HSV-1 strain McKrae. On post inoculation (PI) day 3, rabbits were placed in three balanced groups based on slit-lamp examination (SLE) scores. Treatment began on PI day 3, five times a day for five consecutive days. In addition to the daily, masked SLE scoring, the eyes

© 2013 The Authors. Published by Elsevier B.V. All rights reserved.

Corresponding Author: James M. Hill, PhD, Department of Ophthalmology LSUHSC School of Medicine, 533 Bolivar Street, Suite B, New Orleans, LA 70112.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

were assessed daily for stromal opacity, scleral inflammation, neovascularization, eyelid inflammation, inflammatory discharge, and epiphora. C31G and TFT were very effective in reducing the lesions and pathogenesis associated with HSV-1 ocular keratitis. The vehicle control SLE scores were significantly higher, proving that the vehicle did not effectively treat HSV-1 keratitis. C31G has valuable potential to treat herpetic keratitis as well as other herpetic topical lesions in humans.

Keywords

Anti-herpetic; C31G; HSV-1 keratitis; Novel anti-herpetic; Novel anti-viral; Rabbit

2. Introduction

C31G is a broad spectrum, antimicrobial and antiviral agent (Calis et al. 1992; Corner et al. 1988; Thompson et al. 1996) composed of a mixture of an alkyl dimethylglycine (alkyl betaine) and an alkyl dimethyl amino oxide, buffered with citric acid to pH 5.5–6. This unique agent is potent *in vitro* against gram negative and gram positive bacteria, enveloped viruses, and numerous fungi and yeast (including *Candida*). Since both components in C31G are amphoteric, surface active chemicals, we suggest the mechanism of action involves the binding of the polar head of the group of the amino oxide betaine mixture to the pathogen surface. This subsequently results in a disruption of the membrane by the alkyl portion of the molecules thus killing the infectious agent.

C31G, in addition to antimicrobial activity, was demonstrated in human studies to be an effective, acceptable, and safe spermicidal agent (Burke et al. 2010). Thus, C31G has dual activity, reducing infectivity of enveloped virus and certain other sexually transmitted disease organisms and application as a safe and well-tolerated spermicidal agent.

The present study assessed the safety and tolerance of the buffered formulation of C31G in the rabbit eye model and the eyes of squirrel monkeys. C31G has been shown to be safe and well-tolerated as a vaginal gel in humans (Burke et al. 2010) and in the pig-tailed macaques (Patton et al. 2006). Patton et al. (2006) showed that 1.0% C31G gel applied to the vagina or rectum had no toxicity. However no safety and tolerance studies have been done in any eye model including rabbit or primate eyes. In addition to assessing the safety and tolerance in the rabbit and squirrel monkey eye models, the efficacy of C31G as an anti-herpetic was determined in comparison to commercial trifluorothymidine (TFT). TFT is considered by many to be the drug of choice to treat herpes keratitis in humans (Wellings et al. 1972; Coen and Richman 2007; Hill et al. 2009). Many other anti-herpetics have been tested in the rabbit eye model (Coen and Richman 2007; Toma et al. 2008; Webre et al. 2012). A 1% gel of C31G was tested in pigtailed macaques and found to be safe and well tolerated (Patton et al. 2006).

3. Materials and Methods

3.1. Drugs, Compounds, and Solutions

The antiviral drug trifluorothymidine (TFT) is a 1% ophthalmic solution provided by Falcon Pharmaceutical, Fort Worth, TX. The pH was approximately 6.0 and the osmolarity is approximately 283 oMsm. The C31G was used at 0.25% in 0.5% HPMC at pH 6 provided by BioSyn Inc., Philadelphia, PA. The 0.5% HPMC at pH 6 had no C31G and was provided by Biosyn Inc., Philadelphia. Sterile balanced salt solution (BSS) at pH 6 was a laboratory preparation.

3.2. Approvals

Our studies in rabbits and primates adhered to the policies of the Association for Research and Provision in Ophthalmology (ARVO). In addition to the ARVO policies, we obtained approval for these studies from the LSUHSC Institutional Biosafety Committee (IBC) #7024 and the LSUHSC Institutional Animal Care and Use Committee (IACUC) #2850.

3.3. Animal Models

3.3.1. Safety and Tolerance in Rabbit Eyes—*In vivo* safety and tolerance of C31G was carried out on naïve, uninfected rabbit eyes. A series of studies was performed. Each test group had five rabbits. First, we assessed the safety and tolerance of three types of treatments in rabbit corneas with no injury for five consecutive days, five times per day. These three treatments were: (1) 0.25% C31G; (2) 0.5% HPMC at pH 6; and (3) sterile balanced salt solution (BSS) with no HPMC and no C31G. Each rabbit eye received 50 µl topically.

For the next experiment, rabbit corneas were scarified by mild crosshatch on the corneal surface. This experimental design assessed whether C31G would alter the wound healing and reepithelialization in a small crosshatch area.

A third safety and tolerance test involved removing the entire corneal epithelium in a 5 mm circle marked with a 5 mm trephine with a depth of ~0.2 mm. The three solutions noted above were used to treat the scarified rabbit eyes five times per day for five consecutive days (see Table 1).

3.3.2. Safety and Tolerance in Nonhuman Primate Eyes—We tested the three solutions in eight adult (1.2–2 kg) squirrel monkeys (*Saimiri sciureus*). These monkeys had no corneal defects. To determine any adverse effects in these primates, the right eyes were treated with 0.25% C31G in 0.5% HPMC and the left eyes were treated with BSS. This was done five times per day for five consecutive days. Slit-lamp examinations were performed each day before treatment (see Tables 1 and 2).

3.4. Virus

HSV-1 strain McKrae was used in all experiments. New Zealand white rabbit corneas were scarified in a mild crosshatch fashion. The rabbits were under general anesthesia and the corneas received topical anesthesia. Both corneas were inoculated with 50 µl of HSV-1 strain McKrae at 3×10^5 PFU.

3.5. HSV-1 Keratitis Model/Chemotherapy

Three days after viral inoculation, all 48 rabbit corneas (in 24 rabbits) were scored using ocular parameters and by slit-lamp examinations with fluorescein staining (see Table 2), and then were placed into three balanced groups of eight rabbits per group. The rabbit treatment groups were (1) 0.25% C31G in 0.5% HPMC at pH 6, (2) the vehicle at 0.5% HPMC at pH 6, and (3) 1% TFT (commercially available). Topical drops (50 µl) were applied to all eyes 1.5 hours apart, five times per day for five consecutive days. Administration of drugs was masked (treatment groups unknown by those giving treatments).

The slit-lamp examination and scoring of ocular parameters listed in Table 2 were marked and performed independently by two experienced corneal specialists. We have previously given details of slit-lamp procedures (Kwon et al. 1979; Majumdar et al. 2005; Hill et al. 1998; Toma. et al 2008; Webre et al. 2012).

Table 2 is a description of the 7 ocular clinical parameters. In addition to the 0 to +4 analyses of the corneal lesions by slit-lamp examination, the severity of ocular disease was assessed by evaluating six other clinical parameters: stromal opacity, scleral injection, eyelid inflammation, ocular neovascularization, inflammatory discharge, and epiphora (excessive tearing). Means and standard error of the means of the clinical scores were calculated daily for post inoculation (PI) days 3–7, 7.5, and 10. On PI day 7, scoring was done once before the first treatment and once after the last treatment. All the clinical parameters were masked and assessed. Ocular parameter scores of the three treatment groups are provided for comparison (See Figure 1 and Table 2).

3.6. Statistics

Our statistical analysis was by a full non-parametric repeated mean as described by Dmitreanko et al. 2007 and Brunner et al. 2001.

4. Results

Safety and tolerance tests found that C31G was safe and well-tolerated in both rabbit and squirrel monkey eyes. Based on these experiments, we proceeded with a similar C31G dose and frequency of administration in rabbits displaying HSV-1 keratitis. Table 1 outlines these animal studies.

In the first phase of the safety and tolerance in naïve rabbit eyes, the three drugs noted in Table 1 were given to rabbits five times per day for five consecutive days and rabbits were scored. Ocular parameters and scoring criteria are given in Table 2. Safety and tolerance studies showed no adverse effects in any clinical parameter scored in any eye in any rabbit.

The second phase of safety and tolerance was a mild crosshatch of the corneal surface that mimics the viral inoculation procedure. These were masked analyses and all 5 rabbits (10 eyes) in each group healed completely within 72 hours. We continued treatment until PI day 7 and there were no observable adverse effects in any of the six parameters listed in Table 2.

The most important test of safety and tolerance involved the debridement of the corneal epithelium to evaluate both epithelialization and migration. This mimics an infectious geographic lesion. Three groups noted in Table 1 were employed. Again, the reepithelialization was almost complete (~85%) at 72 hours and was fully completed by day 4 (94 hours) in all eyes in all groups. The experiment was continued for a fifth day and no adverse effects were observed.

We then conducted safety and tolerance testing without injury on normal naïve squirrel monkey eyes. This was the same dosing frequency as in rabbits, but without any injury to the corneal epithelium. Over the five days, the monkeys were examined by fluorescein staining and SLE, and there were no differences in safety and tolerance between the C31G in HPMC and the BSS. This demonstrated that C31G in HPMC is safe and well-tolerated in the non-human primate eye.

Figure 1A shows the average corneal slit-lamp examination (SLE) scores of the three treatment groups, beginning on post-inoculation (PI) day 3. This scoring continued through PI day 10. During the first phase of treatment, there were no statistical differences (PI day 4). In the second phase of treatment, Figure 1A shows C31G and TFT scores significantly ($p < 0.005$) lower than those from the 0.5% HPMC group. This statistical difference continued from PI day 5 through PI day 10.

Another characteristic noted in the SLE scoring is that eyes treated with C31G had geographic lesions (+2 or greater) in two of 112 observations, while eyes treated with the

TFT had 3 geographic lesions out of 112 observations, and eyes treated with vehicle had geographic lesions in 14 of 112 observations. This is another indicator of the efficacy of the two drugs compared to the vehicle. Most studies done on HSV-1 keratitis in animal models and patients emphasize masked slit-lamp examination scores as the most relevant. However, because this study was employing a new drug, we thoroughly assessed other ocular parameters for both safety and tolerance and efficacy in treatment.

Figures 1B–1G are of the individual ocular parameters graphed like those in Figure 1A. Figure 1B is the assessment of the stromal opacity. The vehicle treated eyes had significantly ($p<0.005$) increased stromal opacity scores beginning at PI day 6 after initiation of treatments. The stromal opacity remained constant for the duration of treatments with C31G and TFT. Stromal opacity continues to increase in the 0.5% HPMC group up to PI day 10. Stromal opacity, especially when scores are 1 or greater, is irreversible and is a significant contributor to decreased vision.

Figure 1C is the assessment of scleral injection (bloodshot eyes). The treatments with C31G and TFT were both significantly ($p<0.005$) decreased as compared to the control, and the decline in score began on PI day 6 and continued until PI day 10. All treatments were stopped at PI day 7.

Corneal neovascularization is a major problem in acute and recurrent ocular herpes. In animal models, this condition often occurs late relative to the initial appearance of corneal lesions. Figure 1D provides scores for corneal neovascularization in the 3 treatment groups. This parameter was significantly ($p<0.005$) lower on PI day 7 for both the C31G and TFT groups as compared to the vehicle group. Corneal neovascularization, similar to an increase in stromal opacity, is often irreversible if not aggressively treated in patients and is a significant factor leading to blindness.

Figure 1E is the assessment of eyelid inflammation. A significant ($p<0.005$) reduction in eyelid inflammation began for the C31G and TFT treatment groups on PI day 6. This parameter remained constant through PI day 10, even though treatment stopped at the end of PI day 7. Figure 1E shows that scores remained high for the vehicle group.

Figure 1F is the assessment of inflammatory discharge (primarily mucous). C31G and TFT were effective beginning on PI day 4, 24 hours after initiating treatment. The scores for C31G and TFT were significantly ($p<0.005$) lower than the vehicle.

Figure 1G is the scores of assessment of epiphora (tears/fluid). This was the most rapid response of the six (6) ocular parameters assessed. This response appeared 24 hours after initiation of treatment and continued until PI day 10. At PI days 4–10, C31G and TFT showed significantly less ($p<0.005$) epiphora than the vehicle group.

Figure 1H is a cumulative representation of six ocular parameters indicated in Table 2, excluding the corneal epithelium scores (Figure 1A). The data in Figure 1H show that 48 hours after treatment initiation (PI day 5), a statistical difference ($p<0.005$) was observed between the vehicle group compared to the TFT and C31G groups, and this difference continued to PI day 10.

5. Discussion

A major health problem in industrialized countries, especially the USA, is the increased incidence of acyclovir-resistant and multi drug-resistant strains of HSV-1 in patients with ocular herpes infections (Webre, et al. 2012). Most of the drug resistance is related to the herpes thymidine kinase. We know of no resistance to C31G in any infectious organism.

Although the exact mechanism of action is not known, it may involve pore formation disrupting the viral envelope.

Many antivirals are nucleoside analogs (Coen and Richman 2007). A few potent antivirals are non-nucleoside analogs (Coen and Richman 2007; Himaki et al. 2012). Compounds such as plant extracts have been reported to exhibit anti-herpetic activity *in vitro* and some *in vivo* (Nakama et al. 2012). This is due, in part, to the disruption of the lipid envelope of herpesviruses. Also, numerous proteins (Brandt et al. 2007), peptides (Bultmann and Brandt 2002; Jose et al. 2013) such as the apolipoprotein E mimetic peptide (Bhattacharjee, et al. 2008; Bhattacharjee et al. 2009) and amphoteric compounds (Bultmann and Brandt, 2002) have been reported to exhibit anti-herpetic activity in animal models. Other novel inhibitors have been suggested to block herpes acute infections (Mulik et al. 2012) and viral reactivation epigenetically (Knipe et al. 2013; Kristie, 2012; Liang et al. 2013A; Liang et al. 2013B).

C31G has been reported to have antibacterial, antifungal, and antiviral activity *in vitro* (Wyrick et al. 1997; Corner et al. 1988; Howett et al. 1999; Malamud et al. 1998; Thompson et al. 1996). There are no reported antiherpetic studies of C31G *in vivo*. C31G exhibits potent spermicidal activity and is known to penetrate cervical mucosa (Thompson et al. 1996). C31G was found safe and well-tolerated in the rabbit eye model, and allowed assessment of chemotherapeutic efficacy for HSV-1 keratitis. We evaluated C31G compared to TFT (a potent anti-herpetic agent) and to a vehicle with the same ionic strength and pH as C31G. We found C31G and TFT to have comparable activity. However, C31G is more economical and may be able to inactivate most enveloped viruses, including HIV (Krebs et al. 1999). Furthermore, C31G's mechanism of action is likely to be active in the treatment of HSV-2.

We assessed the corneal epithelium by slit-lamp examination (SLE) and six (6) other ocular parameters known to be associated with HSV-1 lesions and pathogenesis. C31G and TFT both increased healing and decreased the ocular parameters directly associated with herpes keratitis. C31G was proven safe, tolerable, and efficacious at treating herpes keratitis. It is more economical than comparable treatments (TFT) and has no known resistant organisms. Thus, we propose that C31G has promising potential for treatment of ocular herpes keratitis.

Acknowledgments

This research was supported in part by funding from the National Eye Institute grants EY014289 (Richard Bax), R21-EY019144 647 (PSB), R01-EY006311 (JMH); National Cancer Institute R01-CA107974 (ACO), R21-CA162133 (PCR); the NIH National Institute on Aging AG18031 (WJL) and AG038834 (WJL), AG23085 (JMH); P20GMM103501 (ACO, TPF, PCR); National Center for Research Resources (P20RR016456), the National Institute of General Medical Sciences (P20GM103424), Louisiana Cancer Research Consortium and the NIH-RCM1 grant #5G12RR02620 from the National Institute on Minority Health and Health Disparities (HEM). This research was also supported by an unrestricted research grant from the LSU Health Sciences Center (JMH); Transitional Research Initiative grants (WJL and JMH); a research grant from LSUHSC (JMH); a Research to Prevent Blindness (RPB) Senior Scientific Investigator Award (JMH); an unrestricted grant to the LSU Eye Center from RPB, New York, NY; the Louisiana Biotechnology Research Network (WJL, JMH); an Alzheimer Association Investigator-Initiated Research Grant IIRG-09-131729 (WJL); the Louisiana Cancer Research Consortium (PSB); the Louisiana Lions Eye Foundation, New Orleans; the Louisiana Vaccine Center sponsored by the Louisiana Board of Regents (JMH)

References

- Bhattacharjee PS, Neumann DM, Foster TP, Clement C, Singh G, Thompson HW, Kaufman HE, Hill JM. Effective treatment of ocular HSK with a human apolipoprotein E mimetic peptide in a mouse eye model. *Invest Ophthalmol Vis Sci*. 2008; 49:4263–4268. [PubMed: 18515564]

- Bhattacharjee PS, Neumann DM, Hill JM. A human apolipoprotein E mimetic peptide effectively inhibits HSV-1 TK-positive and TK-negative acute epithelial keratitis in rabbits. *Curr Eye Res.* 2009; 34:99–102. [PubMed: 19219680]
- Brandt CR, Akkarawongsa R, Altmann S, Jose G, Kolb AW, Waring AJ, Lehrer RI. Evaluation of a theta-defensin in a Murine model of herpes simplex virus type 1 keratitis. *Invest Ophthalmol Vis Sci.* 2007; 48:5118–5124. [PubMed: 17962464]
- Brunner, E.; Domhof, S.; Langer, F. *Nonparametric Analysis of Longitudinal Data in Factorial Experiments.* 1. Wiley; New York: 2001.
- Bultmann H, Brandt CR. Peptides Containing Membrane-transiting Motifs Inhibit Virus Entry. *J Biol Chem.* 2002; 277:36018–36023. [PubMed: 12130644]
- Burke AE, Barnhart K, Jensen JT, Creinin MD, Walsh TL, Wan LS, Westhoff C, Thomas M, Archer D, Wu H, Liu J, Schlaff W, Carr BR, Blithe D. Contraceptive efficacy, acceptability, and safety of C31G and nonoxynol-9 spermicidal gels: a randomized controlled trial. *Obstet Gynecol.* 2010; 116:1265–1273. [PubMed: 21099590]
- Calis S, Yulug N, Sumnu M, Ayhan A, Hincal AA. A non-antibiotic antimicrobial mixture (C31G): evaluation of the antimicrobial efficiency of C31G on vaginal cultures. *Boll Chim Farm.* 1992; 131:335–338. [PubMed: 1492969]
- Coen, DM.; Richman, DD. Antiviral Agents. In: Knipe, DM.; Howley, PM., editors. *Fields Virology.* 5. Lippincott Williams & Wilkins; Baltimore: 2007. p. 448-486.
- Corner AM, Dolan MM, Yankell SL, Malamud D. C31G, a new agent for oral use with potent antimicrobial and antiadherence properties. *Antimicrob Agents Chemother.* 1988; 32:350–353. [PubMed: 3364952]
- Dmitrienko, A.; Chuang-Stein, C.; D'Agostino, R. *Pharmaceutical Statistics Using SAS: A Practical Guide.* Cary: 2007.
- Hill, JM.; Denham, LV.; Booth, BA.; Friedman, DA.; Hobden, JA.; Murina, AT.; Acierno, MD.; Jacob, JT.; Kaufman, HE.; Bergsma, D. Ocular pharmacology. In: Agarwal, A., editor. *Color Atlas of Ophthalmology.* 2. Jaypee Brothers Medical Publishers; India: 2009. p. 405-429.
- Hill, JM.; Wen, R.; Halford, WP. Pathogenesis and molecular biology of ocular HSV in the rabbit. In: Brown, MS.; MacLean, AR., editors. *Herpes Simplex Virus Protocols.* Humana Press/Wiley; Totowa, NJ: 1998. p. 291-315.
- Himaki T, Masui Y, Chono K, Daikoku T, Takemoto M, Haixia B, Okuda T, Suzuki H, Shiraki K. Efficacy of ASP2151, a helicase-primase inhibitor, against thymidine kinase-deficient herpes simplex virus type 2 infection in vitro and in vivo. *Antiviral Res.* 2012; 93:301–304. [PubMed: 22155691]
- Howett MK, Neely EB, Christensen ND, Wigdahl B, Krebs FC, Malamud D, Patrick SD, Pickel MD, Welsh PA, Reed CA, Ward MG, Budgeon LR, Kreider JW. A broad-spectrum microbicide with virucidal activity against sexually transmitted viruses. *Antimicrob Agents Chemother.* 1999; 43:314–321. [PubMed: 9925525]
- Jose GG, Larsen IV, Gauger J, Carballo E, Stern R, Brummel R, Brandt CR. A cationic peptide, TAT-Cd0, inhibits Herpes Simplex Virus Type 1 Ocular infection in vivo. *Invest Ophthalmol Vis Sci.* 2013; 54:1070–1079. [PubMed: 23341013]
- Knipe DM, Lieberman PM, Jung JU, McBride AA, Morris KV, Ott M, Margolis D, Nieto A, Nevels M, Parks RJ, Kristie TM. Snapshots: chromatin control of viral infection. *Virology.* 2013; 435:141–156. [PubMed: 23217624]
- Krebs FC, Miller SR, Malamud D, Howett MK, Wigdahl B. Inactivation of human immunodeficiency virus type 1 by nonoxynol-9, C31G, or an alkyl sulfate, sodium dodecyl sulfate. *Antiviral Res.* 1999; 43:157–173. [PubMed: 10551374]
- Kristie TM. The rise of epigenetic targets for the development of novel antivirals. *Expert Rev Anti Infect Ther.* 2012; 10:1359–1361. [PubMed: 23253311]
- Kwon B, Gangarosa LP, Park NH, Hull DS, Fineberg E, Wiggins C, Hill JM. Effect of iontophoretic and topical application of antiviral agents in treatment of experimental HSV-1 keratitis in rabbits. *Invest Ophthalmol Vis Sci.* 1979; 18:984–988. [PubMed: 90030]

- Liang Y, Quenelle D, Vogel JL, Mascaro C, Ortega A, Kristie TM. A Novel Selective LSD1/KDM1A Inhibitor Epigenetically Blocks Herpes Simplex Virus Lytic Replication and Reactivation from Latency. *MBio*. 2013A; 4:1–10.
- Liang Y, Vogel JL, Arbuckle JH, Rai G, Jadhav A, Simeonov A, Maloney DJ, Kristie TM. Targeting the JMJD2 Histone Demethylases to Epigenetically Control Herpesvirus Infection and Reactivation from Latency. *Sci Transl Med*. 2013B; 5:167ra5.
- Majumdar S, Nashed YE, Patel K, Jain R, Itahashi M, Neumann DM, Hill JM, Mitra AK. Dipeptide monoester ganciclovir prodrugs for treating HSV-1-induced corneal epithelial and stromal keratitis: in vitro and in vivo evaluations. *J Ocul Pharmacol Ther*. 2005; 21:463–474. [PubMed: 16386088]
- Malamud D, Howett MK, Wigdahl B, Christensen ND, Krebs FC, Weisz J, Kreider JW. Anti-viral activity of microbicides in model systems. *Int Conf AIDS*. 1998; 12:621.
- Mulik S, Xu J, Reddy PB, Rajasagi NK, Gimenez F, Sharma S, Lu PY, Rouse BT. Role of miR-132 in angiogenesis after ocular infection with herpes simplex virus. *Am J Pathol*. 2012; 181:525–534. [PubMed: 22659469]
- Nakama S, Tamaki K, Ishikawa C, Tadano M, Mori N. Efficacy of *Bidens pilosa* Extract against Herpes Simplex Virus Infection In Vitro and In Vivo. *Evid Based Complement Alternat Med*. 2012; 2012:413–453.
- Patton DL, Sweeney YT, Balkus JE, Hillier SL. Vaginal and rectal topical microbicide development: safety and efficacy of 1.0% Savvy (C31G) in the pigtailed macaque. *Sex Transm Dis*. 2006; 33:691–695. [PubMed: 16641825]
- Thompson KA, Malamud D, Storey BT. Assessment of the anti-microbial agent C31G as a spermicide: comparison with nonoxynol-9. *Contraception*. 1996; 53:313–318. [PubMed: 8724622]
- Toma HS, Murina AT, Areaux RG Jr, Neumann DM, Bhattacharjee PS, Foster TP, Kaufman HE, Hill JM. Ocular HSV-1 latency, reactivation, and recurrent Disease. *Semin Ophthalmol*. 2008; 23:249–273. [PubMed: 18584563]
- Webre J, Hill J, Nolan N, Clement C, McFerrin H, Bhattacharjee P, Hsai V, Neumann D, Foster T, Lukiw W, Thompson H. Rabbit and Mouse Models of HSV-1 Latency, Reactivation, and Recurrent Eye Disease. *J Biomed Biotechnol*. 2012; 2012:1–18. [PubMed: 21836813]
- Wellings PC, Awdry PN, Bors FH, Jones BR, Brown DC, Kaufman HE. Clinical evaluation of trifluorothymidine in the treatment of herpes simplex corneal ulcers. *Am J Ophthalmol*. 1972; 73:932–942. [PubMed: 4555825]
- Wyrick PB, Knight ST, Gerbig DG Jr, Raulston JE, Davis CH, Paul TR, Malamud D. The microbicidal agent C31G inhibits *Chlamydia trachomatis* infectivity in vitro. *Antimicrob Agents Chemother*. 1997; 41:1335–1344. [PubMed: 9174195]

Highlights

- We establish the potent anti-herpetic activity of C31G *in vivo* (HSV-1 keratitis)
- We tested 0.25% C31G plus 0.5% hydroxypropyl methylcellulose (HPMC) against 1% trifluorothymidine and a vehicle, 0.5% HPMC
- Rabbits were assessed based on slit-lamp examination scores and 6 other ocular parameters
- C31G has the potential to be used to treat herpetic keratitis as well as other herpetic topical lesions in humans

(A) SLE Scores in HSV-1 Epithelial Keratitis in NZW Rabbits

*TFT and C31G significantly different from Vehicle

(B) Assessment of Stromal Opacity

*TFT and C31G significantly different from Vehicle

(C) Assessment of Scleral Injection

*TFT and C31G significantly different from Vehicle

(D) Assessment of ocular neovascularization

*TFT and C31G significantly different from Vehicle

(E) Assessment of eyelid inflammation

*TFT and C31G significantly different from Vehicle

(F) Assessment of inflammation discharge (mucous)

*TFT and C31G significantly different from Vehicle

(G) Assessment of epiphora

*TFT and C31G significantly different from Vehicle

(H) Cumulative Scores of 6 Ocular Parameters

*TFT and C31G significantly different from Vehicle

Figure 1.
Assessments of 7 Ocular Parameters

Table 1

Outline of Animal Studies

Experiment	Rabbits			Nonhuman Primates	
	Group 1	Group 2	Group 3	Group 1 (Right Eyes)	Group 2 (Left Eyes)
Safety and Tolerance Normal, Naïve Corneas (No Deficit)	C31G (0.25%) + HPMC (0.5%)	HPMC (0.5%)	BSS	C31G (0.25%) + HPMC (0.5%)	BSS
Safety and Tolerance with Crosshatch of Corneal Surface	C31G (0.25%) + HPMC (0.5%)	HPMC (0.5%)	BSS	N/A	N/A
Safety and Tolerance with Corneal Debridement (5 mm circle)	C31G (0.25%) + HPMC (0.5%)	HPMC (0.5%)	BSS	N/A	N/A
Inoculation of HSV-1 *	C31G (0.25%) + HPMC (0.5%)	HPMC (0.5%)	TFT (1%)	N/A	N/A

All treatments were masked. All solutions had a pH of 6.

Treatments were given at 50 µl, 5 times per day at 90 minute intervals.

* Rabbits were placed into three balanced groups based on SLE scores

Table 2

Criteria for scoring of ocular parameters

Ocular parameters	Clinical scores			
	0 (symptomatic of naïve uninfected with no manifestations)	+1 (slight manifestations)	+2 (moderate manifestations)	+3 (severe manifestations)
Stromal opacity	No coloration, clear and transparent	Slight thickening of stromal area, translucent	Whitish-yellow coloration, thickening and scarred	Yellowish coloration of >90% of stromal area, thickening and open sores with blood patches
Scleral injection (hyperemia)	No redness, no prominent blood vessels and no chemosis	Reddish tinge, blood vessels more visible and localized chemosis	Red, prominent blood vessel network and localized chemosis	Hyperemia with individual blood vessels indistinguishable and chemosis
Ocular neovascularization	No visible blood vessels	Developing blood vessel network with <25% of the surface affected	Blood vessel network with 26–75% of the surface affected	Prominent blood vessel network with >75% of the surface vascularized
Eyelid inflammation	Eyelid margins thin with no visible redness	Eyelid margins showing localized swelling and redness	Eyelid margins showing localized swelling and visible blood vessel network	Eyelid margins swollen, red and engorged outer parts of eyelids
Inflammatory discharge	No mucous discharge	Mucous discharge collecting at the caruncle and lacrimal apparatus	Mucous discharge collecting at the caruncle, lacrimal apparatus and eyelid margin	Mucous discharge collecting at the adnexa and causing peri-ocular fur to be matted down (pseudo-membranes)
Epiphora (tears/moisture)	Moist corneal surface	Watery fluid collection mainly at the lower eyelid margin	Watery fluid collection mainly at the lower eyelid margin and wetness of the entire outer part of eyelids	Watery fluid collection mainly at the lower eyelid margin, wetness of the entire outer part of eyelids and pseudo-membranes
Slit-lamp examination (SLE)	No lesions	~15–20% of corneal surface involvement, primarily dendrites	At least one geographic lesion and numerous dendrites	Mostly geographic lesions covering between 50–80% of the corneal surface

* SLE score of +1.5 means that at least 25% up to 40% of the cornea shows dendrites and punctate lesions but no geographic lesion

* SLE score of +4.0 means that at least 90% of the corneal surface is covered with a very large geographic lesion