Published in final edited form as:

Anesthesiology. 2014 March ; 120(3): 601–613. doi:10.1097/ALN.0000000000000080.

A pilot study evaluating pre-surgery neuroanatomical biomarkers for postoperative cognitive decline after total knee arthroplasty in older adults

Catherine C. Price, Ph.D.^{1,2} [Assistant Professor], Jared J. Tanner, Ph.D.¹ [Post-Doctoral Associate], Ilona Schmalfuss, M.D.^{3,4} [Associate Professor], Cynthia Wilson Garvan, Ph.D.⁵ [Statistician], Peter Gearen, M.D.⁶ [Professor], David Dickey, Ph.D.¹ [Post-Doctoral Fellow], Kenneth Heilman, M.D.⁷ [Professor], David L. McDonagh, M.D.⁸ [Associate Professor], David. J. Libon, Ph.D.⁹ [Professor], Christiana Leonard, Ph.D.¹⁰ [Professor], Dawn Bowers, Ph.D.^{1,6} [Professor], and Terri G. Monk, M.D., M.S.⁷ [Professor]

¹Department of Clinical and Health Psychology, University of Florida, Gainesville, Florida

²Joint Appointment, Department of Anesthesiology, University of Florida, Gainesville, Florida

³Department of Radiology, University of Florida, Gainesville, Florida

⁴Department of Radiology, North Florida South Georgia Veteran Association, Gainesville, Florida

⁵Health Science Center, University of Florida, Gainesville, Florida

⁶Department of Orthopedic Surgery, University of Florida, Gainesville, Florida

⁷Department of Neurology, University of Florida, Gainesville, Florida

⁸Department of Anesthesiology, Duke University, Durham, North Carolina

⁹Department of Neurology, Drexel University, Philadelphia, Pennsylvania

¹⁰Department of Neuroscience, University of Florida, Gainesville, Florida

Abstract

Background—Total knee arthroplasty improves quality of life but is associated with postoperative cognitive dysfunction in older adults. This prospective longitudinal pilot study with a parallel control group tested the hypotheses that: 1) nondemented adults would exhibit primary memory and executive difficulties after total knee arthroplasty, and 2) reduced preoperative hippocampus/entorhinal volume would predict postoperative memory change, whereas preoperative leukoaraiosis and lacunae volumes would predict postoperative executive dysfunction.

Methods—Surgery (n = 40) and age–education-matched controls with osteoarthritis (n = 15) completed pre- and postoperative (3 weeks, 3 months, and 1 year) memory and cognitive testing. Hypothesized brain regions of interest were measured in patients completing preoperative magnetic resonance scans (surgery: n = 31, control: n = 12). Analyses used reliable change methods to identify the frequency of cognitive change at each time point.

Results—The incidence of postoperative memory difficulties was shown with delay test indices (i.e., story memory test: 3 weeks = 17%, 3 months = 25%, 1 year = 9%). Postoperative executive

Correspondence: Catherine Price, Ph.D., Department of Clinical and Health Psychology, PO Box 100165, Gainesville, Florida, 32610, Phone: 352-265-0294; Fax: 352-265-0096; cep23@phhp.ufl.edu.

Study's Primary Affiliation: Departments of Clinical and Health Psychology, Anesthesiology, University of Florida

The authors declare no competing interests.

difficulty with measures of inhibitory function (i.e., Stroop Color Word: 3 weeks = 21%, 3 months = 22%, 1 year = 9%). Hierarchical regressions assessing the predictive interaction of group (surgery, control) and preoperative neuroanatomical structures on decline showed that greater preoperative volumes of leukoaraiosis/ lacunae were significant contributors to post-operative executive (inhibitory) declines.

Conclusions—This pilot suggests that executive and memory declines occur in non-demented adults having orthopedic surgery. Severity of preoperative cerebrovascular disease may be relevant for understanding executive decline, in particular.

Introduction

Postoperative cognitive dysfunction (POCD) involves pre- to postoperative reductions in memory, mental flexibility, and information processing. It is distinct from delirium and dementia (1). POCD occurs after non-cardiac surgery (2, 3) to some extent in all age groups at hospital discharge (37% for those aged 18–39, 30% for 40–59 years, and 41% for 60 and older), with longer-term POCD at 3 months for adults over 60 years of age (10%–13%). POCD is associated with early retirement and dependency on social transfer payments (4). It is also associated with increased mortality 1 year after surgery (3).

There are at least three types of POCD. Older non-demented adults with POCD can have isolated difficulty in learning/memory functions, isolated difficulties in executive functions, or a combination of memory and executive difficulties (5). Executive and combined impairments have been associated with functional limitations (5). To date, there have been no prospective investigations examining which cognitive/memory indices may best identify these POCD types, or if there are predictive neuroanatomical risk factors for these impairments. Such studies are necessary because at present, there are no known mechanisms of POCD.

Specific preoperative neuroimaging markers may indicate risk for POCD. It is well established that the entorhinal cortex (ERC) and hippocampus are important in declarative memory processes (6, 7) as measured by delay memory tests (8). These structures change with Alzheimer's disease (AD) (9). ERC volume predicts conversion to AD, with smaller volumes having greater conversion rates (10–12). Because of its involvement in the limbic-hypothalamic-pituitary-adrenal axis, the hippocampus is vulnerable to neuronal degeneration after severe biological/psychological stress (13, 14). Post-traumatic stress disorder associates with reduced hippocampal volumes (15, 16), and hippocampal degeneration also occurs in rats after anoxia and mild hypothermia (17). Preoperative leukoaraiosis and lacunae volume, by contrast, may indicate vulnerability to postoperative executive decline. Leukoaraiosis involves hyperintense white matter regions on brain computed tomography/magnetic resonance images (MRIs) (18) and occurs in 15% to 65% of adults (19). Leukoaraiosis associates with demyelination and hyalinosis narrowing of small brain arterioles. This signifies microvascular burden to frontal-subcortical white matter pathways important for executive functions (20–22). Lacunae suggest ischemic strokes (often considered “silent”). Lacunae often occur in subcortical gray matter nuclei (e.g., thalamus, caudate) necessary for filtering/disengaging attention (23). Leukoaraiosis and lacunae mark chronic small brain vessel disease (24) and are contributors to insidious executive dysfunction (25).

For this pilot investigation, we used a comprehensive neuropsychological protocol to assess whether the learning of new information (memory) and inhibitory functions (executive function) would be dominant forms of postoperative cognitive impairment (1). We then examined the hypothesis that specific presurgical neuroanatomical markers of early disease states (i.e., MRI-based hippocampus/ERC, leukoaraiosis/lacunae volumes) would differentially predict pre-postoperative memory and executive changes. Although

individuals may not present with clinical signs of impairment preoperatively, we hypothesized that preoperative neuroimaging markers might serve as an indication of brain vulnerability to perioperative insult and resulting memory/executive decline. We secondarily examined intraoperative variables (e.g., emboli number, anesthesia duration) as contributors to decline.

Materials and Methods

Standard Protocol Approvals, Registrations, and Patient Consents

The University of Florida Institutional Review Board, Gainesville, Florida, approved this study, and all participants signed consents. Authors followed principles from the Declaration of Helsinki.

Participants

Total knee arthroplasty (TKA) and control participants were recruited through University of Florida orthopedic clinics, screened, and enrolled between the years of 2003 and 2005. Participants in the control group were selected from patients who had chosen to abstain from surgery for at least 1 year. These two groups were recruited over the same time frame and were tested and scanned at the same time intervals. Participants had to meet the following inclusion/exclusion criteria: 1) aged 60 or older, 2) have English as a first language, 3) have osteoarthritis, 4) have intact activities of daily living, and 5) have baseline neuropsychological testing unsupportive for dementia criteria per Diagnostic and Statistical Manual of Mental Disorders – Fourth Edition. Additional exclusion criteria included: another major surgery within the study timeline, history of head trauma/ neurodegenerative illness, documented learning or seizure disorder, substance abuse in the last year, major cardiac disease, chronic medical illness known to induce encephalopathy, implantable device precluding an MRI, and an unwillingness to complete repeat testing. Two neuropsychologists reviewed the baseline data to confirm that test scores met the expected ranges for non-demented individuals.

Procedures

This was a prospective longitudinal study with a TKA surgery and a non-surgery control group. Participants completed a battery of tests that included a comprehensive history/ systems interview, a comorbidity rating (26), activities of daily living (27), and neurocognitive and mood testing. Baseline MRI was performed on eligible participants. Cognitive and mood status were re-evaluated at 3 weeks, 3 months, and 1 year post-baseline (postoperative for the TKA group). Delirium was ruled out postoperatively (28). Cognition was assessed 2 h after pain medication was given.

Anesthesia and Surgery Protocols—Protocols were standardized, with intravenous midazolam for anxiety, and fentanyl, thiopental, and rocuronium for anesthesia induction and intubation. Patients were ventilated with an air/oxygen mixture to maintain an end-tidal carbon dioxide at 35 ± 5 mm; anesthesia was maintained with inhaled isoflurane and intravenous fentanyl and rocuronium. We attempted to record emboli incidents with transcranial Doppler probes over the transtemporal window placed by the same anesthesiologist and technician (29). The same surgeon and anesthesiology teams completed all surgeries.

Neuropsychology Assessment—Baseline measures taken were within 1 week of the brain MRI. Baseline general cognitive (30) and intellectual estimates (31) helped equate groups. Measures of semantic/language fluency, visuoperception, and motor function were added to the protocol to examine the hypothesis that POCD primarily involved changes in

memory and executive measures. Multiple measures were administered as part of the investigation to examine instruments and expected sensitivity to postoperative change. The same neuropsychologist administered all measures at each time point. Tests were scored by the neuropsychologist but were also rescored and double-data entered by two technicians who were blinded to group status. Alternate test forms were used, when applicable, with administration order based on a random generation list. See Ref. 32³² for test citations/descriptions. Primary neuropsychological measures of interest were theoretically grouped:

Memory: Auditory and visual stimuli with immediate and delay indices. Alternate versions were randomly administered for each period of testing.

1. Hopkins Verbal Learning Test - Revised (HVLT-R): Twelve-word list-learning test with three immediate free recall learning trials, one 20-minute delay free recall, and one recognition trial. Dependent variables (DVs): Immediate and delay total correct.
2. Story Memory Test (33): Paragraph recall test with immediate and 30-minute delay recall indices. DVs: Immediate and delay total correct.
3. Brief Visuospatial Memory Test – Revised (BVMT-R): Geometric figural memory with three learning trials, one 20-minute delay free recall, and one recognition trial. DVs: Immediate and delay total correct.

Executive Functions: Tests associated with working memory, word fluency, processing speed, and disengagement-response inhibition were applied:

1. Digit Span Backward Subtest from the Wechsler Memory Scale - Third Edition: Requires listening to an increasing digit series and then repeating those digits in forward and backward sequences. DV: Total backward.
2. Spatial Span Subtest of the Wechsler Memory Scale - Third Edition: A visual analog to the WMS-III Digit Span test. DV: Total backward.
3. Digit Symbol Subtest of the Wechsler Adult Intelligence Scale - Third Edition: A measure of psychomotor processing speed requiring rapid matching of numbers and symbols. DV: Total correct.
4. Controlled Oral Word Association Test: Generating words beginning with a specific letter within 60 seconds, excluding numbers and proper nouns. DV: Total words minus errors.
5. Stroop Color Word Test: Involves selective attention and cognitive control by expecting participants to suppress the automatic tendency to read aloud words rather than name the color of the ink in which the words are printed on the page. DV: Color-word score in 45 seconds.

Other Cognitive Domains: Semantic/language fluency, perceptual-spatial function, and motor measures were administered to better examine the hypothesis that memory and executive dysfunction were primary POCD forms.

1. Category (Animal) Fluency: Requires generating animal names in 60 seconds. DV: Total words generated.
2. Judgment of Line Orientation: Involves matching two lines of varying degree to a spectrum of lines ranging from 0 to 180°. DV = Total correct.
3. Finger Tapping Test: Entails rapid lever pressing for 10 seconds and 10 trials. DVs = Average dominant and non-dominant taps across all trials.

Mood and Pain—Depression and anxiety were evaluated with the Geriatric Depression Scale (34) and the State Trait Anxiety Inventory (35). A visual analog scale gauged pain severity (36).

Neuroimaging—TKA participants were scanned within 2 weeks of surgery and control participants were scanned at their baseline assessment using a Siemens 3T Allegra scanner. T1-weighted, three-dimensional magnetization prepared rapid acquisition gradient-echo (MP-RAGE) sequence (repetition time = 2500 ms, echo time = 4.38 ms, inversion time = 1100 ms, flip angle = 8 degrees, matrix = 256×144) reconfigured to 160 gapless, 1-mm images provided gray and white matter segmentation. Leukoaraiosis volumetrics were acquired from two-dimensional Fluid Attenuated Inversion Recovery sequences (repetition time range across scans = 8,402–12,800; echo time range = 125–147 ms; inversion time range = 1800–2200 ms, flip angle = 90 degrees, gap = 5–7 mm).

Magnetic Resonance Imaging predictor variables—Raters were blind to participant group. Hippocampi (37) were manually segmented using ITK-SNAP (38) by one reliable rater with an excellent Dice Similarity Coefficient (DSC; 39) [grand DSC = 80 ± 0.02 ; intra-rater: grand DSC = 0.81 ± 0.05 , Pearson r range = 0.75–0.83; all $p < 0.001$]. Volume DVs = Left, right, and total hippocampal volumes in cubed millimeters.

ERCs (40) were segmented by highly reliable raters [Intraclass Correlation Coefficient $r > 0.93$ (33) using MEASURE (41)]. Tracings were made on oblique coronal slices with volumes calculated by compiling individual slice measurements. DVs = Left, right, and total volumes in cubed millimeters.

Leukoaraiosis was measured by a reliable rater [DSC inter-rater range of 0.84–0.93, intra-rater range > 0.99] using ImageJ (42) and an in-house macro (43). Using published methods (21), leukoaraiosis voxels were thresholded and created into two-dimensional leukoaraiosis binary masks that were subsequently concatenated to form a single, three-dimensional binary mask for each brain. DV = total volume in cubed millimeters.

Lacunae were measured by a neuroradiologist using volumetric T1 scans. Only well-defined, dark lesions with a diameter ≥ 2 mm that held a stationary position between slices were graded as lacunae with their volume estimated using the formula of a sphere ($\frac{4}{3}\pi r^3$; 44). DV = total volume in cubed millimeters.

Imaging control variables included total brain volume corrected for group total intracranial volume (45) to correct for head size and age-related atrophy. Intracranial volume (brain plus cerebral spinal fluid) and *supratentorial whole brain volume* (gray and white tissue plus ventricular cerebrospinal fluid minus brainstem and cerebellum) were segmented using FSL and BrainSuite methods (46, 47).

Statistical Analysis

A power analysis addressed the primary hypothesis that a baseline neuroanatomical variable of interest would correlate to postoperative cognitive change. Given the pilot nature of the study and that we did not have prior data from which to draw upon to estimate the effect size, we used STPLAN software (1993, University of Texas M.D. Anderson Cancer Center, Houston, TX) and a Fisher Z transformation method for sample estimation. Based on a 0.05 level of significance, a power of 0.80, and a moderate correlation ($r = 0.32$), we expected to enroll 60 participants total for the investigation. All analyses were completed with SPSS, version 11.0 (IBM, New York, NY). The level of significance was set at 0.05.

Assessing POCD Frequency—Independent t- and chi-square tests examined group differences on baseline demographics (e.g., imaging, cognition, mood, and pain variables). A modification of Jacobson and Truax's Reliable Change Index (RCI; see Refs. 1, 3, 48, and

49) assessed frequency of POCD by test: $\frac{\Delta X - \Delta X_c}{SD_{(\Delta X_c)}}$. Change was calculated by subtracting prefrom the postoperative performance (ΔX). The averaged control group change (ΔX_c , which was assumed to represent systematic error) was then subtracted from the individual change, with this value then divided by the control group's standard deviation of the change ($SD_{\Delta X_c}$). Abnormal cognitive decline was a z-score -1.96 . Using this method, we then examined tests for false positives, which is a consideration for POCD test sensitivity (49). Chi-square analyses assessed differences in POCD frequency. An overall doubly repeated longitudinal analysis was performed using the MIXED procedure in SAS, version 9.2 (Cary, NC). The repeated measures were z-scores observed over time (baseline, 2 weeks, 3 months, and 1 year), and cognitive domain: 1) immediate learning and memory (HVLTI, Story I, and BMVT 1), 2) delayed learning and memory (HVLTD, Story D, and BMVT D), 3) attention, processing, and executive function (Dg Span, Sp Span, Dg Symbol, and Stroop C-W), and 4) language, visuospatial, and motor (animals, judgment of line orientation, finger D, and finger ND). Norms were not available to form controlled oral word association z-scores thus controlled oral word association was omitted from this analysis.

Predicting cognitive change—Multivariate analyses were strategic to 1) confirm that there were certain domains that would change with surgery (we expected primary changes in memory and executive function); and 2) investigate the expected sensitivity of certain instruments to detect these changes (i.e., memory test delay indices word list memory measures versus story paragraph memory measures; story memory measures are considered less dependent upon the processing speed (32)). We limited the multivariate analyses to one memory and one executive measure, with this selection based demonstration of change in the surgery group relative to the non-surgery group, and knowledge of which tests may be most sensitivity to anterograde memory changes (e.g., delay index) and inhibitory decline. We did not consider using a composite score approach as we believed that would give an inaccurate assessment of POCD type in our sample, for less sensitive tests have the potential to introduce measurement error. Given the growing field of POCD research and the numerous concerns regarding which tests should be administered, we include information for the separate test measures within the provided tables. We believe that this information is relevant for to guide future investigations. Planned hierarchical regressions examined 1) hypothesized interaction of baseline hippocampal, ERC volumes, and group (surgery, control) on the delay index of the Story Memory Test, and 2) the baseline interaction of lacunae/leukoaraiosis volume and group (surgery, control) on the Stroop Color Word Test Color-Word condition score. The first regression model always included the baseline cognitive variable of interest (e.g., baseline delay memory score) and the imaging control variable of TBVc. Including the preoperative cognitive score in the first regression model renders the DV into a "residual change score," so that the effects of the predictors on the DV may be interpreted as predictors of change. Covariates of education, anesthesia duration, emboli count, and TKA type were analyzed and retained in the model if they were found to be significant. The second regression model included group type (surgery, control), and the third model included the interaction variable of interest [i.e., centered variable (50) of group $\times$ hippocampal volume].

Results

TKA patients (n = 40) and controls (n = 15) were similar regarding general demographic variables, general cognitive status, mood, and baseline pain (Table 1, all $p > 0.05$). Although not significant, the control group had on average two more years of education and were eight points higher on an abbreviated intellectual estimate ($p = 0.05$). All were considered healthy, with low comorbidity, and there were no statistically significant group baseline differences on the memory and cognitive measures (Table 2, all $p > 0.05$). Emboli counts were acquired only on a subset of the surgery participants (n = 15; mean = 14.40; SD = 25.63; range = 1–100 emboli) because of difficulties maintaining transcranial Doppler placement throughout surgery. Surgeries included unilateral TKA (n = 28) and bilateral TKA (n = 12).

A subset of individuals was unable to complete the preoperative brain scan because of new onset claustrophobia (n = 4), the size of the scanner bore, which limited patients with larger chests (n = 5), and poor image quality (n = 3). Because of pilot study timeline enrollment limitations, the final subgroup completing preoperative imaging included 31 surgery individuals and 12 control individuals who were similar regarding demographics, general cognitive status, mood, pain variables (Table 1, $p > 0.05$), baseline neuropsychology variables (Table 2, $p > 0.05$), and baseline brain variables of interest (Table 3, all $p > 0.05$). This MRI subset of controls had approximately two more years in education, on average, and scored six points higher on the intellectual estimate. Emboli were acquired within a subset (n = 12; mean = 14.50; SD = 28.45; range = 1–100 emboli). This MRI subgroup also included unilateral (n = 20) and bilateral (n = 11) TKAs.

The rate of attrition for TKA patients was 0% at 3 weeks, 8% (3/40) at 3 months, and 15% (6/40) at 1 year. Two patients were unavailable at the 3-month time point, but were tested at 1 year. Post-baseline testing was completed at 22 ± 7 days, 3 months ± 29 days, and 1 year ± 81 days.

Frequency of Cognitive Decline

Learning and Memory—POCD rates were more frequent for TKA patients on the delay relative to immediate memory indices [$X^2(1) = 5.98$, $p = 0.01$]. The highest POCD rate involved the Story Memory Test delay and the lowest POCD rate involved the HVLT delay (3 weeks: 17% and 8%; 3 months: 25% and 5%; 1 year: 9% and 15%, respectively) with test comparisons at 3 weeks and 3 months, $p < 0.05$. The visual memory test was accompanied by equally high rates of impairment in the control group at the 3-week and 3-month time points (both $p > 0.05$), suggesting high false-positive rates (Table 4).

Executive Functions—For TKA patients, the highest rates of POCD involved the inhibitory subtest of the Stroop Color Word Test, with 21%, 22%, and 9% at 3 weeks, 3 months and 1 year, respectively. The test rate comparison at 3 weeks and 3 months were all non-significant.

Overall Analysis—The TKA patients had lower z-scores in an overall analysis of time and cognitive domain. The regression coefficient in the doubly repeated model was -0.13 [$t(2893)$, p value = 0.003].

Neuroanatomical Predictors of Cognitive Decline

The two tests with the highest rates of POCD and minimal false positives in the control group were examined as outcome markers of memory and executive function (Memory

Test: Story Memory Test Delay Index; Executive Function Test: Stroop Color Word Test and Color-Word condition).

Preoperative Hippocampal/ERC Volumes and Story Memory Test Performance

—Baseline story memory ability was a predictor for postoperative story performance for each time period (β 's: 0.51–0.71, all $p < 0.01$). Adding in group status (surgery or control) to the model negatively contributed to memory performance (β 's: -0.42 to -0.25 , 3-week and 3-month p values = 0.001, 1-year p value < 0.05). The interaction variable of group (surgery or control) by left hippocampus or ERC volume never significantly contributed to the model over that of the other variables. There were no significant findings for the right hippocampus or ERC (Tables 5 and 6).

Preoperative Leukoaraiosis and Lacunae Volumes and Stroop Color-Word Test Performance

—Baseline Stroop performance and TBVc were significant independent predictors of postoperative Stroop performance for each time period (β 's: 0.76–0.82, all p values < 0.001 and $\beta = -0.23$, $p = 0.03$, respectively). Adding group status alone (surgery or control) to the model was not a significant predictor for any time period (all p values > 0.11). Adding in the interaction variable of group (surgery or control) and leukoaraiosis/lacunae volume significantly improved prediction of executive change for 3 weeks ($\beta = -0.22$, $p = 0.027$; adjusted $R^2 = 0.66$, F change = 5.31) and 1 year ($\beta = -0.27$, $p = 0.01$, adjusted $R^2 = 0.68$, F change = 8.40), with a trend for contribution at three months ($\beta = -0.17$, $p = 0.08$). Post hoc analyses on preoperative leukoaraiosis and lacunae volumes on postoperative memory changes were not found to be significant (Table 7).

Education and Surgical Considerations

Education, anesthesia duration (mean $\pm$ SD = 225.74 $\pm$ 99.39 min), and TKA type (unilateral/bilateral) were examined as covariates but did not contribute significantly to the first step of the regression models (Stroop Color-Word Test: education β 's = -0.03 to 0.01 ; anesthesia β 's = -0.09 to 0.02 ; TKA type β 's = -0.11 to -0.26 ; Story Memory Performance: education β 's = 0.17 – 0.29 ; anesthesia β 's = -0.18 to 0.06 ; TKA type β 's = -0.06 to 0.22) and were therefore not retained in the models. In the subsample of surgery participants with emboli measurement, greater emboli number negatively contributed to the acute 3-week postoperative Stroop Color-Word performance ($n=15$; Stroop Color-Word 3-week $\beta = -0.52$, $p = 0.03$; all other time point β 's < -0.16), but not significantly to delay Story Memory Test performance at any time period ($\beta = -0.05$ to -0.27). Although not statistically significant, emboli number were higher in bilateral ($n = 6$; mean $\pm$ SD = 24.0 $\pm$ 39.06 emboli) than unilateral TKA ($n = 9$; mean $\pm$ SD = 8.0 $\pm$ 8.97 emboli).

Discussion

This is the first prospective pilot study examining the role of presurgical neuroanatomical factors on POCD type after TKA in non-demented older adults. While we acknowledge the pilot nature of the study, our data suggest that memory and executive declines were the primary forms of cognitive change at 3 weeks post-TKA. Five percent or less of the patients exhibited declines in language, perceptual-spatial, or frontal motor function measures. The pilot study found limited value for using presurgery ERC/hippocampal volumes as neuroanatomical predictors for POCD memory decline at any time point (3 weeks, 3 months, or 1 year). In contrast, preoperative neuroimaging evidence of microvascular disease (preoperative leukoaraiosis and lacunae volume) explained a portion of executive function decline at the 3-week and 1-year postoperative sessions, with a trend at 3-months postoperation. We encourage researchers to conduct similar but larger scale prospective studies. Before clinical change can occur, and risk versus surgical benefit can be weighed,

we need more definitive evidence regarding the nature of baseline microvascular disease and executive decline, as well as rigorous examinations on neuroanatomical contributors to postoperative memory decline.

Considerations for Neuropsychological Measures and Potential Perioperative Variables

1. Executive Decline—Leukoaraiosis and lacunae volume accounted for a significant portion of variance on a well-known measure of executive function and the interference condition of the Stroop Color-Word Test at the 3-week and 1-year postoperative intervals. Other frontal system tests [i.e., Digit Symbol subtest (51), which involves processing speed and visual-motor integration (32, 52)] revealed postoperative change, but not to the same extent as the Stroop Color-Word Test. The Stroop Color-Word Test has been sensitive to delirium in similar patient groups (53). Via functional neuroimaging, this test associates with dorsal and medial frontal lobe (54-56), and parietal lobe (57), thereby implicating the involvement of large frontal-parietal and frontal-subcortical white matter networks. Leukoaraiosis and lacunae disrupt these connections (21). Thus, the Stroop Color-Word Test warrants consideration as a key neuropsychological measure in future POCD investigations.

Pilot findings extend upon studies showing leukoaraiosis as a risk factor for high-risk cardiac patients. Leukoaraiosis predicts frequency of POCD 3 months after “on-pump” coronary artery bypass surgery (58), and is associated with cerebral ischemic events following coronary artery bypass grafting (59). Our pilot study is the first to suggest that leukoaraiosis and lacunae volume should be considered for *healthy* non-demented older adults without clinically significant atherosclerotic disease who elect orthopedic surgery.

The control group with similar leukoaraiosis and lacunae burden load allowed us to examine the interaction of baseline microvascular disease and group (surgery/control) status. Findings suggest that the preoperative injuries of leukoaraiosis and lacunae did not contribute significantly to executive dysfunction until “insult (perioperative events) was added to injury.” A larger study needs to replicate these findings. We also need to identify the perioperative events that negatively interact with presurgery evidence of microvascular disease.

Potential perioperative events that may negatively interact with preoperative leukoaraiosis-lacunae volume include acute anemia (60-63), hypotension (64), oxygen desaturation (65-67), and the production of emboli. In our sample, we acquired valid emboli data on 15 of our surgery participants. The number of emboli in our sample (1–100 emboli) is similar to published reports (68). Post hoc findings suggest an association between emboli and acute (3-week) Stroop change score, but did not contribute to the predictive model. Although there are more reports that emboli frequency do not relate to cognitive change (69-71) than otherwise (72), study sample neurovascular burden factors and the interaction with emboli remain poorly understood. Small emboli may lodge in regions with low blood flow (73), and regions of the brain with leukoaraiosis might represent such regions. Support for this postulate comes from the observation that leukoaraiosis contributes to the development of silent stroke after cardiac surgery (74). We did not identify any significant predictors involving the duration of anesthesia, but the interaction with these and other baseline microvascular diseases warrants future examination.

The transient nature of leukoaraiosis and lacunae volume on cognition across our three postoperative time points suggests that perioperative events around TKA may be associated with ischemia. Ischemia, unlike infarction, is potentially reversible. If infarction were the mechanism of the decline, we would have expected that many participants would not have had a full cognitive reversal. The questions of whether transient decline might also be related to changes in other systems (e.g., neurotransmitter systems) needs empirical

assessment. Transient decline is a well-established occurrence in cardiac and non-cardiac surgery (3, 75-78).

2. Memory Decline—Memory functions have been classically associated with three neuroanatomic regions within the brain (and the pathways that interconnect them). These include the medial temporal lobe (hippocampus, entorhinal cortex) (33, 79), the thalamus (dorsomedial, anterior nuclei) (80), and the basal forebrain, which innervates the hippocampus with essential cholinergic neurons (81). In older adults, these regions are most associated with the amnesic form of mild cognitive impairment or clinical presence of AD (11, 82, 83). Despite the current study's high incident rate of postoperative recall impairment on sensitive indices of memory function, preoperative ERC and hippocampal volumes did not significantly contribute to postoperative memory abilities.

We do not consider our inability to support the original ERC-hippocampus hypothesis as a consequence of the neuropsychological or imaging measurement approaches. Rather, our reliable change analyses showed a marked decline on a delay recall index that is a characteristic of individuals with amnesic disorders (84, 85) and functional changes to the entorhinal cortex (8). The greatest rate of decline was also observed on a story recall test, which more closely resembles the everyday memory demands for interpersonal discourse, radio and television programs, as well as material that has been read, such as newspapers (32). We used reliable neuroanatomical measurement approaches (33) with resultant structure volumes that correspond to published values and ranges (40). We controlled for differences in intracranial and brain volume factors that may contribute to ERC/hippocampal variability between participants (86). Given these methodological strengths, we interpret that preoperative macrostructural measurement of the ERC and hippocampus is not sufficiently sensitive for predicting memory decline for non-demented otherwise "healthy" adults having TKA surgery. In vivo molecular-based examinations of medial temporal integrity and associated regions via diffusion and/or spectroscopy methods appear warranted; changes in microstructure may be earlier markers of underlying neurodegenerative disease before alterations in macrostructure such as changes in volume become manifest (87).

Study Considerations

We recognize study limitations. The sample size is small (hence increasing the probability of a Type II error) and unequal within the surgery and control groups. Despite our best attempts at matching the groups on age and education, the surgery group on average had more years of education. Although we restricted our regression analysis on one neuropsychological index, we recognize that our multiple hypothesis testing increased the experiment-wise Type I error rate. An additional limitation was that the neuropsychologist knew of the group (surgery, control) condition. We attempted to rectify this by having all tests rescored and re-entered by trained individuals who were blinded to the groups. Finally, this study was conducted with physically and cognitively well individuals. This limits the applicability of our pilot data to other populations, but does provide some reference for the volume of lacunae and leukoaraiosis in non-demented samples for future comparison purposes. We encourage similar studies on higher risk patients such as those with metabolic syndrome who have shown high rates of POCD after non-cardiac and cardiac surgery (88).

Despite the limitations, the pilot study has design strengths. We targeted control group recruitment during the same time period as the surgery recruitment and attempted to identify individuals similar in age, medical comorbidity, and baseline brain variables of interest. Cognitive change was examined using the RCI, which expressed change relative to the error estimated from a control group matched for age, comorbidity, intelligence, and baseline brain status. The RCI has demonstrated adequate specificity to detect POCD in non-cardiac

samples, with results replicated across studies (2, 3, 48). We further examined our hypotheses regarding POCD types with tests known for their sensitivity in examining memory and executive function. Examining cognitive change over time with a non-surgery control group allowed us to examine for potential false positives with specific cognitive measures (49). This approach was very useful given our small sample size, which prohibited a more powerful confirmatory factor analysis. Now, larger studies are necessary to re-examine our findings with the individual cognitive tests, but also with specific cognitive composites. The neuroanatomical variables were examined relative to brain and intracranial volumes, a technique that reduces inter-individual variability thereby clarifying cognitive-neuroanatomical associations (86).

The pilot findings warrant further consideration and larger empirical study. Investigators are encouraged to consider test protocols that include delay memory test conditions and executive inhibitory measures. In addition to standard structural measurements of leukoaraiosis and lacunae volume, future neuroimaging investigations should include molecular diffusion and functional-based sequences. Diffusion-weighted sequences allow quantification of tissue integrity before changes are seen on traditional clinical sequences (89). Researchers have shown that tensor-based algorithms have relevancy for understanding one-year survival after major cardiac and brain trauma (90, 91), as well as delirium (92). Resting and functional task-based sequences can be incorporated for understanding neuronal network risk profiles. Overall, it is our expectation that similar hypothesis-driven investigations using sensitive neuropsychological tools combined, and molecular- and functional-based tools in addition to standard clinical scans (i.e., FLAIR scans used to calculate leukoaraiosis) will improve our appreciation for pre-surgery neuroanatomical risk profiles and POCD types.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are most grateful to the participants for their time. We appreciate and acknowledge the insightful comments from the Anesthesiology reviewers and associated editors. We are also grateful for the research assistance of Ilcia Shugarman (College of Medicine, University of Florida), who helped with the procedural aspects of the investigation, and to Ria van der Aa, M.S., Katherine Moyses, B.A., and Jacqueline Nicosia, B.S. (Departments of Clinical and Health Psychology or Anesthesiology at the University of Florida), who assisted with double and triple data entry and management. We give special thanks to Jenny Mace, M.D. (Department of Anesthesiology, University of Florida), who assisted with intraoperative monitoring and postoperative management. We acknowledge Stephen Tower, M.S. (University of Florida), for his development of the ImageJ macro and employment of freeware for leukoaraiosis and white matter segmentation. We dedicate this pilot study to the memory of J.S. Gravenstein, M.D. (Department of Anesthesiology, University of Florida), a valued mentor and advocate for research on postoperative cognitive dysfunction.

Disclosure of Funding: National Institutes of Health F32 AG021363 from the National Institute on Aging, Bethesda, Maryland (CP); The University of Florida Opportunity Fund (TM), NIH General Clinical Research Center Grant # UL1 TR000064; I. Heermann Anesthesia Foundation, Gainesville, Florida (TM, CP).

References

1. Rasmussen LS, Larsen K, Houx P, Skovgaard LT, Hanning CD, Moller JT. The assessment of postoperative cognitive function. *Acta Anaesthesiol Scand*. 2001; 45:275–89. [PubMed: 11207462]
2. Moller JT, Cluitmans P, Rasmussen LS, Houx P, Rasmussen H, Canet J, Rabbitt P, Jolles J, Larsen K, Hanning CD, Langeron O, Johnson T, Lauen PM, Kristensen PA, Biedler A, van Beem H, Fraidakis O, Silverstein JH, Beneken JE, Gravenstein JS. Long-term postoperative cognitive

- dysfunction in the elderly ispod1 study. ISPOCD investigators. International study of post-operative cognitive dysfunction. *Lancet*. 1998; 351:857–61. [PubMed: 9525362]
3. Monk TG, Weldon BC, Garvan CW, Dede DE, van der Aa MT, Heilman KM, Gravenstein JS. Predictors of cognitive dysfunction after major noncardiac surgery. *Anesthesiology*. 2008; 108:18–30. [PubMed: 18156878]
 4. Steinmetz J, Christensen KB, Lund T, Lohse N, Rasmussen LS. Long-term consequences of postoperative cognitive dysfunction. *Anesthesiology*. 2009; 110:548–55. [PubMed: 19225398]
 5. Price CC, Garvan CW, Monk TG. Type and severity of cognitive decline in older adults after noncardiac surgery. *Anesthesiology*. 2008; 108:8–17. [PubMed: 18156877]
 6. Squire LR. Memory and the hippocampus: A synthesis from findings with rats, monkeys, and humans. *Psychol Rev*. 1992; 99:195–231. [PubMed: 1594723]
 7. Zola-Morgan S, Squire LR, Amaral DG. Human amnesia and the medial temporal region: Enduring memory impairment following a bilateral lesion limited to field ca1 of the hippocampus. *J Neurosci*. 1986; 6:2950–67. [PubMed: 3760943]
 8. Reitz C, Brickman AM, Brown TR, Manly J, DeCarli C, Small SA, Mayeux R. Linking hippocampal structure and function to memory performance in an aging population. *Archives of neurology*. 2009; 66:1385–92. [PubMed: 19901171]
 9. Braak H, Braak E. morphology of alzheimer disease. *Fortschr Med*. 1990; 108:621–4. [PubMed: 2289729]
 10. Foundas AL, Leonard CM, Mahoney SM, Agee OF, Heilman KM. Atrophy of the hippocampus, parietal cortex, and insula in alzheimer's disease: A volumetric magnetic resonance imaging study. *Neuropsychiatry Neuropsychol Behav Neurol*. 1997; 10:81–9. [PubMed: 9150507]
 11. Jack CR Jr, Petersen RC, Xu YC, O'Brien PC, Smith GE, Ivnik RJ, Boeve BF, Waring SC, Tangalos EG, Kokmen E. Prediction of ad with mri-based hippocampal volume in mild cognitive impairment. *Neurology*. 1999; 52:1397–403. [PubMed: 10227624]
 12. Xu Y, Jack CR Jr, O'Brien PC, Kokmen E, Smith GE, Ivnik RJ, Boeve BF, Tangalos RG, Petersen RC. Usefulness of mri measures of entorhinal cortex versus hippocampus in ad. *Neurology*. 2000; 54:1760–7. [PubMed: 10802781]
 13. Sapolsky RM. Glucocorticoids and hippocampal atrophy in neuropsychiatric disorders. *Arch Gen Psychiatry*. 2000; 57:925–35. [PubMed: 11015810]
 14. Lozovaya N, Miller AD. Chemical neuroimmunology: Health in a nutshell bidirectional communication between immune and stress (limbic-hypothalamic-pituitary-adrenal) systems. *Chembiochem*. 2003; 4:466–84. [PubMed: 12794857]
 15. Shu XJ, Xue L, Liu W, Chen FY, Zhu C, Sun XH, Wang XP, Liu ZC, Zhao H. More vulnerability of left than right hippocampal damage in right-handed patients with post-traumatic stress disorder. *Psychiatry Res*. 2013; 212:237–44. [PubMed: 23149034]
 16. Smith ME. Bilateral hippocampal volume reduction in adults with post-traumatic stress disorder: A meta-analysis of structural mri studies. *Hippocampus*. 2005; 15:798–807. [PubMed: 15988763]
 17. Jia F, Mao Q, Liang YM, Jiang JY. Effect of post-traumatic mild hypothermia on hippocampal cell death after traumatic brain injury in rats. *J Neurotrauma*. 2009; 26:243–52. [PubMed: 19236165]
 18. Hachinski VC, Potter P, Merskey H. Leuko-araiosis: An ancient term for a new problem. *Can J Neurol Sci*. 1986; 13:533–4. [PubMed: 3791068]
 19. Ylikoski A, Erkinjuntti T, Raininko R, Sarna S, Sulkava R, Tilvis R. White matter hyperintensities on mri in the neurologically nondiseased elderly. Analysis of cohorts of consecutive subjects aged 55 to 85 years living at home. *Stroke*. 1995; 26:1171–7. [PubMed: 7604409]
 20. Pantoni L, Garcia JH. Pathogenesis of leukoaraiosis: A review. *Stroke*. 1997; 28:652–9. [PubMed: 9056627]
 21. Price CC, Mitchell SM, Brumback B, Tanner JJ, Schmalfuss I, Lamar M, Giovannetti T, Heilman K, Libon DJ. Mri-leukoaraiosis thresholds and the phenotypic expression of dementia. *Neurology*. 2012; 79:734–40. [PubMed: 22843264]
 22. Brickman AM, Siedlecki KL, Muraskin J, Manly JJ, Luchsinger JA, Yeung LK, Brown TR, Decarli C, Stern Y. White matter hyperintensities and cognition: Testing the reserve hypothesis. *Neurobiol Aging*. 2011; 32:1588–98. [PubMed: 19926168]

23. Libon DJ, Price CC, Davis Garrett K, Giovannetti T. From binswanger's disease to leuokoaraiosis: What we have learned about subcortical vascular dementia. *Clin Neuropsychol*. 2004; 18:83–100. [PubMed: 15595361]
24. Chui HC, Victoroff JJ, Margolin D, Jagust W, Shankle R, Katzman R. Criteria for the diagnosis of ischemic vascular dementia proposed by the state of california alzheimer's disease diagnostic and treatment centers. *Neurology*. 1992; 42:473–80. [PubMed: 1549205]
25. Ishii N, Nishihara Y, Imamura T. Why do frontal lobe symptoms predominate in vascular dementia with lacunes? *Neurology*. 1986; 36:340–5. [PubMed: 3951700]
26. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: Development and validation. *J Chronic Dis*. 1987; 40:373–83. [PubMed: 3558716]
27. Lawton MP, Brody EM. Assessment of older people: Self-maintaining and instrumental activities of daily living. *Gerontologist*. 1969; 9:179–86. [PubMed: 5349366]
28. Inouye SK, van Dyck CH, Alessi CA, Balkin S, Siegel AP, Horwitz RI. Clarifying confusion: The confusion assessment method. A new method for detection of delirium. *Ann Intern Med*. 1990; 113:941–8. [PubMed: 2240918]
29. Markus HS, Ackerstaff R, Babikian V, Bladin C, Droste D, Grosset D, Levi C, Russell D, Siebler M, Tegeler C. Intercenter agreement in reading doppler embolic signals. A multicenter international study. *Stroke*. 1997; 28:1307–10. [PubMed: 9227673]
30. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res*. 1975; 12:189–98. [PubMed: 1202204]
31. Wechsler, D. Wechsler abbreviated scale of intelligence. San Antonio, TX: Harcourt Assessment; 1999 Feb 01.
32. Lezak, MD.; Howieson, DB.; Loring, DW. Neuropsychological assessment. Fourth edition. New York: Oxford University Press; 2004.
33. Price CC, Wood MF, Leonard CM, Towler S, Ward J, Montijo H, Kellison I, Bowers D, Monk T, Newcomer JC, Schmalfuss I. Entorhinal cortex volume in older adults: Reliability and validity considerations for three published measurement protocols. *J Int Neuropsychol Soc*. 2010; 16:846–55. [PubMed: 20937164]
34. Yesavage JA. Geriatric depression scale. *Psychopharmacol Bull*. 1988; 24:709–11. [PubMed: 3249773]
35. Spielberger, CD. Self-evaluation questionnaire. Palo Alto: Cons Psych Press; 1968.
36. Price DD, Bush FM, Long S, Harkins SW. A comparison of pain measurement characteristics of mechanical visual analogue and simple numerical rating scales. *Pain*. 1994; 56:217–26. [PubMed: 8008411]
37. Duvernoy, HM. The human hippocampus : Functional anatomy, vascularization, and serial sections with mri. 3. Berlin ; New York: Springer; 2005.
38. Zijdenbos AP, Dawant BM. Brain segmentation and white matter lesion detection in mr images. *Crit Rev Biomed Eng*. 1994; 22:401–65. [PubMed: 8631195]
39. Zijdenbos AP, Dawant BM, Margolin RA, Palmer AC. Morphometric analysis of white matter lesions in mr images: Method and validation. *IEEE Trans Med Imaging*. 1994; 13:716–24. [PubMed: 18218550]
40. Insausti R, Juottonen K, Soininen H, Insausti AM, Partanen K, Vainio P, Laakso MP, Pitkanen A. Mr volumetric analysis of the human entorhinal, perirhinal, and temporopolar cortices. *AJNR Am J Neuroradiol*. 1998; 19:659–71. [PubMed: 9576651]
41. Barta PE, Dhingra L, Royall R, Schwartz E. Improving stereological estimates for the volume of structures identified in three-dimensional arrays of spatial data. *J Neurosci Methods*. 1997; 75:111–8. [PubMed: 9288642]
42. Abramooff M, Magelhaes P, Ram S. Image processing with imagej. *Biophotonics International*. 2004; 11:36–42.
43. Towler SD, Price CC, Mitchell SM, Libon DJ. White Matter Hyperintensity Quantification in T2 FLAIR MRI: A Reliability Study. *JINS*. 2009; 15:148. [PubMed: 19128539]
44. Gerraty RP, Parsons MW, Alan Barber P, Darby DG, Davis SM. The volume of lacunes. *Stroke*. 2001; 32:1937–8. [PubMed: 11486134]

45. Bigler ED, Tate DF. Brain volume, intracranial volume, and dementia. *Invest Radiol*. 2001; 36:539–46. [PubMed: 11547042]
46. Smith SM, Jenkinson M, Woolrich MW, Beckmann CF, Behrens TE, Johansen-Berg H, Bannister PR, De Luca M, Drobnjak I, Flitney DE, Niazy RK, Saunders J, Vickers J, Zhang Y, De Stefano N, Brady JM, Matthews PM. Advances in functional and structural mr image analysis and implementation as fsl. *Neuroimage*. 2004; 23:S208–19. [PubMed: 15501092]
47. Shattuck DW, Prasad G, Mirza M, Narr KL, Toga AW. Online resource for validation of brain segmentation methods. *Neuroimage*. 2009; 45:431–9. [PubMed: 19073267]
48. Lewis MS, Maruff P, Silbert BS, Evered LA, Scott DA. The sensitivity and specificity of three common statistical rules for the classification of post-operative cognitive dysfunction following coronary artery bypass graft surgery. *Acta Anaesthesiol Scand*. 2006; 50:50–7. [PubMed: 16451151]
49. Lewis M, Maruff P, Silbert B. Statistical and conceptual issues in defining post-operative cognitive dysfunction. *Neurosci Biobehav Rev*. 2004; 28:433–40. [PubMed: 15289007]
50. Aiken, LS.; West, SG. Multiple regression: Testing and interpreting interactions. Thousand Oaks: Sage Publications; 1991.
51. Wechsler, D. Wechsler adult intelligence scale. third edition. San Antonio, Texas: The Psychological Corporation; 1997.
52. Glosser G, Butters N, Kaplan E. Visuo-perceptual processes in brain damaged patients on the digit symbol substitution test. *Int J Neurosci*. 1977; 7:59–66. [PubMed: 893004]
53. Greene NH, Attix DK, Weldon BC, Smith PJ, McDonagh DL, Monk TG. Measures of executive function and depression identify patients at risk for postoperative delirium. *Anesthesiology*. 2009; 110:788–95. [PubMed: 19326494]
54. MacDonald AW 3rd, Cohen JD, Stenger VA, Carter CS. Dissociating the role of the dorsolateral prefrontal and anterior cingulate cortex in cognitive control. *Science*. 2000; 288:1835–8. [PubMed: 10846167]
55. Christensen TA, Lockwood JL, Almryde KR, Plante E. Neural substrates of attentive listening assessed with a novel auditory stroop task. *Front Hum Neurosci*. 2011; 4:236. [PubMed: 21258643]
56. Floden D, Vallesi A, Stuss DT. Task context and frontal lobe activation in the stroop task. *J Cogn Neurosci*. 2011; 23:867–79. [PubMed: 20350183]
57. Roberts KL, Hall DA. Examining a supramodal network for conflict processing: A systematic review and novel functional magnetic resonance imaging data for related visual and auditory stroop tasks. *J Cogn Neurosci*. 2008; 20:1063–78. [PubMed: 18211237]
58. Lund C, Sundet K, Tennoe B, Hol PK, Rein KA, Fosse E, Russell D. Cerebral ischemic injury and cognitive impairment after off-pump and on-pump coronary artery bypass grafting surgery. *Ann Thorac Surg*. 2005; 80:2126–31. [PubMed: 16305857]
59. Andréll C, Jensen C, Norrsell H, Ekre O, Ekholm S, Norrsell U, Eliasson T, Mannheimer C, Blomstrand C. White matter disease in magnetic resonance imaging predicts cerebral complications after coronary artery bypass grafting. *Ann Thorac Surg*. 2005; 79:74–80. [PubMed: 15620918]
60. Smith PL, Treasure T, Newman SP, Joseph P, Ell PJ, Schneidau A, Harrison MJ. Cerebral consequences of cardiopulmonary bypass. *Lancet*. 1986; 1:823–5. [PubMed: 2870314]
61. Floyd TF, McGarvey M, Ochroch EA, Cheung AT, Augoustides JA, Bavaria JE, Acker MA, Pochettino A, Detre JA. Perioperative changes in cerebral blood flow after cardiac surgery: Influence of anemia and aging. *Ann Thorac Surg*. 2003; 76:2037–42. [PubMed: 14667637]
62. Cook DJ, Oliver WC Jr, Orszulak TA, Daly RC, Bryce RD. Cardiopulmonary bypass temperature, hematocrit, and cerebral oxygen delivery in humans. *Ann Thorac Surg*. 1995; 60:1671–7. [PubMed: 8787461]
63. Li M, Bertout JA, Ratcliffe SJ, Eckenhoff MF, Simon MC, Floyd TF. Acute anemia elicits cognitive dysfunction and evidence of cerebral cellular hypoxia in older rats with systemic hypertension. *Anesthesiology*. 2011; 113:845–58. [PubMed: 20808217]

64. Newman MF, Croughwell ND, Blumenthal JA, Lowry E, White WD, Spillane W, Davis RD Jr, Glower DD, Smith LR, Mahanna EP, Reves JG. Predictors of cognitive decline after cardiac operation. *Ann Thorac Surg.* 1995; 59:1326–30. [PubMed: 7733762]
65. Mutch WA, Ryner LN, Kozlowski P, Scarth G, Warrian RK, Lefevre GR, Wong TG, Thiessen DB, Girling LG, Doiron L, McCudden C, Saunders JK. Cerebral hypoxia during cardiopulmonary bypass: A magnetic resonance imaging study. *Ann Thorac Surg.* 1997; 64:695–701. [PubMed: 9307459]
66. Browne SM, Halligan PW, Wade DT, Taggart DP. Postoperative hypoxia is a contributory factor to cognitive impairment after cardiac surgery. *J Thorac Cardiovasc Surg.* 2003; 126:1061–4. [PubMed: 14566247]
67. Goto T, Yoshitake A, Baba T, Shibata Y, Sakata R, Uozumi H. Cerebral ischemic disorders and cerebral oxygen balance during cardiopulmonary bypass surgery: Preoperative evaluation using magnetic resonance imaging and angiography. *Anesth Analg.* 1997; 84:5–11. [PubMed: 8988991]
68. Edmonds CR, Barbut D, Hager D, Sharrock NE. Intraoperative cerebral arterial embolization during total hip arthroplasty. *Anesthesiology.* 2000; 93:315–8. [PubMed: 10910475]
69. Jacobs A, Neveling M, Horst M, Ghaemi M, Kessler J, Eichstaedt H, Rudolf J, Model P, Bonner H, de Vivie ER, Heiss WD. Alterations of neuropsychological function and cerebral glucose metabolism after cardiac surgery are not related only to intraoperative microembolic events. *Stroke.* 1998; 29:660–7. [PubMed: 9506609]
70. O'Brien DJ, Bauer RM, Yarandi H, Knauf DG, Bramblett P, Alexander JA. Patient memory before and after cardiac operations. *J Thorac Cardiovasc Surg.* 1992; 104:1116–24. [PubMed: 1405672]
71. Stygall J, Suvarna S, Harrington J, Hayward M, Walesby RK, Newman SP. Effect on the brain of two techniques of myocardial protection. *Asian Cardiovasc Thorac Ann.* 2009; 17:259–65. [PubMed: 19643849]
72. Pugsley W, Klinger L, Paschalis C, Treasure T, Harrison M, Newman S. The impact of microemboli during cardiopulmonary bypass on neuropsychological functioning. *Stroke.* 1994; 25:1393–9. [PubMed: 8023354]
73. Caplan LR, Hennerici M. Impaired clearance of emboli (washout) is an important link between hypoperfusion, embolism, and ischemic stroke. *Arch Neurol.* 1998; 55:1475–82. [PubMed: 9823834]
74. Floyd TF, Shah PN, Price CC, Harris F, Ratcliffe SJ, Acker MA, Bavaria JE, Rahmouni H, Kuersten B, Wiegers S, McGarvey ML, Woo JY, Pochettino AA, Melhem ER. Clinically silent cerebral ischemic events after cardiac surgery: Their incidence, regional vascular occurrence, and procedural dependence. *Ann Thorac Surg.* 2006; 81:2160–6. [PubMed: 16731147]
75. Selnes OA, Grega MA, Borowicz LM Jr, Royall RM, McKhann GM, Baumgartner WA. Cognitive changes with coronary artery disease: A prospective study of coronary artery bypass graft patients and nonsurgical controls. *Ann Thorac Surg.* 2003; 75:1377–84. [PubMed: 12735550]
76. Treasure T, Smith PL, Newman S, Schneidau A, Joseph P, Ell P, Harrison MJ. Impairment of cerebral function following cardiac and other major surgery. *Eur J Cardiothorac Surg.* 1989; 3:216–21. [PubMed: 2624784]
77. Abildstrom H, Rasmussen LS, Rentowl P, Hanning CD, Rasmussen H, Kristensen PA, Moller JT. Cognitive dysfunction 1-2 years after non-cardiac surgery in the elderly. Ispocd group International study of post-operative cognitive dysfunction. *Acta Anaesthesiol Scand.* 2000; 44:1246–51. [PubMed: 11065205]
78. Gerriets T, Schwarz N, Bachmann G, Kaps M, Kloeveborn WP, Sammer G, Tschernatsch M, Nottbohm R, Blaes F, Schonburg M. Evaluation of methods to predict early long-term neurobehavioral outcome after coronary artery bypass grafting. *Am J Cardiol.* 2010; 105:1095–101. [PubMed: 20381659]
79. Goto M, Abe O, Miyati T, Yoshikawa T, Hayashi N, Takao H, Inano S, Kabasawa H, Mori H, Kunimatsu A, Aoki S, Ino K, Iida K, Yano K, Ohtomo K. Entorhinal cortex volume measured with 3t mri is positively correlated with the wechsler memory scale-revised logical/verbal memory score for healthy subjects. *Neuroradiology.* 2011; 53:617–22. [PubMed: 21455719]

80. Edelstyn NM, Ellis SJ, Jenkinson P, Sawyer A. Contribution of the left dorsomedial thalamus to recognition memory: A neuropsychological case study. *Neurocase*. 2002; 8:442–52. [PubMed: 12529453]
81. Miyamoto M, Kato J, Narumi S, Nagaoka A. Characteristics of memory impairment following lesioning of the basal forebrain and medial septal nucleus in rats. *Brain Res*. 1987; 419:19–31. [PubMed: 3676724]
82. Petersen RC, Smith GE, Waring SC, Ivnik RJ, Tangalos EG, Kokmen E. Mild cognitive impairment: Clinical characterization and outcome. *Arch of Neurol*. 1999; 56:303–8. [PubMed: 10190820]
83. Grothe M, Zaborszky L, Atienza M, Gil-Neciga E, Rodriguez-Romero R, Teipel SJ, Amunts K, Suarez-Gonzalez A, Cantero JL. Reduction of basal forebrain cholinergic system parallels cognitive impairment in patients at high risk of developing alzheimer's disease. *Cereb Cortex*. 20:1685–95. [PubMed: 19889714]
84. Tremont G, Miele A, Smith MM, Westervelt HJ. Comparison of verbal memory impairment rates in mild cognitive impairment. *J Clin Exp Neuropsychol*. 2010; 32:630–6. [PubMed: 20603742]
85. Christidi F, Bigler ED, McCauley SR, Schnelle KP, Merkley TL, Li X, Macleod MC, Chu Z, Hunter JV, Levin H, Clifton G, Wilde EA. Diffusion tensor imaging of the perforant pathway zone and its relation to memory function in patients with severe traumatic brain injury. *J Neurotrauma*. 2011; 28:711–25. [PubMed: 21381986]
86. Bigler ED, Neeley ES, Miller MJ, Tate DF, Rice SA, Cleavinger H, Wolfson L, Tschanz J, Welsh-Bohmer K. Cerebral volume loss, cognitive deficit and neuropsychological performance: Comparative measures of brain atrophy: I. Dementia. *JINS*. 2004; 10:442–52. [PubMed: 15147601]
87. van Norden AG, de Laat KF, Fick I, van Uden IW, van Oudheusden LJ, Gons RA, Norris DG, Zwiers MP, Kessels RP, de Leeuw FE. Diffusion tensor imaging of the hippocampus and verbal memory performance: The run dmc study. *Hum Brain Mapp*. 2012; 33:542–51. [PubMed: 21391278]
88. Hudetz JA, Patterson KM, Amole O, Riley AV, Pagel PS. Postoperative cognitive dysfunction after noncardiac surgery: Effects of metabolic syndrome. *J Anesth*. 2011; 25:337–44. [PubMed: 21516370]
89. Sullivan EV, Pfefferbaum A. Diffusion tensor imaging and aging. *Neurosci Behav Rev*. 2006:749–61.
90. Luyt CE, Galanaud D, Perlberg V, Vanhaudenhuyse A, Stevens RD, Gupta R, Besancenot H, Krainik A, Audibert G, Combes A, Chastre J, Benali H, Laureys S, Puybasset L. for the Neuro Imaging for Coma E, Recovery C. Diffusion tensor imaging to predict long-term outcome after cardiac arrest: A bicentric pilot study. *Anesthesiology*. 2012; 117:1311–21. [PubMed: 23135257]
91. Galanaud D, Perlberg V, Gupta R, Stevens RD, Sanchez P, Tollard E, de Champfleure NM, Dinkel J, Faivre S, Soto-Ares G, Veber B, Cottenceau V, Masson F, Tourdias T, Andre E, Audibert G, Schmitt E, Ibarrola D, Dailler F, Vanhaudenhuyse A, Tshibanda L, Payen JF, Le Bas JF, Krainik A, Bruder N, Girard N, Laureys S, Benali H, Puybasset L. for the Neuro Imaging for Coma E, Recovery C. Assessment of white matter injury and outcome in severe brain trauma: A prospective multicenter cohort. *Anesthesiology*. 2012; 117:1300–10. [PubMed: 23135261]
92. Shioiri A, Kurumaji A, Takeuchi T, Matsuda H, Arai H, Nishikawa T. White matter abnormalities as a risk factor for postoperative delirium revealed by diffusion tensor imaging. *Am J Geriatr Psychiatry*. 2010; 18:743–53. [PubMed: 20220599]