Methamphetamine-Like Discriminative-Stimulus Effects of Nicotinic Agonists

Rajeev I. Desai and Jack Bergman

Preclinical Pharmacology Laboratory, McLean Hospital/Harvard Medical School, Belmont, Massachusetts

Received November 11, 2013; accepted January 2, 2014

ABSTRACT

Nicotine was recently shown to engender *d*-methamphetamine (MA)-like discriminative-stimulus effects in rats, which may be indicative of shared psychomotor stimulant properties. To further investigate such overlapping discriminative-stimulus effects, nicotinic agonists varying in efficacy and selectivity were studied in squirrel monkeys that discriminated a moderate intramuscular dose of MA (0.1 mg/kg) from vehicle. These included $\alpha 4\beta 2$ -selective ligands that may vary in efficacy from relatively high [nicotine, (+)- and (–)-epibatidine] to relatively low [isoarecolone, varenicline, (–)-cytisine, (–)-lobeline] and the $\alpha 7$ -selective ligands anabaseine and anabasine. Results show that nicotine, (+)-epibatidine, and (–)-epibatidine substituted fully for MA, whereas the highest doses of other nicotinic agonists produced intermediate levels of MA-like effects (isoarecolone, anabaseine, anabasine, and varenicline) or did not substitute for MA [(–)-cytisine and (–)-lobeline]. The relative

potencies of nicotinic agonists, based on effective dose₅₀ (ED₅₀) values, corresponded more closely with their relative affinities at $\alpha 4\beta 2$ than at $\alpha 7$ receptors. Regardless of selectivity or efficacy, nicotinic agonists also were observed to produce untoward effects, including salivation and emesis during or after experimental sessions. In pretreatment studies, the $\alpha 4\beta 2$ -selective antagonist dihydro- β -erythroidine hydrobromide (DH β E) (0.032 and 0.1 mg/kg) and the partial agonists varenicline (0.0032–0.1 mg/kg) and (–)-cytisine (0.032 and 0.1 mg/kg) surmountably antagonized (>10-fold rightward shift) nicotine's MA-like effects but were ineffective in blocking nicotine's emetic effects. Overall, our results show that 1) MA-like discriminative-stimulus effects of nicotinic agonists likely are mediated through $\alpha 4\beta 2$ nicotinic acetylcholine receptor actions, and 2) nicotinic $\alpha 4\beta 2$ partial agonists, like the nicotinic antagonist DH β E, can reduce MA-like behavioral effects of nicotine.

Introduction

The discriminative-stimulus effects of nicotine have been widely characterized in laboratory animals (Jutkiewicz et al., 2011; Cunningham et al., 2012; Smith and Stolerman, 2009), and have been related to its subjective effects among users of tobacco or other nicotine delivery devices (Smith and Stolerman, 2009; Benowitz, 2010). Pharmacological studies with selective agonists and antagonists additionally have identified likely mechanisms of action mediating the discriminative-stimulus effects of nicotine. For example, such effects of nicotine are readily mimicked by centrally acting nicotinic agonists with high affinity for the $\alpha 4\beta 2$ nicotinic acetylcholine receptor (nAChR) subtype, but not by drugs that act selectively at other subtypes of nAChR (e.g., $\alpha 3\beta 4$ or $\alpha 7$) or by drugs from other pharmacological classes (e.g., muscarinic agents; see Smith and Stolerman, 2009, for review). Moreover, noncompetitive (mecamylamine) and competitive [dihydro- β -erythroidine hydrobromide (DH β E)] antagonists that block $\alpha 4\beta 2$ nAChRs in the central nervous system attenuate nicotine's discriminative-stimulus effects, whereas peripherally restricted antagonists or antagonists at other nAChR subtypes (e.g., nicotinic $\alpha 7$) are

ineffective. In conjunction, such evidence strongly suggests that the discriminative-stimulus effects of nicotine are centrally mediated, primarily via $\alpha 4\beta 2$ nAChRs (Smith and Stolerman, 2009).

A growing body of evidence also indicates that, as with monoaminergic psychomotor stimulant drugs [e.g., cocaine, *d*-amphetamine, *d*-methamphetamine (MA)], the projection from the ventral tegmental area to the nucleus accumbens in the mesocorticolimbic dopamine (DA) system is a key element in brain circuitry that mediates the neurochemical and behavioral effects of nicotine. Accordingly, increases in DA neurotransmission have been proposed to mediate the reinforcing effects of nicotine and its consumption (Di Chiara, 2000; Smith and Stolerman, 2009). The involvement of common neural substrates also has led to the suggestion that nicotine and monoaminergic psychomotor stimulant drugs might engender overlapping subjective effects and, in laboratory animals, discriminative-stimulus effects (Smith and Stolerman, 2009). Data from some, but not all, previous studies in nicotine- and cocaine- or *d*-amphetamine-trained subjects have supported this suggestion (see Smith and Stolerman, 2009, for review), and have profitably advanced our understanding of the stimulant-like effects of nicotine and other nicotinic ligands. For example, our recent MA-discrimination studies in rats suggest that 1) nicotinic agonists vary in the extent to which they produce

This research was supported by the National Institutes of Health National Institute on Drug Abuse [Grant R21-DA026548] (to R.I.D.).
dx.doi.org/10.1124/jpet.113.211235.

ABBREVIATIONS: DA, dopamine; DH β E, dihydro- β -erythroidine hydrobromide; ED₅₀, effective dose₅₀; FR, fixed-ratio; LEDs, light-emitting diodes; MA, *d*-methamphetamine; R(–)-NPA, R(–)-10,11-dihydroxy-*N*-*n*-propyl noraporphine hydrochloride; SKF82958 hydrobromide, (±)-6-chloro-7,8-dihydroxy-3-allyl-1-phenyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine hydrobromide; TO, time-out; nAChR, nicotinic acetylcholine receptor.

MA-like stimulant effects, 2) $\alpha 4\beta 2$ nAChR-mediated actions play an important role in the MA-like stimulant effects of nicotinic agonists, and 3) varenicline dose-dependently antagonizes the MA-like stimulant actions of nicotine—consistent with the view that nicotinic partial agonists may help manage nicotine addiction and tobacco consumption (Rollema et al., 2007; Desai and Bergman, 2010).

The present research was conducted to further investigate the discriminative-stimulus effects of nicotine and related compounds. The goals of this work were to determine whether the discriminative-stimulus effects of MA and nicotine overlap in primate species and, if so, to examine the pharmacology of that overlap with a range of nAChR ligands. Using standard drug discrimination procedures, squirrel monkeys first were trained to distinguish a moderate dose of 0.1 mg/kg MA from saline. Next, the effects of monoamine uptake inhibitors (MA, cocaine), DA D₁- and D₂-like agonists [SKF82958 (($\pm$)-6-chloro-7,8-dihydroxy-3-allyl-1-phenyl-2,3,4,5-tetrahydro-1*H*-3-benzazepine hydrobromide), R-($-$)-NPA [R-($-$)-10,11-dihydroxy-*N*-*n*-propyl-noraporphine hydrochloride]], and a selective serotonin reuptake inhibitor (citalopram) were tested to confirm the role of dopaminergic mechanisms in these effects of MA (Tidey and Bergman, 1998). This provided a pharmacologically empirical basis for characterizing the effects of nicotinic ligands in MA-trained subjects. Subsequently, substitution tests with a wide range of nicotinic agonists were conducted to evaluate their ability to mimic MA's discriminative-stimulus effects. Drugs studied included $\alpha 4\beta 2$ nAChR subtype-selective ligands previously characterized as either full agonists [nicotine, (+)-epibatidine, ($-$)-epibatidine] or partial agonists [isoarecolone, varenicline, ($-$)-cytisine, ($-$)-lobeline]; Anderson and Arneric, 1994; Badio and Daly, 1994; Hahn et al., 2003; Rollema et al., 2007]. Substitution tests also were conducted with the $\alpha 7$ nAChR subtype-selective agonists anabasine and anabaseine (de Fiebre et al., 1995; Kem et al., 1997). Finally, drug interaction studies were conducted to compare modulation of the MA-like discriminative-stimulus effects of nicotine by the $\alpha 4\beta 2$ competitive antagonist DH β E (Williams and Robinson, 1984) and the partial agonists varenicline and ($-$)-cytisine. Overall, results show overlap in the discriminative-stimulus effects of nicotinic agonists and MA in nonhuman primates, and provide further support for the views that 1) MA-like stimulant effects of nicotinic agonists are primarily mediated through actions at $\alpha 4\beta 2$ nAChRs, and 2) nicotinic partial agonists that attenuate nicotine's stimulant-like discriminable effects may be useful pharmacotherapeutic adjuncts in the management of nicotine addiction.

Materials and Methods

Subjects. Four experimentally naïve adult male squirrel monkeys (*Saimiri sciureus*), weighing 650–900 g were subjects in the present studies. All subjects were individually housed in a climate-controlled vivarium under an automated 12-hour light/dark cycle. Except during testing, monkeys had unlimited access to water and were fed a daily allotment of high protein monkey chow (Purina Monkey Chow, St. Louis, MO), supplemented with fruit and multivitamins in the home cage. All monkeys were weighed daily, food intake was not restricted, and diets were adjusted as needed to maintain recommended body weights. Behavioral experiments were conducted daily (Monday–Friday) between 08:00 AM and 06:00 PM, under protocols that were approved by the Institutional Animal Care and Use Committee at McLean Hospital. Subjects were maintained in the McLean Animal

Care Facility in accordance with guidelines provided by the 2011 Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Animals Resources, Commission on Life Sciences, National Research Council. This facility is licensed by the U.S. Department of Agriculture.

Apparatus. The apparatus and methodology were comparable to those employed previously (Tidey and Bergman, 1998; Kangas et al., 2013). During experimental sessions, monkeys sat in customized Plexiglas chairs (Kelleher and Morse, 1968) that were enclosed in ventilated, sound-attenuating chambers provided with white noise at all times to mask extraneous sounds. While seated, monkeys faced a panel containing two sets of colored stimulus lights. Two response levers extended into the chamber, one below each set of stimulus lights, and were comfortably within the subject's reach. The two response levers were set 15 cm apart. Depression of either lever with a force greater than 0.2 N produced an audible click and was recorded as a response. Prior to each behavioral session, a shaved portion of the monkey's tail was secured under brass electrodes by a small stock and was coated with electrode paste to ensure a low-resistance electrical contact between electrodes and tail. Brief, low-intensity stimuli (200 milliseconds; 3 mA) could be delivered to the electrodes from a 60 Hz transformer. Experimental variables and data collection were controlled by PC computers with Med Associates interfacing equipment and operating software (MED-PC, MedState Notation; Med Associates, Inc., St. Albans, VT).

MA Discrimination. Subjects first were trained to press each of the two response levers under a 10-response fixed-ratio (FR) schedule of stimulus-termination. Under this schedule, a brief, mild electric stimulus (200 milliseconds, 3 mA) was programmed for delivery to the tail every 10 seconds during the illumination of red lights on the front panel. The completion of 10 consecutive lever-press responses (FR10) on one lever within 10 seconds turned off the red lights and the associated program of current delivery. The completion of each FR10 also initiated a 50-second time-out (TO) period, during which all lights in the chamber were extinguished and responding had no programmed consequences. The delivery of four electric stimuli prior to completion of the FR requirement also turned off all lights, terminated the program of stimulus delivery, and initiated the 50-second TO period. Once performance was stable on both levers under the FR10 response requirement, subjects initially were trained to discriminate intramuscular injections of 0.32 mg/kg MA from intramuscular injections of saline (Tidey and Bergman, 1998). Previous studies have indicated that the generalization profiles for many drugs greatly depend on training dose, and that higher doses generally are more pharmacologically restrictive than lower doses. For example, this type of relationship has been exploited by Spealman and colleagues to study the role of different monoaminergic mechanisms in the discriminative-stimulus effects of cocaine (Spealman, 1995). Thus, in the present study, we lowered the training dose of MA to 0.1 mg/kg to better assess the possibility of overlapping discriminative-stimulus effects among compounds from different pharmacological classes. After MA injection, only responses on one lever were reinforced; after saline injection, only responses on the other lever were reinforced. The assignment of MA-associated and saline-associated levers was counterbalanced across monkeys. During all training sessions, responses on the inappropriate lever reset the FR response requirement on the injection-appropriate lever.

When discrimination performance was stable from day to day at or above the criterion (90% accuracy), daily training sessions were extended to encompass one to four components. Each component, which consisted of 10 presentations of the FR10, TO 50-second schedule, was preceded by a 10-minute TO period during which vehicle or MA could be administered. The number of components in daily training sessions varied in a pseudo-random manner, with the stipulation that MA was injected only before the final component of the session. Additionally, sessions in which only saline was administered in all components occurred periodically to avoid an invariant association between injection of MA and the final session component.

Drug testing was initiated when >90% of responses occurred on the injection-appropriate lever during the preceding training session and four of the last five training sessions. Test sessions included four components, during which all schedule parameters and contingencies were identical to those in the training sessions, with the exception that 10 consecutive responses on either lever extinguished the red lights and terminated the associated program of current delivery. Testing was conducted once or twice per week with training sessions on intervening days. During test sessions, incremental doses of a test drug were administered at the beginning of the 10-minute TO periods preceding components of the test session (cumulative dosing). This procedure allowed determination of the effects of up to four cumulative doses during a single test session.

After all drugs were studied for their ability to substitute for MA, drug interaction studies were conducted to determine how pretreatment with selected compounds [varenicline, (–)-cytisine, and DHβE] modified the MA-like effects of nicotine. Pretreatment times were based on data from preliminary experiments and published reports (Stolerman et al., 1995, 1997; Rollema et al., 2007; Desai and Bergman, 2010). Studies were conducted by administering single doses of varenicline (0.0032–0.1 mg/kg), (–)-cytisine (0.032–0.1 mg/kg), or DHβE (0.032–0.1 mg/kg) 5 minutes prior to redetermination of the cumulative dose-response function for nicotine (0.032–1.0 mg/kg), i.e., 15 minutes prior to the first session component. Cumulative doses of nicotinic ligands in substitution studies and pretreatment doses in drug interaction studies were selected on the basis of preliminary dose-ranging experiments. Doses ranged from those without effect to those that: a) engendered nicotine-like effects or b) produced untoward physiologic effects that precluded further increase in either cumulative dose or, in drug interaction studies, pretreatment dose. In the latter case, observed effects were tabulated for presentation in a table.

Drugs. Methamphetamine hydrochloride, cocaine hydrochloride, (±)-SKF82958 hydrobromide, R(–)-NPA, (–)-nicotine hydrogen tartrate, and (–)-lobeline hydrochloride were obtained from Sigma-Aldrich (St. Louis, MO). Citalopram hydrobromide was generously

supplied by Lundbeck (Valby, Denmark). (–)-Cytisine [(1*R*,5*S*)-1,2,3,4,5,6-hexahydro-1,5-methano-8*H*-pyrido[1,2-*a*][1,5]diazocin-8-one], and anabesine hydrochloride [(*S*)-(+)-3-(2-piperidinyl)pyridine hydrochloride] were obtained from Tocris (Minneapolis, MN). Anabaseine dihydrochloride (3,4,5,6-tetrahydro-2,3'-bipyridine dihydrochloride), (+)-epibatidine [(2*R*)-2-(6-chloro-3-pyridinyl)-7-azabicyclo[2.2.1]heptane monohydrochloride], (–)-epibatidine [(2*R*)-2-(6-chloro-3-pyridinyl)-7-azabicyclo[2.2.1]heptane monohydrochloride], isoarecolone hydrochloride (1-methyl-4-acetyl-1,2,3,6-tetrahydropyridine hydrochloride), and dihydro-β-erythroidine hydrobromide [(2*S*,13*bS*)-2,3,5,6,8,9,10,13-octahydro-2-methoxy-1*H*,12*H*-benzo[*i*]pyrano[3,4-*g*]indolizin-12-one hydrobromide] were obtained from the National Institute on Drug Abuse (Bethesda, MD). Varenicline [6,7,8,9-tetrahydro-6,10-methano-6*H*-pyrazino[2,3-*b*][3]benzazepine] was generously donated by Dr. Hans Rollema (Pfizer Global Research and Development). All drugs were dissolved in 0.9% saline or water, and were injected by the intramuscular route of administration. The pH of nicotine and varenicline was adjusted as needed to 7.0 with 0.1 N sodium hydroxide. Doses of each drug are expressed in terms of the free base.

Data Analysis. Data from the test component immediately following injection were used to express the effects of the administered cumulative dose. The percentage MA-associated lever responding in each component of the session was calculated by dividing the number of responses on the injection lever by the total number of responses on both levers. Response rates were calculated for each component by dividing the total number of responses by the duration of the component minus time-out periods. If the mean response rate in a component was less than 0.2 responses per second, data from that component were excluded from further analysis. Mean results for vehicle and each dose of a drug were calculated by averaging data for the four subjects. Complete substitution with a dose of test drug alone or after pretreatment was defined for individual subjects and for the group of subjects as the allocation of ≥90% of total responses to the MA-associated lever. An intermediate level of responding (31 to 89%) on the MA-lever was defined as incomplete substitution, whereas the allocation of ≤30% of responses to the MA-lever was defined as no substitution.

TABLE 1

ED₅₀ values (± S.E.M.) and relative potencies with which nicotinic agonists produce MA-like effects in squirrel monkeys (nicotine = 1) and their affinity and relative affinity at α4β2 and α7 nACh receptors (nicotine = 1)

The in vitro values for inhibiting [³H]nicotine binding at α4β2 nicotinic receptors and [¹²⁵I]α-bungarotoxin binding at α7 nicotinic receptors are the average of values taken from the references cited below.

Drug	Doses	ED ₅₀ ± S.E.M.	Relative Potency (MA-Like Stimulus Effects)	In Vitro Affinity (K _i Values) at α4β2	Relative Affinity at α4β2 Receptors	In Vitro Affinity (K _i values) at α7	Relative Affinity at α7 Receptors
	mg/kg	mg/kg (μmol/kg)		nM		nM	
Nicotine	0.01–0.32	0.032 ± 0.014 (0.198)	1	3.4 ^{a-e,g-j,l}	1	4,895 ⁱ	1
(+)-Epibatidine	0.0001–0.001	0.0005 ± 0.00003 (0.0024)	0.015	0.05 ^{h,j}	0.015	255 ^f	0.052
(–)-Epibatidine	0.0001–0.001	0.0005 ± 0.00002 (0.0024)	0.015	0.06 ^{h,j}	0.018	109 ^f	0.022
Isoarecolone	0.32–10	2.8 ± 0.73 (19.3)	87.5	611 ⁱ	179.7	>100,000 ⁱ	20.43
Anabaseine	0.1–1.0	0.53 ± 0.17 (3.31)	16.6	32 ^m	9.41	58 ^m	0.012
Anabaseine	0.1–1.0	0.74 ± 0.09 (4.31)	23.1	260 ^m	76.5	58 ^m	0.012
Varenicline	0.032–0.18	0.11 ± 0.025 (0.52)	3.44	0.17 ^{k,l}	0.05	620 ^k	0.127
(–)-Cytisine	0.032–1.0	N.D.	N.D.	0.012–1.5 ^{a-e}	0.004–0.44	260–15,000 ^{a,d,n}	0.05–3.06
(–)-Lobeline	0.1–3.2	N.D.	N.D.	1.5–16 ^{b,o,p}	0.44–4.71	11,600–13,100 ^{n,p}	2.37–2.68

N.D., not determined.

^a Jensen et al., 2003.

^b Anderson and Arneric, 1994.

^c Xiao and Kellar, 2004.

^{d,e} Marks et al., 1986, 1996.

^f Sullivan et al., 1994.

^g Decker et al., 1995.

^h Badio and Daly, 1994.

ⁱ Hahn et al., 2003.

^j Damaj et al., 1994.

^{k,l} Rollema et al., 2007, 2010.

^m Kem et al., 1997.

ⁿ de Fiebre et al., 1995.

^o Damaj et al., 1997.

^p Miller et al., 2003.

Data were further analyzed to compare potency and maximum effects among drugs, to evaluate drug interactions, and to examine correspondence between the effects of nicotinic drugs in the present experiments and their published affinities for different subtypes of the nicotinic receptor. As appropriate, analysis of variance followed by Dunnett's *t* test or a paired *t* test was used to evaluate statistical significance of averaged data (defined at the 95% level of confidence; $P < 0.05$). When appropriate, interpolation or linear regression using Bioassay Software (Bioassay version Beta 6.2; MED Associates, Inc.) was used to calculate effective dose₅₀ (ED₅₀) values ($\pm$ S.E.M. for interpolation and 95% confidence limits for regression) from data points on the linear portions of the dose-response functions (Snedecor and Cochran, 1967). In experiments to evaluate varenicline-, (-)-cytisine-, and DH β E-nicotine interactions, ED₅₀ values were determined for nicotine alone and in the presence of each drug; pairs of ED₅₀ values were considered to be significantly different if their 95% confidence limits did not overlap. When significant differences in ED₅₀ values were obtained, relative potency estimates were calculated using standard parallel-line bioassay techniques described by Finney (1964). Finally, correspondence between the effects of drugs in drug discrimination and receptor binding studies was examined by comparing the relative potency of nicotinic drugs in the present experiments (i.e., ED₅₀ values divided by the ED₅₀ value for nicotine) and their published relative affinities for binding $\alpha 4\beta 2$ and $\alpha 7$ nicotinic receptors in vitro (K_i values divided by the K_i value for nicotine). Relative affinity values for each drug were obtained from previously published radioligand binding experiments in rat brain. Data were taken from experiments using [³H]nicotine binding for the $\alpha 4\beta 2$ receptor subtype and [¹²⁵I] α -bungarotoxin (Bgt) binding for the $\alpha 7$ nicotinic receptor subtype, and affinities relative to nicotine were averaged across studies (see Table 1).

Results

MA Discrimination. The 0.1 mg/kg training dose of MA maintained reliable discriminative-stimulus control in all four subjects throughout the present studies (> 18 months). During testing sessions, maximum responding on the MA-associated lever in all subjects occurred following a cumulative dose of 0.1 mg/kg MA (Fig. 1; top panel). MA also produced a dose-related elevation in response rate, which increased to approximately 200% of vehicle control values following the training dose or the cumulative test dose of 0.1 mg/kg; however, differences in the magnitude of effect among individual subjects precluded statistical significance for the averaged data ($F_{4,15} = 2.17$; $P > 0.05$). Neither the position and slope of the cumulative dose-response function for MA discrimination (0.0032–0.1 mg/kg) nor its apparent rate-increasing effects varied significantly over the course of the present studies. Consequently, discrimination and response rate data for MA at the beginning and end of the present studies were averaged for control values and graphic presentation.

Dopaminergic Drugs. The administration of MA (0.0032–0.1 mg/kg) in test sessions produced dose-dependent substitution for the training dose of MA, with full substitution following the cumulative dose of 0.1 mg/kg MA (Fig. 1; top panel). Like MA, the nonselective monoamine transport blocker cocaine and the DA D₁- and D₂-like agonists SKF82958 and R(-)-NPA, respectively, produced dose-dependent and full substitution for 0.1 mg/kg MA, with maxima of 90–94% responding on the MA-associated lever following cumulative doses of 0.32 mg/kg cocaine, 0.1 mg/kg SKF82958, and 0.01 mg/kg R(-)-NPA (Fig. 1; top panel). In contrast, the serotonin-selective reuptake inhibitor citalopram did not substitute for the training dose of MA

Fig. 1. Effects of cumulative administration of the indirect monoamine agonists MA and cocaine, the direct DA D₁- and D₂-like agonists SKF82958 and R(-)-NPA, respectively, and the selective serotonin reuptake inhibitor, citalopram, in squirrel monkeys trained to discriminate 0.1 mg/kg MA from saline under a FR schedule of stimulus termination. Ordinates: percentage MA-associated responding (top), response rates (bottom). Abscissae: cumulative drug dose in mg/kg (log scale). Each data point represents the average of effects in four subjects tested at each dose. The percentage of responses emitted on the MA-associated lever was not plotted if fewer than three subjects responded at that dose. During control sessions, the training dose of MA produced $97.3\% \pm 1.8$ (S.E.M.) responding on the MA-associated lever, and injections of saline produced $1.1\% \pm 0.9$ (S.E.M.) responding on the saline-associated lever. V, vehicle.

(maximum: 2% drug-lever responding after a cumulative dose of 10 mg/kg; Fig. 1, top panel). Higher cumulative doses of citalopram were not studied to avoid untoward effects, e.g., convulsions, previously observed with high doses of serotonin-selective reuptake inhibitors (Spealman, 1995). Although cocaine, like MA, increased responding in a dose-related manner, response rates were not significantly changed from vehicle values by any of the drugs studied ($F_{s,4,15} \leq 1.23$; $P_s > 0.05$; Fig. 1; bottom panels).

Nicotinic Agonists. Nicotine produced dose-dependent increases in responding on the MA-associated lever and fully substituted for the 0.1 mg/kg training dose of MA following cumulative doses of 0.1 (90%) and 0.32 mg/kg (100%) nicotine (Fig. 2, top left panel). The (+)- and (-)-enantiomers of epibatidine also produced dose-dependent increases on the

Fig. 2. Effects of the cumulatively administered nicotinic agonists nicotine and the enantiomers of epibatidine, in squirrel monkeys trained to discriminate 0.1 mg/kg MA from saline. Abscissae and ordinates for the top and bottom panels are as in Fig. 1. See Fig. 1 for other details.

MA-associated lever, and both isomers produced $>85\%$ responding on the MA-associated lever following the cumulative dose of 0.001 mg/kg [mean $\pm$ S.E.M.: $86 \pm 14.3\%$ for (+)-epibatidine and $96.8 \pm 3.3\%$ for (-)-epibatidine; Fig. 2, top middle and right panels, respectively]. Neither nicotine nor the enantiomers of epibatidine significantly altered rates of responding from control values in the group of four monkeys (Fig. 2; bottom panels). However, the highest doses of each drug produced observable effects, including profuse salivation and emesis (see below and Table 2).

The nicotinic agonists isoarecolone, anabaseine, anabasine, and varenicline also produced dose-dependent increases in responding on the MA-associated lever (Fig. 3; top left panel). However, substitution was incomplete following the highest cumulative doses of these agonists, with maximal values ranging from approximately 50 to 65% responding on the MA-associated lever. Among these ligands, a clear plateau in MA-like effects indicative of partial agonist actions was observed only with varenicline: the mean MA-like effects of the highest cumulative pretreatment dose, 0.32 mg/kg, did not exceed those of the immediately preceding cumulative dose (0.18 mg/kg) and rates of responding were comparable to

control values. The two highest doses of isoarecolone also produced comparable effects; however, the 0.5 log unit increase in dose from 3.2 to 10 mg/kg produced a small, albeit nonsignificant, increase in responding on the MA-associated lever and a decrease in response rate to 60% of control values, precluding a more definitive characterization of its efficacy. No indication of a plateau in MA-like effects was apparent with either anabasine or anabaseine. Anabasine did not alter mean rates of responding over the range of doses studied, whereas anabaseine reduced response rate in a dose-dependent manner and, following the highest cumulative dose (3.2 mg/kg), nearly or completely eliminated responding. As with nicotine and regardless of the presence or absence of effects on response rates, the highest cumulative doses of each of these nicotinic agonists produced untoward physiologic signs that precluded further testing (see below and Table 2).

The rank order of potency with which nicotinic agonists produced MA-like effects was: (-)-epibatidine $\approx$ (+)-epibatidine $>$ nicotine $\approx$ varenicline $>$ anabaseine $>$ anabasine $>$ isoarecolone (Table 1). Based on ED_{50} estimates, isoarecolone and the two enantiomers of epibatidine were, respectively, the least and most potent nicotinic ligands in the present studies, and

TABLE 2

Cumulative intramuscular doses (mg/kg) of nicotinic agonist that produced observable untoward effects (excessive salivation, emesis, tremor, or convulsions) during test sessions ($n = 4$ Ss)

One or more effects were observed following injection of the listed cumulative dose either prior to or following the session component in which that dose was studied. The + sign represents the observation of each effect in an individual subject, and the – sign indicates that no untoward effect was observed at that dose in any subjects.

Drug	Doses	Salivation/ Foam	Emesis	Tremor	Convulsions
	mg/kg				
Vehicle	0	–	–	–	–
Nicotine	0.32	+	+	–	–
(+)-Epibatidine	0.001	+	+	–	–
(–)-Epibatidine	0.001	+	+	–	–
Isoarecolone	1.0	+	+	–	–
	3.2	+	+	–	–
	10.0	+	+	+	–
Anabaseine	1.0	+	+	–	–
	3.2	+	+	–	–
Anabasine	1.0	+	+	+	–
Varenicline	0.1	+	+	–	–
	0.18	+	+	–	–
	0.32	+	+	–	–
(–)-Cytisine	0.32	+	+	–	–
	1.0	+	+	+	–
(–)-Lobeline	3.2	+	+	+	+
	10.0	+	+	+	+

approximately 85–80-fold less and more potent than nicotine, respectively. The remaining drugs were approximately 3-fold (varenicline) and 10–20-fold (anabaseine, anabasine) less potent than nicotine in producing MA-like discriminative-stimulus effects (Table 1).

In contrast to the above ligands, (–)-cytisine and (–)-lobeline failed to substitute for the training dose of MA, producing $\leq 30\%$ responding on the MA-associated lever over the range of doses that could be studied (Fig. 3; top right panel). Although response rates were not significantly altered following the highest cumulative doses of (–)-cytisine and (–)-lobeline, untoward observable effects precluded the administration of higher doses (see below; Table 2).

Observable Effects of Nicotinic Agonists. The observable effects of nicotinic agonists in the present study included untoward physiologic signs following one or more cumulative doses of each drug (Table 2). Regardless of the presence or absence of MA-like discriminative-stimulus effects, the highest cumulative doses of each agonist produced profuse salivation in all subjects that often was followed by emesis. In addition to producing excessive salivation and emesis, the highest cumulative doses of isoarecolone, anabaseine, (–)-cytisine, and (–)-lobeline produced tremor in individual subjects (Table 2). Convulsions after cumulatively administered doses of 3.2 or 10 mg/kg lobeline also were noted in two subjects; these signs were quickly and completely attenuated by diazepam (1.0 mg/kg i.m.).

Antagonism of Nicotine's MA-like Effects. Pretreatment with the selective $\alpha 4\beta 2$ nicotinic antagonist, DH β E (0.032 or 0.1 mg/kg) antagonized the MA-like discriminative-stimulus effects of cumulatively administered nicotine (0.01–1.0 mg/kg; Fig. 4, top panel). Both doses of DH β E produced a comparable level of antagonism, evident as an approximately 10-fold rightward shift in the dose-response curve that yielded an approximately 10-fold increase in the ED₅₀ value for nicotine's MA-like effects (Fig. 4, top panel; Table 3). The highest cumulative dose of nicotine after

treatment with DH β E (1.0 mg/kg) produced approximately 70% responding on the MA-associated lever and an approximately 30 to 40% decrease in response rates (Fig. 4; bottom right panel).

Pretreatment with the nicotinic $\alpha 4\beta 2$ partial agonist varenicline (0.0032, 0.032, or 0.1 mg/kg) also antagonized MA-like discriminative-stimulus effects engendered by cumulatively administered nicotine (0.01–1.0 mg/kg). Like DH β E, all doses of varenicline shifted the nicotine dose-response curve rightward to a similar extent (Fig. 5, top left panel). Based on estimated ED₅₀ values, the potency of nicotine for producing MA-associated responding decreased approximately 12-fold in the presence of 0.0032 mg/kg varenicline, and approximately 16-fold following a 30-fold higher pretreatment dose of varenicline (0.1 mg/kg; Table 3). As in experiments with DH β E, the highest cumulative dose of nicotine (1.0 mg/kg) after treatment with the two highest doses of varenicline (0.032 and 0.1 mg/kg) produced a moderate (approximately 30–40%) but statistically non-significant decrease in response rates (Fig. 5; bottom left panel).

Like DH β E and varenicline, the nicotinic $\alpha 4\beta 2$ partial agonist (–)-cytisine (0.032 or 0.1 mg/kg) antagonized the MA-like discriminative-stimulus effects of cumulatively administered nicotine (0.01–1.0 mg/kg) by shifting its dose-effect curve rightward (Fig. 5, top right panel). Based on ED₅₀ values, the effects of nicotine were displaced approximately 7- and 14-fold rightward by pretreatment with 0.032 mg/kg and 0.1 mg/kg (–)-cytisine, respectively (Table 3). Like varenicline, pretreatment doses of (–)-cytisine did not substantively alter nicotine's effects on response rates, and only moderate (<50%) decreases from vehicle-control rates of responding were observed following their combination (Fig. 5, bottom right panel).

Although not studied separately, neither the $\alpha 4\beta 2$ antagonist DH β E nor the $\alpha 4\beta 2$ partial agonists varenicline and (–)-cytisine appeared to attenuate the emetic effects of cumulative doses of nicotine that produced full substitution for MA. As described above, tremor or frank convulsions following treatment with several nicotinic ligands was evident in the present studies. In this regard, higher doses of DH β E, like $\alpha 4\beta 2$ agonists, previously have been reported to produce seizure activity (Damaj et al., 1999; Dobelis et al., 2003). Consequently, higher pretreatment doses of DH β E, varenicline, or (–)-cytisine or higher cumulative doses of nicotine after pretreatment with these drugs were not studied so as to avoid further untoward effects in the present studies.

Discussion

The main objective of the present studies was to characterize discriminative-stimulus effects of nicotine and nicotinic ligands in monkeys that discriminated a moderate training dose of the monoaminergic stimulant MA (0.1 mg/kg). Initial experiments indicated that, as in previous studies with a higher MA training dose (0.32 mg/kg), indirect monoaminergic agonists (MA, cocaine) and direct DA D₁- and D₂-like receptor agonists (SKF82958 and R(–)-NPA, respectively) engendered dose-dependent and full substitution for MA (Tidey and Bergman, 1998). These results, supporting the view that DA-related mechanisms play a prominent role in the discriminative-stimulus effects of MA, provide a pharmacologically empirical basis for evaluating behavioral overlap

Fig. 3. (Left) Effects of the cumulatively administered nicotinic agonists, isoarecolone, anabaseine, anabasine, and varenicline in squirrel monkeys trained to discriminate 0.1 mg/kg MA from saline. (Right) Effects of the cumulatively administered nicotinic agonists, (-)-cytisine and (-)-lobeline in squirrel monkeys trained to discriminate 0.1 mg/kg MA from saline. See Fig. 1 for other details.

in the effects of drugs that act via different (e.g., nicotinic) receptor mechanisms.

Nicotine and the enantiomers of epibatidine also produced dose-dependent increases in MA-associated responding, and fully (or, in the case of (+)-epibatidine, nearly fully) substituted for MA without greatly altering response rates. The comparable effects of these nicotinic full agonists are consistent with their similar nicotinic $\alpha 4\beta 2$ subtype selectivity (see Table 1) and with previous drug discrimination data from nicotine-trained rodents (Reavill et al., 1987; Damaj et al., 1994). They contrast somewhat with data from MA-trained rats in which only nicotine fully substituted for the training dose of MA (Desai and Bergman, 2010). In those studies, 0.001 mg/kg of both (+)-epibatidine and (-)-epibatidine produced approximately 60 to 70% responding on the MA-associated lever, whereas a 3-fold increase in the dose of both enantiomers completely eliminated responding, precluding further testing. Differences in the two studies may reflect species-related differences in vulnerability to the rate-decreasing effects of the epibatidine enantiomers or, alternatively, differences in the resistance of responding to their behaviorally disruptive effects when it is maintained by stimulus-termination (present studies) or food presentation (previous studies). Notwithstanding these

considerations, the present findings clearly show that the discriminative-stimulus effects of nicotine and the enantiomers of epibatidine overlap substantively with those of monoaminergic stimulants like MA in primate species.

The nicotinic receptor ligand varenicline also engendered dose-related MA-like effects but, in contrast to nicotine and epibatidine, the highest doses produced only intermediate levels of substitution. Varenicline previously has been characterized as a nicotinic partial agonist at the $\alpha 4\beta 2$ receptor subtype (Rollema et al., 2007, 2010) and, depending on experimental conditions, may substitute partially or fully for nicotine in nicotine-trained rats and monkeys (Rollema et al., 2007; Smith et al., 2007; LeSage et al., 2009; Jutkiewicz et al., 2011; Cunningham et al., 2012). The plateau in the dose-effect function for varenicline at an intermediate level of responding on the MA-associated lever in the present experiments, in conjunction with its ability to antagonize the stimulant effects of nicotine in MA-trained rodents (Desai and Bergman, 2010), is consistent with its characterization as a nicotinic partial agonist.

Like varenicline, isoarecolone is characterized as an $\alpha 4\beta 2$ -selective ligand and can fully reproduce the discriminative-stimulus effects of nicotine in nicotine-trained rats (Reavill et al., 1987; Damaj et al., 1994). Based upon other behavioral

Fig. 4. Effects of pretreatment with the selective competitive $\alpha 4\beta 2$ nicotinic antagonist DH β E on MA-like responding produced by nicotine in squirrel monkeys trained to discriminate 0.1 mg/kg MA from saline. Abscissae: dose of cumulatively administered nicotine (mg/kg; log scale). See Fig. 1 for other details.

and biochemical findings, however, isoarecolone has been forwarded as a nicotinic partial agonist (Reavill et al., 1987; Whiteaker et al., 1995; Mirza et al., 1996; Hahn et al., 2003; Shoaib, 2006). The present findings that the highest doses of isoarecolone produced only an intermediate level of

substitution for MA might be considered supporting evidence for that view. However, this interpretation remains speculative in the absence of a more definitive characterization of isoarecolone's efficacy, e.g., evaluation of varenicline-like antagonism of nicotine's behavioral effects.

The minor tobacco alkaloids anabasine and anabaseine produced dose-related increases in responding on the MA-associated lever without nicotine-like full substitution or a varenicline-like plateau in MA-like effects. Both drugs previously have displayed $\alpha 7$ and, with lower efficacy, $\alpha 4\beta 2$ receptor-mediated agonist actions (Arendash et al., 1995; Kem et al., 1997; Stevens et al., 1998). Possibly, the limited nicotine-like effects of anabasine and anabaseine in the present experiments reflect relatively low efficacy at the $\alpha 4\beta 2$ receptor (Takada et al., 1989; Brioni et al., 1994; de Fiebre et al., 1995; Stolerman et al., 1995; Desai and Bergman, 2010). Alternatively, $\alpha 7$ -mediated actions of anabasine and anabaseine may have obscured the full expression of their MA-like effects. Although such explanations are speculative in the absence of further information, the present and previous findings in monkeys and rats (Desai and Bergman, 2010) show that anabasine and anabaseine can produce MA-like effects. Although limited, such stimulant-like effects of minor tobacco alkaloids, like those of nicotine, may contribute to the maintenance of tobacco consumption (Clemens et al., 2009; see Hoffman and Evans, 2013, for review).

(-)-Cytisine and (-)-lobeline, which have high nAChR affinity and $\alpha 4\beta 2$ subtype-selectivity (see Table 1), failed to engender MA-like discriminative-stimulus effects in the present studies. Previously, (-)-cytisine was shown to both partially substitute for nicotine in rats and block its discriminative-stimulus effects, consistent with its characterization as a nicotinic partial agonist (Stolerman et al., 1984; Reavill et al., 1990; Brioni et al., 1994; Jutkiewicz et al., 2011; Cunningham et al., 2012). The absence of MA-like effects in the present studies suggests that (-)-cytisine may have less of a stimulant action in primate species than other $\alpha 4\beta 2$ partial agonists such as varenicline. (-)-Lobeline, like (-)-cytisine, is considered a partial agonist at the $\alpha 4\beta 2$ nAChR but, in addition, appears to act through multiple mechanisms, including monoamine uptake inhibition (Damaj et al., 1997; Dwoskin and Crooks, 2002). Thus, (-)-lobeline has been shown to substitute for a low training dose of cocaine, yet

TABLE 3

Doses of nicotine alone and after pretreatment with DH β E, varenicline, or (-)-cytisine that are calculated to produce 50% responding on the MA-associated lever (ED_{50} values with 95% confidence intervals) and relative potencies of nicotine after pretreatment (ED_{50} nicotine/ ED_{50} nicotine after pretreatment)

Data were obtained in squirrel monkeys trained to discriminate intramuscular injections of 0.1 mg/kg (0.67 μ mol/kg) MA from saline.

Pretreatment Drug	Doses	ED_{50}	Relative Potency
	mg/kg	mg/kg (95% CI) (μ mol/kg)	
Nicotine alone	0.01–0.32	0.03 (0.02–0.05) (0.20)	1
DH β E	0.032	0.39 (0.19–1.90) (2.39)	0.08 (0.03–0.20)
	0.1	0.30 (0.12–2.76) (1.85)	0.11 (0.03–0.27)
Varenicline	0.0032	0.39 (0.19–2.55) (2.40)	0.09 (0.02–0.23)
	0.032	0.75 (0.45–1.69) (4.63) ^a	0.06 (0.01–0.14)
	0.1	0.53 (0.26–3.02) (3.25) ^a	0.07 (0.02–0.16) ^a
(-)-Cytisine	0.032	0.23 (-0.19–0.02) ^b (1.39)	0.13 (0.03–0.36)
	0.1	0.44 (0–0.13) ^b (2.72)	0.08 (0.02–0.23)

^aSignificant deviation from linearity.

^bEstimate due to nonsignificant regression.

Fig. 5. Effects of pretreatment with the partial nicotinic agonists varenicline and (–)-cytisine on MA-like responding produced by nicotine in squirrel monkeys trained to discriminate 0.1 mg/kg MA from saline. Abscissae: dose of cumulatively administered nicotine (mg/kg; log scale). See Fig. 1 for other details.

attenuate the effects of a higher training dose of cocaine or MA (Miller et al., 2001; Harrod et al., 2003; Desai et al., 2003; Cunningham et al., 2006). Further indicative of its poorly-understood actions, (–)-lobeline has been reported to produce nicotine-like effects in studies of locomotor activity, but not in place conditioning, self-administration, or drug discrimination studies (Fudala and Iwamoto, 1986; Corrigan and Coen, 1989; Reavill et al., 1990; Stolerman et al., 1995; Harrod et al., 2003). Although the absence of MA-like effects in the present study is not inconsistent with its characterization as a nicotinic partial agonist, additional data showing that (–)-lobeline, like varenicline, can antagonize such effects of nicotine would strengthen this categorization.

Although only isoarecolone and anabaseine decreased response rates, high cumulative doses of all nicotinic agonists produced untoward physiologic signs (emesis, tremor, or convulsions) that precluded the study of higher doses (see Table 2). However, profuse salivation and emesis alone did not appear to interfere with discrimination behavior, e.g., nicotine and epibatidine fully substituted for MA despite profuse salivation and emesis in all subjects. Although the precise mechanism responsible for these adverse physiologic signs remains unclear, it is notable that they can be produced by both $\alpha 4 \beta 2$ nAChR agonists and antagonists as well as by

$\alpha 7$ -selective ligands (Damaj et al., 1999; Dobelis et al., 2003). Thus, it is unlikely that these signs reflect actions at a single subtype of nicotinic receptor.

Pretreatment with the competitive antagonist DH β E and the $\alpha 4 \beta 2$ -selective partial agonists varenicline and (–)-cytisine shifted the dose-effect function for nicotine's MA-like effects rightward, complementing similar results in nicotine-trained rats (Stolerman et al., 1997; Jutkiewicz et al., 2011). Although each drug served as a surmountable antagonist, the range of antagonism was surprisingly limited, i.e., an approximately 1.0–1.25 log unit rightward shift in the nicotine dose-effect function. The limited range of antagonist actions of DH β E and (–)-cytisine might reflect the use of two antagonist doses spanning only a 0.5 log unit range; higher pretreatment doses might have led to additional antagonism. In the case of varenicline, however, the lowest pretreatment dose (0.0032 mg/kg) increased nicotine's ED₅₀ value approximately 10-fold, whereas 10- and 30-fold increases in pretreatment dose produced only a <2-fold further increase in nicotine's ED₅₀ value. The reasons for such limited dose dependence in the nicotine-antagonist effects of varenicline are uncertain but may be partly related to training dose. In previous studies of the same three antagonists in nicotine-trained rats, dose dependence was more evident in subjects that discriminated a low, rather

Fig. 6. Relationship between the relative potencies of nicotinic drugs in the present MA-discrimination studies and their relative affinities at $\alpha 4\beta 2$ and $\alpha 7$ nicotinic receptors in radioligand binding studies (see *Materials and Methods*). Abscissae: affinity relative to nicotine for inhibiting binding of radioligand to $\alpha 4\beta 2$ (left) and $\alpha 7$ (right) nicotinic receptors. Ordinates: potency of nicotinic drugs relative to nicotine, based on ED_{50} values, for engendering MA-associated lever responding (from Table 1). Isoarecolone was excluded from this correlation analysis at the $\alpha 7$ nicotinic receptor subtypes because affinity values obtained at this site are not clearly defined (see Table 1).

than high, dose of nicotine (Jutkiewicz et al., 2011). Possibly, a lower training dose of MA and a concomitant increase in nicotine's potency might also have revealed greater antagonist dose dependence in the present studies.

Comparison of the potencies of nicotinic agonists with their reported binding affinities at $\alpha 4\beta 2$ and $\alpha 7$ nicotinic receptor subtypes reveals some correspondence between their relative behavioral potencies and their relative potencies for inhibiting [3H]nicotine binding at the $\alpha 4\beta 2$ receptors ($r^2 = 0.83$, $P = 0.005$) but not [^{125}I] α -bungarotoxin binding at $\alpha 7$ receptors ($r^2 = 0.01$, $P = 0.83$; Fig. 6; Table 1). These observations parallel a similar analysis in MA-trained rats, and are consistent with the ability of the $\alpha 4\beta 2$ receptor blocker DH β E, but not the $\alpha 7$ receptor blocker MLA, to antagonize nicotine's discriminative-stimulus effects (Brioni et al., 1996; Desai and Bergman, 2010). In concert with the antagonist effects of DH β E and the partial agonists varenicline and (–)-cytisine in the present experiments, such correspondence provides added support for the idea that the stimulant-like effects of nicotine and other nicotinic agonists are predominantly mediated by their actions at the $\alpha 4\beta 2$ nAChR (Reavill et al., 1987, 1988; Stolerman et al., 1995; Desai and Bergman, 2010).

Acknowledgments

The authors thank Elise Trowel for expert technical support, Dr. Hans Rollema (Pfizer, Inc.) for providing varenicline, Drs. B. Kangas and C. A. Paronis for their comments on an earlier version of this manuscript, and the National Institute on Drug Abuse Drug Supply Program for providing enantiomers of epibatidine, isoarecolone, anabaseine, and DH β E.

Authorship Contributions

Participated in research design: Desai, Bergman.

Conducted experiments: Desai.

Performed data analysis: Desai.

Wrote or contributed to the writing of the manuscript: Desai, Bergman.

References

- Anderson DJ and Arneric SP (1994) Nicotinic receptor binding of [3H]cytisine, [3H]nicotine and [3H]methylcarbamylcholine in rat brain. *Eur J Pharmacol* **253**: 261–267.
- Arendash GW, Sengstock GJ, Sanberg PR, and Kem WR (1995) Improved learning and memory in aged rats with chronic administration of the nicotinic receptor agonist GTS-21. *Brain Res* **674**:252–259.
- Badio B and Daly JW (1994) Epibatidine, a potent analgesic and nicotinic agonist. *Mol Pharmacol* **45**:563–569.
- Benowitz NL (2010) Nicotine addiction. *N Engl J Med* **362**:2295–2303.
- Brioni JD, Kim DJ, O'Neill AB, Williams JE, and Decker MW (1994) Clozapine attenuates the discriminative stimulus properties of (–)-nicotine. *Brain Res* **643**:1–9.
- Brioni JD, Kim DJ, and O'Neill AB (1996) Nicotine cue: lack of effect of the $\alpha 7$ nicotinic receptor antagonist methyllycaconitine. *Eur J Pharmacol* **301**:1–5.
- Clemens KJ, Caillé S, Stinus L, and Cador M (2009) The addition of five minor tobacco alkaloids increases nicotine-induced hyperactivity, sensitization and intravenous self-administration in rats. *Int J Neuropsychopharmacol* **12**:1355–1366.
- Cunningham CS, Polston JE, Jany JR, Segert IL, and Miller DK (2006) Interaction of lobeline and nicotinic receptor ligands with the discriminative stimulus properties of cocaine and amphetamine. *Drug Alcohol Depend* **84**:211–222.
- Cunningham CS, Javors MA, and McMahon LR (2012) Pharmacologic characterization of a nicotine-discriminative stimulus in rhesus monkeys. *J Pharmacol Exp Ther* **341**:840–849.
- Corrigall WA and Coen KM (1989) Nicotine maintains robust self-administration in rats on a limited-access schedule. *Psychopharmacology (Berl)* **99**:473–478.
- Damaj MI, Creasy KR, Grove AD, Rosecrans JA, and Martin BR (1994) Pharmacological effects of epibatidine optical enantiomers. *Brain Res* **664**:34–40.
- Damaj MI, Patrick GS, Creasy KR, and Martin BR (1997) Pharmacology of lobeline, a nicotinic receptor ligand. *J Pharmacol Exp Ther* **282**:410–419.
- Damaj MI, Glassco W, Dukat M, and Martin BR (1999) Pharmacological characterization of nicotine-induced seizures in mice. *J Pharmacol Exp Ther* **291**:1284–1291.
- Decker MW, Brioni JD, Bannon AW, and Arneric SP (1995) Diversity of neuronal nicotinic acetylcholine receptors: lessons from behavior and implications for CNS therapeutics. *Life Sci* **56**:545–570.
- Desai RI and Bergman J (2010) Drug discrimination in methamphetamine-trained rats: effects of cholinergic nicotinic compounds. *J Pharmacol Exp Ther* **335**:807–816.
- Desai RI, Barber DJ, and Terry P (2003) Dopaminergic and cholinergic involvement in the discriminative stimulus effects of nicotine and cocaine in rats. *Psychopharmacology (Berl)* **167**:335–343.
- de Fiebre CM, Meyer EM, Henry JC, Muraskin SI, Kem WR, and Papke RL (1995) Characterization of a series of anabaseine-derived compounds reveals that the 3-(4-dimethylaminocinnamylidene) derivative is a selective agonist at neuronal nicotinic $\alpha 7/125I$ - α -bungarotoxin receptor subtypes. *Mol Pharmacol* **47**:164–171.
- Di Chiara G (2000) Role of dopamine in the behavioural actions of nicotine related to addiction. *Eur J Pharmacol* **393**:295–314.
- Dobelis P, Hutton S, Lu Y, and Collins AC (2003) GABAergic systems modulate nicotinic receptor-mediated seizures in mice. *J Pharmacol Exp Ther* **306**: 1159–1166.
- Dwoskin LP and Crooks PA (2002) A novel mechanism of action and potential use for lobeline as a treatment for psychostimulant abuse. *Biochem Pharmacol* **63**:89–98.
- Finney DJ (1964) *Statistical Method in Biological Assay*, Ed. 2nd. Hafner, New York.
- Fudala PJ and Iwamoto ET (1986) Further studies on nicotine-induced conditioned place preference in the rat. *Pharmacol Biochem Behav* **25**:1041–1049.
- Hahn B, Sharples CG, Wonnacott S, Shoaib M, and Stolerman IP (2003) Attentional effects of nicotinic agonists in rats. *Neuropharmacology* **44**:1054–1067.
- Harrod SB, Dwoskin LP, Green TA, Gehrke BJ, and Bardo MT (2003) Lobeline does not serve as a reinforcer in rats. *Psychopharmacology (Berl)* **165**:397–404.
- Hoffman AC and Evans SE (2013) Abuse potential of non-nicotinic tobacco smoke components: acetaldehyde, nornicotine, cotinine, and anabaseine. *Nicotine Tob Res* **15**:622–632.
- Jensen AA, Mikkelsen I, Frølund B, Bräuner-Osborne H, Falch E, and Krosgaard-Larsen P (2003) Carbamoylcholine homologs: novel and potent agonists at neuronal nicotinic acetylcholine receptors. *Mol Pharmacol* **64**:865–875.

- Jutkiewicz EM, Brooks EA, Kynaston AD, Rice KC, and Woods JH (2011) Patterns of nicotinic receptor antagonism: nicotine discrimination studies. *J Pharmacol Exp Ther* **339**:194–202.
- Kangas BD, Delatte MS, Vemuri VK, Thakur GA, Nikas SP, Subramanian KV, Shukla VG, Makriyannis A, and Bergman J (2013) Cannabinoid discrimination and antagonism by CB(1) neutral and inverse agonist antagonists. *J Pharmacol Exp Ther* **344**:561–567.
- Kelleher RT and Morse WH (1968) Determinants of the specificity of behavioral effects of drugs. *Ergeb Physiol* **60**:1–56.
- Kem WR, Mahnir VM, Papke RL, and Lingle CJ (1997) Anabaseine is a potent agonist on muscle and neuronal alpha-bungarotoxin-sensitive nicotinic receptors. *J Pharmacol Exp Ther* **283**:979–992.
- LeSage MG, Shelley D, Ross JT, Carroll FI, and Corrigan WA (2009) Effects of the nicotinic receptor partial agonists varenicline and cytisine on the discriminative stimulus effects of nicotine in rats. *Pharmacol Biochem Behav* **91**:461–467.
- Marks MJ, Stitzel JA, Romm E, Wehner JM, and Collins AC (1986) Nicotinic binding sites in rat and mouse brain: comparison of acetylcholine, nicotine, and alpha-bungarotoxin. *Mol Pharmacol* **30**:427–436.
- Marks MJ, Robinson SF, and Collins AC (1996) Nicotinic agonists differ in activation and desensitization of 86Rb⁺ efflux from mouse thalamic synaptosomes. *J Pharmacol Exp Ther* **277**:1383–1396.
- Miller DK, Crooks PA, Teng L, Witkin JM, Munzar P, Goldberg SR, Acri JB, and Dwoskin LP (2001) Lobeline inhibits the neurochemical and behavioral effects of amphetamine. *J Pharmacol Exp Ther* **296**:1023–1034.
- Miller DK, Harrod SB, Green TA, Wong MY, Bardo MT, and Dwoskin LP (2003) Lobeline attenuates locomotor stimulation induced by repeated nicotine administration in rats. *Pharmacol Biochem Behav* **74**:279–286.
- Mirza NR, Pei Q, Stolerman IP, and Zetterström TS (1996) The nicotinic receptor agonists (-)-nicotine and isoarecolone differ in their effects on dopamine release in the nucleus accumbens. *Eur J Pharmacol* **295**:207–210.
- Reavill C, Spivak CE, Stolerman IP, and Waters JA (1987) Isoarecolone can inhibit nicotine binding and produce nicotine-like discriminative stimulus effects in rats. *Neuropharmacology* **26** (7A):789–792.
- Reavill C, Jenner P, Kumar R, and Stolerman IP (1988) High affinity binding of [3H] (-)-nicotine to rat brain membranes and its inhibition by analogues of nicotine. *Neuropharmacology* **27**:235–241.
- Reavill C, Walther B, Stolerman IP, and Testa B (1990) Behavioural and pharmacokinetic studies on nicotine, cytisine and lobeline. *Neuropharmacology* **29**:619–624.
- Rollema H, Chambers LK, Coe JW, Glowa J, Hurst RS, Lebel LA, Lu Y, Mansbach RS, Mather RJ, and Rovetti CC, et al. (2007) Pharmacological profile of the alpha4beta2 nicotinic acetylcholine receptor partial agonist varenicline, an effective smoking cessation aid. *Neuropharmacology* **52**:985–994.
- Rollema H, Shrikhande A, Ward KM, Tingley FD, 3rd, Coe JW, O'Neill BT, Tseng E, Wang EQ, Mather RJ, and Hurst RS, et al. (2010) Pre-clinical properties of the alpha4beta2 nicotinic acetylcholine receptor partial agonists varenicline, cytisine and dianicline translate to clinical efficacy for nicotine dependence. *Br J Pharmacol* **160**:334–345.
- Shoaib M (2006) Effects of isoarecolone, a nicotinic receptor agonist in rodent models of nicotine dependence. *Psychopharmacology (Berl)* **188**:252–257.
- Smith JW and Stolerman IP (2009) Recognising nicotine: the neurobiological basis of nicotine discrimination. *Handb Exp Pharmacol* **295**:333.
- Smith JW, Mogg A, Tafi E, Peacey E, Pullar IA, Szekeres P, and Tricklebank M (2007) Ligands selective for alpha4beta2 but not alpha3beta4 or alpha7 nicotinic receptors generalise to the nicotine discriminative stimulus in the rat. *Psychopharmacology (Berl)* **190**:157–170.
- Snedecor GW and Cochran WG (1967) *Statistical Methods*, Ed. 6th. Iowa State University Press, Ames, Iowa.
- Spealman RD (1995) Noradrenergic involvement in the discriminative stimulus effects of cocaine in squirrel monkeys. *J Pharmacol Exp Ther* **275**:53–62.
- Stevens KE, Kem WR, Mahnir VM, and Freedman R (1998) Selective alpha7-nicotinic agonists normalize inhibition of auditory response in DBA mice. *Psychopharmacology (Berl)* **136**:320–327.
- Stolerman IP, Garcha HS, Pratt JA, and Kumar R (1984) Role of training dose in discrimination of nicotine and related compounds by rats. *Psychopharmacology (Berl)* **84**:413–419.
- Stolerman IP, Garcha HS, and Mirza NR (1995) Dissociations between the locomotor stimulant and depressant effects of nicotinic agonists in rats. *Psychopharmacology (Berl)* **117**:430–437.
- Stolerman IP, Chandler CJ, Garcha HS, and Newton JM (1997) Selective antagonism of behavioural effects of nicotine by dihydro-beta-erythroidine in rats. *Psychopharmacology (Berl)* **129**:390–397.
- Sullivan JP, Decker MW, Brioni JD, Donnelly-Roberts D, Anderson DJ, Bannon AW, Kang CH, Adams P, Piattoni-Kaplan M, and Buckley MJ, et al. (1994) (+/-)-Epibatidine elicits a diversity of in vitro and in vivo effects mediated by nicotinic acetylcholine receptors. *J Pharmacol Exp Ther* **271**:624–631.
- Takada K, Swedberg MD, Goldberg SR, and Katz JL (1989) Discriminative stimulus effects of intravenous l-nicotine and nicotine analogs or metabolites in squirrel monkeys. *Psychopharmacology (Berl)* **99**:208–212.
- Tidey JW and Bergman J (1998) Drug discrimination in methamphetamine-trained monkeys: agonist and antagonist effects of dopaminergic drugs. *J Pharmacol Exp Ther* **285**:1163–1174.
- Whiteaker P, Garcha HS, Wonnacott S, and Stolerman IP (1995) Locomotor activation and dopamine release produced by nicotine and isoarecolone in rats. *Br J Pharmacol* **116**:2097–2105.
- Williams M and Robinson JL (1984) Binding of the nicotinic cholinergic antagonist, dihydro-beta-erythroidine, to rat brain tissue. *J Neurosci* **4**:2906–2911.
- Xiao Y and Kellar KJ (2004) The comparative pharmacology and up-regulation of rat neuronal nicotinic receptor subtype binding sites stably expressed in transfected mammalian cells. *J Pharmacol Exp Ther* **310**:98–107.

Address correspondence to: Rajeev I. Desai, Preclinical Pharmacology Laboratory, McLean Hospital/Harvard Medical School, 115 Mill Street, Belmont, MA 02478. E-mail: rdesai@mclean.harvard.edu