Published in final edited form as:

J Am Chem Soc. 2014 January 15; 136(2): 582–585. doi:10.1021/ja4099559.

Redox Non–innocence of a *N*–heterocyclic Nitrenium Cation Bound to a Nickel–Cyclam Core

Florian Heims[§], Florian Felix Pfaff[§], Sarah-Luise Abram[§], Erik R. Farquhar[‡], Maurizio Bruschi[✓], Claudio Greco[✓], and Kallol Ray^{§,*}

[§]Humboldt–Universität zu Berlin, Institut für Chemie, Brook–Taylor–Straße 2, D–12489 Berlin, Germany

[‡]Case Western Reserve University Center for Synchrotron Biosciences and Center for Proteomics and Bioinformatics, National Synchrotron Light Source, Brookhaven National Laboratory, Upton, NY 11973, USA

[✓]Department of Environmental and Earth Sciences, University of Milan–Bicocca, Piazza della Scienza 1, 20126 Milan – Italy

Abstract

The redox properties of Ni complexes bound to a new ligand [DMC–nit]⁺, where a *N*–heterocyclic nitrenium group is anchored on a 1,4,8,11–tetraazacyclotetradecane backbone, have been examined using spectroscopic and DFT methods. Ligand based [Ni(DMC–nit)]^{2+/+} reduction and metal based [Ni(DMC–nit)]^{2+/3+} oxidation processes have been established for the [Ni(DMC–nit)]^{+/2+/3+} redox series, which represents the first examples of nitrenium nitrogen (N_{nit}) bound first row transition metal complexes. An unprecedented bent binding mode of N_{nit} in [Ni(DMC–nit)]²⁺ is observed, which possibly results from the absence of any N_{nit}→Ni σ–donation. For the corresponding [Ni(DMC–nit)]³⁺ complex, σ–donation is dominant, and hence a coplanar arrangement at N_{nit} is predicted by DFT. The binding of the triazolium ion to Ni enables new chemistry (formate oxidation) that is not observed in a derivative that lacks this functional group. Thus the *N*–heterocyclic nitrenium ligand is a potential useful and versatile reagent in transition–metal based catalysis.

Transition metal complexes are at the heart of organometallic chemistry, and homogeneous catalysis.¹ Very often the bound ligands are anionic or neutral species. However, cationic ligands ligated to a transition metal are also known in a few cases.^{2,3} Notably, for the positively charged ligands (L⁺) π–acidity rather than σ–basicity of L⁺ must be considered as the decisive factor for the stability of their transition metal complexes, and the efficacy of stabilizing influences such as low formal oxidation state and high atomic number of the metal (M) can be traced to the stabilization of the M→L⁺ π–bond. The weaker π–acceptance ability of a cationic *N*–atom, relative to *P*– and *As*– atoms, may therefore explain why the nitrenium ion has the weakest binding ability amongst all the *N*–heterocyclic carbene–type ligands of the Group 15 *p*–block elements.³ Notably, *N*–heterocyclic nitrenium (NHN) ions were, until very recently, unknown to behave as ligands. By incorporating the triazolium

*Corresponding Author: kallol.ray@chemie.hu-berlin.de.

Notes

Authors declare no competing financial interest.

ASSOCIATED CONTENT

Supporting Information

Additional syntheses and characterization data, kinetic data, X–ray diffraction parameters, crystal data (Tables S2, S9–S10), and computational data (Tables S11–16). This material is available free of charge via the Internet at <http://pubs.acs.org>.

ring within a pincer-type system Gandelman *et al.*^{3a} demonstrated coordination of nitrenium ions to electron-rich transition metals like Rh(I) and Ru(II) in both the solid and solution states. This study filled a missing link in the series of both fundamentally and practically important *N*-heterocyclic carbenes and their analogues, which have become ubiquitous in the fields of transition metal- and organo-catalysis.⁴

Interestingly, all the reported [(NHN)-M] (M=Ru(II), Rh(I)) complexes are found to be planar at the nitrenium nitrogen (N_{nit}) atom.^{3a} However, in the complexes of the *N*-heterocyclic phosphonium (NHP) and arsenium (NHAs) ligands, which are heavier analogues of NHN, both planar and pyramidal geometries at the central *P* or *As* atoms have been observed.³ Whereas, both σ -donation from the filled sp^2 hybrid orbital of *P* or *As* and π -acceptance into the unoccupied p - π orbital ensure planar coordination, the presence of a π -interaction alone results in a pyramidal geometry at *P* or *As*.^{3e} We therefore initiated the present study to investigate whether the planar geometric situation would persist for NHN complexes of the first row transition metals, which are weaker σ -donor and π -acceptors, as compared to their heavier congeners. Accordingly, we designed a new ligand system [DMC-nit]⁺, where a triazolium ring is anchored on a well investigated^{5,6} cyclam (1,4,8,11-tetraazacyclotetradecane) backbone, and studied its coordination chemistry with nickel.

Coupling of the nitrenium precursor 1,3-di-(2'-bromoethyl)-benzotriazolium bromide^{3a} with 1,8-dimethyl-1,4,8,11-tetraazacyclotetradecane in a basic medium led to initial formation of the benzotriazolium-capped cyclam bromide (Figs. S1–S2), which upon anion exchange with KPF₆ or NaBPh₄ yielded the desired [DMC-nit]⁺ ligand cation (Scheme S1) as PF₆[−] or BPh₄[−] salts (Figs. S3–S5). Reactions of equimolar amounts of [DMC-nit]X (X=PF₆ or BPh₄) and Ni(cod)₂ in CH₃CN afforded the purple [(DMC-nit)Ni]X₂ (**1**)-(X)₂; X=PF₆ or BPh₄) complexes in ~35–50% yield (Scheme 1); the formation of **1**-(X)₂ is accompanied by the one-electron oxidation of Ni(0) and concomitant reduction of [DMC-nit]X, which is in agreement with the observation that the yields of **1**-(X)₂ never exceeded 50% of the ligand. Electrospray ionization mass spectrometry (ESI-MS) of **1**-(PF₆)₂ shows a prominent peak at m/z = 458.259 (calc.: 458.254) (Fig. S6). X-ray analysis of single crystals of **1**-(PF₆)₂ obtained from CH₃CN/(C₂H₅)₂O confirmed coordination of [DMC-nit]⁺ as a pentadentate ligand with the cyclam macrocycle adopting a *trans-I* configuration⁶ and the central nitrogen (N_{nit}) of the benzotriazolium residue bound at the apical position of a distorted square pyramid (SQP; τ = 0.48;⁷ Scheme 1 inset). Interestingly, **1**-(PF₆)₂ is pyramidal at the N_{nit} center and has a coordination geometry (sum of all angles (Σ) = 332°)⁷ consistent with a nonbonding lone pair still present on N_{nit} . Thus, in contrast to the previously reported benzotriazolium bound Rh(I) complexes^{3a}, which involved both σ - and π -interactions leading to a coplanar arrangement, the bonding in **1**-(PF₆)₂ can be described as primarily $M \rightarrow L^+$ π -bonding; the lone pair of electrons on N_{nit} , located in the sp^2 hybrid orbital, do not participate in bonding to the metal center (see also results of DFT calculation below).

Cyclic voltammetric (CV) investigations of **1**-(X)₂ (X = PF₆ or BPh₄) performed in CH₃CN solutions at room temperature reveal one reversible oxidation wave centered at −0.51 V versus Fc^+/Fc and a reversible reduction wave centered at −1.71 V (Fig. 1 and Fig. S7) in each case. Coulometric measurements show that both oxidation and reduction correspond to one electron processes. Interestingly, the CV of [DMC-nit]⁺ exhibits a reversible reduction (−1.77 V vs Fc^+/Fc) at a potential very similar to the potential associated with the **1**]^{2+/+} process. The CV of the previously reported [Ni(TMC)]OTf (TMC = 1,4,8,11-tetramethyl-1,4,8,11-tetraazacyclotetradecane) complex⁸ was also measured for comparison, as it lacks the benzotriazolium residue. In this case a single reversible Ni^{I/II} oxidation wave centered at −1.22 V vs. Fc^+/Fc was observed. We therefore assign the reduction in **1**-(X)₂ as ligand centered and the oxidation as metal centered,

which are also corroborated by X-ray absorption spectroscopy (see below). The 700 mV positive shift of the $\text{Ni}^{\text{I/II}}$ potential in $[\mathbf{1}]\text{-(X)}_2$ relative to $[\text{Ni}(\text{TMC})]\text{OTf}$ reflects the strong π -acceptance property of the bound benzotriazolium residue in $[\mathbf{1}]\text{-(X)}_2$.

The absorption spectrum (Fig. 1) of $[\mathbf{1}]\text{-(PF}_6)_2$ displays two intense absorption maxima at 538 nm ($\epsilon = 11500 \text{ M}^{-1}\text{cm}^{-1}$) and 786 nm ($\epsilon = 10000 \text{ M}^{-1}\text{cm}^{-1}$). We tentatively assign these bands to metal to ligand charge transfer (MLCT) type (Ni(I) to benzotriazolium transition), which are absent in $[\text{Ni}(\text{TMC})]\text{OTf}$,⁸ as well as in the electrochemically generated one-electron oxidized $[\mathbf{2}]^{3+}$ and one-electron reduced $[\mathbf{3}]^+$ complexes. The EPR spectrum of a frozen solution of $[\mathbf{1}]\text{-(PF}_6)_2$ (Fig. 1) in CH_3CN at 77 K shows a slightly anisotropic signal ($g_x = 2.22$; $g_y = 2.20$ and $g_z = 2.18$),⁹ which supports the formulation of $[\mathbf{1}]\text{-(PF}_6)_2$ as a Ni(I) complex.

$[\mathbf{1}]\text{-(PF}_6)_2$ can also be chemically oxidized by using 1 eq. of ferrocenium tetrafluoroborate in CH_3CN under anaerobic conditions to yield a green powder of $[(\text{DMC-nit})\text{Ni}(\text{F})](\text{PF}_6)_2$, $[\mathbf{2-F}]\text{-(PF}_6)_2$, in about 73 % yield. $[\mathbf{2-F}]\text{-(PF}_6)_2$ is EPR silent and in its $^1\text{H-NMR}$ spectrum (Fig. S8) twelve paramagnetically shifted resonances are observed, which may indicate the presence of an octahedral (O_h) Ni(II) (d^8 , $S=1$) center (see also XAS studies below). Interestingly, the number and relative integration of these resonances suggest that the *trans-I* configuration of the cyclam macrocycle is preserved in $[\mathbf{2-F}]\text{-(PF}_6)_2$. ESI-MS (Fig. S9) spectrum of a solution of isolated $[\mathbf{2-F}]\text{-(PF}_6)_2$ in CH_3CN shows a prominent peak at $m/z = 238.626$ (calc.: 238.627), with an isotope distribution pattern consistent with its formulation as $[(\text{DMC-nit})\text{Ni}(\text{F})]^{2+}$. Subsequent studies have shown that the presence of fluoride, which is possibly abstracted from the tetrafluoroborate anion, is necessary to stabilize the high positive charge of the $[(\text{DMC-nit})\text{Ni}^{\text{II}}]^{3+}$ core in $[\mathbf{2-F}]\text{-(PF}_6)_2$. Accordingly, $[\mathbf{2-CH}_3\text{CN}]\text{-(BPh}_4)_3$, (O_h Ni^{II} ; $S=1$ from $^1\text{H-NMR}$; Fig S10) generated in the reaction of $[\mathbf{1}]\text{-(BPh}_4)_2$ with ferrocenium tetraphenylborate, is extremely unstable. In the absence of any external fluoride donors and excess oxidant it is spontaneously converted to the starting Ni(I) complex, $[\mathbf{1}]\text{-(BPh}_4)_2$ (Fig S11); the electron necessary for the reduction of $[\mathbf{2-CH}_3\text{CN}]\text{-(BPh}_4)_3$ to $[\mathbf{1}]\text{-(BPh}_4)_2$ is presumably provided by the BPh_4^- anion.¹⁰ EPR silent yellowish-brown solutions of $[(\text{DMC-nit})\text{Ni}]\text{PF}_6$ ($[\mathbf{3}]\text{-(PF}_6)$) could be chemically generated by the reduction of $[\mathbf{1}]\text{-(PF}_6)_2$ with excess KC_8 . $^1\text{H-NMR}$ spectrum of $[\mathbf{3}]\text{-(PF}_6)$ shows all the resonances within $\delta = 0$ to + 10 ppm, confirming an $S=0$ ground state; ESI-MS study shows the molecular ion peak at $m/z = 458.254$ (Fig. S12).

X-ray absorption near-edge spectroscopic studies (XANES; Fig. 2A and Fig. S13) were performed at the Ni K-edge in order to directly probe the metal oxidation states in $[\mathbf{1}]\text{-(PF}_6)_2$, $[\mathbf{2-F}]\text{-(PF}_6)_2$ and $[\mathbf{3}]\text{-(PF}_6)$, and provide support for the electrochemically-derived assignments. $[\mathbf{1}]\text{-(PF}_6)_2$ exhibits an edge inflection energy of ca. 8341.2 eV, with a shoulder along the rising edge at 8336.3 eV corresponding to a $1s \rightarrow 4p$ shakedown transition,¹¹ in accord with the known SQP structure of this complex. A broad $1s \rightarrow 3d$ pre-edge peak is present at 8331.4 eV, with a peak area of approx. 8.1 units. The XANES spectrum of $[\mathbf{2-F}]\text{-(PF}_6)_2$ shows a +2.8 eV blue shift of the edge inflection energy to 8344.0 eV, relative to $[\mathbf{1}]\text{-(PF}_6)_2$, thereby supporting metal-centered oxidation to the Ni(II) state. The $1s \rightarrow 3d$ pre-edge transition has also blue shifted by ~0.5 eV to 8332.0 eV and weakened in intensity. Most notably, $[\mathbf{2-F}]\text{-(PF}_6)_2$ lacks the diagnostic $1s \rightarrow 4p$ shakedown transition shoulder on the rising edge, providing strong evidence for a $6C$ O_h nickel site in this complex.¹¹ Finally, the XANES spectrum of $[\mathbf{3}]\text{-(PF}_6)$ shows an unexpected 1.8 eV *blue-shift* in edge inflection energy to 8343.0 eV relative to $[\mathbf{1}]\text{-(PF}_6)_2$. This suggests that the reducing agent has imparted a significant increase in the Ni effective nuclear charge (Z_{eff}). Most importantly, the existence of a $1s \rightarrow 3d$ pre-edge transition at an identical position (8331.4 eV) relative to $[\mathbf{1}]\text{-(PF}_6)_2$ indicates that $[\mathbf{3}]\text{-(PF}_6)$ *does not* contain a d^{10} Ni^0 center. The Ni site in $[\mathbf{3}]\text{-(PF}_6)$

is likely to still be 5C SQP, given that the $1s \rightarrow 4p$ shakedown transition is present at 8335.9 eV and similar in intensity relative to $[1]-(PF_6)_2$.

Extended X-ray absorption fine structure (EXAFS) analysis reveals further structural details (Figs. 2B, 2C and 2D). For $[1]-(PF_6)_2$ the first coordination sphere could be satisfactorily fitted by 5 nitrogen scatterers at a distance of 2.11 Å (Table S1, fit 8). While the outer-shell features could be accounted for by several single scattering paths involving 14N/C at 2.94 Å, 4C at 3.45 Å, 2C at 3.97 Å, and 2C at 5.34 Å, the fit could be significantly improved by introducing multiple-scattering paths involving elements of the cyclam and triazolium moieties. In general, the metrical parameters obtained for $[1]-(PF_6)_2$ are in agreement with the molecular structure determined by X-ray crystallography, thereby confirming that the solid state structure is retained in frozen CH_3CN solutions of $[1]-(PF_6)_2$. For $[2-F]-(PF_6)_2$ the best fit (Table S1, fit 8) of the Ni EXAFS consists of two subshells of one fluorine scatterer at 1.93 Å and 5 nitrogen scatterers at 2.12 Å, thereby supporting a six coordinate Ni center in $[2-F]-(PF_6)_2$ in agreement with the XANES and DFT (see below) studies. Fits to a single shell of 6 nitrogen scatterers at 2.11 Å produced a significant decrease in fit quality (Table S1, fits 3 and 9). Analysis of the EXAFS data of $[3]-(PF_6)_2$ was less straightforward. $[3]-(PF_6)_2$ has much weaker EXAFS and FT intensity compared to the other two complexes. The best fit (Table S1, fit 10) requires two subshells consisting of one nitrogen scatterer at 1.85 Å and three nitrogen scatterers at 2.06 Å. Notably, the latter shell has a very large Debye-Waller factor, consistent with a disordered distribution of bond lengths rather than a narrowly constrained range. We note that EXAFS supports a 4-coordinate Ni center in $[3]-(PF_6)_2$, and the XANES points to a 5-coordinate Ni site. DFT calculations for $[3]-(PF_6)_2$ (see below) show that both 4- and 5-coordinate states are feasible. Thus, the weak EXAFS intensity of $[3]-(PF_6)_2$ and large bond disorder may be attributable to destructive interference caused by the simultaneous existence of 4- and 5-coordinate Ni sites in the EXAFS sample of $[3]-(PF_6)_2$.

DFT calculations were carried out for the $[1]^{2+}$, $[2-F]^{2+}$ and $[3]^+$ cations in order to obtain deeper insights into their electronic structures. For spin-doublet (Ni^I ; d^9 ; $S_T=1/2$) $[1]^{2+}$ the computed Ni- N_{nit} distance at 2.082 Å and average Ni-N(cyclam) distance at 2.18 Å are in good agreement with the experimentally determined values of 2.021(3) Å and 2.149(3) Å, respectively (Table S2; Fig S14).⁷ Moreover, the distorted SQP ($\tau=0.48$) geometry at Ni and the pyramidal geometry ($\angle_{sum}=332^\circ$) at N_{nit} are well reproduced in the calculation (calculated $\tau=0.48$ and $\angle_{sum}=338$).⁷ Furthermore, the computational prediction of the EPR spectrum of $[1]^{2+}$ is in good accord with the experiment (Table S3, Fig. S15). Interestingly, the bent binding mode of N_{nit} is predicted to persist in the DFT optimized structure of the hypothetical $[Ni^I(cyclam)(benzotriazolium)]^{2+}$ complex, which lacks any ethylene linkers between the cyclam and the benzotriazolium residue and methyl substituents on the cyclam nitrogens (Fig S16). This may indicate that the conformation of N_{nit} in $[1]^{2+}$ is not determined by steric constraints imposed by the $[DMC-nit]^+$ ligand. Further insights were obtained from the nature of the DFT calculated singly occupied molecular orbital (SOMO) of $[1]^{2+}$, which features an antibonding combination of the Ni $d_{x^2-y^2}$ orbital with the N_4 lone pairs of the cyclam ligand (Fig. S17). Thus, the d^9 'hole' of the Ni(I) center is located in the N_4 plane of cyclam, which is orthogonal to the plane of the N_{nit} atom. Hence, $N_{nit} \rightarrow Ni$ σ -donation is negligible, as also indicated in the natural bond orbital (NBO) analysis¹² (Table S4) by the near-identical occupancy of the N_{nit} sp^2 lone pair in $[1]^{2+}$ (1.92 e^-) and free $[DMC-nit]^+$ (1.94 e^-). In the absence of σ -donation, repulsive interactions between electrons of the Ni-centered dz^2 orbital and of the N_{nit} lone pair possibly explains the pyramidal geometry at N_{nit} in $[1]^{2+}$. NBO analysis also reveals that the Ni- N_{nit} bond in $[1]^{2+}$ features a dominating π -interaction, as evidenced by the transfer of 0.29 e^- from the metal to the π -system of the triazolium ring.

A reasonable agreement between the calculated structures of $[2-F]^{2+}$ and $[3]^+$ cations and the EXAFS experiments further validates our theoretical methods (Fig. S14, Table S1). For $[2-F]^{2+}$ the ground state is calculated to be a triplet (Ni^{II} ; d^8 ; $S_T=1$) consistent with experiment. Since, $[2-F]-(PF_6)_2$ was generated in CH_3CN , which is a potential ligand, we also calculated the structure of the $[2-CH_3CN]^{3+}$ species with a bound CH_3CN instead of a fluoride. While DFT calculations also support a $S_T=1$ ground state for $[2-CH_3CN]^{3+}$, the calculated geometry shows a set of four shorter Ni–N distances of 2.17 Å and two long Ni–N distance of 2.34 Å, which is not in agreement with the EXAFS studies (Table S1). Notably, the oxidation of $[1]^{2+}$ to $[2-F]^{2+}$ or $[2-CH_3CN]^{3+}$ involves the loss of an electron from the d_{z^2} orbital, and the additional SOMO in $[2-F]^{2+}$ or $[2-CH_3CN]^{3+}$ is given by an antibonding combination of the d_{z^2} orbital and N_{nit} lone pair (Fig S18). This explains the increased $N_{nit} \rightarrow Ni$ σ -donation in $[2-F]^{2+}$ and $[2-CH_3CN]^{3+}$ with respect to $[1]^{2+}$, as evident from the NBO analysis (Tables S5–S6), which is also reflected in the predicted near coplanar geometry at N_{nit} in $[2-F]^{2+}$ and $[2-CH_3CN]^{3+}$ (Fig. S14). It is also important to note that in contrast to $[1]^{2+}$, no significant π -backdonation from Ni occurs in $[2-F]^{2+}$ and $[2-CH_3CN]^{3+}$.

For the $[3]^+$ cation, the ground state can be best described by a broken-symmetry (BS) solution, where the Ni is in a 5-coordinate geometry (Fig. S14) and a Ni-based d_{xz} orbital is coupled antiferromagnetically to a N_{nit} -based ligand radical (Fig. S19), thereby resulting in a $S_T=0$ state. The closed-shell (CS) calculation for $[3]^+$, however, reveals a 4-coordinate geometry (Fig. S14), with one of the N-atoms of the $[DMC-nit]^+$ macrocycle remaining out of the coordination sphere. Interestingly, the CS state is only 3 kcal/mol higher in energy than the BS state. This may imply that $[3]^+$ may coexist in both 4- and 5-coordinate states, which may account for the structural disorder observed in the EXAFS experiments. Notably, $[3]^+$ involves a reduced ligand in both the CS and BS states in full agreement with the experiment; the population of the π -system of the triazolium ring is calculated to be 0.74 e^- and 0.91 e^- larger in $[3]^+$ for the CS and BS states (Tables S7–S8), respectively, than in the free ligand.

Recently, it has been shown that for a series of nickel(II) complexes the presence of a basic pendant amine on the ligand backbone plays an important role in the oxidation of formate to carbon dioxide and a proton.¹³ With the availability of the compounds $[1]-(PF_6)_2$, $[2-F]-(PF_6)_2$, and $[3]-(PF_6)_2$, we decided to investigate the role of the Lewis-acidic nitrenium cation in the nickel mediated formate oxidation reactions. Although, the nickel(I) centers in $[1]-(PF_6)_2$ and $[3]-(PF_6)_2$ are unable to bind and oxidize formate, the nickel(II) complex $[2-F]-(PF_6)_2$ is reduced to **1** (45% yield) in the presence of $NBu_4HCO_2^+HCO_2H$ or $(HNEt_3)HCO_2$ ¹³ in CH_3CN at 25 °C (Fig. S20); CO_2 is produced in 40% yield. Interestingly, the reaction of the corresponding $[Ni(TMC)]^{2+}$ complex that lacks the benzotriazolium backbone does not lead to formate oxidation; instead a stable $[Ni(TMC)(HCO_2)]^+$ adduct is formed, the molecular structure of which as determined by X-ray crystallography (Fig. S21) shows a 5-coordinate SQP nickel center with the formate bound at the apical position through oxygen.

In summary, we have synthesized and characterized Ni(I) and Ni(II) complexes of a new N-heterocyclic nitrenium ligand $[DMC-nit]^+$, and established an unprecedented bent binding mode of the nitrenium nitrogen to the Ni(I) center that is apparently caused by the presence of a nonbonding electron pair in the N_{nit} sp^2 hybrid orbital. NBO analysis reveals that for the Ni(I) complex π -donation from Ni is the major interaction, which is in contrast for the Ni(II) complex, where σ -donation from $[DMC-nit]^+$ is dominant. $[DMCnit]^+$ is a potential non-innocent ligand, and can be reversibly reduced by one-electron to a neutral ligand π -radical species, both in the free state and when coordinated to a Ni(I) center. Thus, the nitrenium ligands can be considered as NO^+ surrogates,^{2e} not only because they are cationic

and non-innocent, but also because they can adopt planar or pyramidal coordination modes, which are analogous to the linear and bent coordination modes of metal-nitrosyls.^{1a} Furthermore, we have shown that the electrophilic property of a nickel(II) center containing a bound nitrenium ligand can be utilized successfully in the oxidation of formate, thereby, supporting the *N*-heterocyclic nitrenium ligand as a potential useful and versatile reagent in transition metal based catalysis. Further studies of the reactivities of [1]-(PF₆)₂, [2-F]-(PF₆)₂ and [3]-(PF₆)₂, are currently underway in our laboratory, along with possible extension of the coordination chemistry of [DMC-nit]⁺ to other first-row transition metals and their application in catalysis.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We dedicate this manuscript to Dr. Eckhard Bill on the occasion of his 60th birthday. We gratefully acknowledge financial support of this work from the Cluster of Excellence "Unifying Concepts in Catalysis" (EXC 314/1), Berlin. XAS data were obtained on beamline X3B of the National Synchrotron Light Source (Brookhaven National Laboratory, Upton, NY, USA), with support from NIH Grant P30-EB-009998. NSLS is supported by the US DOE Office of Science (OBES), under Contract DE-AC02-98CH10886. FH thanks BIG-NSE for a scholarship. We also thank Corinna Matlachowski for her help with GC experiments.

References

- (a) Crabtree, RH. *The Organometallic Chemistry of the Transition Metals*. 5. Wiley-VCH; Weinheim, Germany: 2009. (b) Duca, G. *Homogenous Catalysis with Metal Complexes*. 7. Springer; 2012.
- (a) Dumrath A, Wu XF, Neumann H, Spannenberg A, Jackstell R, Beller M. *Angew Chem Int Ed*. 2010; 49:8988–8992. (b) Coles MP, Hitchcock PB. *Chem Commun*. 2007:5229–5231. (c) Petušková J, Patil M, Holle S, Lehmann CW, Thiel W, Alcarazo M. *J Am Chem Soc*. 2011; 133:20758–20760. [PubMed: 22136140] (d) Day GS, Pan B, Kellenberger DL, Foxman BM, Thomas CM. *Chem Commun*. 2011; 47:3634–3636. (e) Praneeth VKK, Paulat F, Berto TC, George SD, Näther C, Sulok CD, Lehnert N. *J Am Chem Soc*. 2008; 130:15288–15303. [PubMed: 18942830]
- (a) Tulchinsky Y, Iron MA, Botoshansky M, Gandelman M. *Nat Chem*. 2011; 3:525–531. [PubMed: 21697872] (b) Gudat D. *Coord Chem Rev*. 1997; 163:71–106. (h) Rosenberg L. *Coord Chem Rev*. 2012; 256:606–626. (c) Caputo CA, Jennings MC, Tuononen HM, Jones ND. *Organometallics*. 2009; 28:990–1000. (d) Burck S, Daniels J, Gans-Eichler T, Gudat D, Nättinen K, Nieger M. *Z Anorg Allg Chem*. 2005; 631:1403–1412. (e) Price JT, Lui M, Jones ND, Ragogna PJ. *Inorg Chem*. 2011; 50:12810–12817.
- (a) Crudden CM, Allen DP. *Coord Chem Rev*. 2004; 248:2247–2273. (b) Bourissou D, Guerret O, Gabbai FP, Bertrand G. *Chem Rev*. 2000; 100:39–91. [PubMed: 11749234] (c) Marion N, Nolan SP. *Acc Chem Res*. 2008; 41:1440–1449. [PubMed: 18774825]
- Barefield EK. *Coord Chem Rev*. 2010; 254:1607–1627.
- Bosnich B, Poon CK, Tobe ML. *Inorg Chem*. 1965; 4:1102–1108.
- Addison AWT, Rao N, Reedijk J, Rijn J, Verschoor GC. *J Chem Soc Dalton Trans*. 1984:1349–1356. For the second independent molecule in crystals of [1]-(PF₆)₂ $\theta_{\text{sum}} = 337^\circ$, $\tau = 0.47$, Ni–N_{nit} = 1.986(3) Å, average Ni–N(cyclam) = 2.133(3) Å.
- (a) Kieber-Emmons MT, Schenker R, Yap GPA, Brunold TC, Riordan CG. *Angew Chem, Int Ed*. 2004; 43:6716–6718. (b) Jubran N, Ginzburg G, Cohen H, Koresh Y, Meyerstein D. *Inorg Chem*. 1985; 24:251–258. (c) Ram MS, Riordan CG, Ostrander R, Rheingold AL. *Inorg Chem*. 1995; 34:5884–5892.
- For [1]-(BPh₄)₂ (Fig. S15) $g_x = 2.28$; $g_y = 2.22$ and $g_z = 2.15$.
- Pal PK, Chowdhury S, Drew MGB, Datta D. *New J Chem*. 2002; 26:367–371.

11. Colpas GJ, Maroney MJ, Bagyinka C, Kumar M, Willis WS, Suib SL, Baidya N, Mascharak PK. *Inorg Chem.* 1991; 30:920–928.
12. Reed AE, Curtiss LA, Weinhold F. *Chem Rev.* 1988; 88:899–926.
13. Galan BR, Schöffel J, Linehan JC, Seu C, Appel AM, Roberts JAS, Helm ML, Kilgore UJ, Yang JY, DuBois DL, Kubiak CP. *J Am Chem Soc.* 2011; 133:12767–12779. [PubMed: 21692477]

Figure 1.

Left: Cyclic voltammetry of [DMC-nit]⁺, [1]-(PF₆)₂, and [Ni(TMC)]⁺ complexes recorded in CH₃CN solutions containing 0.10 M NBu₄PF₆. L[•]/₊ represents the [(DMC-nit)]⁺⁰ couple. **Right:** Absorption spectra of [1]²⁺, [2-F]²⁺, and [3]⁺ in CH₃CN at 25 °C. The inset depicts the experimental and simulated X-band EPR spectrum of [1]-(PF₆)₂ in a frozen CH₃CN solution at 77 K. For simulation parameters see text and Fig S15.

Figure 2.

Normalized Ni *K*-edge XANES spectra for $[1]-(PF_6)_2$, $[2-F]-(PF_6)_2$ and $[3]-PF_6$ (A) and the Fourier-transformed Ni *K*-edge EXAFS spectra of $[1]-(PF_6)_2$ (B), $[2-F]-(PF_6)_2$ (C) and $[3]-(PF_6)$ (D). The expansions of the $1s \rightarrow 3d$ and $1s \rightarrow 4p$ transitions are shown in Fig. A inset. The insets of Figs. B, C and D depict the $k^3\chi(k)$ EXAFS data and best fits.

Scheme 1.

Formation of [1]^{2+} , [2-F]^{2+} , and [3]^+ and the reactivity of [2-F]^{2+} with formate. Central nitrogen of the benzotriazolium residue is referred as N_{nit} throughout the manuscript. In the inset is shown the structure of one of the two crystallographically independent molecules in crystals of $\text{[1]}(\text{PF}_6)_2$. (S_T = total spin).