Published in final edited form as:

Rev Neurosci. 2012 ; 23(3): 253–262. doi:10.1515/revneuro-2012-0036.

The role of prefrontal cortex in psychopathy

Michael Koenigs

Department of Psychiatry, University of Wisconsin-Madison, 6001 Research Park Blvd, Madison, Wisconsin, 53719, USA

Michael Koenigs: mrkoenigs@wisc.edu

Abstract

Psychopathy is a personality disorder characterized by remorseless and impulsive antisocial behavior. Given the significant societal costs of the recidivistic criminal activity associated with the disorder, there is a pressing need for more effective treatment strategies, and hence, a better understanding of the psychobiological mechanisms underlying the disorder. The prefrontal cortex (PFC) is likely to play an important role in psychopathy. In particular, the ventromedial and anterior cingulate sectors of PFC are theorized to mediate a number of social and affective decision-making functions that appear to be disrupted in psychopathy. This article provides a critical summary of human neuroimaging data implicating prefrontal dysfunction in psychopathy. A growing body of evidence associates psychopathy with structural and functional abnormalities in ventromedial PFC and anterior cingulate cortex. Although this burgeoning field still faces a number of methodological challenges and outstanding questions that will need to be resolved by future studies, the research to date has established a link between psychopathy and PFC.

Keywords

Neuroimaging; Emotion; Decision-Making; Antisocial Personality Disorder; Crime; Anterior Cingulate Cortex; Ventromedial Prefrontal Cortex

INTRODUCTION

Psychopathy—a personality disorder characterized by callous and impulsive antisocial behavior—has devastating effects on society, particularly through crime. It is estimated that the prevalence of psychopathy is about 1% in the general population, but 15–25% in the U.S. adult prison population. Compared to prisoners without psychopathy, psychopathic prisoners have significantly higher rates of violent crime and recidivism. Given that (1) the societal cost of all criminal behavior in the U.S. is estimated at over a trillion dollars per year, (2) psychopaths constitute up to 25% of the prison population, and (3) psychopaths are known to commit a disproportionate amount of crime, this disorder costs society roughly several hundred billion dollars per year. Despite the dire consequences of the disorder, there are currently no effective treatment strategies. One likely reason for the limited treatment options for psychopathy is that the psychobiological mechanisms of the disorder remain poorly understood. In this regard, neuroscience holds much promise. Identification of reliable neural correlates of psychopathy could serve to refine diagnostic criteria for the disorder, help predict the likelihood of future offense, locate potential biological targets for pharmacological treatment, and identify neuropsychological dysfunction that may be addressed through novel cognitive-behavioral therapies. The aim of this review article is to evaluate the evidence linking psychopathy to a particular area of the brain with diverse roles in cognitive and affective function—the prefrontal cortex (PFC).

PSYCHOPATHY AND HOW IT IS MEASURED

In order to assess the putative role of PFC in psychopathy, it is first necessary to detail the principal characteristics of the disorder and how they are typically measured for research purposes. The concept of psychopathy, in its modern formulation, is based largely on the personality disorder described at length by Hervey Cleckley in “The Mask of Sanity” (Cleckley 1941, Cleckley 1976). To develop his conception of psychopathy, Cleckley cataloged a series of cases he had observed in his psychiatric practice—individuals with no apparent cognitive defect or traditional mental illness, yet a severe and pervasive disregard for the rights and welfare of others. In summary of these individuals, Cleckley states, “My concept of the psychopath’s functioning postulates a selective defect or elimination which prevents important components of normal experience from being integrated into the whole human reaction, particularly an elimination or attenuation of those strong affective components that ordinarily arise in major personal and social issues” (Cleckley 1976, p. 374). Distilling the common personality and behavioral traits that he observed in these cases, Cleckley outlined a number of hallmark characteristics (Table 1, left column).

Building on Cleckley’s conception, Robert Hare subsequently developed a formal assessment procedure for psychopathy in forensic settings, the Psychopathy Checklist-Revised (PCL-R; Hare 2003), which has become the most widely used and most extensively validated measure of psychopathy. The PCL-R assessment involves a 60–90 minute semi-structured interview and file review to obtain information used to rate 20 psychopathy-related items as 0, 1, or 2, depending on the degree to which the individual exhibits each trait (Table 1, middle column). As will later be shown in the review of neuroimaging studies of psychopathy, total scores on the PCL-R can be treated categorically in order to classify inmates as non-psychopathic (e.g., PCL-R 20) or psychopathic (e.g., PCL-R 30) for between-group analyses, or continuously, in order to provide a quantity for correlation or regression analyses. Furthermore, the PCL-R has been shown to consist of interrelated clusters of items, or factors. The conventional two-factor model consists of an interpersonal/affective cluster (Factor 1) and an antisocial/lifestyle cluster (Factor 2) (Hare 2003, Harpur et al 1989). Each factor score can be entered separately in regression analyses.

While forensic studies of psychopathy have almost uniformly employed the PCL-R, studies of noncriminal samples have commonly used self-report scales to measure psychopathy-related traits. The most widely used such instrument, the Psychopathic Personality Inventory (PPI) (Lilienfeld & Andrews 1996), includes 187 first-person statements (e.g., “I am easily flustered in pressured situations”), which are rated on a 4-point scale (1=false, 2=mostly false, 3=mostly true, 4=true). Total scores are obtained by summing across items after reversing specified items. The test consists of eight factors or subscales (Table 1, right column). With the cardinal features of psychopathy (and their measurement) in mind, we will now turn to a discussion of how this disorder may relate to dysfunction in the PFC.

FUNCTIONAL ANATOMY OF PREFRONTAL CORTEX RELEVANT TO PSYCHOPATHY

Ventromedial PFC and pseudopsychopathy

Perhaps the earliest evidence for a critical role of PFC in psychopathic personality is the famous case of Phineas Gage, the 19th century railroad construction foreman who had an iron rod shot through his face and out the top of his head in a freak excavation accident (Harlow 1868). Remarkably, Gage survived the blast, but his personality was permanently altered. Whereas before the accident Gage had been dependable and trustworthy, afterward he became disrespectful, profane, impulsive, and unreliable—traits reminiscent of psychopathy. Based on posthumous measures of his skull, Gage’s brain injury was projected

to primarily involve the medial PFC (Damasio et al 1994). Throughout the 20th century, a number of similar cases of personality change following brain injury were reported. In 1975, Blumer and Benson coined the term “pseudopsychopathy” to encapsulate the personality changes (“the lack of adult tact and restraints”) observed in a subset of neurological patients with brain damage involving ventral PFC (Blumer & Benson 1975). The connection between PFC injury and pseudopsychopathic changes was subsequently elaborated by Antonio Damasio and colleagues. Reports from Damasio’s group associated damage to a specific sector of the PFC—the ventromedial prefrontal cortex (vmPFC) (Figure 1)—with an array of emotion and decision-making deficits, including conspicuously diminished guilt, shame, and empathy; irritability; poor planning; irresponsibility; and failure to learn from punishment (Barrash et al 2000, Bechara et al 1997, Damasio 1994, Eslinger & Damasio 1985). Experimentally, both psychopaths and vmPFC lesion patients have demonstrated reduced autonomic arousal to emotionally charged stimuli (Damasio et al 1990, Hare et al 1978, Patrick et al 1993, Patrick et al 1994) and deficits in reversal learning (Budhani et al 2006, Hornak et al 2004). More recent studies have linked vmPFC damage to changes in moral judgment (Ciaramelli et al 2007, Koenigs et al 2007, Young et al 2010), economic decision-making (Koenigs & Tranel 2007, Krajcich et al 2009, Moretti et al 2009), and reductions in negative affect (Koenigs et al 2008a, Koenigs et al 2008b). In some cases, comparable moral and economic decision-making profiles have been observed in primary (low-anxious) psychopaths (Koenigs et al 2010, Koenigs et al 2011). Overall, this collection of findings hints strongly that vmPFC dysfunction may underlie certain aspects of psychopathy (Blair 2007, Blair 2008). Accordingly, the review of neuroimaging results for psychopathy below will specifically address the extent to which this neuroanatomical hypothesis has been supported.

Before evaluating neuroimaging studies of vmPFC dysfunction in psychopathy, it is perhaps first worthwhile to consider the normal function of vmPFC, as it is pertinent to emotion, social/moral behavior, and decision-making. Although there is no clear consensus on the essential function of vmPFC, there are several proposals with significant empirical support. The seminal theory of vmPFC function is Damasio’s “somatic marker hypothesis” (Damasio 1994, Damasio 1996), which was initially developed to explain the personality and behavioral changes and decision-making deficits associated with vmPFC damage. This theory, supported by neuropsychological and electrodermal data (Bechara et al 1997), posits that vmPFC links potential decision outcomes with an associated emotion, or somatic state, based on previous experiences of reward or punishment, which effectively marks the outcome as something to pursue or avoid. A second, related proposal supported by more recent functional neuroimaging data is that vmPFC plays a key role in representing the value of goal-directed outcomes and options (Grabenhorst & Rolls 2011, O’Doherty 2011). A third proposal is that vmPFC serves to regulate negative affect by exerting top-down inhibition of amygdala, which is noted for its role in processing emotionally and socially relevant information, particularly fear and anxiety (Milad et al 2006, Phelps et al 2004, Rauch et al 2006). This model is supported by human and animal data from fear conditioning paradigms, as well as human imaging data of anxiety disorders (but see also Myers-Schulz & Koenigs 2012). However, this amygdala-based model appears to conflict with the longstanding theory that psychopathy arises from an underlying deficit in processing fear—the “low fear hypothesis” (Lykken 1957, Lykken 1995). If vmPFC serves to regulate negative affect through top-down inhibition of the amygdala, then vmPFC dysfunction would be associated with disinhibition of the amygdala, and consequently higher levels of fear and anxiety. According to this model, vmPFC dysfunction would seemingly not be expected to yield psychopathic personality traits. Finally, a fourth model suggests that vmPFC may underlie aspects of self-processing, such as self-reflection and rumination, which facilitate social emotions such as empathy, guilt, and embarrassment (Beer et al 2006, Mitchell et al 2005, Qin & Northoff 2011). Regardless of the precise

function (or functions) of vmPFC, it is brain area that warrants particular attention in the neuroscientific study of psychopathy.

Despite the intriguing degree of similarity between psychopathic personality characteristics and personality changes accompanying vmPFC damage, it is important to note that vmPFC lesion patients do not typically exhibit the full-blown disorder, particularly with respect to antisocial and criminal behavior. Moreover, there are additional regions of PFC whose functions are also likely germane to psychopathy.

Anterior cingulate cortex and behavioral control

A second area of PFC that may play an especially important role in psychopathy is the anterior cingulate cortex (ACC) (Figure 1). The ACC is widely regarded to be a key brain area for motivating and regulating behavior through cognitive and affective mechanisms. Activity in the ACC has been related to a number of functions potentially relevant to psychopathy, including reward, punishment, pain, negative affect, empathy, error detection, performance monitoring, and cognitive control (for reviews see Bush et al 2000, Devinsky et al 1995, Etkin et al 2011, Ridderinkhof et al 2004, Rushworth & Behrens 2008, Shackman et al 2011). Moreover, like the vmPFC, lesion studies of the ACC confirm its importance for emotion and social behavior; cingulotomy patients typically exhibit blunted affect and reduced motivation but greater irritability and social disinhibition (Tow & Whitty 1953). The functional relationship between ACC and vmPFC is reflected in brain structure, as the two regions are anatomically adjacent with dense reciprocal connections (Carmichael & Price 1996, Ongur & Price 2000).

Although this section has highlighted two particular subregions of PFC—the vmPFC and ACC—that figure to play a role in psychopathic behavior, it is important to keep in mind that these regions interact and interconnect with more dorsal and lateral sectors of PFC as well. The following section will summarize neuroimaging data associating psychopathy with structural and functional deficits in PFC.

NEUROIMAGING EVIDENCE FOR PFC DYSFUNCTION IN PSYCHOPATHY

Scope of review

Before summarizing the neuroimaging results, this section will first outline the scope of the studies that were evaluated for this article. This review covers original published reports of human neuroimaging data wherein the authors associate abnormalities in PFC structure or function with psychopathy in adults (in particular, neuroimaging reports with “psychopathy”, “psychopaths”, or “psychopathic” in the title; see Table 2). This approach omits two important related lines of research, which are briefly mentioned here. One is the study of the neural correlates of antisocial traits commonly associated with, but not limited to, psychopathy. Examples include violence (Raine et al 1997, Volkow et al 1995), antisocial personality disorder (Barkataki et al 2006, Raine et al 2000), aggressive/impulsive behavior (Dolan et al 2002), and pathological lying (Yang et al 2005a). Although these traits may commonly overlap with psychopathy, none are unique to psychopathy. Accordingly, neuroimaging findings associated with these traits may not specifically inform the neural basis of psychopathy, and so such studies are omitted in this review. (For a recent review on neuroimaging of antisocial behavior see Yang & Raine 2009). The other line of research omitted here is the neuroimaging of children and adolescents with psychopathic tendencies (e.g., De Brito et al 2009, Jones et al 2009, Marsh et al 2008). Research in children and adolescents is of course critical for understanding the development of antisocial behavior. However, the comparison of imaging data from adult and child/adolescent studies can be challenging for a number of reasons. One reason is that the diagnostic criteria for antisocial behavior in children/adolescents (such as conduct disorder) are necessarily somewhat

different than the criteria for adult psychopathy, reflecting the considerable differences in life circumstances for children, adolescents, and adults. A second reason is that the brain undergoes substantial structural and functional development throughout childhood and adolescence, such that neuroimaging findings vary significantly across pre-adult age groups, even among neurologically and psychologically healthy individuals (Giedd et al 2009). Given these important differences, the child/adolescent literature warrants its own separate review and evaluation. (For a recent review on neuroimaging findings related to antisocial behavior in children see Crowe & Blair 2008).

PFC structure in psychopathy: Gray matter

First we consider studies of PFC gray matter (GM) structure. These studies are largely based on T1 magnetic resonance imaging (MRI) data, from which one can compute measures of GM volume, thickness, and density/concentration. Study designs include between-group analyses (i.e., comparison of psychopathic and non-psychopathic groups) as well as correlation/regression analyses that relate GM measures to psychopathy severity. Together, the studies described in this section converge to associate psychopathy with reductions in PFC GM.

A series of studies by Yang, Raine, and colleagues advanced this area of research by recruiting subjects with varying degrees of psychopathy from temporary employment agencies. The first such study (Yang et al 2005b) compared gray matter volume across the entire PFC in three subject types: “unsuccessful” psychopaths (those with one or more criminal convictions; $n=16$), “successful” psychopaths (no criminal convictions; $n=13$) and a non-psychopathic comparison group ($n=23$). Overall, PFC GM volume correlated negatively with total PCL-R score, as well as with each factor/subscale of psychopathy. Unsuccessful psychopaths had lower PFC volumes than both successful psychopaths and non-psychopaths, although successful psychopaths and non-psychopaths did not significantly differ with respect to PFC volume. In a follow-up study, Yang et al found reduced GM volume specifically in lateral PFC and ventral PFC/orbitofrontal cortex (OFC) in unsuccessful psychopaths, but not in successful psychopaths, as compared to non-psychopaths (Yang et al 2010). In a separate study, Yang et al used cortical pattern matching to examine regional cortical thickness in a group of $n=27$ psychopathic adult males to a group of $n=32$ non-psychopathic males, matched for age, gender, and substance abuse (Yang et al 2009a). The psychopathic group had reduced cortical thickness in right lateral PFC, and to a lesser extent, in right vmPFC and rostral ACC. In these areas, cortical thickness was negatively correlated with the affective facet of psychopathy (a subset of Factor 1).

Focusing on brain areas involved in moral cognition, de Oliveira-Souza et al used voxel-based morphometry (VBM) to evaluate gray matter volume (GMV) and concentration (GMC) in a group of $n=15$ adult neuropsychiatric patients diagnosed with Antisocial Personality Disorder (ASPD) and a group of $n=15$ neuropsychiatrically healthy adults matched for age, sex, education, and cognitive ability (de Oliveira-Souza et al 2008). The ASPD group had significant GMV reductions in OFC and frontopolar cortex, including parts of vmPFC. In addition, GMC in an area of posterior OFC/vmPFC significantly correlated with Factor 1 scores in the patient group.

While the aforementioned studies were performed in non-forensic settings, a number of structural imaging studies of psychopathy have involved incarcerated subject samples. One advantage to studying incarcerated samples is that it is typically easier to recruit individuals with especially high psychopathy scores (see Methodological Caveats below). In one of the first such studies, Muller et al used VBM to compare GM in a group of $n=17$ adult male forensic inpatients diagnosed with psychopathy (based on the PCL-R) to a group of $n=17$ non-incarcerated, healthy age-matched males (Muller et al 2008). The psychopathic group

had significant GM reductions in right dorsal ACC and dorsolateral PFC bilaterally. In a subsequent study, Boccardi et al used a cortical pattern matching technique to compare gray matter density in a group of $n=26$ adult male violent offenders to a group of $n=25$ adult male age-matched non-offenders (Boccardi et al 2011). The greatest group differences in PFC were observed in vmPFC and ACC—in both areas the psychopathic group had lower gray matter density. Recently, Ly et al compared cortical thickness in a group of adult male psychopathic prison inmates ($n=21$) to a group of adult male non-psychopathic prison inmates ($n=31$) matched for age, IQ, and substance abuse history (Ly et al in press). The psychopathic inmates exhibited significantly thinner GM in the left dorsal ACC and adjacent dorsomedial PFC as well as left dorsolateral PFC. In by far the largest study to date, Ermer et al used VBM and multiple regression analysis in a group of $n=254$ adult incarcerated men to examine the relationship between psychopathy severity and GM structure, while covarying for age, substance use, and overall brain volume (Ermer et al 2011). They found a significant negative relationship between total PCL-R score and GMV/GMC in OFC and vmPFC. In addition, they found a negative relationship between GMC and Factor 2 score in left OFC/vmPFC.

In sum, across all of the aforementioned studies, psychopathy was associated with significant reductions in prefrontal gray matter, with the most commonly reported subregions being vmPFC/OFC and ACC.

PFC structure in psychopathy: White matter

There have been comparatively fewer studies examining PFC white matter (WM) structure in psychopathy. Diffusion tensor imaging (DTI) is an MRI technique for assessing microstructural integrity of WM tracts (Pierpaoli & Basser 1996). To date, only two studies have used DTI to study psychopathy (Craig et al 2009, Motzkin et al 2011), and both associate psychopathy with reduced integrity of the uncinate fasciculus, a major white matter tract connecting PFC with subcortical structures. Importantly, the uncinate fasciculus includes fibers connecting vmPFC and amygdala. Despite the remarkable convergence of findings between these two studies, there is a need for subsequent studies to confirm the initial findings, which are based on relatively small samples. The first study (Craig et al 2009) included $n=9$ psychopaths and $n=9$ non-psychopaths recruited from a forensic psychiatric hospital, while the second study (Motzkin et al 2011) included $n=14$ psychopaths and $n=13$ non-psychopaths recruited from a medium-security prison. Nonetheless, these initial studies of WM integrity are consistent with PFC dysfunction in psychopathy.

PFC function in psychopathy

Functional imaging studies of psychopathy can be divided into two groups based on the type of fMRI analysis: resting-state functional connectivity MRI (rs-fcMRI) or task-related fMRI. rs-fcMRI measures the degree of spontaneously correlated BOLD activity between brain regions while the subject is at rest (e.g., staring at a fixation cross with no task demands). The degree of spontaneously correlated activity at rest is thought to reflect the extent to which macroscopic brain areas are functionally interconnected. Two studies have identified reduced PFC functional connectivity in criminal psychopaths using rs-fcMRI. One study of $n=20$ psychopathic and $n=20$ non-psychopathic adult male prison inmates found reduced functional connectivity between the amygdala and anterior vmPFC in psychopathy (Motzkin et al 2011), while a second study of the same subject sample found reduced functional connectivity between the dorsal ACC and insula (Ly et al in press).

Results from task-related fMRI studies are somewhat more difficult to summarize, as they have employed a range of stimulus and response paradigms to test hypotheses regarding particular patterns of brain activation. Despite a variety of task paradigms, a considerable

number of studies associate psychopathy with abnormal activity within vmPFC and/or ACC. Such findings have been obtained in studies of fear conditioning (Birbaumer et al 2005), cognitive control (Sadeh et al 2011), speech production (Sheng et al 2010), affective theory of mind (Sommer et al 2010), anticipation and receipt of reward (Bjork et al 2012), social competition and aggression (Veit et al 2009), social cooperation and defection (Rilling et al 2007), recollection of emotionally salient words (Kiehl et al 2001), viewing emotionally salient scenes (Harenski et al 2009, Muller et al 2003), viewing facial expressions of emotion (Dolan & Fullam 2009), and viewing pictures of moral violations (Harenski et al 2010). Psychopathy has also been associated with PFC abnormalities outside vmPFC/ACC, including during tasks involving social-affective processing, such as moral decision-making (Glenn et al 2009) and recognition of facial expressions of emotion (Gordon et al 2004).

Overall, the available functional imaging data suggest abnormal PFC function in psychopathy, particularly within vmPFC and ACC.

CAVEATS

Methodological caveats

When considering the results of neuroimaging studies of psychopathy, it is important to keep in mind some of the methodological challenges associated with studying this disorder. One significant issue is how “psychopathy” is defined. As described above, most studies of criminal populations rely on the Hare Psychopathy Checklist-Revised (PCL-R) to define psychopathy. The PCL-R manual advises standard cutoff scores for grouping subjects: total scores of 30 or greater indicate psychopathy, scores of 20 or less indicate non-psychopathy, and scores of 21–29 are considered intermediate (Hare 2003). However, because subjects with high PCL-R scores (i.e., 30) are typically difficult to recruit for imaging studies, especially outside forensic settings, researchers have employed a variety of minimum PCL-R total scores to define psychopathy. Cutoff scores in the mid-20s (or even lower) are fairly common (see Table 2). The issue with such lenient classification criteria is that the imaging findings in the “psychopathic” group may pertain more generally to ASPD or externalizing behavior, rather than specifically to psychopathy, *per se*.

A second methodological challenge for studies of psychopathic criminals is assembling a group of non-psychopathic comparison subjects that are well-matched to the psychopathic subjects on important variables that can affect brain structure and function (e.g., age, education, cognitive ability, substance abuse history). In several of the studies described above, the psychopathic group and non-psychopathic comparison group differed dramatically on one or more of these variables (e.g., Boccardi et al 2011, Muller et al 2008).

Finally, it is important to make a distinction between studies that involve criminal offenders who meet the diagnostic criteria for psychopathy (e.g., Harenski et al 2010, Ly et al in press, Motzkin et al 2011, Muller et al 2003), versus those studies that involve community samples of relatively normal individuals with varying degrees of “psychopathic” personality traits (e.g., Bjork et al 2012, Harenski et al 2009, Rilling et al 2007, Sadeh et al 2011, Sheng et al 2010). The neurobiological features that correlate with PPI scores in normal functioning individuals (e.g., college undergraduates) may not necessarily correspond to the same features associated with the categorical diagnosis of psychopathy. Future neuroimaging research will be necessary to resolve this question.

Neurobiological caveats

Although this review has focused primarily on subregions of PFC, there are of course regions outside PFC that may play important roles in psychopathy as well. In particular, the amygdala has received considerable attention, with numerous studies associating

psychopathy with abnormal size, shape, or activity of this subcortical structure (Birbaumer et al 2005, Boccardi et al 2011, Dolan & Fullam 2009, Glenn et al 2009, Harenski et al 2010, Harenski et al 2009, Yang et al 2010, Yang et al 2009b). Furthermore, the anterior temporal cortex, which overlies and densely interconnects with the amygdala, has also been repeatedly associated with structural and functional deficits in psychopathy (Ermer et al 2011, Kiehl et al 2004, Ly et al in press, Yang et al 2009a). Additional areas that warrant mention are the hippocampus (Boccardi et al 2010, Laakso et al 2001), insula (de Oliveira-Souza et al 2008, Ly et al in press), and striatum (Buckholz et al 2010, Glenn et al 2010). The role of PFC in psychopathy likely depends in large part on its interactions with these areas outside PFC; therefore, future studies of PFC functional and structural connectivity will help elucidate the neurobiological mechanisms of the disorder.

CONCLUSION

The neuroscientific study of psychopathy has advanced rapidly in recent years, with an increasing appreciation for the potential of the results to impact law, psychiatric medicine, and public policy. Although a growing number of studies are converging to implicate the prefrontal cortex (and specific subsectors thereof) as key pathophysiological substrates for the disorder, the field continues to face a number of methodological challenges and limitations that will need to be addressed in future studies. Moreover, a comprehensive understanding of prefrontal dysfunction in psychopathy will no doubt depend on further refinement of the basic systems-level understanding of normal prefrontal function. Pending future progress in these areas of research, the sober assessment of Hervey Cleckley from “The Mask of Sanity” still applies: “I do not believe that the cause of the psychopath’s disorder has yet been discovered and demonstrated. Until we have more and better evidence than is at present available, let us admit the incompleteness of our knowledge and modestly pursue our inquiry.”

References

- Barkataki I, Kumari V, Das M, Taylor P, Sharma T. Volumetric structural brain abnormalities in men with schizophrenia or antisocial personality disorder. *Behav Brain Res*. 2006; 169:239–47. [PubMed: 16466814]
- Barrash J, Tranel D, Anderson SW. Acquired personality disturbances associated with bilateral damage to the ventromedial prefrontal region. *Dev Neuropsychol*. 2000; 18:355–81. [PubMed: 11385830]
- Bechara A, Damasio H, Tranel D, Damasio AR. Deciding advantageously before knowing the advantageous strategy. *Science*. 1997; 275:1293–5. [PubMed: 9036851]
- Beer JS, John OP, Scabini D, Knight RT. Orbitofrontal cortex and social behavior: integrating self-monitoring and emotion-cognition interactions. *J Cogn Neurosci*. 2006; 18:871–9. [PubMed: 16839295]
- Birbaumer N, Veit R, Lotze M, Erb M, Hermann C, et al. Deficient fear conditioning in psychopathy: a functional magnetic resonance imaging study. *Arch Gen Psychiatry*. 2005; 62:799–805. [PubMed: 15997022]
- Bjork JM, Chen G, Hommer DW. Psychopathic tendencies and mesolimbic recruitment by cues for instrumental and passively obtained rewards. *Biol Psychol*. 2012; 89:408–15. [PubMed: 22178441]
- Blair RJ. The amygdala and ventromedial prefrontal cortex in morality and psychopathy. *Trends Cogn Sci*. 2007; 11:387–92. [PubMed: 17707682]
- Blair RJ. The amygdala and ventromedial prefrontal cortex: functional contributions and dysfunction in psychopathy. *Philos Trans R Soc Lond B Biol Sci*. 2008; 363:2557–65. [PubMed: 18434283]
- Blumer, D.; Benson, DF. Personality changes with frontal and temporal lesions. In: Benson, DF.; Blumer, D., editors. *Psychiatric Aspects of Neurological Disease*. New York: Stratton; 1975.

- Boccardi M, Frisoni GB, Hare RD, Cavedo E, Najt P, et al. Cortex and amygdala morphology in psychopathy. *Psychiatry Res.* 2011; 193:85–92. [PubMed: 21676597]
- Boccardi M, Ganzola R, Rossi R, Sabattoli F, Laakso MP, et al. Abnormal hippocampal shape in offenders with psychopathy. *Hum Brain Mapp.* 2010; 31:438–47. [PubMed: 19718651]
- Buckholtz JW, Treadway MT, Cowan RL, Woodward ND, Benning SD, et al. Mesolimbic dopamine reward system hypersensitivity in individuals with psychopathic traits. *Nat Neurosci.* 2010; 13:419–21. [PubMed: 20228805]
- Budhani S, Richell RA, Blair RJ. Impaired reversal but intact acquisition: probabilistic response reversal deficits in adult individuals with psychopathy. *J Abnorm Psychol.* 2006; 115:552–8. [PubMed: 16866595]
- Bush G, Luu P, Posner MI. Cognitive and emotional influences in anterior cingulate cortex. *Trends Cogn Sci.* 2000; 4:215–22. [PubMed: 10827444]
- Carmichael ST, Price JL. Connectional networks within the orbital and medial prefrontal cortex of macaque monkeys. *J Comp Neurol.* 1996; 371:179–207. [PubMed: 8835726]
- Ciaromelli E, Muccioli M, Ladavas E, di Pellegrino G. Selective deficit in personal moral judgment following damage to ventromedial prefrontal cortex. *Soc Cogn Affect Neurosci.* 2007; 2:84–92. [PubMed: 18985127]
- Cleckley, H. *The Mask of Sanity.* St. Louis: Mosby; 1941.
- Cleckley, H. *The Mask of Sanity.* 5. St. Louis: Mosby; 1976.
- Craig MC, Catani M, Deeley Q, Latham R, Daly E, et al. Altered connections on the road to psychopathy. *Mol Psychiatry.* 2009; 14:946–53. 07. [PubMed: 19506560]
- Crowe SL, Blair RJ. The development of antisocial behavior: what can we learn from functional neuroimaging studies? *Dev Psychopathol.* 2008; 20:1145–59. [PubMed: 18838035]
- Damasio, AR. *Descartes' Error.* New York: Putnam; 1994.
- Damasio AR. The somatic marker hypothesis and the possible functions of the prefrontal cortex. *Philos Trans R Soc Lond B Biol Sci.* 1996; 351:1413–20. [PubMed: 8941953]
- Damasio AR, Tranel D, Damasio H. Individuals with sociopathic behavior caused by frontal damage fail to respond autonomically to social stimuli. *Behav Brain Res.* 1990; 41:81–94. [PubMed: 2288668]
- Damasio H, Grabowski T, Frank R, Galaburda AM, Damasio AR. The return of Phineas Gage: clues about the brain from the skull of a famous patient. *Science.* 1994; 264:1102–5. [PubMed: 8178168]
- De Brito SA, Mechelli A, Wilke M, Laurens KR, Jones AP, et al. Size matters: increased grey matter in boys with conduct problems and callous-unemotional traits. *Brain.* 2009; 132:843–52. [PubMed: 19293245]
- de Oliveira-Souza R, Hare RD, Bramati IE, Garrido GJ, Azevedo Ignacio F, et al. Psychopathy as a disorder of the moral brain: fronto-temporo-limbic grey matter reductions demonstrated by voxel-based morphometry. *Neuroimage.* 2008; 40:1202–13. [PubMed: 18289882]
- Devinsky O, Morrell MJ, Vogt BA. Contributions of anterior cingulate cortex to behaviour. *Brain.* 1995; 118 (Pt 1):279–306. [PubMed: 7895011]
- Dolan MC, Deakin JF, Roberts N, Anderson IM. Quantitative frontal and temporal structural MRI studies in personality-disordered offenders and control subjects. *Psychiatry Res.* 2002; 116:133–49. [PubMed: 12477598]
- Dolan MC, Fullam RS. Psychopathy and functional magnetic resonance imaging blood oxygenation level-dependent responses to emotional faces in violent patients with schizophrenia. *Biol Psychiatry.* 2009; 66:570–7. [PubMed: 19446795]
- Ermer E, Cope LM, Nyalakanti PK, Calhoun VD, Kiehl KA. Aberrant paralimbic gray matter in criminal psychopathy. *J Abnorm Psychol.* 2011
- Eslinger PJ, Damasio AR. Severe disturbance of higher cognition after bilateral frontal lobe ablation: patient EVR. *Neurology.* 1985; 35:1731–41. [PubMed: 4069365]
- Etkin A, Egner T, Kalisch R. Emotional processing in anterior cingulate and medial prefrontal cortex. *Trends Cogn Sci.* 2011; 15:85–93. [PubMed: 21167765]

- Giedd JN, Lalonde FM, Celano MJ, White SL, Wallace GL, et al. Anatomical brain magnetic resonance imaging of typically developing children and adolescents. *J Am Acad Child Adolesc Psychiatry*. 2009; 48:465–70. [PubMed: 19395901]
- Glenn AL, Raine A, Schug RA. The neural correlates of moral decision-making in psychopathy. *Mol Psychiatry*. 2009; 14:5–6. [PubMed: 19096450]
- Glenn AL, Raine A, Yaralian PS, Yang Y. Increased volume of the striatum in psychopathic individuals. *Biol Psychiatry*. 2010; 67:52–8. [PubMed: 19683706]
- Gordon HL, Baird AA, End A. Functional differences among those high and low on a trait measure of psychopathy. *Biol Psychiatry*. 2004; 56:516–21. [PubMed: 15450788]
- Grabenhorst F, Rolls ET. Value, pleasure and choice in the ventral prefrontal cortex. *Trends Cogn Sci*. 2011; 15:56–67. [PubMed: 21216655]
- Hare, RD. The Hare psychopathy checklist-revised. 2. Toronto: Multi-Health Systems; 2003.
- Hare RD, Frazelle J, Cox DN. Psychopathy and physiological responses to threat of an aversive stimulus. *Psychophysiology*. 1978; 15:165–72. [PubMed: 652911]
- Harenski CL, Harenski KA, Shane MS, Kiehl KA. Aberrant neural processing of moral violations in criminal psychopaths. *J Abnorm Psychol*. 2010; 119:863–74. [PubMed: 21090881]
- Harenski CL, Kim SH, Hamann S. Neuroticism and psychopathy predict brain activation during moral and nonmoral emotion regulation. *Cogn Affect Behav Neurosci*. 2009; 9:1–15. [PubMed: 19246323]
- Harlow JM. Recovery from the passage of an iron bar through the head. Publications of the Massachusetts Medical Society. 1868; 2:327–47.
- Harpur TJ, Hare RD, Hakstian R. A two factor conceptualization of psychopathy: Construct validity and implications for assessment. *Psychological Assessment*. 1989; 1:6–17.
- Hornak J, O'Doherty J, Bramham J, Rolls ET, Morris RG, et al. Reward-related reversal learning after surgical excisions in orbito-frontal or dorsolateral prefrontal cortex in humans. *J Cogn Neurosci*. 2004; 16:463–78. [PubMed: 15072681]
- Jones AP, Laurens KR, Herba CM, Barker GJ, Viding E. Amygdala hypoactivity to fearful faces in boys with conduct problems and callous-unemotional traits. *Am J Psychiatry*. 2009; 166:95–102. [PubMed: 18923070]
- Kiehl KA, Smith AM, Hare RD, Mendrek A, Forster BB, et al. Limbic abnormalities in affective processing by criminal psychopaths as revealed by functional magnetic resonance imaging. *Biol Psychiatry*. 2001; 50:677–84. [PubMed: 11704074]
- Kiehl KA, Smith AM, Mendrek A, Forster BB, Hare RD, Liddle PF. Temporal lobe abnormalities in semantic processing by criminal psychopaths as revealed by functional magnetic resonance imaging. *Psychiatry Res*. 2004; 130:27–42. [PubMed: 14972366]
- Koenigs M, Huey ED, Calamia M, Raymond V, Tranel D, Grafman J. Distinct regions of prefrontal cortex mediate resistance and vulnerability to depression. *J Neurosci*. 2008a; 28:12341–48. [PubMed: 19020027]
- Koenigs M, Huey ED, Raymond V, Cheon B, Solomon J, et al. Focal brain damage protects against post-traumatic stress disorder in combat veterans. *Nat Neurosci*. 2008b; 11:232–7. [PubMed: 18157125]
- Koenigs M, Kruepke M, Newman JP. Economic decision-making in psychopathy: a comparison with ventromedial prefrontal lesion patients. *Neuropsychologia*. 2010; 48:2198–204. [PubMed: 20403367]
- Koenigs M, Kruepke M, Zeier J, Newman JP. Utilitarian moral judgment in psychopathy. *Soc Cogn Affect Neurosci*. 2011
- Koenigs M, Tranel D. Irrational economic decision-making after ventromedial prefrontal damage: evidence from the Ultimatum Game. *J Neurosci*. 2007; 27:951–6. [PubMed: 17251437]
- Koenigs M, Young L, Adolphs R, Tranel D, Cushman F, et al. Damage to the prefrontal cortex increases utilitarian moral judgements. *Nature*. 2007; 446:908–11. [PubMed: 17377536]
- Krajibich I, Adolphs R, Tranel D, Denburg NL, Camerer CF. Economic games quantify diminished sense of guilt in patients with damage to the prefrontal cortex. *J Neurosci*. 2009; 29:2188–92. [PubMed: 19228971]

- Laakso MP, Vaurio O, Koivisto E, Savolainen L, Eronen M, et al. Psychopathy and the posterior hippocampus. *Behav Brain Res.* 2001; 118:187–93. [PubMed: 11164516]
- Lilienfeld SO, Andrews BP. Development and preliminary validation of a self-report measure of psychopathic personality traits in noncriminal populations. *J Pers Assess.* 1996; 66:488–524. [PubMed: 8667144]
- Ly M, Motzkin JC, Philippi C, Kirk GR, Newman JP, et al. Cortical thinning in psychopathy. *American Journal of Psychiatry.* in press.
- Lykken DT. A study of anxiety in the sociopathic personality. *J Abnorm Psychol.* 1957; 55:6–10. [PubMed: 13462652]
- Lykken, DT. *The antisocial personalities.* Mahwah: Erlbaum; 1995.
- Marsh AA, Finger EC, Mitchell DG, Reid ME, Sims C, et al. Reduced amygdala response to fearful expressions in children and adolescents with callous-unemotional traits and disruptive behavior disorders. *Am J Psychiatry.* 2008; 165:712–20. [PubMed: 18281412]
- Milad MR, Rauch SL, Pitman RK, Quirk GJ. Fear extinction in rats: implications for human brain imaging and anxiety disorders. *Biol Psychol.* 2006; 73:61–71. [PubMed: 16476517]
- Mitchell JP, Banaji MR, Macrae CN. The link between social cognition and self-referential thought in the medial prefrontal cortex. *J Cogn Neurosci.* 2005; 17:1306–15. [PubMed: 16197685]
- Moretti L, Dragone D, di Pellegrino G. Reward and social valuation deficits following ventromedial prefrontal damage. *J Cogn Neurosci.* 2009; 21:128–40. [PubMed: 18476758]
- Motzkin JC, Newman JP, Kiehl KA, Koenigs M. Reduced prefrontal connectivity in psychopathy. *J Neurosci.* 2011; 31:17348–57. [PubMed: 22131397]
- Muller JL, Ganssbauer S, Sommer M, Dohnel K, Weber T, et al. Gray matter changes in right superior temporal gyrus in criminal psychopaths. Evidence from voxel-based morphometry. *Psychiatry Res.* 2008; 163:213–22. [PubMed: 18662867]
- Muller JL, Sommer M, Wagner V, Lange K, Taschler H, et al. Abnormalities in emotion processing within cortical and subcortical regions in criminal psychopaths: evidence from a functional magnetic resonance imaging study using pictures with emotional content. *Biol Psychiatry.* 2003; 54:152–62. [PubMed: 12873805]
- Myers-Schulz B, Koenigs M. Functional anatomy of ventromedial prefrontal cortex: implications for mood and anxiety disorders. *Mol Psychiatry.* 2012; 17:132–41. [PubMed: 21788943]
- O'Doherty JP. Contributions of the ventromedial prefrontal cortex to goal-directed action selection. *Ann N Y Acad Sci.* 2011; 1239:118–29. [PubMed: 22145881]
- Ongur D, Price JL. The organization of networks within the orbital and medial prefrontal cortex of rats, monkeys and humans. *Cereb Cortex.* 2000; 10:206–19. [PubMed: 10731217]
- Patrick CJ, Bradley MM, Lang PJ. Emotion in the criminal psychopath: startle reflex modulation. *J Abnorm Psychol.* 1993; 102:82–92. [PubMed: 8436703]
- Patrick CJ, Cuthbert BN, Lang PJ. Emotion in the criminal psychopath: fear image processing. *J Abnorm Psychol.* 1994; 103:523–34. [PubMed: 7930052]
- Phelps EA, Delgado MR, Nearing KI, LeDoux JE. Extinction learning in humans: role of the amygdala and vmPFC. *Neuron.* 2004; 43:897–905. [PubMed: 15363399]
- Pierpaoli C, Basser PJ. Toward a quantitative assessment of diffusion anisotropy. *Magn Reson Med.* 1996; 36:893–906. [PubMed: 8946355]
- Qin P, Northoff G. How is our self related to midline regions and the default-mode network? *Neuroimage.* 2011; 57:1221–33. [PubMed: 21609772]
- Raine A, Buchsbaum M, LaCasse L. Brain abnormalities in murderers indicated by positron emission tomography. *Biol Psychiatry.* 1997; 42:495–508. [PubMed: 9285085]
- Raine A, Lencz T, Bihle S, LaCasse L, Colletti P. Reduced prefrontal gray matter volume and reduced autonomic activity in antisocial personality disorder. *Arch Gen Psychiatry.* 2000; 57:119–27. discussion 28–9. [PubMed: 10665614]
- Rauch SL, Shin LM, Phelps EA. Neurocircuitry models of posttraumatic stress disorder and extinction: human neuroimaging research--past, present, and future. *Biol Psychiatry.* 2006; 60:376–82. [PubMed: 16919525]

- Ridderinkhof KR, Ullsperger M, Crone EA, Nieuwenhuis S. The role of the medial frontal cortex in cognitive control. *Science*. 2004; 306:443–7. [PubMed: 15486290]
- Rilling JK, Glenn AL, Jairam MR, Pagnoni G, Goldsmith DR, et al. Neural correlates of social cooperation and non-cooperation as a function of psychopathy. *Biol Psychiatry*. 2007; 61:1260–71. [PubMed: 17046722]
- Rushworth MF, Behrens TE. Choice, uncertainty and value in prefrontal and cingulate cortex. *Nat Neurosci*. 2008; 11:389–97. [PubMed: 18368045]
- Sadeh N, Spielberg JM, Heller W, Herrington JD, Engels AS, et al. Emotion disrupts neural activity during selective attention in psychopathy. *Soc Cogn Affect Neurosci*. 2011
- Shackman AJ, Salomons TV, Slagter HA, Fox AS, Winter JJ, Davidson RJ. The integration of negative affect, pain and cognitive control in the cingulate cortex. *Nat Rev Neurosci*. 2011; 12:154–67. [PubMed: 21331082]
- Sheng T, Gheytanchi A, Aziz-Zadeh L. Default network deactivations are correlated with psychopathic personality traits. *PloS one*. 2010; 5:e12611. [PubMed: 20830290]
- Sommer M, Sodian B, Dohnel K, Schwerdtner J, Meinhardt J, Hajak G. In psychopathic patients emotion attribution modulates activity in outcome-related brain areas. *Psychiatry Res*. 2010; 182:88–95. [PubMed: 20417065]
- Tow PM, Whitty CW. Personality changes after operations on the cingulate gyrus in man. *J Neurol Neurosurg Psychiatry*. 1953; 16:186–93. [PubMed: 13085202]
- Veit R, Lotze M, Sewing S, Missenhardt H, Gaber T, Birbaumer N. Aberrant social and cerebral responding in a competitive reaction time paradigm in criminal psychopaths. *Neuroimage*. 2009
- Volkow ND, Tancredi LR, Grant C, Gillespie H, Valentine A, et al. Brain glucose metabolism in violent psychiatric patients: a preliminary study. *Psychiatry Res*. 1995; 61:243–53. [PubMed: 8748468]
- Yang Y, Raine A. Prefrontal structural and functional brain imaging findings in antisocial, violent, and psychopathic individuals: a meta-analysis. *Psychiatry Res*. 2009; 174:81–8. [PubMed: 19833485]
- Yang Y, Raine A, Colletti P, Toga AW, Narr KL. Abnormal temporal and prefrontal cortical gray matter thinning in psychopaths. *Mol Psychiatry*. 2009a; 14:561–2. 55. [PubMed: 19455172]
- Yang Y, Raine A, Colletti P, Toga AW, Narr KL. Morphological alterations in the prefrontal cortex and the amygdala in unsuccessful psychopaths. *J Abnorm Psychol*. 2010; 119:546–54. [PubMed: 20677843]
- Yang Y, Raine A, Lencz T, Bihle S, Lacasse L, Colletti P. Prefrontal white matter in pathological liars. *Br J Psychiatry*. 2005a; 187:320–5. [PubMed: 16199789]
- Yang Y, Raine A, Lencz T, Bihle S, LaCasse L, Colletti P. Volume reduction in prefrontal gray matter in unsuccessful criminal psychopaths. *Biol Psychiatry*. 2005b; 57:1103–8. [PubMed: 15866549]
- Yang Y, Raine A, Narr KL, Colletti P, Toga AW. Localization of deformations within the amygdala in individuals with psychopathy. *Arch Gen Psychiatry*. 2009b; 66:986–94. [PubMed: 19736355]
- Young L, Bechara A, Tranel D, Damasio H, Hauser M, Damasio A. Damage to ventromedial prefrontal cortex impairs judgment of harmful intent. *Neuron*. 2010; 65:845–51. [PubMed: 20346759]

Figure 1. Mid-sagittal view of the brain illustrating vmPFC (red) and ACC (blue). The vmPFC also includes the medial sector of the orbital surface.

Table 1

Measures of psychopathy.

Cleckley characteristics	Hare PCL-R items (Factor 1 or 2)	PPI subscales
Superficial charm and good intelligence	Glibness/Superficial Charm (F1)	Social Potency
Pathologic egocentricity and incapacity for love	Grandiose Sense of Self-Worth (F1)	Machiavellian Egocentricity
	Need for Stimulation/Proneness to Boredom (F2)	Fearlessness
Untruthfulness and insincerity	Pathological Lying (F1)	
	Conning/Manipulative (F1)	
Lack of remorse or shame	Lack of Remorse or Guilt (F1)	
General poverty in major affective reactions	Shallow Affect (F1)	
Unresponsiveness in general interpersonal relations	Callous/Lack of Empathy (F1)	Coldheartedness
	Parasitic Lifestyle (F2)	
Poor judgment and failure to learn by experience	Poor Behavioral Controls (F2)	
Sex life impersonal, trivial, and poorly integrated	Promiscuous Sexual Behavior (F2)	
	Early Behavior Problems (F2)	
Failure to follow any life plan	Lack of Realistic, Long-Term Goals (F2)	Carefree Nonplanfulness
Inadequately motivated antisocial behavior	Impulsivity (F2)	Impulsive Nonconformity
Unreliability	Irresponsibility (F2)	
	Failure to Accept Responsibility for Own Actions (F1)	Blame Externalization
	Many Short-term Marital Relationships (neither)	
	Juvenile Delinquency (F2)	
	Revocation of Conditional Release (F2)	
	Criminal Versatility (neither)	
Absence of delusions and irrational thinking		
Absence of "nervousness"		Stress Immunity
Specific loss of insight		
Fantastic and uninviting behavior with drink or sometimes without		
Suicide rarely carried out		

Similar items between measures are placed in the same row.

Table 2
Neuroimaging studies associating psychopathy with abnormal PFC structure or function.

First Author	Year	Title	Type of imaging	Type of analysis	PCL-R cutoff for P	Mean PCL-R for Ps
Birbaumer	2005	Deficient fear conditioning in psychopathy : a functional magnetic resonance imaging study	F	BG	15	24.9
Bjork	2012	Psychopathic tendencies and mesolimbic recruitment by cues for instrumental and passively obtained rewards	F	C/R	n/a	n/a
Boccardi	2011	Cortex and amygdala morphology in psychopathy	S	BG	21	29.9
Craig	2009	Altered connections on the road to psychopathy	S	BG, C/R	25	28.4
de Oliveira-Souza	2008	Psychopathy as a disorder of the moral brain: fronto-temporo-limbic grey matter reductions demonstrated by voxel-based morphometry	S	BG, C/R	n/a	n/a
Dolan	2009	Psychopathy and functional magnetic resonance imaging blood oxygenation level-dependent responses to emotional faces in violent patients with schizophrenia	F	BG, C/R	n/a	n/a
Erner	2011	Aberrant paralimbic gray matter in criminal psychopathy	S	C/R	n/a	n/a
Glenn	2009	The neural correlates of moral decision-making in psychopathy	F	C/R	n/a	n/a
Gordon	2004	Functional differences among those high and low on a trait measure of psychopathy	F	BG	n/a	n/a
Harenski	2009	Neuroticism and psychopathy predict brain activation during moral and nonmoral emotion regulation	F	C/R	n/a	n/a
Harenski	2010	Aberrant neural processing of moral violations in criminal psychopaths	F	BG, C/R	30	31.8
Intrator	1997	A brain imaging (single photon emission computerized tomography) study of semantic and affective processing in psychopaths	F	BG	25	29.9
Kiehl	2001	Limbic abnormalities in affective processing by criminal psychopaths as revealed by functional magnetic resonance imaging	F	BG	24	32.8
Ly	2012	Cortical thinning in psychopathy	F,S	BG	30	31.8
Motzkin	2011	Reduced prefrontal connectivity in psychopathy	F,S	BG	30	31.9
Muller	2003	Abnormalities in emotion processing within cortical and subcortical regions in criminal psychopaths : evidence from a functional magnetic resonance imaging study using pictures with emotional content	F	BG	31	36.8
Muller	2008	Gray matter changes in right superior temporal gyrus in criminal psychopaths . Evidence from voxel-based morphometry	S	BG	28	33.4
Muller	2008	Disturbed prefrontal and temporal brain function during emotion and cognition interaction in criminal psychopathy	F	BG	28	30.5
Rilling	2007	Neural correlates of social cooperation and non-cooperation as a function of psychopathy	F	C/R	n/a	n/a
Sadeh	2011	Emotion disrupts neural activity during selective attention in psychopathy	F	C/R	n/a	n/a
Sheng	2010	Default network deactivations are correlated with psychopathic personality traits	F	C/R	n/a	n/a

First Author	Year	Title	Type of imaging	Type of analysis	PCL-R cutoff for P	Mean PCL-R for Ps
Sommer	2010	In psychopathic patients emotion attribution modulates activity in outcome-related brain areas	F	BG	28	28.6
Veit	2009	Aberrant social and cerebral responding in a competitive reaction time paradigm in criminal psychopaths	F	C/R	n/a	n/a
Yang	2005	Volume reduction in prefrontal gray matter in unsuccessful criminal psychopaths	S	BG, C/R	23	28.4
Yang	2009	Abnormal temporal and prefrontal cortical gray matter thinning in psychopaths	S	BG, C/R	n/a	n/a
Yang	2010	Morphological alterations in the prefrontal cortex and the amygdala in unsuccessful psychopaths	S	BG	23	n/a

P= psychopathy; S= structural; F = functional; C/R = correlation or regression analysis; BG = between-group analysis