CASE REPORT

Huge trichobezoar causing obstructive jaundice

Ankit Verma,¹ Sourabh Sharma,¹ Gaurav Tyagi,¹ Sugandha Singh²¹Department of General Surgery, Sawai Man Singh Hospital, Jaipur, Rajasthan, India²MGM Medical College, Aurangabad, Maharashtra, India**Correspondence to**Dr Ankit Verma,
ankit.doc2004@gmail.com

Accepted 3 February 2014

SUMMARY

Trichobezoars are composed of hair or hair-like fibres that are commonly observed in the gastrointestinal tract of children and young women with psychiatric illnesses. Presentation of trichobezoars can be diverse. We report a case of a 29-year-old woman who presented with obstructive jaundice and a large upper abdominal lump. She was diagnosed with post-traumatic stress disorder on psychiatric evaluation. She underwent exploratory laparotomy and a large trichobezoar was removed. Her postoperative course was uneventful and she was referred to the psychiatric department for further management.

BACKGROUND

Trichobezoar may present with an upper abdominal lump, anaemia, weight loss, malnutrition, perforation and rarely as obstructive jaundice. Our case is important for its rarity. The possibility of trichobezoar should be considered in every case of obstructive jaundice, especially those presenting as epigastric lumps in patients with known psychiatric health problems.

CASE PRESENTATION

A 29-year-old married woman presented to the outpatient department with abdominal pain and epigastric fullness for 4 months along with jaundice and pruritus. She also reported occasional non-bilious, non-bloody vomiting. There was no history of any drug intake or repeated attacks of pain. She was asthenic, with yellowish discoloration of the eyes.

Abdominal examination revealed a single large, mobile lump approximately 16×12 cm in the epigastric region, which was extending up to the umbilicus. It was firm in consistency (figures 1 and 2).

On psychiatric consultation, she was diagnosed as a case of post-traumatic stress disorder, considering a change in her social circumstances induced by repeated abuse by her husband. On enquiry, she reported a history of 'flashbacks' with emotional blunting, numbness and insomnia.

Figure 2 Clinical picture showing the extent of the abdominal lump.

She had a disturbed personal life, as her husband was a chronic alcoholic, construction labourer who used to physically assault her. She has been living with her parents for the past 6–8 months during which she was observed by family members to ingest her own hair.

INVESTIGATIONS

Her haemoglobin was 8.2 g/dL, and total bilirubin was 0.854 mmol/L with direct bilirubin 0.566 mmol/L and indirect bilirubin 0.288 mmol/L. Serum alkaline phosphatase was raised 496 IU/L (figure 3). Serum aspartate aminotransferase and alanine aminotransferase were within normal limits (43 and 50 µ/L). All other routine blood investigations were normal. Ultrasonography revealed a grossly dilated stomach containing echogenic material with extension into the first part of the duodenum. Contrast-enhanced CT abdomen showed a grossly distended stomach showing a hypodense, whorled lesion filling the lumen with extension beyond the first part of the duodenum suggestive of bezoar (figures 4 and 5). Upper gastrointestinal (UGI) endoscopy revealed trichobezoar occupying the whole of the stomach, extending from the gastro-oesophageal junction to the first part of the duodenum. The stomach was not distensible. D1 was full of trichobezoar and D2 could not be visualised (figures 6 and 7).

DIFFERENTIAL DIAGNOSIS

- ▶ Periapillary mass
- ▶ Pancreatic cancer
- ▶ Gastric tumour
- ▶ Gallbladder fossa mass
- ▶ Gastrointestinal stromal tumours
- ▶ Large choledochal cyst
- ▶ Pseudopancreatic cyst

CrossMark

Figure 1 Clinical picture showing lump abdomen.

To cite: Verma A, Sharma S, Tyagi G, et al. *BMJ Case Rep* Published online: [please include Day Month Year] doi:10.1136/bcr-2013-201667

Figure 3 Patient's photograph showing severe icterus.

TREATMENT

Considering a diagnosis of a huge trichobezoar obstructing the ampulla, a laparotomy was planned. The trichobezoar, which measured 28 × 20 cm, was taken out from the stomach after vertical gastrotomy (figures 8 and 9). The postoperative course was uneventful and the patient was discharged on the fifth post-operative day. She was referred to the psychiatric department for treatment, and was prescribed selective serotonin reuptake inhibitors (SSRIs) tab. Sertraline 50 mg twice daily initially for 1 month.

OUTCOME AND FOLLOW-UP

Six months after follow-up, the patient seems to be psychiatrically stable and was on antianxiety medication.

DISCUSSION

Trichobezoars are composed of hair or hair-like fibres and are commonly found in children or adolescent women younger than 30 years,¹ often with an underlying psychiatric or social problem. Clinical presentation of these patients may be confusing as often they are not forthcoming with a history of trichophagia either due to embarrassment or the unintentional nature of the problem. Most bezoars are located in the stomach, but they may be encountered in the oesophagus, small intestines and even rectum.² Bezoars can also be found distally in the gastrointestinal tract without continuity with the stomach bezoar due to breakage and distal propulsion. Trichobezoars continue to grow in size with continued ingestion of hair and this increases the risk of severe complications. The ingested hair

Figure 4 Contrast-enhanced CT abdomen showing mass in stomach suggesting bezoar.

Figure 5 Contrast-enhanced CT abdomen showing extension beyond the first part of the duodenum.

gets trapped in gastric mucosal folds and is typically black regardless of the colour of fibres ingested. Previous gastric surgery, diabetic gastroparesis, hypothyroidism and gut motility disorders have all been associated with trichobezoar formation. Most patients present with symptoms of vague epigastric discomfort. Other symptoms include nausea, vomiting, anorexia, early satiety and weight loss. The most common complications that are reported over the years include gastric mucosal erosion, ulceration and perforation of the stomach or small intestine, gastric outlet obstruction and intussusception. Obstructive jaundice, pancreatitis and even death, although rare, have also been reported.^{3–5}

In our case, irritation by the bezoar tail caused oedematous obstruction at the ampulla of Vater. It possibly explains the cause of the raised alkaline phosphatase and bilirubin levels. The supportive evidence can be derived from a case report of obstructive jaundice due to trichobezoar.³

Trichobezoars can be demonstrated on plain X-ray, barium films, ultrasonography and CT, but up to 75% of them may be missed on plain radiography. UGI endoscopy is considered as the gold standard investigation for the diagnosis. On endoscopy, the appearance is of a black, hard and concrete-like material.

Figure 6 Upper gastrointestinal endoscopy showing trichobezoar showing reflective material.

Figure 7 Upper gastrointestinal endoscopy showing trichobezoar going into the duodenum.

Management options include endoscopic removal, laparoscopic removal or laparotomy. Gorter *et al*, in a retrospective review of 108 cases of trichobezoar, evaluated the available management options; they noted that whereas 5% of attempted endoscopic removals were successful, 75% of attempted laparoscopies were successful. However, laparotomy was 100% successful and thus favoured as their management of choice.⁶ Surgery is usually required after failure of endoscopic therapy, or complications such as bleeding, perforation or obstruction. Among all bezoars, trichobezoars are the ones most likely to require surgical management.^{7, 8} The lack of invasiveness of these endoscopic techniques does not seem to outweigh the disadvantages and the complexity of surgical procedures.

In addition to the acute surgical treatment, psychiatric consultation is of paramount importance in order to prevent relapses.⁸ It is emphasised that the majority of these patients have an underlying psychiatric or social disorder. Our patient was diagnosed with post-traumatic stress disorder as a consequence of repeated abuse by her husband.

Patients with underlying psychiatric disorders require specific therapy which may include SSRIs, hypnosis or play therapies.⁹ Therefore, a multidisciplinary approach is essential to prevent recurrence of the problem.¹⁰

Figure 8 Trichobezoar being delivered out of gastrotomy.

Figure 9 Dimensions of trichobezoar.

Learning points

- ▶ In the presence of mental health problems, bezoars should always be kept in the differential diagnosis of patients with lump abdomen.
- ▶ Obstructive jaundice can be a rare presentation of bezoars.
- ▶ They can be easily treated with minimal morbidity if diagnosed and treated early.
- ▶ Upper gastrointestinal endoscopy is the gold standard investigation.
- ▶ Complete removal is a must and should be without spillage during extraction.
- ▶ Psychiatric evaluation and treatment is as important as any intervention for trichobezoar.

Acknowledgements The authors would like to thank Dr Prabha Om.

Contributors AV participated in the conception and design and also in the acquisition, analysis and interpretation of data. SSH and SSI participated in the conception and design and also in the drafting and critical revision of the article. GT participated in the drafting and critical revision of the article.

Competing interests None.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- 1 Debakey M, Oschner A. Bezoars: classification, pathophysiology and treatment. *Am J Gastroenterol* 1988;83:476–8.
- 2 Byrne WJ. Foreign bodies, bezoars and caustic ingestion. *Gastrointest Endos Clin North Am* 1994;4:99–104.
- 3 Schreiber H, Filston HC. Obstructive jaundice due to gastric trichobezoars. *J Pediatr Surg* 1976;11:103–14.
- 4 Shawis RN, Doig CM. Gastric trichobezoar associated with transient pancreatitis. *Arch Dis Child* 1986;59:994–5.
- 5 Deitrich NA, Gauf C. Postgastrectomy phytobezoars: endoscopic diagnosis and treatment. *Arch Surg* 1985;120:432–5.
- 6 Gorter RR, Kneepkens CMF, Mattens ECJL, *et al*. Management of trichobezoar: case report and literature review. *Pediatr Surg Int* 2010;26:457–63.
- 7 Phillips MR, Zaheer S, Drugas GT. Gastric trichobezoar: case report and literature review. *Mayo Clin Proc* 1998;73:653–6.
- 8 Siriwardana HPP, Ammori BJ. Laproscopic removal of a large gastric trichobezoar in a mentally retarded patient with pica. *Surg Endosc* 2003;17:834.
- 9 Christenson GA, Crow SJ. The characterization and treatment of trichotillomania. *J Clin Psychiatry* 1996;57:42–9.
- 10 Mewa Kinoo S, Singh B. Gastric trichobezoar: an enduring intrigue. *Case Rep Gastrointest Med* 2012;2012:136963.

Copyright 2014 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow