Published in final edited form as:

Eur J Appl Physiol. 2014 March ; 114(3): 561–578. doi:10.1007/s00421-013-2761-1.

Blood pressure regulation X: What happens when the muscle pump is lost? Post-exercise hypotension and syncope

John R. Halliwill, Dylan C. Sieck, Steven A. Romero, Tahisha M. Buck, and Matthew R. Ely
Department of Human Physiology, University of Oregon, Eugene, OR 97403–1240 USA

Abstract

Syncope which occurs suddenly in the setting of recovery from exercise, known as post-exercise syncope, represents a failure of integrative physiology during recovery from exercise. We estimate that between 50 and 80% of healthy individuals will develop pre-syncopal signs and symptoms if subjected to a 15-min head-up tilt following exercise. Post-exercise syncope is most often neurally mediated syncope during recovery from exercise, with a combination of factors associated with post-exercise hypotension and loss of the muscle pump contributing to the onset of the event. One can consider the initiating reduction in blood pressure as the tip of the proverbial iceberg. What is needed is a clear model of what lies under the surface; a model that puts the observational variations in context and provides a rational framework for developing strategic physical or pharmacological countermeasures to ultimately protect cerebral perfusion and avert loss of consciousness. This review summarizes the current mechanistic understanding of post-exercise syncope and attempts to categorize the variation of the physiological processes that arise in multiple exercise settings. Newer investigations into the basic integrative physiology of recovery from exercise provide insight into the mechanisms and potential interventions that could be developed as countermeasures against post-exercise syncope. While physical counter maneuvers designed to engage the muscle pump and augment venous return are often found to be beneficial in preventing a significant drop in blood pressure after exercise, countermeasures that target the respiratory pump and pharmacological countermeasures based on the involvement of histamine receptors show promise.

Keywords

Syncope; vasovagal; Post-exercise hypotension; Tilt-table test; Hypotension; Orthostatic; Regional Blood Flow; Exercise; Recovery; Receptors; Histamine; Baroreflex; Sympathetic Nervous System; Athletic Performance

INTRODUCTION

Post-exercise syncope can be defined as loss of consciousness or development of pre-syncopal signs and symptoms during recovery from a bout of physical activity or exercise. It is an alarming response to exercise which can occur in apparently healthy individuals, including athletes, or in individuals with autonomic disorders, and represents a failure of integrative physiology in the setting of recovery from exercise. It is associated with regulated reductions in blood pressure known as post-exercise hypotension, which can be

ADDRESS FOR CORRESPONDENCE: John R. Halliwill, PhD, 122 Esslinger Hall, 1240 University of Oregon, Eugene, OR 97403-1240, (541) 346-2841 fax, (541) 346-5425 phone, halliwil@uoregon.edu.

AUTHOR CONTRIBUTIONS

All authors contributed directly to the writing of this review. Each drafted a substantial section, assisted in revising the entire review, and approved the final version.

large enough in magnitude to become symptomatic (Kenny & Seals, 1993; Halliwill, 2001; Halliwill *et al.*, 2013). However, most reports of post-exercise syncope are likely to be incidents of neurally mediated syncope (formally called vasovagal syncope) (Freeman *et al.*, 2011) that have occurred during recovery from exercise, with the underlying changes associated with post-exercise hypotension contributing to the onset of the event (Kosinski *et al.*, 2000; Krediet *et al.*, 2004; O'Connor *et al.*, 2009).

Several case studies of individuals who have reportedly fainted following physical activity have been examined, yet the physiology remains poorly understood (see Table 1 below for an extensive listing). Important from these case studies is the recognition that cessation of exercise is a key step in the cascade leading to neurally mediated syncope in healthy individuals (Krediet *et al.*, 2004), whereas hypotension during exercise is more likely to occur in individuals with autonomic disorders (Low *et al.*, 2012). Characteristically, post-exercise syncopal events occur when an individual is standing motionless during the first 5 to 10 min after exercise (Krediet *et al.*, 2004), at time when the muscle pump (with its implicit role of maintaining central venous pressure during exercise in the upright position) is no longer engaged.

Prior work on “the postural problem” had suggested a key role of postural muscles (activated tonically to offset gravity and rhythmically during postural sway) in aiding venous return to the heart during quiet standing, as highlighted by Amberson (1943). However, it was classic studies in the 1940’s (Hellebrandt & Franseen, 1943; Pollack & Wood, 1949; Hellebrandt & Cary, 1949; Barcroft & Dornhorst, 1949) that coined the term “muscle pump” and demonstrated the profound power of skeletal muscle contractions to reduce both venous pressure and blood volume within the dependent exercising limbs, as well as the rapidity with which this beneficial effect of skeletal muscle contraction is lost when transitioning from exercise to quiet standing. Figure 1 illustrates some of these classic demonstrations. Indeed, this early work, as well as studies by Stegall (1966), show that rhythmic muscle contractions in the dependent limbs are capable of moving blood toward the heart against significant pressure gradients (i.e., greater than 90 mmHg) so long as venous valves are competent.

Thus, with quiet standing after exercise, one can anticipate a significant rise in dependent limb venous pressure and blood volumes, and a consequent fall in venous return to the heart, when this second generator of forward blood flow is switched off. This shift in blood volume, along with sustained post-exercise vasodilation and other responses associated with post-exercise hypotension, set the stage for syncope in individuals who attempt to remain stationary and upright. Along these lines, we estimate that between 50 and 80% of healthy individuals will develop pre-syncopal signs and symptoms if they are subjected to a head-up tilt for 15 minutes following a range of exercise activities (see table 2 below). This is markedly higher than the incidence of ~ 8% pre-syncope during similar head-up tilt in the general population (Kosinski & Grubb, 1994; Natale *et al.*, 1995)

This topic was last explored from a mechanistic or integrative physiology perspective a decade ago (Kosinski *et al.*, 2000; Halliwill, 2001; Krediet *et al.*, 2004), but has been reviewed from a clinical assessment perspective in more recent years (Link & Estes, 2007; O'Connor *et al.*, 2009; Asplund *et al.*, 2011; Hastings & Levine, 2012). We believe that recent investigations into the basic integrative physiology of recovery from exercise provide new insight into the mechanisms (Halliwill *et al.*, 2013) and potential interventions (Lacewell *et al.*, 2013) that could be developed as countermeasures against post-exercise syncope.

One can consider the reduction in blood pressure associated with post-exercise hypotension and syncope as the tip of the proverbial iceberg. What we have endeavored to create is a clear model of what lies under the surface; a model that puts the observational variations in context and provides a rational framework for developing strategic physical or pharmacological countermeasures. Part one of this review summarizes the current understanding of the mechanisms that contribute to post-exercise hypotension and syncope in humans. Part two of this review explores and attempts to categorize the variations on these physiological processes that arise in the settings of 1) moderate-intensity exercise, 2) high-intensity exercise, 3) exercise in the heat, 4) exercise at altitude, and 5) resistance exercise. Part three of this review explores and attempts to categorize targets of intervention including physical and pharmacological countermeasures against post-exercise syncope.

PART 1: REVIEW OF THE MECHANISMS UNDERLYING POST-EXERCISE HYPOTENSION AND SYNCOPE

The observation that “arterial pressure becomes depressed below normal resting pressure after severe muscular work” (Hill, 1898) is nearly as old as the sphygmomanometer itself. There are two distinct phenomena to consider in this context. Post-exercise hypotension represents a well-regulated reduction in arterial pressure lasting several hours following a single bout of exercise. It is well-regulated in that the hemodynamic state is stable, the reduction in pressure is similar in magnitude whether the individual is studied upright or supine, and it is driven by modest quantitative changes in autonomic control of arterial pressure. In contrast, post-exercise syncope represents a failure of the normal regulation of pressure and or cerebral perfusion such that symptoms develop which are syncopal in nature. It is an unstable hemodynamic state in which there is either a failure to compensate for symptomatic hypotension or in which a qualitative change in autonomic control of arterial pressure results in paradoxical vasodilatory and bradycardic responses. With a basic definition of these two phenomena, let us explore the known mechanisms which link them to recovery from exercise.

As recently reviewed (Halliwill *et al.*, 2013), the hemodynamic patterns most readily observed following exercise (e.g., reduced blood pressure, increased heart rate) are part of a larger pattern of hemodynamic responses and underlying mechanisms. Work over the last few years has generated a number of important mechanistic insights into post-exercise hypotension and identified the related phenomenon of sustained post-exercise vasodilation and its primary cause. During the exercise recovery period, the combination of centrally mediated decreases in sympathetic nerve activity, in addition to a reduced signal transduction from sympathetic nerve activation into vasoconstriction, and local vasodilator mechanisms contribute to the fall in arterial blood pressure seen after exercise, which in some cases may contribute to the development of pre-syncopal signs or symptoms.

Post-exercise baroreflex resetting and blunted transduction

After exercise, the arterial baroreflex is reset to defend lower blood pressures in humans (Halliwill *et al.*, 1996) and in rats (Kajekar *et al.*, 2002; Miki *et al.*, 2003), resulting in reduced sympathetic outflow (Floras *et al.*, 1989; Halliwill *et al.*, 1996; Kulics *et al.*, 1999; Kajekar *et al.*, 2002; Miki *et al.*, 2003). In this discussion, baroreflex resetting is used to indicate a shift in the operating point of the baroreceptor reflex in absence of any change in gain or sensitivity of the control of sympathetic outflow. In humans, the sensitivity of baroreflex control of sympathetic outflow is unaltered during post-exercise resetting (Halliwill *et al.*, 1996) but it appears blunted in rats (Kajekar *et al.*, 2002; Miki *et al.*, 2003). Baroreflex control of heart rate has proven to be more variable in its response to exercise, perhaps due to differences in methodology, idiosyncrasies resulting from the dual autonomic

control of the heart, or thermal influences on heart rate. The current model for the mechanism of post-exercise baroreflex resetting is based on studies conducted by Chen and colleagues (reviewed in Chen & Bonham, 2010), in which muscle afferents to the nucleus tractus solitarius (NTS) provide the key modulation of sympathetic output from the cardiovascular control centers in the medulla. In brief, muscle afferents activated in response to muscle contraction release substance-P at neurokinin-1 receptors on GABAergic interneurons in the caudal (NTS). These GABAergic interneurons inhibit 2nd order barosensitive neurons within the NTS which convey information from baroreceptor afferents to the caudal ventrolateral medulla. GABA reduces their excitability, resulting in less inhibition of sympathetic neurons in the rostral medulla, greater firing of sympathetic vasoconstrictor neurons during exercise, and resetting of the baroreflex to higher pressures during exercise. As exercise continues, neurokinin-1 receptors internalize on the GABA interneuron so that after exercise, neurokinin-1 receptors are less available for binding (Chen *et al.*, 2009). As a result, the GABAergic interneurons exert less inhibitory influence on the 2nd order barosensitive neurons, leading to an overall decrease in sympathetic outflow from the rostral ventrolateral medulla after exercise. When neurokinin-1 receptors are blocked prior to exercise, post-exercise hypotension is attenuated (Chen *et al.*, 2002). Likewise, GABA_A receptor antagonism in the rostral ventrolateral medulla prevents post-exercise hypotension (Kajekar *et al.*, 2002).

In young healthy humans exercised in thermoneutral conditions, neither the splanchnic nor renal vasculatures demonstrate changes in vascular conductance from resting pre-exercise levels to post-exercise recovery (although both vasoconstrict during exercise). This is an interesting contrast to the well known vasoconstriction that occurs in these vascular beds in response to hypotension in general, and has been interpreted to be another representation of the central resetting of the arterial baroreflex (Pricher *et al.*, 2004). Hence, these vascular beds do not directly contribute to post-exercise hypotension, yet they do not attenuate it. Likewise, cutaneous vascular conductance quickly returns to pre-exercise levels despite continued elevations in body core temperature (Wilkins *et al.*, 2004). This appears to be an expression of a shift in threshold of the thermoreflexes which allow core temperature to remain higher during recovery from exercise.

In addition to the sympathoinhibition that results from baroreflex resetting, post-exercise hypotension is associated with a blunted transduction of sympathetic outflow into vasoconstriction. For comparable levels of muscle sympathetic nerve activity, there is a reduced vascular resistance after exercise in the previously active skeletal muscles (Halliwill *et al.*, 1996). It is unlikely that a post-junctional mechanism is responsible for this blunted sensitivity to sympathetic outflow, as both α_1 - and α_2 -adrenergic responsiveness remain intact following exercise (Halliwill *et al.*, 2003). Presynaptic inhibition of norepinephrine release or increased reuptake of norepinephrine are possible explanations for the blunted transduction.

Post-exercise Vasodilation

There are two recognized vasodilatory phenomena during recovery from exercise: 1) immediate post-exercise hyperemia, and 2) sustained post-exercise vasodilation of the previously active skeletal muscle vascular beds (Laughlin *et al.*, 2012). The immediate post-exercise hyperemia can last from several seconds up to 20 min. The magnitude and duration of the hyperemia is dependent on the time, type, and intensity of exercise, but is not fully explained by mechanisms related to oxygen consumption within the previously active skeletal muscle (Morganroth *et al.*, 1975; Bangsbo & Hellsten, 1998). It has been suggested that the same vasodilator signals that drive exercise hyperemia contribute to immediate post-

exercise hyperemia (Bangsbo & Hellsten, 1998; Halliwill *et al.*, 2013), but there is no consensus on its cause which is likely to be multi-factorial.

In contrast to the short-lived immediate post-exercise hyperemia, sustained post-exercise vasodilation typically lasts upwards of two hours following moderate intensity aerobic exercise. It is dependent on the activation of histamine H₁- and H₂-receptors, as combined H₁- and H₂-receptor antagonism reduces post-exercise vasodilation by ~ 80% following 60 min of moderate-intensity cycle ergometry in both sedentary and endurance-trained athletes (Lockwood *et al.*, 2005; McCord & Halliwill, 2006; McCord *et al.*, 2006). Further, sustained post-exercise vasodilation following 60 min of moderate-intensity unilateral dynamic knee-extension exercise is abolished by H₁- and H₂-receptor antagonism (Barrett-O'Keefe *et al.*, 2013). Several possible mechanisms may increase intramuscular histamine during recovery from exercise. Mast cells located within the connective tissue layer surrounding skeletal muscle fascicles or near blood vessels may degranulate, releasing histamine locally (Metcalf *et al.*, 1997). Exercise-related factors that have been associated with mast cell degranulation in other contexts include reactive oxygen species (Son *et al.*, 2006), a variety of cytokines (and perhaps myokines), increases in temperature, and vibration (Grabbe, 2001). Alternatively, histamine can be formed *de novo* without storage in mast cells through histidine decarboxylase (Watanabe & Ohtsu, 2002). Along these lines, histidine decarboxylase mRNA expression (Endo *et al.*, 1998) and enzyme activity increase with prolonged exercise in mice (but not rats) (Graham *et al.*, 1964; Ayada *et al.*, 2000). There appear to be links between oxidative stress (Höcker *et al.*, 1998), hypoxia inducible factor-1 α (Jeong *et al.*, 2009), and histidine decarboxylase transcription. There is also older evidence that increased shear stress promotes histamine formation in large vessels such as the aorta (DeForrest & Hollis, 1978). Future studies are needed to define the cascade of events that result in post-exercise histamine receptor activation. In particular, studies must identify the exercise-related signal (e.g., oxidative stress, increased muscle temperature) that activates the histaminergic pathway, determine the source of histamine released by this signal (i.e., mast cell degranulation and/or *de novo* formation), and the timing or dose-response of this system in relation to exercise.

Obligatory versus Situational Influences

As a way to comprehend varied outcomes during recovery from exercise, we identified mechanisms that appear to be obligatory versus situational influences that vary from case to case. Along these lines, prolonged muscle afferent activation and consequent resetting of the baroreflex, resetting of thermoreflexes (Journeay *et al.*, 2006), pre-synaptic inhibition of sympathetic vasoconstrictor nerves, and activation of H₁- and H₂-receptors in the previously active muscle all appear to be obligatory components of post-exercise hypotension. Of these factors, histaminergic vasodilation may be the most important contributor to the fall in pressure in the normotensive and recreationally active population, as blockade of histamine receptors (Lockwood *et al.*, 2005; McCord & Halliwill, 2006; McCord *et al.*, 2006; Barrett-O'Keefe *et al.*, 2013), but not removal of the sympathetic component (Halliwill *et al.*, 2000), reduces post-exercise hypotension. Situational influences include the fluid status of the individual, heat balance with the environment, and the presence or absence of gravitational stress. We have explored how combinations of obligatory mechanisms and situational influences produce the integrated responses observed, as illustrated conceptually in Figure 2 (reproduced from Halliwill *et al.*, 2013).

The majority of studies report an elevated cardiac output concurrent with post-exercise hypotension, at least when subjects are studied in the supine position (Halliwill, 2001); however, this is not always the case. Reductions in cardiac output are commonly reported following exercise in endurance trained men even in the supine position (but not endurance

trained women, for unknown reasons)(Senitko *et al.*, 2002). Heart rate is consistently elevated following exercise, and this is likely the product of higher temperatures affecting the SA node pacemaker. This may confound many of the studies exploring baroreflex control of heart rate following exercise.

Stroke volume is generally well maintained, despite reductions in central venous pressure (Halliwill *et al.*, 2000). Stroke volume appears highly sensitive to both core temperature and fluid status but seems to be supported by enhanced contractility of unknown origin. Again, endurance trained men stand alone as a group that exhibits a reduced stroke volume following exercise (Senitko *et al.*, 2002; Lynn *et al.*, 2009). Ironically, it may be that the greater heat storage seen in less fit men (a response which allows for heat dissipation via physical mechanisms when physiological mechanisms are limited), leads to an enhanced cardiac contractility which is absent in highly fit men, and the absence of this effect leads to the marked fall in stroke volume in endurance trained men after exercise (Lynn *et al.*, 2009).

It is worth noting that the majority of studies reporting cardiac output and stroke volume have been performed with subjects in a supine position, and one would predict that upright recovery, via the increase in gravitational pooling of blood, would shift the system toward lower stroke volumes and cardiac outputs. While this is often observed (McCord *et al.*, 2008), there are also examples in which cardiac output remained surprisingly high during upright tilt after exercise (Lacewell *et al.*, 2013). These differences are likely related to the different exercise stimuli in these studies which may result in differing thermal, autonomic, and hormonal influences on the heart. Further, the impact of changes in plasma volume, which decreases ~ 10% in the upright position, may be markedly impacted by exercise or thermal stress, and may change in response to the vascular responses in liver, skin, and muscle vascular beds.

Despite all these potential variations in the underlying physiology, in most cases where measurements are made upright following moderate intensity exercise, post-exercise blood pressure remains lower than pre-exercise, yet is still well regulated in the upright position (Senitko *et al.*, 2002).

Post-exercise Syncope

Post-exercise syncope has an unknown rate of occurrence. There is likely an overlap between severe post-exercise hypotension, the development of post-exercise syncope, and the phenomenon of exercise-induced collapse, particularly when collapse occurs immediately after exercise and/or completion of athletic performance (e.g., crossing the finish line of a marathon)(Noakes, 1988; Holtzhausen *et al.*, 1994; Holtzhausen & Noakes, 1995, 1997; Kenefick & Sawka, 2007). While it is likely that the regulated reductions in blood pressure known as post-exercise hypotension can be large enough in magnitude as to become symptomatic under some circumstances, most reports of post-exercise syncope are likely incidences of neurally mediated syncope (formerly called vasovagal syncope) during recovery from exercise, with the underlying changes associated with post-exercise hypotension contributing to the onset of the event (Krediet *et al.*, 2004). Generally speaking, when standing upright following exercise, elevations in skeletal muscle blood flow combined with an inactive muscle pump can lead to marked reductions in venous return to the heart, a response which can be exacerbated in the heat (Krediet *et al.*, 2004; McCord *et al.*, 2008; Kenney *et al.*, 2013). By mechanisms which remain surprisingly murky (Noakes, 2003; Hainsworth, 2003; Joyner, 2009), the situation of high cardiac contractility and compromised venous return may trigger both paradoxical reductions in sympathetic vasoconstrictor nerve activity and increases in parasympathetic cardiac nerve activity. The mechanism or trigger for such neurally mediated syncope is often attributed to ventricular mechanoreceptors firing in response to a misinterpreted tissue distortion and producing a

response that is akin to the Bezold-Jarish reflex. Others have argued that the trigger is more likely to be generated within the central nervous system rather than within the ventricles (Hainsworth, 2003), and may involve serotonin receptors in the rostral ventrolateral medulla (Dean & Bago, 2002). This is a neglected topic of investigation. Regardless of where the origins of the response resides, the shift toward greater vasodilation (due to sympathetic withdrawal) and a relative or frank bradycardia leads to a rapid collapse of arterial pressure and cerebral perfusion.

It is possible that some cases of post-exercise syncope are related to alterations in cerebral blood flow independent of vasovagal reactions or frank hypotension. For example, hypocapnia induced by hyperventilation during recovery from exercise could produce a cerebral vasoconstriction and diminish cerebral oxygen delivery in the absence of any change in perfusion pressure. Along these lines, Rasmussen et al. (2006) showed that cerebral CO₂ vascular reactivity following strenuous exercise is increased such that arterial CO₂ becomes the primary factor influencing changes in cerebral blood flow during and after prolonged exercise.

The potential for changes in cerebral autoregulation following exercise has also been explored. While cerebral autoregulation is well-maintained following moderate intensity exercise (Ogoh *et al.*, 2007; Murrell *et al.*, 2007; Willie *et al.*, 2013), two studies suggest that more intense exercise may produce a lasting cerebral autoregulatory deficit. Specifically, Ogoh *et al.* (2005) found reduced dynamic cerebral autoregulation during heavy cycle exercise and Bailey *et al.* (2011) found reduced dynamic cerebral autoregulation during recovery from a recumbent cycle peak test. Thus, it seems likely that cerebral perfusion is less protected from sudden onset hypotension following more intense exercise.

Table 1 summarizes case reports and case series of post-exercise syncope in which individual patient information on age and sex were provided. It is worth noting that many cases include asystole, evidence of the vagal component of neurally mediated syncope. However, depending on the timing of events and the progression toward symptoms, bradycardia is not always evident.

In reviewing the case reports and case series, some interesting patterns are evident regarding age and sex of individual patients, as indicated in Figure 3. First, in contrast to conventional wisdom, the majority of cases of post-exercise syncope occur in men. In general, women have decreased orthostatic tolerance compared to men, and this holds true in response to heat stress (Meendering *et al.*, 2005), but not recovery from exercise (Lacewell *et al.*, 2013). It is reasonable to suggest that the higher number of cases in men is a spurious result, perhaps due to greater historical participation in physical activities, but the pattern holds true in several prospective studies (Fleg & Lakatta, 1986; Lacewell *et al.*, 2013). Second, the cases represent a broad range of ages, with nearly every decade of life represented with a higher proportion of cases below the age of 50.

There are several other notable case series related to post-exercise syncope. First, as part of the Baltimore Longitudinal Study on Aging, Fleg & Lakatta (1986) analyzed post-exercise blood pressure responses in a large cohort of healthy volunteers. During seated recovery from a graded maximal treadmill test, 3.1% of subjects younger than 55 yrs exhibited a marked hypotension that was associated with pre-syncope symptoms. The incidence was lower in women than men, and was only 0.3% for subjects older than 55 yrs. Long-term investigation of subjects who became symptomatically hypotensive following exercise did not indicate any increased health risks, supporting the notion that the prognosis related to post-exercise syncope is generally benign. Second, Sakaguchi *et al.* (1995) reported on a group of 12 patients who were referred for exercise-related syncope. They noted a general

susceptibility to neurally mediated syncope in these individuals, fostering the concept that most cases of syncope following exercise are incidences of neurally mediated syncope. Third, Holtzhausen et al. (1994) reported on 46 male athletes who collapsed during or after a 56-km ultramarathon, finding that 85% of these cases occurred after the finish line (i.e., post-exercise), and that hydration status (a surrogate for circulating blood volume) and core temperature in those who collapsed post-race are no different than control athletes who do not collapse. In other words, traditional contributing factors such as dehydration, hypovolemia, and hyperthermia are not necessary for post-exercise syncope following longer duration athletic events. Finally, an interesting report by Eckart et al. (2010) suggests that the use of ergogenic supplements (e.g., stimulants such as ephedra and ephedra-substitutes) can increase the risk of post-exercise syncope three-fold.

While the long-term prognosis related to post-exercise syncope is generally benign, it is important that other more serious causes be ruled out by thorough clinical evaluation, as syncope after exercise can be a sign of a significant pathophysiology (Kramer *et al.*, 1988; Smith *et al.*, 1993; Krediet *et al.*, 2004; O'Connor *et al.*, 2009; Hastings & Levine, 2012).

PART 2: EXPLORATION OF DIFFERENT EXERCISE MODELS THAT GENERATE POST-EXERCISE SYNCOPES

Much of our understanding of the mechanisms of post-exercise hypotension come from a single model of exercise, namely, moderate-intensity exercise of medium-duration (e.g., up to 60 min) performed in a thermoneutral environment. However, the cases of post-exercise syncope do not fall neatly into this model, but come from a range of exercise activities which may differentially affect arterial pressure regulation or control of cerebral oxygen delivery. Let us explore five archetypal exercise models, with the understanding that when individuals exercise, it may be on a spectrum that combines elements of these five models. Table 2 summarizes prospective studies on post-exercise syncope, as they relate to these models, in which pre-syncopal rates during head-up tilt (or tolerance to lower body negative pressure) were assessed during recovery from exercise.

1) Moderate-intensity exercise

In young healthy normotensive individuals, moderate-intensity exercise for 30–60 min produces modest post-exercise hypotension via the mechanisms described above, but is generally insufficient to elicit post-exercise syncope during seated recovery or brief (5 min) head-up tilt when performed in a thermoneutral environment (Raine *et al.*, 2001; Senitko *et al.*, 2002). However, the incidence of orthostatic intolerance can be surprisingly high in response to even mild exercise when orthostatic stress is more prolonged (15 min) (Mayerson, 1944). Further, when moderate-intensity exercise is of extended duration, or under moderately warm conditions, the pre-syncopal incidence can reach as high as 54 to 81% (Eichna *et al.*, 1947a; Murrell *et al.*, 2009, 2011), as shown in Table 2.

This model of post-exercise syncope is ostensibly explained by the effects of the sustained post-exercise vasodilation on reducing total peripheral resistance and reducing venous return to the heart. Noakes and colleagues (Noakes, 1988, 2007; Holtzhausen & Noakes, 1997) have argued that hyperthermia and dehydration play little or no role in the collapse of athletes at the end of marathons, whereas sustained peripheral vasodilation and loss of the muscle pump appear sufficient to explain many cases of post-exercise syncope following moderate-intensity exercise of extended duration. It is worth noting that the early work by Eichna (1947a) found the increased incidence of pre-syncope during head-up tilt persists for 1 to 2 hours following exercise. This is a timeline that is similar to the duration of the

sustained post-exercise vasodilation that is mediated by histamine-receptor activation (Emhoff *et al.*, 2011).

2) High-intensity exercise

In contrast to moderate-intensity exercise, there are fewer studies investigating post-exercise hypotension and related phenomenon following short-duration high-intensity exercise. However, there is anecdotal evidence that the Wingate test of anaerobic power, a staple of undergraduate exercise physiology labs, results in pre-syncopal symptoms in many individuals. Likewise, the classic study by Eichna *et al.* (1947a) found that more than half of healthy individuals will become pre-syncopal during a 5-min head-up tilt after completion of maximal exercise. More recently, Lacewell *et al.* (2013) developed a modified Wingate test which produces post-exercise syncope in many individuals. Such maximal or supra-maximal efforts when followed by head-up tilt or active standing elicit pre-syncopal responses in 58 to 100% of individuals, as shown in Table 2.

What is interesting in the study by Lacewell *et al.* (2013) is that arterial pressure was often well-maintained at or above pre-exercise levels until a sudden shift to hypotension and development of pre-syncopal symptoms. In other words, arterial pressure did not demonstrate post-exercise hypotension prior to post-exercise syncope, rather pressure was well-maintained (until it was not).

One of the characteristics of recovery from high-intensity exercise is marked hyperventilation which can reduce arterial CO₂ to values that would likely induce prominent cerebral vasoconstriction. For example, after a modified Wingate test, end-tidal CO₂ decreased 14 mmHg to levels that would be associated with a ~ 40% reduction in cerebral perfusion (Lacewell *et al.*, 2013). There are cases of athletes developing syncope as a result of brief vigorous hyperventilation (Buja *et al.*, 1989).

The potential role of autoregulation in this model also merits discussion. While cerebral autoregulation is well-maintained following moderate intensity exercise (Ogoh *et al.*, 2007; Murrell *et al.*, 2007; Willie *et al.*, 2013), two studies suggest that more intense exercise may produce a lasting cerebral autoregulatory deficit. Specifically, Ogoh *et al.* (2005) found reduced dynamic cerebral autoregulation (as assessed by transfer function) during heavy cycle exercise and Bailey *et al.* (2011) found reduced dynamic cerebral autoregulation (as assessed by thigh-cuff release) during recovery from a recumbent cycle peak test. Thus, it seems likely that cerebral perfusion is less protected from sudden onset hypotension following high-intensity exercise.

3) Exercise in the heat

Exercise in the heat adds another level of complexity to these integrated responses. This model is exemplified by moderate-intensity exercise of moderate to long duration that is performed in a hot environment, which could be either arid or humid. Thus, the situational component that adds complexity to this model is the greater heat load, secondary to the reduced ability to dissipate metabolic heat to the environment. When individuals are exercising hard enough or long enough to generate significant increases in core temperature, then the physiological accommodations to heat stress such as cutaneous vasodilation and sweating can lead to reductions in central venous pressure, dehydration, and hypovolemia.

As recently reviewed by Kenney *et al.* (2013), there is a commensal use of cardiac output to serve both the metabolic demands of the exercising muscle and thermoregulatory need for heat dissipation (Rowell, 1977). While skin blood flow may increase to more than 7 L/min (more than half of available cardiac output) during passive heat stress (Rowell, 1986;

Crandall *et al.*, 2010), in normal exercising individuals the blood flow needed to maintain heat dissipation and muscular activity is not a great enough demand to limit either of these vasculatures or to compromise blood pressure regulation (Kenney *et al.*, 2013). That said, the non-competing division of cardiac output may deteriorate into competition for limited cardiac output at the end of exercise (when the muscle pump is lost), but some surprising observations have been made.

While the greater fluid loss associated with exercise in the heat can be predicted to decrease cardiac preload, Lynn *et al.* (2009) showed that greater elevations in heart rate following exercise in the heat are associated with an augmentation of cardiac output that is as great as what can be produced by fluid replacement, at least when subjects are supine. However, it is clear that the skin, when vasodilated in response to passive heat stress, is insufficiently vasoconstricted with orthostatic stress to support cardiac preload and, therefore, predisposes individuals to the development of syncope in the heat (Crandall, 2008; Crandall *et al.*, 2010; Pearson *et al.*, 2013).

Following exercise, other vascular beds likely vasoconstrict to maintain pressure in response to orthostatic challenge. While the arterial baroreflex is reset to maintain arterial pressure at lower levels following exercise, sensitivity of baroreflex control of sympathetic outflow is augmented after exercise in the heat, possibly due to greater reductions in central venous pressure that result in decreased cardiopulmonary baroreceptor activation (Charkoudian *et al.*, 2003). However, it is likely that the vasoconstrictor reserve is reached for important vasculatures such as the splanchnic region during this combination of stressors (Crandall, 2008; Crandall *et al.*, 2010; Pearson *et al.*, 2013), leaving little if any margin for maintaining cerebral perfusion in the upright position. Further, there is some evidence that exercise in the heat may produce a lasting deficit in the cerebral autoregulation which is observable after the return to resting body temperatures (Carter *et al.*, 2006). Thus, it is not surprising that classic studies documented a decreased orthostatic tolerance following exercise in the heat (Greenleaf *et al.*, 1974; Shvartz *et al.*, 1977), and brief high-intensity (Greenleaf *et al.*, 1974), prolonged low-intensity (Shvartz *et al.*, 1977), and medium-duration moderate-intensity (McCord *et al.*, 2008) exercise in the heat have all been used as prospective models for studying post-exercise syncope.

4) Exercise at altitude

Anecdotes of altitude-related syncope both during and after exercise abound, but few (if any) have been documented in the literature (Hultgren, 1997). That said, it is well established that neurally mediated syncope can be produced in most individuals by having them breathe low O₂ levels (< 8 %), and that some individuals will become syncopal while breathing moderately hypoxic O₂ mixtures (13 – 14 % O₂) similar to the O₂ levels at altitudes of 3000–4000 m. At these altitudes, orthostatic tolerance is reduced (Halliwill, 2003).

Early hemodynamic studies highlighted a potential role of exaggerated circulating epinephrine levels in precipitating hypoxic syncope (Halliwill, 2003; Halliwill & Minson, 2005). Dinunno *et al.* (2003) were able to obtain an arterial blood sample from a hypoxic individual at the onset of hypotension and bradycardia and found 4-fold higher epinephrine than other subjects exposed to the same degree of hypoxia. This observation is consistent with the notion of exaggerated circulating epinephrine in hypoxic syncope. In this individual, high epinephrine was associated with progressive skeletal muscle vasodilation, but part of the hypotension may have been linked to hypoxic vasodilation of the splanchnic circulation (Rowell & Blackmon, 1989; Westendorp *et al.*, 1997). One case report suggests that β -adrenergic blockade may prevent hypoxic syncope (Freitas *et al.*, 1996).

With this background on hypoxic syncope, one can surmise that post-exercise syncope at altitude is multi-factorial, combining the elements of the other exercise models with epinephrine-induced vasodilation and a possible splanchnic steal of blood flow. These factors, which acting alone can compromise cerebral perfusion, would act in combination with reduced arterial oxygen content to diminish cerebral oxygenation.

5) Resistance exercise

While most of the focus of this review is on the response to dynamic whole-body exercise, it is worth mentioning that resistance exercise is also associated with post-exercise syncopal episodes. Often referred to as “weightlifter’s blackout,” post-resistance exercise syncope occurs during exercise or within seconds of completion (Compton *et al.*, 1973).

Unlike the moderate blood pressure increase that occurs during aerobic exercise, arterial blood pressure is profoundly elevated during resistance exercise. Arterial blood pressure has been reported to reach 320/250 mmHg during maximal double-leg press (Macdougall *et al.*, 1985), although the rise in pressure during submaximal resistance exercise does not approach these levels (Edwards *et al.*, 2002; Romero & Cooke, 2007). In the face of increased perfusion pressure, cerebral perfusion (measured via transcranial Doppler) is unchanged (Edwards *et al.*, 2002) or slightly elevated (Romero & Cooke, 2007) during moderate intensity resistance exercise, but is reduced during high intensity resistance exercise (Dickerman *et al.*, 2000; Perry *et al.*, 2013) perhaps as protection against hypertension (Tzeng & Ainslie, 2013).

Subsequent to the marked elevation in blood pressure during resistance exercise, pressure falls rapidly to below pre-exercise levels upon the completion of the lift (Macdougall *et al.*, 1985; Edwards *et al.*, 2002; Pott *et al.*, 2003; Romero & Cooke, 2007). As a result, cerebral perfusion falls below pre-exercise levels immediately following exercise (Edwards *et al.*, 2002; Romero & Cooke, 2007). The precipitous fall in arterial blood pressure immediately following resistance exercise may be a sufficient challenge to the cerebrovascular regulatory mechanisms to produce syncope (i.e., the large and rapid “swing” in pressure may be beyond the upper and lower limits of cerebral autoregulation)(Macdougall *et al.*, 1985; Dickerman *et al.*, 2000).

Several additional mechanisms may exacerbate this state and contribute to post-resistance exercise syncope. First, voluntary or involuntary hyperventilation before or during resistance exercise may induce cerebral vasoconstriction (Compton *et al.*, 1973; Romero & Cooke, 2007), similar to what we have discussed above for high-intensity exercise. Second, Valsalva straining during lifting may produce sudden blood pressure transitions both during and immediately after straining (Compton *et al.*, 1973). Third, dehydration is a common practice among competitive weightlifters and other weight-class athletes (e.g., wrestling and boxing), and the reduced circulating blood volume associated with hypohydration can further reduce tolerance to post-exercise orthostatic challenges (Hand, 1997; Moralez *et al.*, 2012).

In summary, one can consider these five archetypal exercise models as situational variations on the underlying physiology that may increase or decrease the predisposition toward post-exercise syncopal events. Some of these models are well developed, but some remain poorly investigated.

PART 3: CLASSIFICATIONS OF POTENTIAL COUNTERMEASURES

The majority of evidence indicates that post-exercise syncope, in the absence of other medical conditions, is a nuisance. One to three percent of emergency department visits in the

United States, with an estimated cost of \$5400 per patient and totaling \$2.5 billion a year, are due to syncope (Sun *et al.*, 2005). It is important to recognize that post-exercise syncope can be situationally life-threatening (e.g., cycling, climbing, firefighting, and military operations), and several case reports on post-exercise syncope involve firefighters (Osswald *et al.*, 1994; van den Berg *et al.*, 1995; Krediet *et al.*, 2006). Thus, the development of countermeasures which can protect against post-exercise syncope should focus on ways to protect the individual from loss of consciousness during recovery from exercise. The current arsenal of countermeasures falls into three basic categories: behavioral, pharmacological, and pacemakers. The evidence for these existing countermeasures is almost exclusively anecdotal. Table 3 summarizes case reports of post-exercise syncope which included an intervention or treatment and follow-up.

To the best of our knowledge, there is one randomized controlled trial for post-collapse treatment (but not prevention) (Anley *et al.*, 2011), one randomized controlled trial for prevention in athletes with post-exercise syncope (Privett *et al.*, 2010b), and two randomized controlled trials for prevention in models of post-exercise syncope performed in healthy subjects (McCord *et al.*, 2008; Lacewell *et al.*, 2013), as summarized in Table 4 and Figure 4.

Behavioral

It is disappointing that one of the most common medical recommendations to patients with post-exercise syncope is to simply avoid vigorous exercise. Considering the numerous health benefits of routine vigorous exercise, this is not an acceptable remedy. Rather, avoidance of exercise should be seen as a last resort when all other countermeasures have failed to protect the individual.

In contrast to preaching avoidance, a number of easily learned physical countermeasures designed to engage the muscle pump and augment venous return are often found to be beneficial in preventing a significant drop in blood pressure after exercise. A classic study by Eichna *et al.* (1947a) showed that alternate movement of the legs during orthostatic hypotension following exhaustive physical work in young, healthy soldiers raised blood pressure markedly. Likewise, Krediet *et al.* (2005) demonstrated that a simple squatting maneuver could ameliorate pre-syncopal symptoms and normalize blood pressure in a patient in whom neurally mediated syncope could be consistently triggered by moderate cycle exercise. One can consider such countermeasures, along with the advice to “keep moving” after exercise, as a simple and effective means to avoid many cases of post-exercise syncope, as active recovery has been reported to be beneficial in some cases (Tamura *et al.*, 1990; Krediet *et al.*, 2006). However, there are likely limitations inherent in asking an exhausted athlete to rely on their fatigued muscles to protect them from post-race collapse, and it is likely that the finish-line chute standstill has contributed to some cases of syncope (Holtzhausen & Noakes, 1995; Anon, 2004). If active recovery is not practical, then use of the Trendelenberg position (or simply lifting the legs of the supine individual) may be an effective way to rectify venous return to the heart (Anley *et al.*, 2011).

Lacewell *et al.* (2013) recently demonstrated the benefits of using an inspiratory resistance device to alleviate post-exercise syncope following high-intensity exercise, as shown in Figure 4B. Originally developed for the treatment of severe hemorrhage and shock, an impedance threshold device generates resistance during inspiration which may augment the function of the respiratory pump in mobilizing venous blood toward the heart. However, it is possible that the main benefit of the device following high-intensity exercise is the attenuation of hypocapnic cerebral vasoconstriction or enhancement of a cerebral siphon effect.

Another behavioral approach with promise is the use of water ingestion (Thijs *et al.*, 2003), not in the prevention of dehydration, but owing to the pressor response elicited by ingestion of modest to large volumes of hypotonic fluid. In other studies, fluid ingestion to replace that lost during exercise has been shown to reduce cardiovascular strain during lower body negative pressure (Davis & Fortney, 1997). However, when treating post-competition collapse following endurance events Anley *et al.* (2011) found that use of the Trendelenberg position was as effective as intravenous fluids.

Lastly, compression garments can be of benefit in reducing the likelihood of post-exercise syncope (Privett *et al.*, 2010b). Use of compression garments of various types are seeing more widespread use by the athletic community, so routine use of such garments in an individual with recurring symptoms during recovery from exercise may be an appealing alternative to some of the more invasive countermeasures described below.

Pharmacological

It is clear that many of the clinical approaches that are employed to treat patients with post-exercise syncope has been adapted from those tools that have also seen routine use in treating neurally mediated syncope. Thus, the use of α -adrenergic blocking drugs is the most common pharmacological countermeasure (see Table 3), followed by the antiarrhythmic agent disopyramide (which is used in part because of its vagolytic effects). These standard pharmacological approaches remain controversial in treatment of neurally mediated syncope, where anecdotal evidence and case reports are promising but evidence from controlled trials is often inconclusive (Kaufmann & Freeman, 2004).

It is important to recognize that none of the common pharmacological approaches, such as α -adrenergic blocking drugs, act directly on the underlying physiology related to post-exercise hypotension or sustained post-exercise vasodilation (except perhaps in the setting of altitude, where circulating epinephrine is an issue). However, McCord *et al.* (2008), tested the effectiveness of H_1 -receptor antagonism in preventing post-exercise syncope induced by head-up tilt following exercise in the heat. They found the magnitude of the arterial pressure drop during head-up tilt was blunted by administration of an H_1 -receptor antagonist, and subjects were able to stand longer before the development of pre-syncope signs and symptoms, as shown in Figure 4A. In essence, this is the first pharmacological countermeasure which directly targets the underlying physiology of recovery from exercise.

Pacemakers

In a number of cases, pacemakers have been implanted in patients with post-exercise syncope. Like the use of α -adrenergic blocking drugs, this is a countermeasure adapted from the clinical toolbox routinely used to deal with neurally mediated syncope. While a number of case reports (see Table 3) suggest that pacemakers are effective, such anecdotes are far from definitive and the use of pacemakers for the treatment of neurally mediated syncope remains controversial (Petersen & Sutton, 1997; Romme *et al.*, 2011). As reviewed here, knowledge of the physiology of post-exercise hypotension, sustained post-exercise vasodilation, and post-exercise syncope do not provide a rationale for the use of implanted pacemakers, when other less-invasive countermeasures are likely to be beneficial.

Other directions

We believe there are other potential directions for interventions, and that a systematic view of the underlying physiology of post-exercise hypotension and syncope can be used as a roadmap. Specifically, each of the five archetypal exercise models may respond to targeted interventions with greater efficacy than a “one size fits all” approach, but some interventions may prove universally beneficial, but perhaps to lesser effect. The ideal solution, if it exists,

would be a non-invasive countermeasure that does not require the individual to medicate daily, and that can be deployed “as needed” without the need to carry equipment or wear special clothing.

One approach would be countermeasures that increase total peripheral resistance. In hot conditions, rapid skin cooling may prove an effective means of reducing cutaneous vascular conductance and improving hemodynamic stability. Along these lines, Wilson *et al.* (2002) demonstrated that, in a model of passive heat stress, rapid skin cooling protected arterial pressure, cerebral perfusion, and decreased pre-syncope symptoms during head-up tilt. In other situations, sympathoexcitatory maneuvers such as bolus fluid ingestion of water or hand immersion in ice water to elicit a cold pressor response might be sufficient to increase total peripheral resistance and provide benefit. Pharmacological interventions that target specific vascular beds such as the highly compliant splanchnic circulation may prove effective in enhancing total peripheral resistance (Jarvis *et al.*, 2012), and could be very advantageous for exercise at altitude.

Another approach would be countermeasures designed to augment cardiac preload, beyond administering fluids and engaging the muscle pump (Tamura *et al.*, 1990; Krediet *et al.*, 2006). Augmentation of the respiratory pump via inspiratory resistance shows some promise in this area (Lacewell *et al.*, 2013), and it is possible that some of the sympathoexcitatory maneuvers or rapid skin cooling also function to augment cardiac preload. External compression by medical compression garments or other means may be viable, but perhaps less than ideal.

The least explored approach would be countermeasures designed to protect cerebral blood flow directly, by either causing cerebral vasodilation or boosting cerebral perfusion pressure gradients. Along these lines, when hypocapnia is a likely contributing factor (high-intensity exercise or resistance exercise), it is possible that something as simple as rebreathing from a paper bag might help protect cerebral blood flow by maintaining higher arterial CO₂ levels. To the extent that the cerebral siphon effect is present in upright humans (which remains highly debated), breathing maneuvers which engage the respiratory pump may prove effective in augmenting the cerebral perfusion pressure gradient (as discussed in Lacewell *et al.*, 2013). Lastly, supplemental oxygen could also be beneficial, but it seems to be seldom deployed during recovery from exercise aside from NFL games played at modest altitudes (~1610m).

One interesting potential countermeasure that may need to be re-explored is the role of acclimatization to heat. Early work by Greenleaf *et al.* (1974) found that orthostatic tolerance following exercise in the heat was, surprisingly, made worse by prior heat acclimatization. It is unclear how acclimatization, which greatly improves the ability to exercise in the heat and is associated with plasma volume expansion, would prove to be deleterious in the setting of post-exercise syncope.

SUMMARY AND CONCLUSIONS

Blood pressure regulation after exercise is about much more than a simple loss of the muscle pump. Newer investigations on the recovery from exercise provide insight into the mechanisms of baroreflex resetting, the sustained histamine-mediated vasodilation, and changes in cerebral autoregulation. Thus, obligatory alterations in the control of sympathetic vascular tone and skeletal muscle blood flow interact with situational effects like cutaneous vasodilation in the heat, dehydration, hypoxic vasodilation, and hypocapnic cerebral vasoconstriction to produce the post-exercise hypotension, and in some cases post-exercise syncope. Mining of case and prospective studies hints at additional influences regarding who

is likely to suffer from pre-syncope symptoms after exercise and under what circumstances. Together, this information creates a framework for challenging some of the traditional treatment strategies and for developing novel countermeasures against post-exercise syncope.

Acknowledgments

This research was supported in part by grants from the American Heart Association, Grant-in-Aid 11GRNT5490000, and National Institutes of Health Grant HL115027.

References

- Amberson WR. Physiological adjustments to the standing position. *Univ Md Sch Med Bull.* 1943; 27:127–145.
- Anley C, Noakes TD, Collins M, Schwellnus MP. A comparison of two treatment protocols in the management of exercise-associated postural hypotension: a randomised clinical trial. *Br J Sports Med.* 2011; 45:1113–1118. [PubMed: 20584756]
- Runner's World. 2004 Mar. A knockout finish.
- Arad M, Solomon A, Roth A, Atsmon J, Rabinowitz B. Postexercise syncope: evidence for increased activity of the sympathetic nervous system. *Cardiology.* 1993; 83:121–123. [PubMed: 8261477]
- Asplund CA, O'Connor FG, Noakes TD. Exercise-associated collapse: an evidence-based review and primer for clinicians. *Br J Sports Med.* 2011; 45:1157–1162. [PubMed: 21948122]
- Ayada K, Watanabe M, Endo Y. Elevation of histidine decarboxylase activity in skeletal muscles and stomach in mice by stress and exercise. *Am J Physiol Regul Integr Comp Physiol.* 2000; 279:R2042–7. [PubMed: 11080067]
- Bailey DM, Evans KA, McEneny J, Young IS, Hullin DA, James PE, Ogoh S, Ainslie PN, Lucchesi C, Rockenbauer A, Culcasi M, Pietri S. Exercise-induced oxidative-nitrosative stress is associated with impaired dynamic cerebral autoregulation and blood-brain barrier leakage. *Exp Physiol.* 2011; 96:1196–1207. [PubMed: 21841038]
- Bangsbo J, Hellsten Y. Muscle blood flow and oxygen uptake in recovery from exercise. *Acta Physiol Scand.* 1998; 162:305–312. [PubMed: 9578376]
- Barcroft H, Dornhorst AC. Demonstration of the muscle pump in the human leg. *J Physiol.* 1949; 108:39P.
- Barrett-O'Keefe Z, Kaplon RE, Halliwill JR. Sustained postexercise vasodilatation and histamine receptor activation following small muscle-mass exercise in humans. *Exp Physiol.* 2013; 98:268–277. [PubMed: 22848080]
- Van den Berg MP, Crijns HJ, Bouwmeester TR, Smit AJ, Lie KI. Cardiac asystole post-exercise: a report of two cases. *Int J Cardiol.* 1995; 51:296–300. [PubMed: 8586479]
- Bjurstedt H, Rosenhamer G, Balldin U, Katkov V. Orthostatic reactions during recovery from exhaustive exercise of short duration. *Acta Physiol Scand.* 1983; 119:25–31. [PubMed: 6650203]
- Brogdon E, Hellebrandt FA. Post-exercise orthostatic collapse. *Am J Physiol.* 1940; 129:P318.
- Buja G, Folino AF, Bittante M, Canciani B, Martini B, Miorelli M, Tognin D, Corrado D, Nava A. Asystole with syncope secondary to hyperventilation in three young athletes. *PACE.* 1989; 12:406–412. [PubMed: 2466265]
- Carter RI, Cheuvront SN, Vernieuw CR, Sawka MN. Hypohydration and prior heat stress exacerbates decreases in cerebral blood flow velocity during standing. *J Appl Physiol.* 2006; 101:1744–1750. [PubMed: 16916922]
- Charkoudian N, Halliwill JR, Morgan BJ, Eisenach JH, Joyner MJ. Influences of hydration on post-exercise cardiovascular control in humans. *J Physiol.* 2003; 552:635–644. [PubMed: 14561843]
- Chen CY, Bechtold AG, Tabor J, Bonham AC. Exercise reduces GABA synaptic input onto nucleus tractus solitarius baroreceptor second-order neurons via NK1 receptor internalization in spontaneously hypertensive rats. *J Neurosci.* 2009; 29:2754–2761. [PubMed: 19261870]
- Chen CY, Bonham AC. Postexercise hypotension: central mechanisms. *Exerc Sport Sci Rev.* 2010; 38:122–127. [PubMed: 20577060]

- Chen CY, Munch PA, Quail AW, Bonham AC. Postexercise hypotension in conscious SHR is attenuated by blockade of substance P receptors in NTS. *Am J Physiol Heart Circ Physiol.* 2002; 283:H1856–62. [PubMed: 12384463]
- Compton D, Hill PM, Sinclair JD. Weight-lifters' blackout. *Lancet.* 1973; 2:1234–1237. [PubMed: 4128562]
- Crandall CG. Heat stress and baroreflex regulation of blood pressure. *Med Sci Sports Exerc.* 2008; 40:2063–2070. [PubMed: 18981943]
- Crandall CG, Shibasaki M, Wilson TE. Insufficient cutaneous vasoconstriction leading up to and during syncopal symptoms in the heat stressed human. *Am J Physiol Heart Circ Physiol.* 2010; 299:H1168–H1173. [PubMed: 20693394]
- Crisafulli A, Melis F, Orrù V, Lener R, Lai C, Concu A. Hemodynamic during a postexertional asystolia in a healthy athlete: a case study. *Med Sci Sports Exerc.* 2000; 32:4–9. [PubMed: 10647522]
- Davis JE, Fortney SM. Effect of fluid ingestion on orthostatic responses following acute exercise. *Int J Sports Med.* 1997; 18:174–178. [PubMed: 9187970]
- Dean C, Bago M. Renal sympathoinhibition mediated by 5-HT_{1A} receptors in the RVLM during severe hemorrhage in rats. *Am J Physiol Regul Integr Comp Physiol.* 2002; 282:R122–R130. [PubMed: 11742830]
- DeForrest JM, Hollis TM. Shear stress and aortic histamine synthesis. *Am J Physiol Heart Circ Physiol.* 1978; 3:H701–H705.
- Dickerman RD, McConathy WJ, Smith GH, East JW, Rudder L. Middle cerebral artery blood flow velocity in elite power athletes during maximal weight-lifting. *Neurol Res.* 2000; 22:337–340. [PubMed: 10874679]
- Dinunno FA, Joyner MJ, Halliwill JR. Failure of systemic hypoxia to blunt alpha-adrenergic vasoconstriction in the human forearm. *J Physiol.* 2003; 549:985–994. [PubMed: 12730336]
- Dockery BK, Newman KP. Exercise-induced asystole with syncope in a healthy young man. *Am J Med Sci.* 2007; 334:145–148. [PubMed: 17700209]
- Eckart RE, Gentlesk PJ, Shry EA. Differential manifestation of cardiovascular complaints as a function of utilization of ergogenic supplements. *PACE.* 2010; 33:286–289. [PubMed: 20015135]
- Edwards MR, Martin DH, Hughson RL. Cerebral hemodynamics and resistance exercise. *Med Sci Sports Exerc.* 2002; 34:1207–1211. [PubMed: 12131264]
- Eichna LW, Horvath SM, Bean WB. Post-exertional orthostatic hypotension. *Am J Med Sci.* 1947a; 213:641–654. [PubMed: 20242611]
- Eichna LW, Horvath SM, Bean WB. Cardiac asystole in a normal young man following physical effort. *Am Heart J.* 1947b; 33:254–262. [PubMed: 20283567]
- Emhoff CA, Barrett-O'Keefe Z, Padgett RC, Hawn JA, Halliwill JR. Histamine-receptor blockade reduces blood flow but not muscle glucose uptake during postexercise recovery in humans. *Exp Physiol.* 2011; 96:664–673. [PubMed: 21478258]
- Endo Y, Tabata T, Kuroda H, Tadano T, Matsushima K, Watanabe M. Induction of histidine decarboxylase in skeletal muscle in mice by electrical stimulation, prolonged walking and interleukin-1. *J Physiol.* 1998; 509:587–598. [PubMed: 9575306]
- Fleg JL, Asante AV. Asystole following treadmill exercise in a man without organic heart disease. *Arch Int Med.* 1983; 143:1821–1822. [PubMed: 6615110]
- Fleg JL, Lakatta EG. Prevalence and significance of postexercise hypotension in apparently healthy subjects. *Am J Cardiol.* 1986; 57:1380–1384. [PubMed: 3717041]
- Floras JS, Sinkey CA, Aylward PE, Seals DR, Thoren PN, Mark AL. Postexercise hypotension and sympathoinhibition in borderline hypertensive men. *Hypertension.* 1989; 14:28–35. [PubMed: 2737735]
- Freeman R, et al. Consensus statement on the definition of orthostatic hypotension, neurally mediated syncope and the postural tachycardia syndrome. *Clin Auto Res.* 2011; 21:69–72.
- Freitas J, Costa O, Carvalho MJ, de Freitas AF. High altitude-related neurocardiogenic syncope. *Am J Cardiol.* 1996; 77:1021. [PubMed: 8644629]
- Grabbe J. Pathomechanisms in physical urticaria. *J Invest Dermatol Symp Proc.* 2001; 6:135–136.

- Graham P, Kahlson G, Rosengren E. Histamine formation in physical exercise, anoxia and under the influence of adrenaline and related substances. *J Physiol.* 1964; 172:174–188. [PubMed: 14205015]
- Gratz G, Mayer H, Skrabal F. Sympathetic reserve, serum potassium, and orthostatic intolerance after endurance exercise and implications for neurocardiogenic syncope. *Eur Heart J.* 2008; 29:1531–1541. [PubMed: 18480094]
- Greenleaf JR, Bosco JS, Matter M. Orthostatic tolerance in dehydrated, heat-acclimated men following exercise in the heat. *Aerospace Med.* 1974; 45:491–497. [PubMed: 4831605]
- Hainsworth R. Syncope: what is the trigger? *Heart.* 2003; 89:123–124. [PubMed: 12527651]
- Halliwill JR. Mechanisms and clinical implications of post-exercise hypotension in humans. *Exerc Sport Sci Rev.* 2001; 29:65–70. [PubMed: 11337825]
- Halliwill JR. Hypoxic regulation of blood flow in humans. Skeletal muscle circulation and the role of epinephrine. *Adv Exper Med Biol.* 2003; 543:223–236. [PubMed: 14713125]
- Halliwill JR, Buck TM, Lacewell AN, Romero SA. Postexercise hypotension and sustained postexercise vasodilation: what happens after we exercise? *Exp Physiol.* 2013; 98:7–18. [PubMed: 22872658]
- Halliwill JR, Dinunno FA, Dietz NM. Alpha-adrenergic vascular responsiveness during postexercise hypotension in humans. *J Physiol.* 2003; 550:279–286. [PubMed: 12766237]
- Halliwill JR, Minson CT. Cardiovascular regulation during combined hypoxic and orthostatic stress: fainters vs. nonfainters. *J Appl Physiol.* 2005; 98:1050–1056. [PubMed: 15531565]
- Halliwill JR, Minson CT, Joyner MJ. Effect of systemic nitric oxide synthase inhibition on postexercise hypotension in humans. *J Appl Physiol.* 2000; 89:1830–1836. [PubMed: 11053333]
- Halliwill JR, Taylor JA, Eckberg DL. Impaired sympathetic vascular regulation in humans after acute dynamic exercise. *J Physiol.* 1996; 495:279–288. [PubMed: 8866370]
- Hand J. Exercise-induced vasodepressor syncope in a collegiate wrestler: a case study. *J Ath Train.* 1997; 32:359–362.
- Hastings JL, Levine BD. Syncope in the Athletic Patient. *Prog Cardiovasc Dis.* 2012; 54:438–444. [PubMed: 22386295]
- Hellebrandt FA, Cary MK. Relative importance of the muscle pump in the prevention of gravity shock. *Phys Ther Rev.* 1949; 29:12–22. [PubMed: 18108783]
- Hellebrandt FA, Franseen EB. Physiological study of the vertical stance of man. *Physiol Rev.* 1943; 23:220–255.
- Hill L. Arterial pressure in man while sleeping, resting, working and bathing. *J Physiol.* 1898; 22:xxvi–xxx.
- Hirata T, Yano K, Okui T, Mitsuoka T, Hashiba K. Asystole with syncope following strenuous exercise in a man without organic heart disease. *J Electrocardiol.* 1987; 20:280–283. [PubMed: 3309112]
- Höcker M, Rosenberg I, Xavier R, Henihan RJ, Wiedenmann B, Rosewicz S, Podolsky DK, Wang TC. Oxidative Stress Activates the Human Histidine Decarboxylase Promoter in AGS Gastric Cancer Cells. *J Biol Chem.* 1998; 273:23046–23054. [PubMed: 9722530]
- Holtzhausen LM, Noakes TD. The prevalence and significance of post-exercise (postural) hypotension in ultramarathon runners. *Med Sci Sports Exerc.* 1995; 27:1595–1601. [PubMed: 8614313]
- Holtzhausen LM, Noakes TD. Collapsed ultraendurance athlete: proposed mechanisms and an approach to management. *Clin J Sport Med.* 1997; 7:292–301. [PubMed: 9397327]
- Holtzhausen LM, Noakes TD, Kroning B, de Klerk M, Roberts M, Emsley R. Clinical and biochemical characteristics of collapsed ultra-marathon runners. *Med Sci Sports Exerc.* 1994; 26:1095–1101. [PubMed: 7808242]
- Hultgren, H. High Altitude Medicine. Hultgren Publications; Stanford, CA: 1997. Syncope; p. 419–423.
- Huycke EC, Card HG, Sobol SM, Nguyen NX, Sung RJ. Postexertional cardiac asystole in a young man without organic heart disease. *Ann Int Med.* 1987; 106:844–845. [PubMed: 3555201]

- Jarvis SS, Florian JP, Curren MJ, Pawelczyk JA. A somatostatin analog improves tilt table tolerance by decreasing splanchnic vascular conductance. *J Appl Physiol.* 2012; 112:1504–1511. [PubMed: 22345429]
- Jeong HJ, Moon PD, Kim SJ, Seo JU, Kang TH, Kim JJ, Kang IC, Um JY, Kim HM, Hong SH. Activation of hypoxia-inducible factor-1 regulates human histidine decarboxylase expression. *Cell Mol Life Sci.* 2009; 66:1309–1319. [PubMed: 19266161]
- Journeay WS, Carter R, Kenny GP. Thermoregulatory control following dynamic exercise. *Aviat Space Environ Med.* 2006; 77:1174–1182. [PubMed: 17086774]
- Joyner MJ. Orthostatic stress, haemorrhage and a bankrupt cardiovascular system. *J Physiol.* 2009; 587:5015–5016. [PubMed: 19770188]
- Kajekar R, Chen CY, Mutoh T, Bonham AC. GABA(A) receptor activation at medullary sympathetic neurons contributes to postexercise hypotension. *Am J Physiol Heart Circ Physiol.* 2002; 282:H1615–24. [PubMed: 11959623]
- Kapoor WN. Syncope with abrupt termination of exercise. *Am J Med.* 1989; 87:597–599. [PubMed: 2816980]
- Karadag B, Aliyev F, Duman D, Ongen Z, Vural VA. Post-exercise cardiac asystole in persons without underlying heart disease: management and follow-up of three cases. *J Cardiol Med.* 2007; 8:855–858.
- Kaufmann H, Freeman R. Pharmacological treatment of reflex syncope. *Clin Auto Res.* 2004; 14(Suppl 1):71–75.
- Kenefick RW, Sawka MN. Heat exhaustion and dehydration as causes of marathon collapse. *Sports Med.* 2007; 37:378–381. [PubMed: 17465613]
- Kenney WL, Stanhewicz AE, Bruning RS, Alexander LM. Blood pressure regulation III: what happens when one system must serve two masters: temperature and pressure regulation? *Eur J Appl Physiol.* 2013;1007/s00421-013-2652-5
- Kenny MJ, Seals DR. Postexercise hypotension: key features, mechanisms, and clinical significance. *Hypertension.* 1993; 22:653–664. [PubMed: 8225525]
- Kosinski D, Grubb BP, Karas BJ, Frederick S. Exercise-induced neurocardiogenic syncope: clinical data, pathophysiological aspects, and potential role of tilt table testing. *Europace.* 2000; 2:77–82. [PubMed: 11225599]
- Kosinski DJ, Grubb BP. Neurally mediated syncope with an update on indications and usefulness of head-upright tilt table testing and pharmacologic therapy. *Curr Opin Cardiol.* 1994; 9:53–64. [PubMed: 8199371]
- Kramer R, Drori Y, Lev B. Sudden death in young soldiers. High incidence of syncope prior to death. *Chest.* 1988; 93:345–347. [PubMed: 3338301]
- Krediet CTP, Wilde AAM, Halliwill JR, Wieling W. Syncope during exercise, documented with continuous blood pressure monitoring during ergometer testing. *Clin Auto Res.* 2005; 15:59–62.
- Krediet, CTP.; Wilde, AAM.; Halliwill, JR.; Wieling, W. Post-exercise vasovagal syncope. In: Civera, R.; Esquivias, G.; Bland, J.; Brignole, M.; Mitjans, A.; Granell, R.; Wieling, W., editors. *Syncope Cases.* Blackwell; Malden, MA: 2006. p. 46-48.
- Krediet CTP, Wilde AAM, Wieling W, Halliwill JR. Exercise related syncope, when it's not the heart. *Clin Auto Res.* 2004; 14 (Suppl):25–36.
- Kulics JM, Collins HL, DiCarlo SE. Postexercise hypotension is mediated by reductions in sympathetic nerve activity. *Am J Physiol Heart Circ Physiol.* 1999; 45:H27–H32.
- Lacewell AN, Buck TM, Romero SA, Halliwill JR. Post-exercise syncope: Wingate syncope test and effective countermeasure. *Exp Physiol.* 2013;108/1113/expphysiol.2013.075333
- Laughlin MH, Davis MJ, Secher NH, van Lieshout JJ, Arce-esquivel AA, Simmons GH, Bender SB, Padilla J, Bache RJ, Merkus D, Duncker DJ. Peripheral Circulation. *Compr Physiol.* 2012; 2:321–447. [PubMed: 23728977]
- Link MS, Estes NAM. How to manage athletes with syncope. *Cardiol Clin.* 2007; 25:457–466. [PubMed: 17961800]
- Lockwood JM, Wilkins BW, Halliwill JR. H1 receptor-mediated vasodilatation contributes to postexercise hypotension. *J Physiol.* 2005; 563:633–642. [PubMed: 15618271]

- Low DA, da Nóbrega AC, Mathias CJ. Exercise-induced hypotension in autonomic disorders. *Auton Neurosci*. 2012; 171:66–78. [PubMed: 23040841]
- Lowe MD, Petch MC, Everard P. Syncope after effort. *Postgrad Med J*. 2000; 76:164–165. [PubMed: 10684329]
- Luft, UC.; Myhre, LG.; Loeppky, JA.; Venters, MD. A study of factors affecting tolerance of gravitational stress simulated by lower body negative pressure. Lovelace Foundation; Albuquerque, NM: 1976. NASA Contract NAS 9-14472
- Lynn BM, Minson CT, Halliwill JR. Fluid replacement and heat stress during exercise alter post-exercise cardiac haemodynamics in endurance exercise-trained men. *J Physiol*. 2009; 587:3605–3617. [PubMed: 19491249]
- Macdougall JD, Tuxen D, Sale DG, Moroz JR, Sutton JR. Arterial blood pressure response to heavy resistance exercise. *J Appl Physiol*. 1985; 58:785–790. [PubMed: 3980383]
- Mayerson HS. A cardiovascular “blackout” test. *J Aviation Med*. 1944; 15:304–315.
- McCord JL, Beasley JM, Halliwill JR. H2-receptor-mediated vasodilation contributes to postexercise hypotension. *J Appl Physiol*. 2006; 100:67–75. [PubMed: 16141376]
- McCord JL, Halliwill JR. H1 and H2 receptors mediate postexercise hyperemia in sedentary and endurance exercise-trained men and women. *J Appl Physiol*. 2006; 101:1693–1701. [PubMed: 16888049]
- McCord JL, Pellingier TK, Lynn BM, Halliwill JR. Potential benefit from an H1-receptor antagonist on postexercise syncope in the heat. *Med Sci Sports Exerc*. 2008; 40:1953–1961. [PubMed: 18845973]
- Meendering JR, Torgrimson BN, Houghton BL, Halliwill JR, Minson CT. Menstrual cycle and sex affect hemodynamic responses to combined orthostatic and heat stress. *Am J Physiol Heart Circ Physiol*. 2005; 289:H631–H642. [PubMed: 15778279]
- Metcalfe D, Baram D, Mekori Y. Mast cells. *Physiol Rev*. 1997; 77:1033–1079. [PubMed: 9354811]
- Miki K, Yoshimoto M, Tanimizu M. Acute shifts of baroreflex control of renal sympathetic nerve activity induced by treadmill exercise in rats. *J Physiol*. 2003; 548:313–322. [PubMed: 12562953]
- Morales G, Romero SA, Rickards CA, Ryan KL, Convertino VA, Cooke WH. Effects of dehydration on cerebrovascular control during standing after heavy resistance exercise. *J Appl Physiol*. 2012; 112:1875–1883. [PubMed: 22461441]
- Morganroth M, Mohrman D, Sparks H. Prolonged vasodilation following fatiguing exercise of dog skeletal muscle. *Am J Physiol*. 1975; 229:38–43. [PubMed: 1147054]
- Murrell C, Cotter JD, George K, Shave R, Wilson L, Thomas K, Williams MJA, Lowe T, Ainslie PN. Influence of age on syncope following prolonged exercise: differential responses but similar orthostatic intolerance. *J Physiol*. 2009; 587:5959–5969. [PubMed: 19884316]
- Murrell C, Wilson L, Cotter JD, Lucas S, Ogoh S, George K, Ainslie PN. Alterations in autonomic function and cerebral hemodynamics to orthostatic challenge following a mountain marathon. *J Appl Physiol*. 2007; 103:88–96. [PubMed: 17379746]
- Murrell CJ, Cotter JD, George K, Shave R, Wilson L, Thomas K, Williams MJA, Ainslie PN. Syncope is unrelated to supine and postural hypotension following prolonged exercise. *Eur J Appl Physiol*. 2011; 111:469–476. [PubMed: 20886226]
- Natale A, Akhtar M, Jazayeri M, Dhala A, Blanck Z, Deshpande S, Krebs A, Sra JS. Provocation of hypotension during head-up tilt testing in subjects with no history of syncope or presyncope. *Circulation*. 1995; 92:54–58. [PubMed: 7788917]
- Noakes TD. Why marathon runners collapse. *S Afr Med J*. 1988; 73:569–571. [PubMed: 3375898]
- Noakes TD. The forgotten Barcroft/Edholm reflex: potential role in exercise associated collapse. *Br J Sports Med*. 2003; 37:277–282. [PubMed: 12782559]
- Noakes TD. Reduced peripheral resistance and other factors in marathon collapse. *Sports Med*. 2007; 37:382–385. [PubMed: 17465614]
- O'Connor FG, Levine BD, Childress MA, Asplundh CA, Oriscello RG. Practical management: a systematic approach to the evaluation of exercise-related syncope in athletes. *Clin J Sport Med*. 2009; 19:429–434. [PubMed: 19741318]

- Ogoh S, Dalsgaard MK, Yoshiga CC, Dawson EA, Keller DM, Raven PB, Secher NH. Dynamic cerebral autoregulation during exhaustive exercise in humans. *Am J Physiol Heart Circ Physiol*. 2005; 288:H1461–H1467. [PubMed: 15498819]
- Ogoh S, Fisher JP, Purkayastha S, Dawson EA, Fadel PJ, White MJ, Zhang R, Secher NH, Raven PB. Regulation of middle cerebral artery blood velocity during recovery from dynamic exercise in humans. *J Appl Physiol*. 2007; 102:713–721. [PubMed: 17068217]
- Osswald S, Brooks R, O’Nunain SS, Curwin JH, Roelke M, Radvany P, Ruskin JN, McGovern BA. Asystole after exercise in healthy persons. *Ann Int Med*. 1994; 120:1008–1011. [PubMed: 8185132]
- Pearson J, Lucas RAI, Crandall CG. Elevated local skin temperature impairs cutaneous vasoconstrictor responses to a simulated haemorrhagic challenge while heat stressed. *Exp Physiol*. 2013; 98:444–450. [PubMed: 22903981]
- Pedersen WR, Janosik DL, Goldenberg IF, Stevens LL, Redd RM. Post-exercise asystolic arrest in a young man without organic heart disease: utility of head-up tilt testing in guiding therapy. *Am Heart J*. 1989; 118:410–413. [PubMed: 2568746]
- Perry BG, Schlader ZJ, Barnes MJ, Cochrane DJ, Lucas SJE, Mündel T. Hemodynamic Response to Upright Resistance Exercise: Effect of Load and Repetition. *Med Sci Sports Exerc*. 2013;10.1249/MSS.0b013e3182a7980f
- Petersen ME, Sutton R. Cardiac pacing for vasovagal syncope: a reasonable therapeutic option? *PACE*. 1997; 20:824–826. [PubMed: 9080519]
- Pollack AA, Wood EH. Venous pressure in the saphenous vein at the ankle in man during exercise and changes in posture. *J Appl Physiol*. 1949; 1:649–662. [PubMed: 18124797]
- Pott F, Van Lieshout JJ, Ide K, Madsen P, Secher NH. Middle cerebral artery blood velocity during intense static exercise is dominated by a Valsalva maneuver. *J Appl Physiol*. 2003; 94:1335–1344. [PubMed: 12626468]
- Pricher MP, Holowatz LA, Williams JT, Lockwood JM, Halliwill JR. Regional hemodynamics during postexercise hypotension. I. Splanchnic and renal circulations. *J Appl Physiol*. 2004; 97:2065–2070. [PubMed: 15310744]
- Privett SE, George KP, Middleton N, Whyte GP, Cable NT. The effect of prolonged endurance exercise upon blood pressure regulation during a postexercise orthostatic challenge. *Br J Sports Med*. 2010a; 44:720–724. [PubMed: 18981045]
- Privett SE, George KP, Whyte GP, Cable NT. The effectiveness of compression garments and lower limb exercise on post-exercise blood pressure regulation in orthostatically intolerant athletes. *Clin J Sport Med*. 2010b; 20:362–367. [PubMed: 20818194]
- Raine NM, Cable NT, George KP, Campbell IG. The influence of recovery posture on post-exercise hypotension in normotensive men. *Med Sci Sports Exerc*. 2001; 33:404–412. [PubMed: 11252067]
- Rasmussen P, Stie H, Nielsen B, Nybo L. Enhanced cerebral CO₂ reactivity during strenuous exercise in man. *Eur J Appl Physiol*. 2006; 96:299–304. [PubMed: 16284788]
- Romero SA, Cooke WH. Hyperventilation before resistance exercise: cerebral hemodynamics and orthostasis. *Med Sci Sports Exerc*. 2007; 39:1302–1307. [PubMed: 17762363]
- Romme JJ, Reitsma JB, Black CN, Colman N, Scholten RJ, Wieling W, Van Dijk N. Drugs and pacemakers for vasovagal, carotid sinus and situational syncope. *Cochrane Database Syst Rev*. 2011;CD004194. [PubMed: 21975744]
- Rowell, LB. Competition between skin and muscle for blood flow during exercise. In: Nadel, ER., editor. *Problems with temperature regulation during exercise*. Academic Press; New York, NY: 1977. p. 49-77.
- Rowell, LB. *Human circulation: Regulation during physical stress*. Oxford University Press; New York, NY: 1986.
- Rowell LB, Blackmon JR. Hypotension induced by central hypovolaemia and hypoxaemia. *Clin Physiol*. 1989; 9:269–277. [PubMed: 2743745]
- Sakaguchi S, Shultz JJ, Remole SC, Adler SW, Lurie KG, Benditt DG. Syncope associated with exercise, a manifestation of neurally mediated syncope. *Am J Cardiol*. 1995; 75:476–481. [PubMed: 7863992]

- Schlesinger Z. Letter: Life-threatening “vagal reaction” to physical fitness test. *JAMA*. 1973; 226:1119. [PubMed: 4800477]
- Senitko AN, Charkoudian N, Halliwill JR. Influence of endurance exercise training status and gender on postexercise hypotension. *J Appl Physiol*. 2002; 92:2368–2374. [PubMed: 12015349]
- Shvartz E, Meroz A, Magazanik A, Shoenfeld Y, Shapiro Y. Exercise and heat orthostatism and the effect of heat acclimation and physical fitness. *Aviat Space Environ Med*. 1977; 48:836–842. [PubMed: 907595]
- Smith GD, Bannister R, Mathias CJ. Post-exertion dizziness as the sole presenting symptom of autonomic failure. *Br Heart J*. 1993; 69:359–361. [PubMed: 8489871]
- Son A, Nakamura H, Kondo N, Matsuo Y, Liu W, Oka S, Ishii Y, Yodoi J. Redox regulation of mast cell histamine release in thioredoxin-1 (TRX) transgenic mice. *Cell research*. 2006; 16:230–239. [PubMed: 16474438]
- Stegall HF. Muscle pumping in the dependent leg. *Circ Res*. 1966; 19:180–190.
- Sun BC, Emond JA, Camargo CA. Direct medical costs of syncope-related hospitalizations in the United States. *Am J Cardiol*. 2005; 95:668–671. [PubMed: 15721118]
- Tamura Y, Onodera O, Kodera K, Igarashi Y, Miida T, Aizawa Y, Izumi T, Shibata A, Takano S. Atrial standstill after treadmill exercise test and unique response to isoproterenol infusion in recurrent postexercise syncope. *Am J Cardiol*. 1990; 65:533–535. [PubMed: 2305697]
- Thijs RD, Reijntjes RHAM, van Dijk JG. Water drinking as a potential treatment for idiopathic exercise-related syncope: a case report. *Clin Auto Res*. 2003; 13:103–105.
- Tse HF, Lau CP. Exercise-associated cardiac asystole in persons without structural heart disease. *Chest*. 1995; 107:572–576. [PubMed: 7842800]
- Tsutsumi E, Hara H, RPB. Syncope after running. *Br Med J*. 1979; 280:1480. [PubMed: 526822]
- Tzeng Y-C, Ainslie PN. Blood pressure regulation IX: cerebral autoregulation under blood pressure challenges. *Eur J Appl Physiol*. 2013
- Ueyama T, Hano T, Kasamatsu K, Mohara O, Shima H, Ueshima K, Fukuda K, Ueno Y, Nishio I. A case of orthostatic and postexertional hypotension with ischemic electrocardiographic changes. *Jap Heart J*. 1994; 35:375–382. [PubMed: 7933554]
- Watanabe T, Ohtsu H. L-histidine decarboxylase as a probe in studies on histamine. *Chem Rec*. 2002; 2:369–376. [PubMed: 12469348]
- Westendorp RG, Blauw GJ, Frölich M, Simons R. Hypoxic syncope. *Aviat Space Environ Med*. 1997; 68:410–414. [PubMed: 9143751]
- Wilkins BW, Minson CT, Halliwill JR. Regional hemodynamics during postexercise hypotension. II. Cutaneous circulation. *J Appl Physiol*. 2004; 97:2071–2076. [PubMed: 15322061]
- Willie CK, Ainslie PN, Taylor CE, Eves ND, Tzeng Y-C. Maintained cerebrovascular function during post-exercise hypotension. *Eur J Appl Physiol*. 2013; 113:1597–1604. [PubMed: 23314684]
- Wilson TE, Cui J, Zhang R, Witkowski S, Crandall CG. Skin cooling maintains cerebral blood flow velocity and orthostatic tolerance during tilting in heated humans. *J Appl Physiol*. 2002; 93:85–91. [PubMed: 12070190]
- Ziegelstein RC. Near-syncope after exercise. *JAMA*. 2004; 292:1221–1226. [PubMed: 15353535]

FIGURE 1.

Early observations of the muscle pump. *Panel A:* Changes in calf volume with muscle contraction as measured by plethysmography. E denotes the onset of exercise (plantar flexion at 1 Hz for 10 s); R denotes onset of rest; CP denotes inflation of a cuff above the knee to 90 mmHg for remainder of tracing. This early study shows the considerable volume of blood that can be mobilized from the calf in response to modest muscle contractions, as well as showing that the muscle pump is capable of moving blood against a considerable pressure gradient. (Reproduced from Barcroft & Dornhorst, 1949). *Panel B:* Changes in saphenous vein pressure with walking on a treadmill at 1.7 mph. Step-by-step changes in minimum and maximum venous pressure at the level of the ankle during quiet standing (control), slow walking, and quiet standing after walking. This early study shows that dependent limb venous pressure can be markedly reduced within a few steps by engagement of the muscle pump, as well as the rate at which venous pressure will return that that predicted by gravitational forces when muscle pump activity ceases. (Reproduced from Pollack & Wood, 1949).

FIGURE 2. Integrated hemodynamic responses following exercise

Regulated reductions in blood pressure known as post-exercise hypotension can be large enough in magnitude to induce pre-syncope symptoms and lead to post-exercise syncope. The expression of post-exercise hypotension is the integration of a variety of obligatory and situational components. Obligatory components (indicated in pale blue) include 1) a sustained histaminergic vasodilation of the previously exercise skeletal muscle vascular beds, 2) resetting of the baroreflex (which generally results in sympathoinhibition of sympathetic nerves to muscle vascular beds), 3) pre-synaptic inhibition of norepinephrine release from sympathetic nerves to the exercised muscle, and 4) resetting of thermoreflexes. These changes manifest as a rise in vascular conductance of the previously exercised muscle, inhibition of sympathetic vasoconstrictor nerve activity to previously active muscle vascular beds, and little or no change in cutaneous vascular conductance despite higher core temperatures. Situational components (indicated in light yellow) include the impact of whether or not significant fluid loss, gravitational pooling of blood, and hyperthermia are present. These components can greatly impact on the extent to which cardiac output is elevated. (Reproduced from Halliwill *et al.*, 2013).

FIGURE 3. Sex and age of patients evaluated with post-exercise syncope
Panel A: The percentage of male and female patients in case reports. *Panel B:* The distribution of case reports on post-exercise syncope by age. Both figures are based on data from references listed in Table 1 (n = 27).

FIGURE 4. Survival time during head-up tilt following exercise - two models and two countermeasures

The proportion of subjects remaining in the tilted position as shown by survival function curve. *Panel A*: Effect of H₁-receptor blockade on head-up tilt after 45 min running in the heat. (Modified from McCord *et al.*, 2008). *Panel B*: Effect of inspiratory resistance on head-up tilt after 1 min high-intensity exercise. (Modified from Lacewell *et al.*, 2013). Solid line represents the control day; dotted line, countermeasure day.

TABLE 1
COMPILATION OF CASE REPORTS ON POST-EXERCISE SYNCOPÉ

Case reports and case series in which pre-syncope or syncope developed during recovery from exercise. Excluded were cases of syncope that occurred during exercise, rather than post-exercise. Clinical tests show the variety of exercise tests subjects underwent to evaluate syncopal episodes.

Reference	Age	Sex	History	Clinic test	Onset time
Eichna <i>et al.</i> , 1947b	22	Male	Syncope while standing after performance of the Harvard Pack Test	Harvard Pack Test	3 min post-exercise
Schlesinger, 1973	35	Male	Pre-syncopal episodes with prolonged standing	6 min cycling @ 100 watts	Immediately after cessation of exercise
Tsutsumi <i>et al.</i> , 1979	48	Male	History of syncopal episodes after exercise [running]	Master's augmented 2-step test	Immediately after cessation of exercise
Fleg & Asante, 1983	52	Male	Pre-syncopal episode after maximal exercise; brief history of mild hypertension	Graded maximal treadmill test [Balke protocol]	10 min post-exercise
Hirata <i>et al.</i> , 1987	29	Male	6 episodes of syncope after strenuous exercise over last 18 years	Stair-stepping exercise test	Immediately after cessation of exercise
Huycke <i>et al.</i> , 1987	22	Male	Syncopal episode several seconds after maximal exertion [cycling uphill]	Graded maximal treadmill test [Bruce protocol]	Immediately after cessation of exercise
Kapoor, 1989	34	Male	2 syncopal episodes after exercise	Graded maximal treadmill test [Bruce protocol]	Immediately after cessation of exercise
Pedersen <i>et al.</i> , 1989	29	Male	No history of syncope	Graded maximal treadmill test [Bruce protocol]	3 min post-exercise
Tamura <i>et al.</i> , 1990	45	Male	Recurrent syncopal episodes after strenuous exercise [rope skipping]	Graded maximal treadmill test [Bruce protocol]	1 min post-exercise
Arad <i>et al.</i> , 1993	36	Male	Hypertension and syncope after a graded maximal cycle test	Graded maximal cycle test	2 min after cessation of exercise
Ueyama <i>et al.</i> , 1994	62	Male	Two episodes of syncope after exertion; history of hypertension; 50 mg atenolol daily	Graded maximal treadmill test	1 min post-exercise
Osswald <i>et al.</i> , 1994	34	Male	Recurrent syncopal and pre-syncopal episodes after weight lifting	Submaximal exercise test	2 min post-exercise
Osswald <i>et al.</i> , 1994	22	Male	Four syncopal episodes during or after playing basketball	Maximal exercise test	1 min post-exercise
Osswald <i>et al.</i> , 1994	35	Male	Two pre-syncopal episodes after strenuous exercise and after fighting a fire	Undescribed exercise test	8 min post-exercise [seated]
Tse & Lau, 1995	41	Male	Three syncopal episodes associated with abdominal pain or micturition	Graded maximal treadmill test [Bruce protocol]	4 min post-exercise
van den Berg <i>et al.</i> , 1995	37	Male	Recurrent syncopal episodes after exercise	Graded maximal cycle test	6 min post-exercise [during active recovery]
van den Berg <i>et al.</i> , 1995	36	Male	Prolonged asystole and syncopal episode after a graded maximal exercise test	Graded maximal cycle test	4 min post-exercise [during active recovery]
Crisafulli <i>et al.</i> , 2000	18	Male	Syncopal episode after a graded maximal cycle test	Graded maximal cycle test	Immediately after cessation of active recovery

Reference	Age	Sex	History	Clinic test	Onset time
Lowe <i>et al.</i> , 2000	29	Male	Syncopal episode after strenuous physical activity; two pre-syncopal episodes after sexual activity.	Graded maximal treadmill test [Bruce protocol]	1 min post-exercise
Thijjs <i>et al.</i> , 2003	20	Male	Recurrent syncopal and pre-syncopal episodes after exercise	Cycle exercise	Immediately after cessation of exercise
Ziegelstein, 2004	72	Male	Recurrent pre-syncopal episodes after exercise	n/a	n/a
Krediet <i>et al.</i> , 2005	27	Female	Recurrent syncopal and pre-syncopal episodes during exercise and sexual activity	Graded maximal cycle test	During maximal or immediately after cessation of exercise
Krediet <i>et al.</i> , 2006	28	Male	Syncopal episode after a graded maximal cycle test	Graded maximal cycle test	3 min post-exercise
Dockery & Newman, 2007	40	Male	Syncopal episode after playing basketball; four pre-syncopal episodes after exercise	Graded maximal treadmill test [Bruce protocol]	30 s post-exercise
Karadag <i>et al.</i> , 2007	44	Male	History of syncope and chest pain	Graded maximal treadmill test [Bruce protocol]	2 min post-exercise
Karadag <i>et al.</i> , 2007	41	Male	Bodybuilder with recurrent syncopal and pre-syncopal episodes associated with strenuous exercise	Graded maximal treadmill test [Bruce protocol]	3 min post-exercise
Karadag <i>et al.</i> , 2007	44	Male	Atypical chest pain with no history of syncope	Graded maximal treadmill test [Bruce protocol]	4 min post-exercise

TABLE 2
COMPILATION OF PROSPECTIVE STUDIES ON POST-EXERCISE SYNCOPE

Prospective studies in which pre-syncopal rates during head-up tilt or tolerance to lower body negative pressure were assessed during recovery from exercise.

Reference	Exercise	Environment	Incidence
1 Moderate-intensity exercise			
Mayerson, 1944	10 min cycling @ 45 w	24 to 32°C	54% pre-syncopal; 60 to 75 head-up tilt [15 min]
Gratze <i>et al.</i> , 2008	42.2 km marathon	20°C, 78% humidity	27% pre-syncopal; active standing [6 min]
Privett <i>et al.</i> , 2010a	42.2 km marathon	10 to 11°C, light rain	10% pre-syncopal; active standing [3 min]
Murrell <i>et al.</i> , 2009	4.5 h running @ 70–80% maximal heart rate	15°C, 72% humidity	81% pre-syncopal; 60 head-up tilt [15 min]
Eichna <i>et al.</i> , 1947a	51.5 km hike with pack	“Moderately warm”	54% pre-syncopal; 70 head-up tilt [5 min]
Holtzhausen & Noakes, 1995	80 km ultra-marathon	14 to 25°C, 44 to 24% humidity	23% pre-syncopal; active standing [10 min]
2 High-intensity exercise			
Brogdon & Hellebrandt, 1940	Brief “violent” exercise	n/a	2 out of 3 pre-syncopal; active standing
Eichna <i>et al.</i> , 1947a	Maximal exercise for up to 5 min	23°C	58% pre-syncopal; 70 head-up tilt [5 min]
Bjurstedt <i>et al.</i> , 1983	Graded maximal cycle test	21°C	83% pre-syncopal; 70 head-up tilt [6 min]
Privett <i>et al.</i> , 2010b	Graded maximal treadmill test	23°C	100% pre-syncopal; active standing [10 min]
Lacewell <i>et al.</i> , 2013	Supra-maximal exercise for 1 min [modified Wingate test]	23°C, 40% humidity	58% pre-syncopal; 60 head-up tilt [15 min]
3 Exercise in the heat			
Luft <i>et al.</i> , 1976	2 h intermittent cycling @ 30% VO ₂ max	50°C, 15% humidity	Lower body negative pressure tolerance was reduced 31%
Shvartz <i>et al.</i> , 1977	3 h bench stepping @ 40 w	39°C, 50% humidity	46% pre-syncopal; leaning upright against wall [20 min]
McCord <i>et al.</i> , 2008	45 min running @ 50% VO ₂ max	35°C, 30% humidity	53% pre-syncopal; 60 head-up tilt [15 min]
4 Exercise at altitude - No prospective studies			
5 Resistance exercise - No prospective studies			

TABLE 3

CASE STUDIES WITH INTERVENTIONS AND FOLLOW-UP

Reference	Treatment/Intervention	Outcome/Follow-up
Behavioral		
Fleg & Asante, 1983	Avoid strenuous exercise	No further episodes at 3 months
Hirata <i>et al.</i> , 1987	Avoid strenuous exercise	n/a
Tamura <i>et al.</i> , 1990	Cardiac pacing and active walking recovery	No further episodes with use of active recovery
Tse & Lau, 1995	Avoid strenuous exercise	No further episodes at 12 months
van den Berg <i>et al.</i> , 1995	Avoid strenuous exercise	n/a
Crisafulli <i>et al.</i> , 2000	Active recovery after exercise	n/a
Thijs <i>et al.</i> , 2003	Ingest 1000 ml water, 15 min prior to exercise	No further episodes with use of countermeasure at 2 months
Krediet <i>et al.</i> , 2005	Instruction to try squatting, leg-crossing, leg-buttock-abdominal muscle tensing, and drinking water	No further episodes with use of physical countermeasures at 4 months
Krediet <i>et al.</i> , 2006	Active recovery after exercise	No further episodes at 5 years
Krediet <i>et al.</i> , 2006	Avoid strenuous exercise	No further episodes at 9 months
Karadag <i>et al.</i> , 2007	Avoid strenuous exercise	No further episodes at 3 years
Pharmacological		
Tsutsumi <i>et al.</i> , 1979	β -blocker [20 mg pindolol]	n/a
Huycke <i>et al.</i> , 1987	β -blocker [14 mg propranolol IV, then 20 mg every 6 hours for 3 days before exercise testing]; avoid strenuous exercise	No further episodes with avoidance at 1 year
Pedersen <i>et al.</i> , 1989	β -blocker [100 mg metoprolol twice daily]	No further episodes at long-term follow up
Osswald <i>et al.</i> , 1994	Anti-arrhythmic/anti-muscarinic [disopyramide]	No further episodes at 38 months
Osswald <i>et al.</i> , 1994	β -blocker [25 mg atenolol daily]	No further episodes at 9 months
Pacemaker		
Kapoor, 1989	Dual-chamber permanent pacemaker	No further episodes at 4 year
Lowe <i>et al.</i> , 2000	Dual-chamber permanent pacemaker	Some pre-syncopal symptoms after exercise, but no measurable hypotension
Ziegelstein, 2004	Dual-chamber permanent pacemaker	No further episodes at 2 years
Dockery & Newman, 2007	Dual-chamber permanent pacemaker	No further episodes
van den Berg <i>et al.</i> , 1995	Rate-responsive [QT] VVI pacemaker	Asymptomatic with exercise
Karadag <i>et al.</i> , 2007	Dual-chamber permanent pacemaker	No further episodes at 26 months
Karadag <i>et al.</i> , 2007	Dual-chamber permanent pacemaker	No further episodes at 12 months

TABLE 4
COUNTERMEASURES AGAINST POST-EXERCISE SYNCOPE

Prospective studies in which pre-syncope rates during head-up tilt were assessed during recovery from exercise with a countermeasure designed to prevent syncope.

Reference	Exercise and orthostatic test	Countermeasure	Outcome
Eichna <i>et al.</i> , 1947a	High-intensity exercise; 70 head-up tilt [5 min]	Alternate movement of the legs; vascular occlusion of the legs prior to tilt	Reduced incidence of pre-syncope
McCord <i>et al.</i> , 2008	Exercise in the heat; 60 head-up tilt [15 min]	Histamine H ₁ -receptor blockade	Reduced incidence of pre-syncope from 53% to 27%
Privett <i>et al.</i> , 2010b	High-intensity exercise; active standing [10 min]	Heel raises	Reduced incidence of pre-syncope from 100% to 33%
Privett <i>et al.</i> , 2010b	High-intensity exercise; active standing [10 min]	Compression stockings	Reduced incidence of pre-syncope from 100% to 16%
Lacewell <i>et al.</i> , 2013	High-intensity exercise; 60 head-up tilt [15 min]	Inspiratory resistance	Reduced incidence of pre-syncope from 58% to 33%