Published in final edited form as:

Int J Dev Neurosci. 2014 April ; 33: 1–7. doi:10.1016/j.ijdevneu.2013.10.006.

Cell size anomalies in the auditory thalamus of rats with hypoxic-ischemic injury on postnatal day 3 or 7

Michelle Alexander¹, Haley Garbus¹, Amanda L. Smith¹, and R. Holly Fitch¹

¹University of Connecticut, Department of Psychology, 406 Babbidge Road, Unit 1020, Storrs, CT 06269

Abstract

Children born prematurely (<37 weeks gestational age) or at very low birth weight (VLBW; <1500 grams) are at increased risk for hypoxic ischemic (HI) brain injuries. Term infants can also suffer HI from birth complications. In both groups, blood/oxygen delivery to the brain is compromised, often resulting in brain damage and later cognitive delays (e.g., language deficits). Literature suggests that language delays in a variety of developmentally impaired populations (including specific language impairment (SLI), dyslexia, and early HI-injury) may be associated with underlying deficits in rapid auditory processing (RAP; the ability to process and discriminate brief acoustic cues). Data supporting a relationship between RAP deficits and poor language outcomes is consistent with the “magnocellular theory,” which purports that damage to or loss of large (magnocellular) cells in thalamic nuclei could underlie disruptions in temporal processing of sensory input, possibly including auditory (medial geniculate nucleus; MGN) information. This theory could be applied to neonatal HI populations that show subsequent RAP deficits. In animal models of neonatal HI, persistent RAP deficits are seen in postnatal (P)7 HI injured rats (who exhibit neuropathology comparable to term birth injury), but not in P1–3 HI injured rodents (who exhibit neuropathology comparable to human pre-term injury). The current study sought to investigate the mean cell size, cell number, and cumulative probability of cell size in the MGN of P3 HI and P7 HI injured male rats that had previously demonstrated behavioral RAP deficits. Pilot data from our lab (Alexander et al., 2011) previously revealed cell size abnormalities (a shift towards smaller cells) in P7 but not P1 HI injured animals when compared to shams. Our current findings support this result, with evidence of a significant shift to smaller cells in the experimental MGN of P7 HI but not P3 HI subjects. P7 HI animals also showed significantly fewer cells in the affected (right) MGN as compared P3 HI and shams animals. Moreover, cell number in the right hemisphere was found to correlate with gap detection (fewer cells = worse performance) in P7 HI injured subjects. These findings could be applied to clinical populations, providing an anatomic marker that may index potential long-term language disabilities in HI injured infants and possibly other at-risk populations.

1.1 Introduction

Hypoxic Ischemic (HI) injuries are commonly diagnosed in infants born prematurely (< 37 weeks gestational age) and/or at very low birth weight (VLBW; < 1500 grams; Annibale &

© 2013 ISDN. Published by Elsevier Ltd. All rights reserved.

Corresponding Author: Michelle Alexander, PhD, 516 Delaware Street SE, Minneapolis MN 55455, 01-612-625-2672, alex0213@umn.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Hill, 2008; Volpe, 2001, 2009). Additionally, HI injuries can arise in term infants suffering from birth complications (e.g., cord prolapse, cord asphyxia, or fetal distress; de Vries & Cowan, 2009; Johnston, et al., 2001; Lai & Yang, 2011; Volpe 2001). In both populations, HI injury reflects a reduction in blood and/or oxygen delivery to the brain. In premature infants, HI brain injuries are commonly related to the presence of intraventricular (IVH)/periventricular hemorrhage (PVH), or periventricular leukomalacia (PVL; a non-hemorrhagic ischemic injury; Volpe 2001), with characteristic patterns of neuropathology including white matter (subcortical) damage and disruption of white matter fiber tracts (e.g., cortico-spinal tracts and internal capsule; Nagy et al., 2010). Conversely, among term infants with HI, hypoxic ischemic encephalopathy (HIE) is more commonly diagnosed. HIE is characterized by global gray matter damage to the cortex, hippocampus, basal ganglia and/or thalamus (Jyoti et al., 2006; Huang & Castilla, 2008; Marinez-Biagant et al., 2012; Nosarti et al., 2008).

Although the timing and location of brain injury differs between premature and term populations, both groups show an elevated rate of learning and language deficits compared to matched controls. In one study, children born prematurely were found to require more support in spelling, reading and writing, and had deficits in receptive and expressive language skills (Luu et al., 2009; Ortiz-Mantilla, et al., 2008; van Lierde et al., 2009). At two and four years old, VLBW infants had significantly lower scores on language comprehension tasks, and these scores correlated with performance on language comprehension, naming, and auditory discrimination tasks at 4 years (Jansson-Verkassiao et al., 2004). At age 6, these subjects showed alterations on mismatched negativity during naming tasks, and difficulty in preattentively discriminating changes in syllables (Jansson-Verkassiao et al., 2004). Full term infants diagnosed with moderate to severe HIE also show receptive language, reading and spelling deficits compared to healthy controls (Badawi et al., 2001), and degree of HIE injury was found to correlate significantly with verbal IQ scores (Steinman et al., 2009).

Previous research has suggested that deficits in rapid auditory processing (RAP; processing mechanisms required to distinguish phonemes) may underlie later disruptions in development of complex language and reading (Bradley et al., 1983; Farmer et al., 1985; Hari et al., 1996, 2001; Kraus et al., 1996; McAnally et al., 1996; McCrosky et al., 1980; Stein 1993; Neville et al., 1993; Robin et al., 1989; Paterson et al., 2006; Tallal et al., 1980, 1985, 1993; Witton et al., 1998; Wright et al., 1997; Benasich et al., 2002; Choudhury et al., 2007). Specifically regarding HI injured populations, VLBW children diagnosed with severe PVL scored significantly lower on auditory temporal processing tasks, with path analysis supporting a relationship between auditory temporal processing and later reading ability (Downie et al., 2002).

To date, among neonatal HI injured populations, no specific anatomical marker has been found to correlate with language delays. However, in other populations of developmentally language disordered populations (e.g., dyslexia), *post mortem* studies have demonstrated significant reductions in the size of magnocellular cells in the lateral geniculate nucleus (LGN), which encode temporal components of visual information (Livingstone et al., 1991; Galaburda et al., 1985; Galaburda & Livingstone, 1993; Lehmkuhle et al., 1993; Lovegrove et al., 1990; Slaghuis et al., 1993; Stein 2001). Thus the anatomical abnormalities were suggested to relate to behavioral deficits in magnocellular aspects of visual information processing (i.e., low contrast motion; Livingstone et al., 1991; Lovegrove et al., 1990; Slaghuis et al., 1993; Lehmkuhle et al., 1993). Subsequently, these findings were extended to cell populations in the medial geniculate nucleus (MGN) from the same dyslexic brains. Here, *post mortem* histological analysis again revealed a shift toward smaller cells in the MGN (Galaburda & Livingstone et al., 1993). It is hypothesized that the large cells in the

MGN might preferentially process temporal aspects of acoustic information (much like functional divisions seen in the LGN), and disruptions to these cells might relate to behavioral impairments in RAP (Galaburda & Livingstone, 1993). The latter results are consistent with the general magnocellular theory, which hypothesizes that behavioral deficits in temporal processing *across various sensory systems* are associated with reductions in large or “magno” cells of thalamic nuclei (Chase & Stein, 2003; Leonard, 1998; Livingstone et al., 1991; Stein 2001). However, it should be noted that a magnocellular sub-division of the MGN has not been anatomically identified (as seen in the LGN).

Animal models can provide further insight into cellular abnormalities following early disruptions in brain development (e.g., see Fitch et al., 2004; Szalkowski & Fitch, 2012). For example, the Rice-Vannucci model can be used to study neonatal HI injury in rodent models, and involves cauterization of the right common carotid artery followed by a period of time in a hypoxic chamber (8% oxygen; Vannucci & Vannucci, 2005). When these injuries are induced early in life in rats (between postnatal (P) 1–5), resulting pathologies roughly correspond to those seen in preterm HI infants (Silbereis et al., 2010). Injuries induced later in life (P7–12) roughly correspond to those of full term infants (i.e., HIE-like pathology; Vannucci & Vannucci 2005). These age translations (rat P1–3 to human GA 24–26; rat P7–10 to human near-term) are also supported by a cross-species comparison of neurodevelopmental events (including auditory system), where many prenatal human neural events occur postnatally in rats (Workman et al., 2013). Previous studies have demonstrated persistent RAP deficits in P7 HI male rats – similar to those identified in language-disabled populations (Alexander et al., 2013; Alexander et al., in review; McClure et al., 2005; McClure et al., 2006; McClure et al., 2006; McClure et al., 2007; Hill et al., 2011). However, other studies found that when injuries were induced earlier in life (on P1 or P3), male rats showed deficits on RAP tasks only in juvenile periods but not adulthood (i.e., the deficits were transient; McClure et al., 2006; Alexander et al., in review). Authors interpreted this as a “recovery of function” in the early but not late HI injured rodents (McClure et al., 2006; Alexander et al., in review). In addition, pilot data indicated significantly smaller cells and abnormal cell size distributions in the MGN of the rats with a P7 HI injury, though P1 HI injured animals showed no changes in cell size or cell distribution compared to shams (Alexander et al., 2011). Taken together, these data suggest that persistent RAP deficits as seen in a term HI model may be associated with permanent cellular abnormalities in the MGN (see also Herman et al., 1997).

Given behavioral evidence of language delays in human neonatal HI populations, behavioral deficits on RAP tasks in late but not early HI injured rats, and pilot indications of cell size abnormalities in the MGN of P7 but not P1 HI injured rodents, the current study sought to evaluate MGN neuronal size distribution in P3 and P7 HI injured rodents that had previously shown rapid auditory processing deficits (see Alexander et al., in review, for behavioral data).

1.2 Methods

Brain tissue from a prior study of P3 and P7 HI injured animals was used for the current study (see Alexander et al., in review). In brief, time-mated pregnant dams were ordered from Charles River Laboratories and shipped to the Psychology Department at the University of Connecticut on embryonic day 5 (E5). At birth, litters were culled to 8 males and 2 females. Only male pups were used based on evidence of sex differences in HI-induced behavioral deficits (Hill et al., 2012). On P3 or P7, pups were anesthetized with isoflurane (2.5%) and received a longitudinal mid-line neck incision and the right common carotid artery was permanently cauterized. Sham animals received the same incision, but no

manipulation of the artery. Animals were randomly selected to be in one of the three treatment conditions (P3 HI, P7 HI, or sham), and all treatment conditions were evenly balanced within and across litters. All pups were marked with a footpad injection and returned to the dam for two hours to feed. HI subjects were then placed in a heated airtight container with flowing 8% humidified oxygen (120 min). Sham animals were placed in a similar open container (also on P3 or P7), and were exposed to normal air oxygen (120 min). At P145 (following behavioral testing), animals were anesthetized with ketamine (100 mg/kg) and xylazine (15 mg/kg) and were transcardially perfused with .9% saline solution followed by 10% buffered formalin. Brains were removed and placed in 10% buffered formalin. Twenty-four hours prior to slicing, brains were placed in 30% sucrose solution for cryoprotection. Brains were sectioned at 60 μm on a cryostat, with every 3rd section saved and mounted. Cell bodies were stained using cresyl violet (CV) staining procedures. Assessment of cellular measures (see below) revealed that data from P3 and P7 shams ($n = 6$ each) could be pooled, yielding final group n 's of: P3 HI $n=14$; P7 HI $n=14$; sham $n=12$.

Cells size in the left and right MGN was measured using Stereo Investigator MicroBrightField software on an Axio 2 Zeiss Microscope set using the 100X objective (Figure 1). With the optical fractionator and nucleator probe, the cell's nucleus was marked, and the distance to six different rays bisecting the perimeter of the cell membrane were measured. On average, 5–6 brain sections including the MGN (per animal/per hemisphere) were used for measurement, using a grid size of 250 $\mu\text{m} \times 250 \mu\text{m}$ and a counting frame of 25 $\mu\text{m} \times 25 \mu\text{m}$. The average cell size for the right and left MGN were thus calculated for each subject, and analyzed as a function of Treatment. Additionally, the cumulative probability for the MGN cell size (μm^2) was analyzed for the right and left MGN. Here, cell size values were ordered from smallest to largest within each Treatment group (for each side), and the cumulative probability was calculated by dividing "1" by the total number of cells counted in each Treatment group. Thus we were able to represent the frequency of data points equal to or smaller than a given point across the X-axis (cell size), ranging from 1 to 400 μm^2 . This cumulative frequency distribution was generated separately for each Treatment group, for each side. Additionally, mean cell area and number of cells per section (total number of cells divided by the number of sections counted) for the left and right MGN were analyzed as a function of Treatment. Finally, bivariate correlations were used in order to analyze a possible relationship between average RAP behavioral scores on a 0–100 ms silent gap detection task (see Alexander et al., in review, for task details), and mean size and number of cells per section in the right MGN. Scores from silent gap detection tasks administered in both juvenile and adult periods were used. For the juvenile task, raw measures included performance scores by day (across four days of testing), and gap duration (2–100ms). For the adult silent gap detection task, multiple gap durations from one day of testing (2–100ms) were assessed. To reduce the number of correlations run, performance scores were averaged across gaps for each subject at each age, and for each day of testing, to obtain four scores for each subject in juvenile period, and one adult gap detection score. Bivariate correlations were then run.

For mean cell size and number of cells per section, data was analyzed using a one-way analysis of variance (ANOVA), and follow-up comparisons were performed using independent samples t-tests. Cumulative distribution of cells between all three groups was analyzed using a non-parametric Kruskal-Wallis test. Specific comparisons between groups were made using nonparametric Kolmogorov-Smirnov tests. Finally, bivariate correlations were run between number/mean size of cells per section and behavioral scores (4 juvenile and 1 adult measure). These correlations were performed within each Treatment group separately, to avoid confounding from mean group differences.

1.3 Results

Cell measurements were taken from a total of 40 brains. As noted above, for the two sub-groups of shams (P3 and P7), no significant differences were seen for average cell size, number of cells per section, cell distribution, nor in previously obtained behavioral RAP scores (Alexander et al., in review). Thus shams were pooled to create final n's of: P3 HI=14, P7 HI=14, Sham=12.

1.3.1 Average Cell Area

A one-way ANOVA revealed no significant overall Treatment effect on average cell area for the left MGN [$F(2,39) = .166, p > .05$] nor right MGN [$F(2,39) = 1.766, p > .05$]. Although this analysis failed to reach significance, scores on the right were in the expected direction, with P7 HI animals showing smaller average cell area (P3 HI = $81.45 \mu\text{m}^2$, P7 HI = $65.58 \mu\text{m}^2$, and sham = $77.55 \mu\text{m}^2$; Table 1A).

1.3.2 Number of Cells per Section

A one-way ANOVA revealed no significant Treatment effect on number of cells for the left MGN [$F(2,39) = .421, p > .05$]. However, analysis did reveal a significant Treatment effect for right MGN [$F(2,39) = 14.423, p < .001$] (Figure 2). Specific follow-up independent samples t-tests revealed a significant difference between P7 HI and sham animals ($t(24) = 5.445, p < .001$) and between P3 HI and P7 HI animals ($t(26) = 4.540, p < .001$), with P7 HI animals showing significantly fewer cells per section compared to both groups. There was no significant difference between P3 HI and sham groups ($t(24) = .104, p > .05$; Table 1B).

1.3.3 Cumulative Distribution

For the left MGN (Figure 3a), the results of the Kruskal Wallis non-parametric analysis revealed a significant overall Treatment effect ($H(2) = 10.655, p < .005$). Specific follow-up comparisons using the two sample Kolmogorov-Smirnov test revealed a significant difference between P7 HI and sham groups ($p < .001$), with P7 HI animals showing a shift towards *larger* cells. There was no difference between P3 HI and sham groups ($p > .05$), nor between P3 HI and P7 HI ($p > .05$).

For the right MGN (Figure 3b) the results of the Kruskal Wallis non-parametric analysis again revealed a significant overall Treatment effect ($H(2) = 86.233, p < .001$). Specific follow-up comparisons using the two sample Kolmogorov-Smirnov test revealed a significant difference between P3 HI and sham groups ($p < .001$), with P3 HI animals showing a shift towards larger cells. Additionally, results showed a significant difference between P7 HI and shams groups ($p < .005$), and between P3 HI and P7 HI ($p < .001$), with P7 HI animals showing a shift towards *smaller* cells as compared to both groups.

1.3.4 Behavioral and Cell Size/Number Correlation

When analyzing histological and behavioral correlations within individual groups, P7 HI animals did show a significant correlation between raw RAP scores in the juvenile period (on the 3rd consecutive day of testing) and number of cells per section in the right MGN ($r = -.577, p < .05$; a negative correlation was expected because higher scores reflect worse performance on the discrimination task). Similar correlations for P3 HI and sham animals were in the expected direction, but failed to reach significance. Finally, there were no significant correlations in any group between cell measures and adult RAP scores, although scores again reflected a trend for fewer/smaller cells to be related to worse performance.

1.4 Discussion

Using the HI rodent model, the current study demonstrated that the number of cells in the right MGN of P7 HI injured animals was significantly lower compared to both P3 HI and sham animals. Furthermore, the current study demonstrated that P7 HI injured animals showed a significant shift towards smaller cells in the right MGN as compared to both P3 HI and sham animals. Interestingly, there was a change in the right MGN for P3 HI versus sham animals, with P3 HI animals showing a significant shift towards *larger* cells. Furthermore, P7 HI animals *also* showed a significant shift towards larger cells compared to shams in the left MGN. Finally, bivariate correlation analysis revealed a significant relationship between juvenile RAP scores and average number of cells per section in the right MGN in P7 HI animals (with fewer cells associated with worse performance).

The current findings are consistent with prior pilot evidence of abnormal cell size distributions in the right MGN of late (P7) HI but not early (P1 or P3) HI animals (Alexander et al., 2011). Results are also consistent with a related study where P1 focal ischemic lesions in male rats (resulting in the formation of cortical “microgyria”) led to a shift towards smaller cells in the MGN (Herman et al., 1997). As in the current study, microgyric animals showed concurrent deficits on RAP tasks (Herman et al., 1997). Our findings are also consistent with human *post mortem* evidence demonstrating a shift towards smaller cells in the MGN in brains of dyslexics, coupled with behavioral evidence of RAP deficits in a wide range of language impaired populations (Galaburda et al., 1994; Bradley & Bryant, 1983; Downie et al., 2002; Farmer & Klein, 1985; Hari & Kiesla, 1996; Hari & Renvall, 2001; Kraus et al., 1996; McAnally & Stein, 1996; McCrosky & Kidder, 1980; Neville, et al., 1993; Robin et al., 1989; Paterson, et al., 2006; Stein, 1993; Tallal, 1980; Tallal et al., 1985, 1993; Witton, et al., 1998; Wright et al., 1997; Benasich et al., 2002; Choudhury et al., 2007). The current results support and extend the magnocellular theory, which purports that deficits in rapid auditory processing (or RAP) in language-impaired populations may reflect, in part, anomalies in large or “magno” thalamic MGN cells. This theory originated through evidence of deficits in magnocellular processing of visual information, coupled with an overall reduction of neuronal cell area in the magnocellular division in the LGN of human dyslexic brains (Lehmkuhle et al., 1993; Livingstone et al., 1991; Lovegrove et al., 1990; Slaghuys et al., 1993). Notably, since magnocellular/parvocellular divisions have not been clearly delineated in the MGN (as in the LGN), it can only be suggested that large cells may preferentially encode temporal components of acoustic information (Galaburda & Livingstone, 1993; Stein, 2001).

Interestingly and unexpectedly, the current study also revealed alterations in cells of the right MGN of P3 HI animals, specifically a significant shift towards *larger* cells compared to shams. Alterations in cell size were also seen in the left MGN (contralateral hemisphere to injury) for P7 HI animals, again with a significant shift towards *larger* cells compared to shams. These changes could reflect reorganization and plasticity (e.g., compensatory) effects following the unilateral neonatal HI injury (though it must be noted that both hemispheres received hypoxia). This assertion is consistent with evidence that altered expression of multiple cell death mediators have been reported in both the ipsi- and contralateral hemisphere of six-week old rats following a unilateral P12 HI injury. Specifically, alterations in cytokines, hypoxia inducing factor-1 α , and phosphorylated-Akt levels in both the ipsi- and contralateral hemisphere were seen following HI injury, supporting bilateral reorganization following unilateral HI injury in the neonatal period (Van Den Tweel et al., 2006). The behavioral significance of larger MGN cells remains unclear, but could reflect partial developmental compensation in the P7 HI subjects. Specifically, although it is not possible to assess the behavioral effects of a bi-lateral HI injury (these are largely lethal), it is possible that subjects would have performed much worse on the task without putative

compensation evidenced in the left MGN. In the P3 HI subjects, evidence of an increase in MGN cell size on the right could relate to more effective compensation, since RAP deficits were no longer evident in these subjects by adulthood. Finally, it must be noted that all interpretations are tempered by the fact that MGN measures were obtained long after behaviors were assessed, thus it is not possible to know the characteristics of MGN cells at the time behavioral scores were obtained. Overall, findings of pervasive (bilateral) reorganization following early unilateral HI injury are generally consistent with the notion that, in direct contrast to adult populations where focal brain damage can be clearly tied to specific and largely predictable functional deficits, disruption of the brain during early development does not lead to “focal” injury but rather, a subtle and pervasive reorganization (Thomas & Karmiloff-Smith, 2002; Van Den Tweel et al., 2006). Although this phenomenon may explain the greater resiliency of neonates with regards to maintaining overall function following severe brain injury, it also explains the stark difficulties in identifying a neural signature for specific types of behavioral deficits in developmentally impaired populations.

The current study also reported a decrease in number of cells per section in the right MGN of P7 HI injured rats. This, coupled with a shift towards smaller cells in the right MGN of P7 animals, suggest that large “magno” cells might selectively be dying. This could lead to an interpretation that the reduction in cell number in the right MGN of P7 HI rats simply reflects retrograde degeneration subsequent to cortical atrophy. Notably, a recently published paper also showed a shift in cell size towards smaller cells in the MGN of rats who received embryonic ventricular transfection of RNAi for the dyslexic candidate gene *Dyx1c1* (Szalkowski et al., 2013). Those results support the view that current alterations in MGN cells did not simply occur through retrograde degeneration, since RNAi manipulation was at the genetic level and not injury-based. Notably, the transfected animals in that study also showed evidence of acoustic processing deficits (Szalkowski et al., 2013). In fact, apoptosis of a selective subset of thalamic cells in P7 but not P3 HI animals could be partially explained by the differences in brain development at the time of HI injury. For example, the development of glutamatergic receptor subtypes involved in glutotoxicity (AMPA and NMDA) might contribute to enhanced vulnerability of the MGN at P7 (since increases in the NMDA receptor in areas such as hippocampus and cortex occur later in postnatal rodent development (P7–12), and these populations continue to rise through postnatal development in the human infant; Rice et al., 2000). Alternately, larger MGN cells might require more oxygen and nutritional support, again rendering them more vulnerable to neonatal HI injury.

The timing and development of the central auditory system might also play a crucial role in the impact of HI injury on later language processes (reviewed in Fitch et al., 2013). In typical development, the auditory nerve projects to the cochlear nucleus of the brain stem and sends both contralateral and ipsilateral ascending fibers to the superior olive (also located in the brain stem). From here, the superior olive sends ipsilateral projections to the inferior colliculus which projects to the MGN and on to the primary auditory cortex (A1). Establishment of these connections in the rodent seems to precede functional activation via sound. For example, synapses between the ganglion hair cells, the cochlear nucleus, and the superior olive are apparent as early as E10–14 (Hoffpauir et al., 2009), and spontaneous activation (which is crucial for proper circuitry formation) also occurs prior to the onset of hearing, which occurs postnatally around P11–12 (Tritsch & Bergles, 2010). In human neonates, central auditory maturation also occurs prenatally, with an onset of hearing around GA 27–29 based on speech recognition and fetal auditory brain stem responses (ABRs; Graven & Brown, 2008; Sininger et al., 1997). Although the onset of human hearing occurs prenatally and the rat postnatally, the relative timing for neurodevelopmental events for these two species (including for CN, IC and MGN) is consistent with the models used here.

That is, neurodevelopmental events in the rat are shifted relative to birth when compared to humans, thus allowing postnatal day 3 in rats to provide an appropriate human early pre-term model (Workman et al., 2013).

In both humans and rats, there is still a high degree of behavioral plasticity that occurs postnatally. For example, ABRs and auditory evoked potential response patterns do not reach adult-like patterns until 16–18 years of age in humans for more complex stimuli (Fischer & Hartnegg, 2004), when full maturation of A1 is seen (Moore & Linthicum, 2007). Similarly for rodent development, detectable ABRs associated with hearing do not come online until P12–14, with adult like patterns not emerging until P22 (Zhang et al., 2001). For both the rodent and human neonate, functional connectivity of the auditory system undergo a high degree of postnatal maturation which can be highly affected by developmental insults, such as HI.

In closing, to our knowledge, this is the first study to demonstrate a shift in the cell size distribution and a reduction in cell number in the right MGN (ipsilateral to HI injury) in P7 HI injured rodents that also showed persistent rapid auditory processing deficits at the behavioral level (Alexander et al., in review). Furthermore, P3 HI injured rodents that previously showed a recovery of function on RAP tasks by adulthood did *not* show the same shift towards smaller cells in the right MGN (Alexander et al., in review). Future studies are needed to make a stronger association between other cellular abnormalities in the MGN of HI injured rodents and RAP deficits, perhaps by identifying a more global pattern of altered circuitry. Additionally, studies investigating the use of neuroprotectants following neonatal HI injury could be performed using MGN cell size at an outcome measure. For example, previous studies have demonstrated the therapeutic benefit of erythropoietin (McClure et al., 2006; McClure et al., 2007; Alexander et al., 2013) on RAP deficits in P7 HI injured rodents. It is possible that these therapeutic benefits manifest in part through preserving cellular integrity in the MGN. Finally, improved characterization of apoptotic markers and other cellular features characterizing early and late neonatal HI injury, starting at time-points immediately following injury induction (and throughout development), could help better characterize the pattern and nature of changes in large MGN cells. Overall, an improved understanding of how early injury -- such as that associated with term-birth incidents that cause HIE -- can in turn lead to anatomic changes that can be tied to functional deficits will better serve the at-risk neonatal population through improved early screening. A more delineated focus on specific “brain changes” that follow these early injuries, and lead to the dramatic increases in cognitive disabilities (and language disabilities) evidenced in these vulnerable populations, is also indicated for future research.

References

- Alexander ML, Hill CA, Rosenkrantz TS, Fitch RH. Evaluation of the therapeutic benefit of delayed administration of erythropoietin following early hypoxic-ischemic injury in rodents. *Dev Neurosci*. 2013; 34:515–524. [PubMed: 23328535]
- Alexander, M. Masters Thesis. University of Connecticut; 2011. Cell size distribution in the MGN and associated auditory temporal processing deficits in P1 and P7 HI injured rodents.
- Alexander, M.; Garbus, H.; Smith, AL.; Rosenkrantz, TS.; Fitch, RH. Behavioral and histological outcomes following neonatal HI injury in a preterm (P3) and term (P7) rodent model. In review
- Annibale, D.; Hill, J. Periventricular hemorrhage-intraventricular hemorrhage. 2008. Retrieved from www.emedicine.medscape.com
- Badawi N, Keogh JM, Dixon G, Kurinczuk JJ. Developmental outcomes of newborn encephalopathy in the term infant. *Indian J Pediatr*. 2001; 68:527–530. [PubMed: 11450384]
- Banasia K, Xia Y, Haddad GG. Mechanisms underlying hypoxia-induced neuronal apoptosis. *Progress in Neurobiology*. 2000; 62:215–249. [PubMed: 10840148]

- Benasich AA, Tallal P. Infant discrimination of rapid auditory cues predicts later language impairment. *Behav Brain Res.* 2002; 136:31–49. [PubMed: 12385788]
- Bradley L, Bryant P. Categorizing sounds and learning to read—A causal connection. *Nature.* 1983; 301:419–421.
- Chase C, Stein J. Visual magnocellular deficits in dyslexia. *Brain.* 2003; 126(9):E2. [PubMed: 12937067]
- Choudhury N, Leppanen PH, Leevers HJ, Benasich AA. Infant information processing and family history of specific language impairment: Converging evidence for RAP deficits from two paradigms. *Dev Sci.* 2007; 10:213–236. [PubMed: 17286846]
- de Vries LS, Jongmans MJ. Long-term outcome after neonatal hypoxic-ischemic encephalopathy. *Arch Dis Child Fetal Neonatal Ed.* 2010; 95:F220–F224. [PubMed: 20444814]
- Dekaban A. Human thalamus; an anatomical, developmental and pathological study. II. Development of the human thalamic nuclei. *J Comp Neurol.* 1954; 100:63–97. [PubMed: 13130709]
- Downie AL, Jakobson LS, Frisk V, Ushycky I. Auditory temporal processing deficits in children with periventricular brain injury. *Brain and Lang.* 2002; 80:208–225.
- Farmer ME, Klein RM. The evidence for a temporal processing deficit linked to dyslexia: A review. *Psychon Bull Rev.* 1995; 2:460–493. [PubMed: 24203785]
- Fischer B, Hartnegg. In the development of low-level auditory discrimination and deficits in dyslexia. *Dyslexia.* 2004; 10:105–118. [PubMed: 15180392]
- Fitch RH, Tallal P, Brown CP, Galaburda AM, Rosen GD. Induced microgyria and auditory temporal processing in rats: a model for language impairment? *Cerebral Cortex.* 1994; 4:260–270. [PubMed: 8075531]
- Fitch RH, Alexander M, Threlkeld SW. Early neural disruption and auditory processing outcomes in a rat model. *Auditory Cognitive Neuroscience.* epub ahead of print.
- Galaburda AM, Sherman GF, Rosen GD, Aboitiz F, Geschwind N. Developmental dyslexia: four consecutive patients with cortical anomalies. *Ann Neurology.* 1985; 18:222–233.
- Galaburda, AM.; Livingstone, M. Evidence for a magnocellular deficit in developmental dyslexia, ppabcxyzpp. The magnocellular theory of developmental dyslexia. In: Stein, J., editor. *Dyslexia.* Vol. 7. 1993. p. 65-73.p. 12-36.
- Graven SN, Brown JV. Auditory development in the fetus and neonate. *Newborn and infant nursing reviews.* 2008; 8:187–193.
- Hari R, Kiesla P. Deficit of temporal auditory processing in dyslexic adults. *Neurosci Lett.* 1996; 205:138–140. [PubMed: 8907336]
- Hari R, Renvall H. Impaired processing of rapid stimulus sequences in dyslexia. *Tends Cogn Sci.* 2001; 5:525–532.
- Hill CA, Threlkeld SW, Fitch RH. Early testosterone modulated sex differences in behavioral outcome following neonatal hypoxia ischemia in rats. *Int J Dev Neurosci.* 2011; 29:381–388. [PubMed: 21473905]
- Hill CA, Fitch RH. Sex differences in mechanisms and outcome of neonatal hypoxia-ischemia in rodent models: Implications for sex-specific neuroprotection in clinical neonatal practice. *Neurol Res Int.* 2012; 2012:867531. [PubMed: 22474588]
- Herman AE, Galaburda AM, Fitch RH, Carter AR, Rosen GD. Cerebral microgyria, thalamic cell size and auditory temporal processing in male and female rats. *Cereb Cortex.* 1997; 7:453–464. [PubMed: 9261574]
- Hoffpauir BK, Marrs GS, Mathers PH, Spirou GA. Does the brain connect before the periphery can detect? A comparison of three sensory systems in mice. *Brain Res.* 2009; 1277:115–129. [PubMed: 19272365]
- Huang B, Castillo M. Hypoxic-ischemic brain injury: Imaging findings from birth to adulthood. *Radiographics.* 2008; 28:417–39. [PubMed: 18349449]
- Jansson-Verkasalo E, Valkama M, Vainionpaa L, Paakko E, Ilkko E, Lehtihalmes M. Language development in very low birth weight preterm children: a follow-up study. *Folia Phoniatr Logo.* 2004; 56:108–119.

- Jansson-Verkasalo E, Korpilahti P, Jantti V, Valkama M, Vainionpaa L, Alku P, Suominen K, Naatanen R. Neurophysiologic correlates of deficient phonological representations and object naming in prematurely born children. *Clin Neurophysiol.* 2004; 115:179–187. [PubMed: 14706486]
- Johnston MV, Trescher WM, Ishda A, Nakajima W. Neurobiology of hypoxic-ischemic injury in the developing brain. *Pediatr Res.* 2001; 49:735–741. [PubMed: 11385130]
- Jyoti R, O-Neil R, Hurion E. Predicting outcome in term neonates with hypoxic-ischemic encephalopathy using simplified MR criteria. *Pediatr Radiol.* 2006; 36:38–42. [PubMed: 16284761]
- Kraus N, McGee TJ, Carrell TD, Zecker SG, Nicol TG, Koch DB. Auditory neurophysiologic responses and discrimination deficits in children with learning problems. *Science.* 1996; 273:971–973. [PubMed: 8688085]
- Lai MC, Yang SN. Perinatal hypoxic-ischemic encephalopathy. *J Biomed Biotechnol.* 2011; 2011:609813. [PubMed: 21197402]
- Lehmukuhle S, Garzia RP, Turner L, Hash T, Baro JA. A defective visual pathway on children with reading disability. *N Endl J Med.* 1993; 8:989–996.
- Leonard, LB. Children with specific language impairment. Cambridge, MA: MIT Press; 1998.
- Livngstone MS, Rosen GD, Drislane FW, Galaburda AM. Physiological and anatomical evidence for a magnocellular deficit in developmental dyslexia. *PNAS.* 1991; 88:7943–7947. [PubMed: 1896444]
- Lovegrove WJ, Garzia RP, Nicholson SB. Experimental evidence for a transient system deficit in specific reading disability. *Optometry.* 1990; 61:137–146.
- Luu TM, Ment LR, Schneider KC, Katz KH, Allan WC, Vohr BR. Lasting effects of preterm birth and neonatal brain hemorrhage at 12 years of age. *Pediatrics.* 2009; 123:1037–1044. [PubMed: 19255037]
- Martinez-Biarge M, Diez-Sebastian J, Kapellou O, Gindner D, Allsop JM, Rutherford MA, Cowan FM. Predicting motor outcome and death in term hypoxic-ischemic encephalopathy. *Neurology.* 2011; 76:2055–2061. [PubMed: 21670434]
- McAnally KI, Stein JF. Auditory temporal coding in dyslexia. *Proceedings of the Royal Society of London.* 1996; 263:961–965.
- McClure MM, Peiffer AM, Rosen GD, Fitch RH. Auditory processing deficits in rats with neonatal hypoxic-ischemic injury. *Int J Dev Neurosci.* 2006; 23:351–362. [PubMed: 15927759]
- McClure MM, Threlkeld SW, Rosen GD, Fitch RH. Auditory processing deficits in unilaterally and bilaterally injured hypoxic-ischemic rats. *Neuroreport.* 2005; 16:1309–1312. [PubMed: 16056130]
- McClure MM, Threlkeld SW, Rosen GD, Fitch RH. Rapid auditory processing and learning deficits in rats with P1 versus P7 neonatal hypoxic-ischemic injury. *Behav Brain Res.* 2006; 172:114–121. [PubMed: 16765458]
- McCrosky R, Kidder H. Auditory fusion among learning disabled, reading disabled and normal children. *J Learn Disabil.* 1980; 13:69–76. [PubMed: 7391678]
- Moore JK, Linthicum HR Jr. The human auditory system: a timeline of development. *Int J Audiol.* 2007; 46:460–478. [PubMed: 17828663]
- Nagy Z, Ashburner J, Anderson J, Jbabdi S, Dranganski B, Skare S, Bohm B, Smedler AC, Frossberg H, Lagercrantz H. Structural correlates of preterm birth in the adolescent brain. *Pediatrics.* 2009; 124:964–972.
- Neville H, Coffey S, Holcomb P, Tallal P. The neurobiology of sensory and language processing in language impaired children. *J Cogn Neurosci.* 1993; 5:235–253. [PubMed: 23972156]
- Nosarti C, Giouroukou E, Healy E, Rifkin L, Walshe M, Reichenberg A, Chitnis X, Williams SC, Murray RM. Grey and white matter distribution in very preterm adolescents mediates neurodevelopmental outcomes. *Brain.* 2008; 131:205–217. [PubMed: 18056158]
- Ortiz-Mantilla S, Choudhury N, Leever H, Benasich AA. Understanding language and cognitive deficits in very low birth weight children. *Dev Psychobiol.* 2008; 50:107–126. [PubMed: 18286580]

- Paterson SJ, Heim S, Friedman JT, Choudhury N, Benasich AA. Development of structure and function in the infant brain: Implications for cognition, language and social behaviour. *Neurosci Biobehav Rev.* 2006; 39:1087–1105. [PubMed: 16890291]
- Rice D, Barone S. Critical periods of vulnerability in the developing nervous system: Evidence from human and animal models. *Perspect.* 2000; 31:511–533.
- Robin D, Tomblin JB, Kearney A. Auditory temporal pattern learning in children with severe speech and language impairment. *Brain Lang.* 1989; 36:604–613. [PubMed: 2720373]
- Silbereis JC, Huang EJ, Back SA, Rowitch DH. Towards improved animal models of neonatal white matter injury associated with cerebral palsy. *Disease Models & Mechanisms.* 2010; 31:678–688. [PubMed: 21030421]
- Sininger YS, Abdala C, Cone-Wesson B. Auditory threshold sensitivity of the human neonate as measured by the auditory brainstem response. *Hear Res.* 1997; 104:27–38. [PubMed: 9119764]
- Slaghuys WL, Lovegrove WJ, Davidson JA. Visual and language processing deficits are concurrent in dyslexia. *Cortex.* 1993; 29:1–15.
- Stein J. Dyslexia-impaired temporal information processing? *Ann N Y Acad Sci.* 1993; 682:83–86. [PubMed: 8323162]
- Stein J. The magnocellular theory of developmental dyslexia. *Dyslexia.* 2001; 7:12–36. [PubMed: 11305228]
- Steinman KJ, Gorno-Tempini ML, Glidden DV, Kramer JH, Miller SP, Barkovich AJ, Ferriero DM. Neonatal watershed brain injury on magnetic resonance imaging correlates with verbal IQ at 4 years. *Pediatrics.* 2009; 123:1025–1030. [PubMed: 19255035]
- Szalkowski, CE.; Fitch, RH. Candidate dyslexia susceptibility genes and disorders of neuronal migration: Behavioral effects of cortical dysgenesis in a rodent model. In: Girard, A.; Moreau, L., editors. *Neuronal Migration: Disorders, Genetic Factors and Treatment Options.* Nova Science Publishers; 2012.
- Szalkowski CE, Booker AB, Truong DT, Threlkeld SW, Rosen GD, Fitch RH. Knockdown of the Candidate Dyslexia Susceptibility Gene Homolog *Dyx1c1* in Rodents: Effects on Auditory Processing, Visual Attention, and Cortical and Thalamic Anatomy. *Dev Neurosci.* 2013 epub ahead of print.
- Tallal P. Auditory temporal perception, phonics and reading disabilities in children. *Brain Lang.* 1980; 9:182–198. [PubMed: 7363063]
- Tallal P, Stark RE, Mellits D. The relationship between auditory temporal analysis and receptive language development: Evidence from studies of developmental language disorder. *Neuropsychologia.* 1985; 23:527–534. [PubMed: 2412182]
- Tallal, P.; Miller, S.; Fitch, RH. Neurobiological basis of speech: A case for the preeminence of temporal processing. Temporal information processing in the nervous system: Special reference to dyslexia and dysphasia. In: Tallal, P.; Galaburda, AM.; Llins, RR.; Von Euler, C., editors. *Ann N Y Acad Sci.* Vol. 682. 1993. p. 27–47.
- Thomas M, Kamiloff-Smith A. Developmental disorders like cases of adult brain damage? Implications from connectionist modeling. *Behav Brain Res.* 2002; 25:727–750.
- Tritsch NX, Bergles DE. Developmental regulation of spontaneous activity in the mammalian cochlea. *J Neurosci.* 2010; 30:1539–1550. [PubMed: 20107081]
- Van Lierde KM, Roeyers H, Boerjan S, De Groote I. Expressive and receptive language characteristics in three year old preterm children with extremely low birth weight. *Folia Phoniatr Logop.* 2009; 61:296–299. [PubMed: 19776605]
- Vannucci RC, Vannucci SJ. A model of perinatal hypoxic-ischemic brain damage. *Ann N Y Acad Sci.* 1997; 19 (835):234–249. [PubMed: 9616778]
- Vannucci RC, Vannucci SJ. Perinatal hypoxic-ischemic brain damage: evolution of an animal model. *Dev Neurosci.* 2005; 27:81–86. [PubMed: 16046840]
- Van Den Tweel E, Kavelaara A, Lombardi S, Nijboer CHA, Groenendaal F, Van Bel F, Hieijnen CJ. Bilateral molecular changes in a neonatal rat model of unilateral hypoxic-ischemic brain damage. *Pediatr Res.* 2006; 59:343–349. [PubMed: 16492969]
- Volpe J. Neurobiology of periventricular leukomalacia in the premature infant. *Pediatr Res.* 2001; 50:553–562. [PubMed: 11641446]

- Volpe J. Brain injury in premature infants: A complex amalgam of destructive and developmental disturbances. *Lancet Neurol.* 2009; 8:110–124. [PubMed: 19081519]
- Workman AD, Charvet CJ, Clancy B, Darlington RB, Finlay BL. Modeling transformations of neurodevelopmental sequences across mammalian species. *J Neurosci.* 2013; 33:7368–7383. [PubMed: 23616543]
- Witton C, Talcott J, Hansen P, Richardson A, Griffiths T, Rees A, Stein JF, Green GG. Sensitivity to dynamic auditory and visual stimuli predicts nonword reading ability in both dyslexic and normal readers. *Curr Biol.* 1998; 8:791–797. [PubMed: 9663387]
- Zhang LI, Bao S, Merzenich MM. Persistent and specific influences of early acoustic environment on primary auditory cortex. *Nat Neurosci.* 2001; 4:1123–1130. [PubMed: 11687817]

Highlights

P7 neonatal HI injured rodent show persitent RAP deficits. P3 HI injured rodents show a recovery of function on the same RAP task

P7 HI show a shift towards smaller cells in the MGN in post mortem tissue. P3 HI animals do not show this shift.

P7 HI animals show a significant decrease in the number of cells in the MGN. P3 HI injured animals do not show any decrease.

Representative Photographs of the MGN

Figure 1.

Microphotographs of MGN

(A) Microphotograph of the nissl-stained MGN as outlined with Microbrightfield software using a 2.5X objective. (B) Microphotograph of nissl stained MGN cells using a 100X objective. (C) Counting frame overlaid on the 100X objective MGN image via Microbrightfield software.

Figure 2.

Number of cells per section.

A one-way ANOVA revealed a significant Treatment effect in the right MGN for the number of cells per section [$F(2,39) = 14.423$, $p < .001$]. Follow-up independent samples t-test revealed a significant difference between P7 HI and shams ($t(24) = 5.445$, $p < .001$) and between P7 HI and P3 HI ($t(26) = 4.540$, $p < .001$). There was no difference between P3 HI and sham ($t(24) = .104$, $p > .05$). There was no Treatment effect in the left MGN [$F(2,39) = .421$, $p > .05$].

Figure 3. Cumulative Probability

(A) A Kruskal-Wallis non-parametric test revealed a significant Treatment effect between P3 HI, P7 HI and shams in the left MGN ($H(2) = 10.655$, $p < .005$). Follow up two-sample Kolmogorov-Smirnov analysis revealed significant differences between P7 HI and sham ($p < .001$) with P7 HI showing a shift towards larger cells. There was no difference between P3 HI and shams ($p > .05$) or P3 HI or P7 HI ($p > .05$). (B) A Kruskal-Wallis non-parametric test revealed a significant Treatment effect between P3 HI, P7 HI and shams in the right MGN ($H(2) = 86.233$, $p < .001$). Follow up two-sample Kolmogorov-Smirnov analysis revealed significant differences between P7 HI and sham ($p < .003$), and P3 HI and P7 HI ($p < .001$) with P7 HI showing a shift towards smaller cells compared to both groups. There was also a significant difference between P3 HI and shams ($p < .001$) with P3 HI showing a shift towards larger cells compared to shams.

Table 1

(A) Average Cell Area - Table of means for the left and right MGN. (B) Number of Cell per Section - Table of means for the left and right MGN.

Average Cell Area		
A	Right MGN	Left MGN
P3 HI, n=14	81.9456	82.5139
P7 HI, n=14	65.5892	80.4919
Sham, n=12	77.5573	76.9701

Number of Cells per Section		
B	Right MGN	Left MGN
P3 HI, n=14	30.4243	28.8229
P7 HI, n=14	19.1643*	30.9186
Sham, n=12	30.1425	29.8408