CORRESPONDENCE

Nosocomial Infection and Antibiotic Use: A Second National Prevalence Study in Germany

by Dr. rer. medic. Michael Behnke, Dr. med. Sonja Hansen, Dr. med. Rasmus Leistner, Dipl.-Inform. Luis Alberto Peña Díaz, Stud.-Inform. Alexander Gropmann, Dr. rer. nat. Dorit Sohr, Prof. Dr. med. Petra Gastmeier, Dr. med. Brar Piening in volume 38/2013

Controversial Measures

Alarming news about nosocomial—hospital acquired—infections have unsettled all participants in the healthcare system, met with enormous interest among the public, and triggered substantial countermeasures. Comprehensive disinfection and early administration of antibiotics serve as the generalized response to the problem, although it is especially these that have equally been considered as its cause. Screening all hospital patients for pathogens posing a special risk contributes further to the explosion of the associated costs.

Questions regarding hygiene have in recent years also included quality management in doctors' practices. Disinfection measures for door handles, exercise bikes, and ultrasound equipment have been recommended, although no scientific basis exists for this whatsoever. Except for special cases, the wearing of surgical gloves when performing phlebotomy in the surgery also ranks among the controversial measures. In addition to the enormous costs owing to the seeming prophylaxis, allergies and skin problems when wearing gloves, comprehensive disinfection resulting in selection of pathogens, and antibiotic resistance are among the problems of the future (1–3).

I have run my private practice for cardiopulmonary medicine for 15 years now, and not a single case of “nosocomial” infection has occurred. This is in spite of the fact that we treat patients with sputum smear positive tuberculosis and tumors, as well as a large number of multimorbid patients. Fortunately, the occupational health physician responsible for the practice has left it to our own discretion whether we wish our staff members to wear gloves when performing phlebotomy or to undertake comprehensive disinfection of the practice equipment. The evidence on which recommendations considering the actual benefit of such hygiene standards are usually based seems rather scarce as far as doctors' private practices are concerned. DOI: 10.3238/arztebl.2014.0066a

REFERENCES

1. Turner S, McNamee R, Agius R, Wilkinson SM, Carder M, Stockes SJ: Evaluating interventions aimed at reducing occupational exposure to latex and rubber glove allergens. *Occup Environ Med* 2012; 69: 925–31.
2. Carlet J, Jalier V, Harbarth S, Voss A, Goossens H, Pittet D: Ready for a world without antibiotics? The Pensières Antibiotic Resistance Call to Action. *Antimicrob Resist Infect Control* 2012; 1: 11.
3. Hsueh PR, Chen WH, Luh KT: Relationships between antimicrobial use and antimicrobial resistance in Gram-negative bacteria causing nosocomial infections from 1991–2003 at a university hospital in Taiwan. *Int J Antimicrob Agents* 2005; 26: 463–72.

4. Behnke M, Hansen S, Leistner R, et al.: Nosocomial infection and antibiotic use: A Second National Prevalence Study in Germany. *Dtsch Arztebl Int* 2013; 110(38): 627–33.

Prof. Dr. med. Damian Franzen
Köln
prof-dr-franzen@damianfranzen.de

In Reply:

Although comprehensive data on the prevalence of nosocomial infection and the transmission of multiresistant pathogens in hospitals are available in Germany, data are scarce on infections associated with outpatient medical treatment. This has two major reasons: on the one hand, the risk of infection in the outpatient setting is estimated to be much lower, because the patients—except for those having outpatient dialysis, endoscopies, or outpatient surgery—experience very few invasive diagnostic and therapeutic measures. On the other hand, it is methodologically complex to follow up the cohorts of patients after outpatient treatment in a systematic fashion, in order to identify such infections.

This is also the reason why there are hardly any intervention studies of medically sensible preventive measures in the outpatient setting; for this reason, preventive recommendations from the inpatient setting are usually transferred to the outpatient setting, which in our opinion is not necessarily a perfect match for the specific problems that may occur. However, certain preventive measures undoubtedly have great importance in the outpatient setting too. Hand disinfection is without doubt one of these. Many doctors working in the outpatient setting have not yet accepted the indications recommended by the World Health Organization (WHO) for hand disinfection for themselves and therefore do not regularly implement them. This is confirmed by data from the “AKTION Saubere Hände [national hand hygiene campaign in Germany]”: 622 German hospitals provided data on their consumption of hand disinfectants, whereas only 40 doctors' practices participated in the campaign. The average use of alcohol based hand disinfectant per treated patient in the practices was only 3.8 mL per case, barely more than is required for disinfecting hands once for every patient. The participating practices showed wide variation in this respect.

DOI: 10.3238/arztebl.2014.0066b

REFERENCES

1. Behnke M, Hansen S, Leistner R, et al.: Nosocomial infection and antibiotic use: A Second National Prevalence Study in Germany. *Dtsch Arztebl Int* 2013; 110(38): 627–33.

Dr. rer. medic. Michael Behnke
Institut für Hygiene und Umweltmedizin
Charité Berlin
michael.behnke@charite.de

Conflict of interest statement

The authors of both contributions declare that no conflict of interest exists.