CORRESPONDENCE

Common Causes of Poisoning—Etiology, Diagnosis, and Treatment

by Dr. med. Dieter Müller, Dipl. Chem. Dr. rer. nat. Herbert Desel
in volume 41/2013

Gastric Lavage in Cases of Poisoning

Once again a claim is made that gastric lavage is not safe and should be considered only in life-threatening cases and within minutes after the ingestion (1). This “vagueness” demands a closer look at the relevant reference literature.

The cited consensus article (2) updates the data on the basis of animal studies, studies of volunteers, experimental studies in patients, case studies, and clinical studies. Animal studies certainly have nothing in common with the clinical setting. Interestingly, several case reports showed positive outcomes for gastric lavage and endoscopy. With regard to clinical studies: the cited studies are from 1942 to 1966 and did not consider the intoxication dose and the factor of time. Further citations (review of 56 small and heterogeneous, exclusively Chinese, studies) produce a controversial picture. Numerous studies reporting the known possible complications of gastric lavage were mentioned. An unprotected respiratory tract is listed as a contraindication. The role of endoscopy in detoxification (which can be undertaken without protecting the respiratory tract) is not mentioned at all. In spite of all this, six out of 11 poison information centers supported gastric lavage in an international study using the example of acetaminophen (paracetamol) intoxication (3). But enough about the evidence. Recommendations have been copied from one another for years. In cases of life-threatening intoxication, all possible options for detoxification should be considered. Gastric lavage is one such option. In such a situation, a benefit not supported by controlled studies is not a counterargument. The factor of time is no argument either, in substances with an enterohepatic circulation. We have repeatedly encountered situations where the lack of, or delay in, gastric lavage resulted in poor outcomes. Especially from the perspective of continuing medical education, a more careful differentiation of the facts is needed in review articles and in the advice provided by poison information centers.

DOI: 10.3238/arztebl.2014.0100a

REFERENCES

1. Müller D, Desel H: Common causes of poisoning-etiology, diagnosis and treatment. *Dtsch Arztebl Int* 2013; 110: 690–700.
2. Benson BE, Hoppu K, Troutman WG, et al.: Position paper update: gastric lavage for gastrointestinal decontamination. *Clin Toxicol (Phila)* 2013; 51: 140–6.
3. Good AM, Kelly CA, Bateman DN: Differences in treatment advice for common poisons by poisons centres—an international comparison. *Clin Toxicol (Phila)* 2007; 45: 234–9.

Dr. med. Christian Marx

Innere Medizin/Nephrologie, Notfallmedizin, Nordhausen
c.marx@dialyse-nordhausen.de

Dr. med. Manfred Marx

Innere Medizin/Nephrologie, Internistische Intensivmedizin, Nordhausen

Conflict of interest statement

The authors declare that no conflict of interest exists.

In Reply:

We thank Marx and Marx for their critical response to the consensus article on gastric lavage we cited in our CME article. Their comment is likely to find agreement with many readers.

Especially experienced colleagues will remember successful cases of gastric lavage, which in the past was often applied in treatment of oral intoxication. We also reviewed many impressive individual case reports. It seems obvious that the removed poison cannot harm the patient any longer—and that gastric lavage represents an effective treatment strategy.

On the other hand, however, we have received many case reports of poor outcomes after gastric lavage.

The consensus article under criticism was compiled by toxicologists representing two medical societies, which conducted a careful and systematic evidence-based evaluation including various sources. Their goal was to describe the best therapeutic recommendation for all poisoned patients. The position paper on gastric lavage was reviewed several times by the working groups—newly formed each time—on the basis of recent literature and was confirmed almost without changes. The recommendation to perform gastric lavage only within 60 minutes after ingestion of a life-threatening dose of a toxic agent is based on a consensus that has remained stable for 16 years.

The cited article by Good et al. was an important contribution to the discussion. The data collected in 2002 about inconsistent advice given by poisons information centers in different countries—in spite of published recommendations from medical specialty societies—have underlined the need for harmonization/standardization of the treatment recommendations. This was also the intention of our article.

Poisons information centers provide advice if gastric lavage is under consideration. They register case reports to be added to the continuing discussion of the subject and may serve as a basis for future alignments of therapeutic recommendations.

DOI: 10.3238/arztebl.2014.0100b

REFERENCES

1. Müller D, Desel H: Common causes of poisoning-etiology, diagnosis and treatment. *Dtsch Arztebl Int* 2013; 110: 690–700.

Dr. med. Dieter Müller

Dipl. Chem. Dr. rer. nat. Herbert Desel
University Medical Center Göttingen
GIZ-Nord Poisons Center
dmueller@giz-nord.de

Conflict of interest statement

Dr Müller declares that no conflict of interest exists.

Dr Desel has received payment for serving as a consultant for the International Association for Soaps, Detergents and Maintenance Products (AISE, Brussels) and for expert opinions for Henkel. AISE and Industrieverband Körperpflege und Waschmittel (German Personal Care Products and Detergents Industry Association, Frankfurt/Main) are sponsoring the multicenter research project MAGAM II coordinated by GIZ-Nord Poisons Center, managed by Dr Desel.