6. Grevitt M, Khazim R, Webb J, Mulholland R, Shepperd J (1997) The Short Form-36 health survey questionnaire in spine surgery. *J Bone Joint Surg [Br]* 79B: 48–52
7. Main CJ (1983) The modified somatic perception questionnaire. *J Psychosom Res* 27: 503–514
8. Main CJ, Wood PL, Hollis S, Spanswick CS, Waddell G (1992) The distress and risk assessment method – a simple patient classification to identify distress and evaluate the risk of poor outcome. *Spine* 17: 41–52
9. Parker H, Woods PLR, Main CJ (1995) The use of the pain drawing as a screening measure to predict psychological distress in chronic low back pain. *Spine* 20: 236–243
10. Patrick DL, Deyo RA, Atlas SJ, Singer DE, Chapin A, Keller RB (1995) Assessing health related quality of life in patients with sciatica. *Spine* 20: 1899–1909
11. Polatin PB, Kirney RK, Gatchel RJ, Lillo E, Mayer TG (1985) Psychiatric illness and chronic low back pain. The mind and the spine – which goes first? *Spine* 10: 932–936
12. Ransford AO, Cairns D, Mooney V (1976) The pain drawing as an aid to the psychologic evaluation of patients with low back pain. *Spine* 1: 127–134
13. Turner RS, Lerding WC (1985) Correlation of the MMPI with lumbosacral spine fusion results: a prospective study. *Spine* 10: 932–936
14. Waddell G, Kummell EG, Lotto WN et al (1979) Failed lumbar disc surgery and repeat surgery following industrial injuries. *J Bone Joint Surg [Am]* 61A: 201–207
15. Waddell G, Main CJ, Morris EW, Di Paola M, Gray ICM (1984) Chronic low back pain, psychological distress and illness behaviour. *Spine* 9: 209–213
16. Wiltse LL, Rocchio PD (1975) Preoperative psychological tests as predictors of success of chemonucleolysis in the treatment of the low back pain syndrome. *J Bone Joint Surg [Am]* 57A: 478–483
17. Zdeblick TA (1995) The treatment of degenerative lumbar disorders – a critical review of the literature. *Spine* 20: 126–137 S
18. Zung WWK (1965) A self-rating depression scale. *Arch Gen Psychiatry* 32: 63–70

REVIEWER'S COMMENT

The authors present results from a specialist back clinic in a secondary care setting, finding 35 patients (28%) to be psychologically distressed. This figure is similar to the 23% found by Greenough and Fraser [1] and the 29% found by Main et al. [2]. The experienced clinicians in the study by Grevitt et al., using a subjective evaluation, diagnosed distress in only 13 patients (10%) and were significantly more likely to miss depression than somatisation ($P < 0.05$).

This important study tells clinicians that relying on subjective assessment alone will lead to an under-diagno-

sis of distress (an extremely important prognostic factor in the treatment of low back pain) and particularly an under-diagnosis of depression, which is a treatable condition. Failure to diagnose depression may be an aetiological factor in failed back surgery. All clinicians involved in the treatment of low back pain should use an instrument to formally assess psychological distress in their patients.

References

1. Greenough CG, Fraser RF (1991) Comparison of eight psychometric instruments in unselected patients with back pain. *Spine* 16: 1068–1074
2. Main CJ, Wood PLR, Hollis S, Spanswick CC, Waddell G (1992) The Distress and Risk Assessment Method. A simple patient classification to identify distress and evaluate risk of poor outcome. *Spine* 17: 42–52

C. G. Greenough
Consultant Orthopaedic Surgeon
Middlesbrough General Hospital
Ayresome Green Lane
Middlesbrough, Cleveland TS5 5AZ, UK