Staphylococcus aureus colonization in Brazilian children

Marcelo J. Mimica^{1,*}, Maria Fernanda Bádue-Pereira²

To the editor,

We read with interest the paper authored by Preotescu LL and Streinu-Cercel O,¹ and would like to input more data into their brief review on the subject of *Staphylococcus aureus* colonization in children. We have previously performed and published² a colonization study, from March to November 2009, in 197 children in São Paulo, Brazil. Samples were obtained using swabs from both the anterior nares and the oropharynx of each individual.

The overall prevalence of *S. aureus* was 65% and prevalence of methicillin-resistant isolates was 5.9%. In 51 patients (25.9%), *S. aureus* was detected in both sites. Oropharyngeal colonization without nasal colonization occurred in 42 cases (21.3%), whereas nasal colonization without colonization of the oropharynx was the case for 35 children (17.8%). Of the eleven MRSA isolates, six (3.2%) were found only in the oropharynx and three (1.6%) only in nares.

Our results have shown, in addition to a high prevalence of colonization, the importance of investigating other sites in addition to the anterior nares.

References

1. Preotescu LL, Streinu-Cercel O. Prevalence of nasal carriage of *S aureus* in children. GERMS. 2013;3:49-51. [\[CrossRef\]](#) [\[PubMed\]](#) [\[FullText\]](#)
2. Bádue Pereira MF, Mimica MJ, de Lima Bigelli Carvalho R, Scheffer DK, Berezin EN. High rate of *Staphylococcus aureus* oropharyngeal colonization in children. J Infect. 2012;64:338-40. [\[CrossRef\]](#) [\[PubMed\]](#)

Please cite this article as:
Mimica MJ, Bádue-Pereira MF. *Staphylococcus aureus* colonization in Brazilian children. GERMS. 2014;4(1):22. doi: 10.11599/germs.2014.1051

Received: 28 February 2014; accepted: 01 March 2014

¹MD, PhD, Division of Microbiology (Department of Pathology) and Division of Infectious Diseases (Department of Pediatrics), Santa Casa School of Medicine, São Paulo, Brazil; ²MD, MSc, Division of Infectious Diseases (Department of Pediatrics), Santa Casa School of Medicine, São Paulo, Brazil.

*Corresponding author: Marcelo J. Mimica, MD, PhD, Division of Microbiology (Department of Pathology) and Division of Infectious Diseases (Department of Pediatrics), Santa Casa School of Medicine, Rua Cesário Motta Jr., 112, São Paulo, Brazil. mjmimica@hotmail.com

Article downloaded from www.germs.ro

Published March 2014

© GERMS 2014

ISSN 2248 - 2997

ISSN - L = 2248 - 2997