Radium-223 exposes tumor cells within the skeleton to α radiation [8]. Other novel agents, such as cabozantinib, inhibit the tyrosine kinases c-Met and VEGFR2 pathways with an indirect effect on the AR [9]. Furthermore, many new treatment options are in development (e.g., ARN509, TAK-700, TOK-001, custirsén), targeting various pathways involved in CRPC.

Despite the increasing number of drugs in the armamentarium against PCa, the optimal sequence of drugs is not yet known and is likely to vary for individual patients given the heterogeneous nature of the disease. Cross-resistance is also an issue: docetaxel is a taxane targeting microtubules, but recent data suggest that the drug also affects the AR and may result in cross-resistance with androgen biosynthesis inhibitors [10]. In the future, biomarkers to better select effective agents will eventually lead to rational drug selection and drug combinations based on the biology of the disease. Many of the drugs used to treat PCa target more than one pathway, and several mechanisms of pathogenesis are at play within an individual patient.

We acknowledge that some patients find the term CRPC offensive. Dr. Luzzatto proposes to differentiate prostate cancer as androgen deprivation- (ADPRC-a) or deletion-resistant (ADPRC-b) PCa rather than CRPC. To further differentiate between the state of the AR would be confusing and could lead to several categories ADPRC-a, ADPRC-b, or even ADPRC-c and ADPRC-d when further AR alterations/mechanisms of action are detected. The challenge is to find an alternative term that is accurate, inoffensive, and not too much of a mouthful. To our knowledge, no good alternative had been proposed.

AXEL S. MERSEBURGER

Department of Urology and Urologic Oncology, Hannover Medical School, Hannover, Germany

JOAQUIM BELLMUNT

Medical Oncology Service, University Hospital del Mar-Institut de Investigació Mèdica, Barcelona, Spain

CHERYL JENKINS

Rocket Science Medical Communications, Wantage, United Kingdom

CHRIS PARKER

The Royal Marsden National Health Service Foundation Trust, Sutton, United Kingdom

JOHN M. FITZPATRICK

Department of Surgery, Mater Misericordiae Hospital University College and Irish Cancer Society, Dublin, Ireland

Disclosures

Axel S. Merseburger: AstraZeneca, Pfizer, Janssen, Astellas, Teva, Ipsen (C/A); AstraZeneca, Pfizer, Janssen, Astellas, Teva, Ipsen, Pierre Fabre, Novartis, GlaxoSmithKline (H); Bayer, Astellas, Janssen, Teva (RF); **Cheryl Jenkins:** Janssen (C/A); **Chris Parker:** Bayer, BNIT (C/A); Astellas, Bayer, Janssen, Sanofi-Aventis, Takeda (H); **John M. Fitzpatrick:** Sanofi, Astellas, Janssen (C/A); Sanofi, Astellas, Janssen (H). The other author indicated no financial relationships.

(C/A) Consulting/advisory relationship; (RF) Research funding; (E) Employment; (ET) Expert testimony; (H) Honoraria received; (OI) Ownership interests; (IP) Intellectual property rights/inventor/patent holder; (SAB) Scientific advisory board

REFERENCES

- Shah RB, Mehra R, Chinnaiyan AM et al. Androgen-independent prostate cancer is a heterogeneous group of diseases: Lessons from a rapid autopsy program. *Cancer Res* 2004;64:9209–9216.
- Aryee MJ, Liu W, Engelmann JC et al. DNA methylation alterations exhibit intraindividual stability and interindividual heterogeneity in prostate cancer metastases. *Sci Transl Med* 2013;5:169ra10.
- Heidenreich A, Bastian PJ, Bellmunt J et al. European Association of Urology: Prostate Cancer Guidelines. 2013. Available at http://www.uroweb.org/gls/pdf/09_Prostate_Cancer_LR.pdf. Accessed October 2013.
- Cookson MS, Roth BJ, Dahm P et al. Castration-resistant prostate cancer: AUA guideline. *J Urol* 2013;190:429–438.
- Horwich A, Parker C, de Reijke T et al. Prostate cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2013 [Epub ahead of print].
- Merseburger AS, Bellmunt J, Jenkins C et al. Perspectives on treatment of metastatic castration-resistant prostate cancer. *The Oncologist* 2013;18:558–567.
- Merseburger AS, Kuczyk MA, Wolff JM. Pathophysiology and therapy of castration-resistant prostate cancer [in German]. *Urologe A* 2013;52:219–225.
- Parker C, Nilsson S, Heinrich D et al. Alpha emitter radium-223 and survival in metastatic prostate cancer. *N Engl J Med* 2013;369:213–223.
- Lee RJ, Smith MR. Targeting MET and vascular endothelial growth factor receptor signaling in castration-resistant prostate cancer. *Cancer J* 2013;19:90–98.
- Fitzpatrick JM, de Wit R. Taxane mechanisms of action: Potential implications for treatment sequencing in metastatic castration-resistant prostate cancer. *Eur Urol* 2013 [Epub ahead of print].

<http://dx.doi.org/10.1634/theoncologist.2013-0395>

Tyrosine Kinase Inhibitors Adverse Events

In their recent article, Lenihan and Kowey (“Overview and Management of Cardiac Adverse Events Associated With Tyrosine Kinase Inhibitors” [1]) do not cite dasatinib among tyrosine kinase inhibitors (TKIs) with cardiovascular toxicity. Dasatinib is a TKI with markedly higher affinity for BCR-ABL kinase in comparison with imatinib, and it inhibits a large number of kinases including the Src family kinase. Dasatinib is approved as first- or second-line treatment for chronic myeloid leukemia. From 2009 on, several cases of pulmonary arterial hypertension (PAH) associated with dasatinib therapy have been reported [2, 3]. In 2012, Montani et al. [4]

published a series of results in nine patients in therapy with dasatinib and severe PAH, and they estimate that the incidence of severe/moderate PAH occurring in patients exposed to dasatinib in France is 0.45%. Dasatinib-associated PAH is described as a precapillary dose-related hypertension that is partially reversible after dasatinib withdrawal. Moreover, we evaluated four patients in therapy with dasatinib 100 mg/day for more than 12 months; in all patients, we observed an increased level of pulmonary pressure (CTCAE 4.03 grade I) in absence of any risk factors for pulmonary hypertension. It is our opinion that dasatinib should be included among TKIs with cardiovascular adverse effects for the risk of PAH. Every patient who needs dasatinib treatment

should be evaluated for pulmonary pressure, and a careful therapy strategy should be evaluated before the occurrence of severe/moderate PAH.

LEONARDO CAMPIOTTI

ANDREA MARESCA

LUIGINA GUASTI

ANNA MARIA GRANDI

Department of Clinical and Experimental Medicine, Università dell'Insubria, Varese, Italy

Disclosures

The authors indicated no financial relationships.

In Reply

Thank you for the opportunity to respond to the letter from Dr. Leonardo Campiotti et al. We very much appreciate the careful reading of our review [1] and thank Dr. Campiotti and his colleagues for their comments. In essence, Dr. Campiotti and colleagues suggest that dasatinib should be included in the list of tyrosine kinase inhibitors (TKIs) associated with cardiac adverse events. In particular, the authors cite the recently reported data that dasatinib is associated with pulmonary arterial hypertension (PAH) [2] in selected patients and furthermore that they are reporting similar findings in four patients. We agree with these authors that PAH is an important cardiovascular event and should be included in the complete description of all cardiovascular events that might be noted with TKI therapy. A thorough discussion of the incidence, monitoring, and treatment are certainly warranted in our opinion. However, we previously chose to limit the discussion in our review to the most common cardiac events encountered, which included systemic hypertension, left ventricular dysfunction, and QT prolongation [1]. The purpose of this review was to cover the major items in some detail but not be completely inclusive because that would have required a much longer article. With that concept in mind, we left out other cardiovascular events potentially encountered with TKI therapy that probably also deserve a careful discussion, including the potential development of atherosclerosis associated with nilotinib [3] or arterial thrombotic events

REFERENCES

1. Lenihan DJ, Kowey PR. Overview and management of cardiac adverse events associated with tyrosine kinase inhibitors. *The Oncologist* 2013;18:900–908.
2. Sano M, Saotome M, Urushida T et al. Pulmonary arterial hypertension caused by treatment with dasatinib for chronic myeloid leukemia—Critical alert. *Intern Med* 2012;51:2337–2340.
3. Orlandi EM, Rocca B, Pazzano AS et al. Reversible pulmonary arterial hypertension likely related to long-term, low-dose dasatinib treatment for chronic myeloid leukaemia. *Leuk Res* 2012;36:e4–e6.
4. Montani D, Bergot E, Günther S et al. Pulmonary arterial hypertension in patients treated by dasatinib. *Circulation* 2012;125:2128–2137.

<http://dx.doi.org/10.1634/theoncologist.2013-0397>

associated with ponatinib [4]. Thank you so much for your consideration of these points.

DANIEL J. LENIHAN

Vanderbilt Heart and Vascular Institute, Nashville, Tennessee, USA

PETER R. KOWEY

Jefferson Medical College and the Lankenau Institute for Medical Research, Main Line Health Heart Center, Wynnewood, Pennsylvania, USA

Disclosures

Daniel J. Lenihan: Roche, Astra Zeneca, Singulex (C/A); Acorda, Inc. (RF); **Peter R. Kowey:** Astra Zeneca, Johnson & Johnson, Novartis (H).

(C/A) Consulting/advisory relationship; (RF) Research funding; (E) Employment; (ET) Expert testimony; (H) Honoraria received; (OI) Ownership interests; (IP) Intellectual property rights/inventor/patent holder; (SAB) Scientific advisory board

REFERENCES

1. Lenihan DJ, Kowey PR. Overview and management of cardiac adverse events associated with tyrosine kinase inhibitors. *The Oncologist* 2013;18:900–908.
2. Montani D, Bergot E, Günther S et al. Pulmonary arterial hypertension in patients treated by dasatinib. *Circulation* 2012;125:2128–2137.
3. Quintás-Cardama A, Kantarjian H, Cortes J. Nilotinib-associated vascular events. *Clin Lymphoma Myeloma Leuk* 2012;12:337–340.
4. Cortes JE, Kim DW, Pinilla-Ibarz J et al. A phase 2 trial of ponatinib in Philadelphia chromosome-positive leukemias. *N Engl J Med* 2013;369:1783–1796.

<http://dx.doi.org/10.1634/theoncologist.2013-0433>