Published in final edited form as:

Photochem Photobiol. 2014 March ; 90(2): 448–454. doi:10.1111/php.12204.

The Contribution of Reactive Oxygen Species to the Photobleaching of Organic Fluorophores†

Qinsi Zheng^{#a,b}, Steffen Jockusch^{#c}, Zhou Zhou^a, and Scott C. Blanchard^{a,b,#}

^aDepartment of Physiology and Biophysics, Weill Medical College of Cornell University

^bTri-Institutional Training Program in Chemical Biology, 1300 York Avenue, New York, NY 10065, USA

^cDepartment of Chemistry, Columbia University, New York, NY 10027, USA

[#] These authors contributed equally to this work.

Abstract

Photo-excitation of fluorophores commonly used for biological imaging applications generates reactive oxygen species (ROS) which can cause bleaching of the fluorophore and damage to the biological system under investigation. In this study we show that singlet oxygen contributes relatively little to Cy5 and ATTO 647N photobleaching at low concentrations in aqueous solution. We also show that Cy5 generates significantly less ROS when covalently linked to the protective agents, cyclooctatetraene (COT), nitrobenzyl alcohol (NBA), or Trolox. Such fluorophores exhibit enhanced photostability both in bulk solutions and in single-molecule fluorescence measurements. While the fluorophores ATTO 647N and ATTO 655 showed greater photostability than Cy5 and the protective-agent linked Cy5 derivatives investigated here, both of ATTO 647N and ATTO 655 generated singlet oxygen and hydroxyl radicals at relatively rapid rates, suggesting that they may be substantially more phototoxic than Cy5 and its derivatives.

INTRODUCTION

Fluorescence microscopy has seen revolutionary advancements in both sensitivity and resolution over the past decade (1). Progress in this field is highlighted by recent developments in single-molecule and super-resolution fluorescence imaging techniques that have yielded new insights into a variety of biological systems in unprecedented detail (2-4). However, photo-induced degradation (photobleaching) of the fluorophores used remains a key obstacle that limits the temporal and spatial resolutions of imaging (5). Photo-induced fluorophore toxicity (phototoxicity) may also lead to unwanted perturbations to the biological system (6-8) that can obscure the signal of interest. Further advancements in fluorescence imaging would therefore be greatly aided by a deeper understanding of the mechanisms of fluorophore photobleaching and the development of fluorophores exhibiting slow photobleaching and low-toxicity.

Molecular oxygen plays a critical role in fluorophore photobleaching and phototoxicity (5, 7, 9, 10) as it can participate in two probable reactions with an excited fluorophore. First, energy transfer can occur from a fluorophore in the triplet excited state to molecular oxygen

†This paper is part of the Special Issue honoring the memory of Nicholas J. Turro

[#]To whom correspondence should be addressed: scb2005@med.cornell.edu.

SUPPLEMENTARY MATERIAL

Supplementary material has been posted online on DOI: xxxxx

leading to the formation of energetically excited singlet oxygen. Second, electron transfer can occur from the triplet fluorophore to molecular oxygen leading to the formation of a superoxide radical (Scheme 1) (9, 11). Singlet oxygen and superoxide radical, along with other oxidizing species formed subsequently (Scheme 1), are collectively termed reactive oxygen species (ROS) which can degrade fluorophores (10-14) and damage biomolecules in various contexts (6, 7, 15-17). However, the role of specific ROS in fluorophore photobleaching is not well understood. In particular, while it is widely held that singlet oxygen is a direct participant in the photobleaching pathway (9-11, 13, 18, 19), at least one early study provided evidence suggesting that it does not directly participate in the photobleaching of fluorescein isothiocyanate (FITC) (20). Here, we aim to elucidate the contribution of singlet oxygen to the photobleaching of Cy5 and ATTO 647N, two organic fluorophores commonly used in single-molecule and super-resolution measurements.

In order to mitigate fluorophore photobleaching and phototoxicity, molecular oxygen, normally at 0.3 mM in aqueous solution (21), can be depleted by enzymatic oxygen scavenging system (18, 22). Moreover, small-molecule protective agents (PA), such as cyclooctatetraene (COT), nitrobenzyl alcohol (NBA), and Trolox (23, 24), can be added to the imaging solution to mitigate the transient dark state caused by the triplet and radical state species (5, 10). Yet, the use of oxygen scavenging systems and PA in solution can be incompatible with, or perturbing to, biological systems, particularly for measurements involving lipid membranes (25) or live cells.

Recently, we demonstrated that the covalent linkage of a single COT, NBA and Trolox to cyanine fluorophores substantially reduces the frequency of transient dark state and the rate of photobleaching, thereby circumventing the requirements of PA and, in some cases, of oxygen scavenger system in solution (26, 27). The protective effect for Cy5-COT results from its capacity to reduce the lifetime of the triplet state by up to 50-fold through energy transfer quenching, while the distinct mechanism for Cy5-NBA and Cy5-Trolox was unknown (28). Here, we show that COT-, NBA- and Trolox-linked Cy5 derivatives (Cy5-PA) (Scheme 2) exhibit reduced rates of ROS generation compared to the parent Cy5 molecule. These findings imply that PA-conjugated fluorophores may exhibit reduced phototoxicity compared to the fluorophores commonly used for fluorescence imaging, and suggest that the Cy5-NBA and Cy5-Trolox in aqueous solution are also protected by triplet state quenching.

Materials and Methods

Materials

N-hydroxysuccinimide (NHS) ester activated fluorophores Cy5 (GE Healthcare), ATTO 647N (ATTO-TEC GmbH), and ATTO 655 (ATTO-TEC-GmbH) were used as received. Cy5-COT, Cy5-NBA, and Cy5-Trolox were synthesized as previously described (28).

Fluorophore bleaching experiments

Fluorophores (5 μ M or 26 μ M for Cy5; 5 μ M for Cy5-COT, Cy5-NBA, and Cy5-Trolox; 10 μ M for ATTO 655 and ATTO 647N) were dissolved in solvent (H_2O , D_2O , $CHCl_3$, $CDCl_3$, acetonitrile, or 100 mM Tris-acetate aqueous solution at pH 7.5). Samples (2ml) were examined in a 1 cm cuvette, illuminated with a 300W Tungsten halogen lamp, using an RG570 longpass filter to block short-wavelength light. Absorbance spectra of each fluorophore were measured using a UV-vis spectrometer (Agilent 8453). The decrease of the absorption was used to estimate the relative rate of fluorophore photobleaching over the first two minutes using a linear fit.

Singlet oxygen measurements

Singlet Oxygen Sensor Green (SOSG; Invitrogen) experiments were performed in aqueous buffer as previously described (29) in a 1 cm cuvette containing 2 mL of 100 mM Tris-acetate (pH=7.5), SOSG (2 μ M) fluorophore (5 or 10 μ M). Samples were illuminated for fixed periods of time with a 300 W Tungsten halogen lamp in conjugation with a RG570 long-pass filter. After each photolysis period, the fluorescence of SOSG was recorded over a range of 510-630 nm using 504 nm for excitation.

Due to the incompatibility of SOSG with organic solvent, 9,10-diphenylanthracene (DPA) was used to detect singlet oxygen in acetonitrile. DPA exhibits strong absorbance at 373 and 393 nm, and forms 9,10-endoperoxide, a compound that has no absorbance above 350 nm (Fig. S4a). The absorption of DPA was measured with a UV-vis spectrometer (Agilent 8453).

Single-molecule fluorescence measurements

Single-molecule fluorescence measurements were performed using a wide-field, prism-based total internal reflection fluorescence microscope as previously described (26, 28).

Fluorescence quantum yield measurements

Fluorescence spectra of solutions containing 0.7 μ M Cy5, Cy5-COT, Cy5-NBA, or Cy5-Trolox were recorded with a spectrofluorometer (Fluorolog 3, HORIBA Jobin Yvon), using an excitation wavelength of either 649 (aqueous Tris buffer, pH = 7.5) or 654 nm (acetonitrile). The fluorescence quantum yield was calculated using Cy5 in aqueous solution ($\phi = 0.20$ (30)) as a standard.

Fluorescence lifetime measurements

Fluorescence lifetimes were measured by time correlated single photon counting (TC-SPC) with an OB920 spectrometer (Edinburgh Analytical Instruments) in conjunction with a pulsed diode laser emitting at 659 nm. Fluorescence decay traces were monitored at 670 nm.

RESULTS AND DISCUSSIONS

The organic fluorophores Cy5, ATTO 647N, and ATTO 655 used for the present investigation represent the three main categories of fluorophore structures (cyanines for Cy5, carbopyronines for ATTO 647N, and oxazines for ATTO 655) that are widely used for single-molecule and super-resolution fluorescence imaging (31-33). Removal of molecular oxygen from the imaging solution substantially reduces the photobleaching rate of each fluorophore (20, 31), implying that photobleaching involves molecular oxygen. Here we show that the photobleaching rates of these dyes follow the order: Cy5>ATTO 647N>ATTO 655 (Fig. 1), inversely correlating with the oxidation potential of the fluorophores ($E_{OX,Cy5}=0.97V$, $E_{OX,ATTO647N}=1.11V$, $E_{OX,ATTO655}=1.31V$ (32, 34), where an increased oxidation potential implies a decreased propensity to be oxidized. Given the distinct structures and properties of these fluorophores (Scheme 2), this inverse correlation strongly supports that photobleaching is dominated by photo-induced oxidation reactions in the presence of oxygen.

To test the involvement of singlet oxygen in fluorophore photobleaching, we examined whether fluorophore photobleaching exhibits a solvent isotope effect (9). Given that the lifetime of singlet oxygen is extended by more than 20-fold in D₂O (68 μ s vs. 3.1 μ s in H₂O) (35), if fluorophore photobleaching is principally mediated by singlet oxygen, a substantial enhancement of the rate of photobleaching should be observed in D₂O (9). However, as observed previously for the organic fluorophore fluorescein isothiocyanate (FITC) (20), in

an initial test the rates of Cy5 (5 μ M) and ATTO 647N (10 μ M) photobleaching were only slightly higher in D₂O than in H₂O (Fig. 1). By contrast, a significant solvent isotope effect was observed in aqueous conditions when the Cy5 concentration was increased 5 fold to 26 μ M (Fig. 2), consistent with the involvement of singlet oxygen in the photobleaching reaction. Experiments at high concentration were not possible for ATTO 647N due to aggregation of the fluorophore. The absence of a solvent isotope effect at low concentration, and its presence at high Cy5 concentration, can be explained by the fact that encounter rate between singlet oxygen and fluorophore increases with increasing fluorophore concentration. At low fluorophore concentrations, singlet oxygen decays much faster than it encounters and reacts with fluorophores. Under such conditions singlet oxygen contributes negligibly to Cy5 photobleaching compared to other photobleaching reactions (scheme 1). At high fluorophore concentration, singlet oxygen can react with neighboring fluorophores to contribute to photobleaching to a greater extent. Consistent with the model that singlet oxygen decays too rapidly in aqueous solutions to contribute meaningfully to the bleaching of the fluorophore that led to its generation, a significant solvent isotope was observed for Cy5 at 5 μ M in chloroform (CHCl₃) and deuterated chloroform (CDCl₃) (Fig. S1), where singlet oxygen has much longer lifetime (approximately 230 μ s in CHCl₃ and 7 ms in CDCl₃ (35)) than it has in water. The significant solvent isotope effects observed in water with high Cy5 concentration, and in chloroform, demonstrate that singlet oxygen can contribute to fluorophore photobleaching under certain conditions. In line with previous results (20), the absence of any significant of solvent isotope effect at low fluorophore concentration in aqueous solution, suggests that singlet oxygen contributes negligibly to fluorophore photobleaching under the conditions of most biological studies at the single-molecule scale.

To examine whether the PA-conjugated fluorophores produce less singlet oxygen, we attempted to spectroscopically measure singlet oxygen via its phosphorescence emission band at 1270 nm (36) upon fluorophore illumination. However, under the conditions of the experiment (5 μ M fluorophore at room temperature), the yield of singlet oxygen was too low to be reliably detected using this approach (data not shown). We therefore used Singlet Oxygen Sensor Green (SOSG) as an indirect probe of singlet oxygen generation. SOSG itself is weakly-fluorescent due to intramolecular electron transfer quenching and becomes highly fluorescent following cycloaddition of singlet oxygen (Fig. 3a) (29).

Initially, singlet oxygen and photobleaching resulting Cy5 illumination (>570nm) were tracked by monitoring fluorophore absorption and SOSG emission spectra as a function of illumination time (647 nm and 528 nm, respectively) (Fig. 3b, d). As expected, the observed increase of SOSG fluorescence emission paralleled the decrease of Cy5 absorption. The increase in SOSG fluorescence was confirmed to arise from its specific reaction with singlet oxygen by performing the same measurement in 90% deuterated water, where the lifetime of singlet oxygen is significantly extended (see discussion above). In contrast to the negligible solvent isotope effect observed for photobleaching of Cy5 at low concentrations (Fig. 1), the rate of SOSG fluorescence increase was enhanced by approximately 8 fold in solution in 90% D₂O (Fig.S2).

In parallel, the rates of singlet oxygen generation were also measured for Cy5-COT, Cy5-NBA, Cy5-Trolox, ATTO 655 and ATTO 647N. As observed for the experiment of Cy5, SOSG fluorescence increased upon illumination of each fluorophore (Fig. 3c, e). All three PA-conjugated fluorophores exhibited lower rates of singlet oxygen generation, which paralleled their increased photostability compared to Cy5. Specifically, Cy5-Trolox, Cy5-COT and Cy5-NBA exhibited 2.5-fold, 5-fold and 20-fold reductions in the rate of singlet oxygen generation, respectively (Fig. 3c; Table 1). By contrast, ATTO 655 exhibited rates of singlet oxygen generation that were similar to those observed for Cy5, while ATTO 647N

generated singlet oxygen at a rate that was approximately 5-fold greater than Cy5 (Fig. 3c). These data imply that Cy5-PA fluorophores may be significantly less phototoxic than Cy5, ATTO 647N, and ATTO 655 in biological contexts.

To further explore the generation of ROS, we examined the rates of hydroxyl radical ($\text{HO}\cdot$) formation using the probe Aminophenyl fluorescein (APF) (37). Upon reaction with hydroxyl radicals, APF converts to fluorescein, a highly fluorescent species (37) (Fig. S3a). Aqueous solutions containing APF and individual fluorophores showed marked increases in fluorescein fluorescence ($\lambda_{\text{max}} = 518 \text{ nm}$) as a function of illumination time (Fig. S3b). Consistent with their reduced rates of singlet oxygen generation, the PA-conjugated fluorophores (Cy5-COT, Cy5-NBA and Cy5-Trolox) exhibited significantly reduced rates of hydroxyl radical generation (Fig. S3c). These data provide supporting evidence that the Cy5-PA fluorophores may be less phototoxic than the Cy5, ATTO 647N, and ATTO 655.

To directly examine the performance of fluorophores in single-molecule fluorescence measurement in ambient oxygen conditions, each fluorophore was conjugated to a double-stranded DNA oligonucleotide and imaged under a total internal reflection fluorescence microscope (26). Visual inspection of fluorescence traces from individual molecules revealed that most of ATTO 647N and ATTO 655 molecules showed substantial transient dark states and intensity fluctuations, while Cy5, Cy5-COT, Cy5-NBA, and Cy5-Trolox rarely blinked or fluctuated (Fig. 4). By tracking the fluorescence of many single molecules (>1000) over time (Fig. 4), the fluorescence intensity and the photobleaching rates were quantified. To compare each fluorophore's photobleaching rate, μ (the average numbers of photons detected from individual fluorophore prior to photobleaching) was calculated. As anticipated from prior investigations (26), the Cy5-COT, Cy5-NBA, and Cy5-Trolox fluorophores exhibited significantly reduced rates of photobleaching (Table 1), emitting up to 4-fold more photons than Cy5 prior to photobleaching (Fig. 4). As previously reported (26), the COT-conjugated fluorophore displayed a roughly 30% increase in brightness relative to Cy5 (Fig. 4). By contrast, both Cy5-NBA and Cy5-Trolox conjugates investigated here were substantially dimmer than Cy5 ($\sim 30\%$ and 50% , respectively; Fig. 4), likely due to the electron transfer between the PA and Cy5 singlet excited state. The slow-photobleaching, non-blinking, and non-fluctuating properties of the Cy5 derivatives investigated here render them suitable for single-molecule quantitative measurements, such as fluorescence resonance energy transfer, anisotropy, and stoichiometry. By contrast, the fluctuating emission behaviors of ATTO 647N and ATTO 655 make them more suitable for single-molecule tracking.

In previous studies (28), we showed that Cy5-COT, but not Cy5-NBA and Cy5-Trolox, exhibits shortened triplet state lifetime compared to Cy5 (Table 1). However, our previous measurements were exclusively performed in acetonitrile. Here we have shown that each of the Cy5-PA fluorophores exhibit reductions in photobleaching rates and in ROS generation, implying that in aqueous solution Cy5-COT, Cy5-NBA, and Cy5-Trolox possess substantially shorter triplet state lifetime than Cy5. These data suggest that the effects of NBA and Trolox are strongly solvent-dependent: in aqueous solution they effectively quench the triplet state of Cy5, likely through an electron-transfer mechanism (38, 39), whereas in acetonitrile they do not quench the triplet state due to the solvent-dependent nature of the electron transfer where radical states form (9). The rate of electron transfer is known to be affected by solvent polarity (40). By contrast, COT can quench the Cy5 triplet state in both solvents because of the solvent-independent nature of triplet-triplet energy transfer (9, 28). This mechanism is supported by a further comparison of fluorophore properties in aqueous solution and in acetonitrile: the fluorescent lifetime, the fluorescent quantum yield, the photobleaching rate and the singlet oxygen generation rate for Cy5-NBA

and Cy5-Trolox were strongly solvent-dependent, while these properties for Cy5 and Cy5-COT were only weakly solvent-dependent (Table 1; Fig. S4).

CONCLUSION

The results presented suggest that the contribution of singlet oxygen to fluorophore photobleaching is negligible at low fluorophore concentrations ($<5 \mu\text{M}$) in aqueous buffers. Singlet oxygen does, however, play a significant role under solvent where its lifetime is significantly extended (e.g. chloroform) or at higher fluorophore concentrations ($>20 \mu\text{M}$) where the collision frequency with neighboring fluorophore molecules is greatly increased. These observations explain discrepancies in the literature regarding the role of singlet oxygen in fluorophore photobleaching (13, 20) and suggest that singlet oxygen is unlikely to contribute to fluorophore photobleaching under most *in vitro* single-molecule imaging conditions. Cy5 derivatives covalently linked to a protective agent exhibit enhanced photostability as a direct consequence of distinct mechanisms of triplet state quenching that result in lower rates of singlet oxygen and hydroxyl radical generation. Correspondingly, these fluorophores may also be less perturbing and exhibit reduced phototoxicity when used for live cell imaging applications.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the National Institutes of Health (GM098859-01A1 to SCB) and the National Science Foundation (CHE 11-11392 to SJ). We are grateful to Roger B. Altman and Daniel S. Terry for critical discussions and comments on the manuscript.

References

1. Lakowicz, J. Principles of Fluorescence Spectroscopy. Springer Science + Business Media, LLC; New York, NY: 2006.
2. Moerner WE. A Dozen Years of Single-Molecule Spectroscopy in Physics, Chemistry, and Biophysics. The Journal of Physical Chemistry B. 2002; 106:910–927.
3. Joo C, Balci H, Ishitsuka Y, Buranachai C, Ha T. Advances in Single-Molecule Fluorescence Methods for Molecular Biology. Annual Review of Biochemistry. 2008; 77:51–76.
4. Huang B, Bates M, Zhuang X. Super-Resolution Fluorescence Microscopy. Annual Review of Biochemistry. 2009; 78:993–1016.
5. Ha T, Tinnefeld P. Photophysics of Fluorescent Probes for Single-Molecule Biophysics and Super-Resolution Imaging. Annual Review of Physical Chemistry. 2012; 63:595–617.
6. Sheetz MP, Koppel DE. Membrane damage caused by irradiation of fluorescent concanavalin A. Proceedings of the National Academy of Sciences. 1979; 76:3314–3317.
7. Dixit R, Cyr R. Cell damage and reactive oxygen species production induced by fluorescence microscopy: effect on mitosis and guidelines for non-invasive fluorescence microscopy. The Plant Journal. 2003; 36:280–290. [PubMed: 14535891]
8. Dobrucki JW, Feret D, Noatynska A. Scattering of Exciting Light by Live Cells in Fluorescence Confocal Imaging: Phototoxic Effects and Relevance for FRAP Studies. Biophysical Journal. 2007; 93:1778–1786. [PubMed: 17416613]
9. Turro, NJ.; Ramamurthy, V.; Scaiano, JC. Modern Molecular Photochemistry of Organic Molecules. University Science Books; 2010.
10. Levitus M, Ranjit S. Cyanine dyes in biophysical research: the photophysics of polymethine fluorescent dyes in biomolecular environments. Q Rev Biophys. 2011; 44:123–51. [PubMed: 21108866]

11. Bonnett R, Martínez G. Photobleaching of sensitizers used in photodynamic therapy. *Tetrahedron*. 2001; 57:9513–9547.
12. Chen P, Li J, Qian Z, Zheng D, Okasaki T, Hayami M. Study on the photooxidation of a near-infrared-absorbing benzothiazolone cyanine dye. *Dyes and Pigments*. 1998; 37:213–222.
13. Byers GW, Gross S, Henrichs PM. Direct and Sensitized Photooxidation of Cyanine Dyes. *Photochemistry and Photobiology*. 1976; 23:37–43. [PubMed: 1265127]
14. Kuramoto N, Kitao T. The contribution of singlet oxygen to the photofading of triphenylmethane and related dyes. *Dyes and Pigments*. 1982; 3:49–58.
15. Vigers GP, Coue M, McIntosh JR. Fluorescent microtubules break up under illumination. *The Journal of Cell Biology*. 1988; 107:1011–1024. [PubMed: 3417772]
16. Davies MJ. Reactive species formed on proteins exposed to singlet oxygen. *Photochemical & Photobiological Sciences*. 2004; 3:17–25. [PubMed: 14743273]
17. Sies H, Menck CFM. Singlet oxygen induced DNA damage. *Mutation Research/DNAging*. 1992; 275:367–375.
18. Aitken CE, Marshall RA, Puglisi JD. An oxygen scavenging system for improvement of dye stability in single-molecule fluorescence experiments. *Biophys J*. 2008; 94:1826–35. [PubMed: 17921203]
19. Touthkine A, Nguyen D-V, Hahn KM. Merocyanine Dyes with Improved Photostability. *Organic Letters*. 2007; 9:2775–2777. [PubMed: 17583344]
20. Johnson GD, Davidson RS, McNamee KC, Russell G, Goodwin D, Holborow EJ. Fading of immunofluorescence during microscopy: a study of the phenomenon and its remedy. *J Immunol Methods*. 1982; 55:231–42. [PubMed: 6819318]
21. Montalti, M.; Credi, A.; Prodi, L.; Gandolfi, MT. *Handbook of photochemistry*. CRC/Taylor & Francis; Boca Raton: 2006.
22. Benesch RE, Benesch R. Enzymatic Removal of Oxygen for Polarography and Related Methods. *Science*. 1953; 118:447–448. [PubMed: 13101775]
23. Rasnik I, McKinney SA, Ha T. Nonblinking and long-lasting single-molecule fluorescence imaging. *Nat Meth*. 2006; 3:891–893.
24. Dave R, Terry DS, Munro JB, Blanchard SC. Mitigating Unwanted Photophysical Processes for Improved Single-Molecule Fluorescence Imaging. *Biophys J*. 2009; 96:2371–2381. [PubMed: 19289062]
25. Alejo, Jose L, Blanchard Scott C. Andersen Olaf S. Small-Molecule Photostabilizing Agents are Modifiers of Lipid Bilayer Properties. *Biophysical Journal*. 2013; 104:2410–2418. [PubMed: 23746513]
26. Altman RB, Terry DS, Zhou Z, Zheng Q, Geggier P, Kolster RA, Zhao Y, Javitch JA, Warren JD, Blanchard SC. Cyanine fluorophore derivatives with enhanced photostability. *Nature Methods*. 2012; 9:68–71. [PubMed: 22081126]
27. Altman RB, Zheng Q, Zhou Z, Terry DS, Warren JD, Blanchard SC. Enhanced photostability of cyanine fluorophores across the visible spectrum. *Nature Methods*. 2012; 9:428–9. [PubMed: 22543373]
28. Zheng Q, Jockusch S, Zhou Z, Altman RB, Warren JD, Turro NJ, Blanchard SC. On the Mechanisms of Cyanine Fluorophore Photostabilization. *The Journal of Physical Chemistry Letters*. 2012; 3:2200–2203. [PubMed: 22984636]
29. Gollmer A, Arnbjerg J, Blaikie FH, Pedersen BW, Breitenbach T, Daasbjerg K, Glasius M, Ogilby PR. Singlet Oxygen Sensor Green®: Photochemical Behavior in Solution and in a Mammalian Cell. *Photochemistry and Photobiology*. 2011; 87:671–679. [PubMed: 21272007]
30. Mujumdar RB, Ernst LA, Mujumdar SR, Lewis CJ, Waggoner AS. Cyanine dye labeling reagents: Sulfoindocyanine succinimidyl esters. *Bioconjugate Chem*. 1993; 4:105–111.
31. Roy R, Hohng S, Ha T. A practical guide to single-molecule FRET. *Nat Methods*. 2008; 5:507–16. [PubMed: 18511918]
32. Vogelsang J, Cordes T, Forthmann C, Steinhauer C, Tinnefeld P. Controlling the fluorescence of ordinary oxazine dyes for single-molecule switching and superresolution microscopy. *Proc Natl Acad Sci U S A*. 2009; 106:8107–12. [PubMed: 19433792]

33. Dempsey GT, Vaughan JC, Chen KH, Bates M, Zhuang X. Evaluation of fluorophores for optimal performance in localization-based super-resolution imaging. *Nature Methods*. 2011; 8:1027–1036. [PubMed: 22056676]
34. Vogelsang J, Kasper R, Steinhauer C, Person B, Heilemann M, Sauer M, Tinnefeld P. A reducing and oxidizing system minimizes photobleaching and blinking of fluorescent dyes. *Angew Chem Int Ed Engl*. 2008; 47:5465–9. [PubMed: 18601270]
35. Schweitzer C, Schmidt R. Physical Mechanisms of Generation and Deactivation of Singlet Oxygen. *Chemical Reviews*. 2003; 103:1685–1758. [PubMed: 12744692]
36. Khan AU, Kasha M. Direct spectroscopic observation of singlet oxygen emission at 1268 nm excited by sensitizing dyes of biological interest in liquid solution. *Proceedings of the National Academy of Sciences*. 1979; 76:6047–6049.
37. Setsukinai, K.-i.; Urano, Y.; Kakinuma, K.; Majima, HJ.; Nagano, T. Development of Novel Fluorescence Probes That Can Reliably Detect Reactive Oxygen Species and Distinguish Specific Species. *Journal of Biological Chemistry*. 2003; 278:3170–3175. [PubMed: 12419811]
38. Tinnefeld P, Cordes T. ‘Self-healing’ dyes: intramolecular stabilization of organic fluorophores. *Nat Meth*. 2012; 9:426–427.
39. Blanchard SC. Reply to “‘Self-healing’ dyes: intramolecular stabilization of organic fluorophores”. *Nat Meth*. 2012; 9:427–428.
40. Chandler, D. Electron transfer in water and other polar environments, how it happens. In: Berne, BJ.; Ciccotti, G.; Coker, DF., editors. *Classical and Quantum Dynamics in Condensed Phase Simulations*. World Scientific; Singapore: 1998. p. 25-49.

Figure 1. Photobleaching of Cy5 (5 μ M), ATTO 647N (10 μ M), and ATTO 655 (10 μ M) in air-saturated H_2O (red) or D_2O (blue), where k_B is the photobleaching rate.

Figure 2. Photobleaching of Cy5 at 26 μM in air-saturated H_2O (solid) or D_2O (open), where k_B is the photobleaching rate.

Figure 3.

(a) The mechanism of singlet oxygen ($^1\text{O}_2$) induced fluorescence of SOSG. (b) The emission spectra of SOSG upon illumination of Cy5. (c) $^1\text{O}_2$ generation, reported by the increase of SOSG fluorescence, for distinct fluorophores at different illumination time. The relative rates of $^1\text{O}_2$ generation compared to the Cy5 fluorophore are shown in parentheses. (d) The absorption spectrum of Cy5 observed at different illumination time. (e) The decrease of Cy5, Cy5-COT, Cy5-NBA, Cy5-Trolox, ATTO 647N, and ATTO 655 absorption at 647nm as a function of illumination time. Experiments were performed in 100 mM Tris-acetate buffered aqueous solution, pH=7.5.

Figure 4.

Representative single-molecule fluorescence traces for Cy5, Cy5-COT, Cy5-NBA, Cy5-Trolox, ATTO 647N, and ATTO 655 covalently linked to DNA oligonucleotides and imaged in air-saturated aqueous solution using a total internal reflection microscope under continuous laser excitation (641 nm). μ is the average number of detected photons before fluorophore photobleaching.

Scheme 1.

Some probable steps to ROS via the reactions between molecular oxygen ($^3\text{O}_2$) and the fluorophore. S_0 : fluorophore in ground state; S_1 : fluorophore in the singlet excited state; T_1 : fluorophore in the triplet state; R^+ : fluorophore in the radical cationic state.

Scheme 2.
Structures of the compounds used in this study.

Table 1

A summary of the data. 100mM Tris acetate buffer (pH=7.5) was used in all experiments in aqueous buffer. All of the rates were normalized to the corresponding numbers for Cy5.

Solvent		Cy5	Cy5-COT	Cy5-NBA	Cy5-Trolox
aqueous	Relative rate of single-molecule photobleaching	1±0.2	0.29±0.02	0.27±0.02	0.42±0.04
	Relative rate of ensemble photobleaching	1±0.2	0.45±0.03	0.24±0.04	0.78±0.06
	1 Relative rate of $^1\text{O}_2$ generation	1±0.05	0.18±0.01	0.05±0.01	0.41±0.01
	Relative rate of hydroxyl radical generation	1±0.03	0.43±0.01	0.235±0.004	0.367±0.009
	Fluorescence lifetime (ns)	0.92±0.02	0.95±0.02	0.59±0.02	0.76±0.03
	Fluorescence quantum yield	0.20 ^a	0.20	0.07	0.11
acetonitrile	Relative rate of ensemble photobleaching	1±0.08	0.50±0.01	1.0±0.1	0.9±0.1
	1 Relative rate of $^1\text{O}_2$ generation	1±0.08	0.57±0.06	0.92±0.06	0.86±0.06
	Fluorescence lifetime (ns)	1.11±0.03	1.18±0.05	0.93±0.04	1.02±0.03
	Fluorescence quantum yield	0.22	0.22	0.18	0.19
	Triplet state lifetime (μs)	63±3	1.1±0.1	62±3	60±4