Back pain in patients with degenerative spine disease and intradural spinal tumor: what to treat? when to treat?

David Bellut · Urs M. Mutter · Martin Sutter ·
Andreas Eggspuehler · Anne F. Mannion ·
François Porchet

Received: 28 January 2013 / Revised: 12 December 2013 / Accepted: 13 December 2013 / Published online: 31 December 2013
© Springer-Verlag Berlin Heidelberg 2013

Abstract

Background Back pain is common in industrialized countries and one of the most frequent causes of work incapacity. Successful treatment is, therefore, not only important for improving the symptoms and the quality of life of these patients but also for socioeconomic reasons. Back pain is frequently caused by degenerative spine disease. Intradural spinal tumors are rare with an annual incidence of 2–4/1,00,000 and are mostly associated with neurological deficits and radicular and nocturnal pain. Back pain is not commonly described as a concomitant symptom, such that in patients with both a tumor and degenerative spine disease, any back pain is typically attributed to the degeneration rather than the tumor.

Objective The aim of the present retrospective investigation was to study and analyze the impact of microsurgery on back/neck pain in patients with intradural spinal tumor in the presence of degenerative spinal disease in adjacent spinal segments.

Methods Fifty-eight consecutive patients underwent microsurgical, intradural tumor surgery using a standardized protocol assisted by multimodal intraoperative neuromonitoring. Clinical symptoms, complications and surgery characteristics were documented. Standardized questionnaires were used to measure outcome from the surgeon's and the patient's perspectives (Spine Tango Registry and Core Outcome Measures Index). Follow-up included clinical and neuroradiological examinations 6 weeks, 3 months and 1 year postoperatively.

Results Back/neck pain as a leading symptom and coexisting degenerative spine disease was present in 27/58 (47 %) of the tumor patients, and these comprised to group under study. Patients underwent tumor surgery only, without addressing the degenerative spinal disease. Remission rate after tumor removal was 85 %. There were no major surgical complications. Back/neck pain as the leading symptom was eradicated in 67 % of patients. There were 7 % of patients who required further invasive therapy for their degenerative spinal disease.

Conclusions Intradural spinal tumor surgery improves back/neck pain in patients with coexisting severe degenerative spinal disease. Intradural spinal tumors seem to be the only cause of back/neck pain more often than appreciated. In these patients suffering from both pathologies, there is a higher risk of surgical overtreatment than undertreatment. Therefore, elaborate clinical and radiological examinations should be performed preoperatively and the indication for stabilization/fusion should be discussed carefully in patients foreseen for first time intradural tumor surgery.

Parts of these results have been presented at the 14th EANS Congress 2011 in Rome, Italy.

D. Bellut (✉) · U. M. Mutter · F. Porchet
Spine Center, Schulthess Clinic Zürich, Lengghalde 2,
8008 Zürich, Switzerland
e-mail: dbellut@gmail.com

M. Sutter · A. Eggspuehler
Department of Neurology, Schulthess Clinic Zürich,
Lengghalde 2, 8008 Zürich, Switzerland

A. F. Mannion
Spine Center Division, Department of Research and
Development, Schulthess Clinic Zürich, Lengghalde 2,
8008 Zürich, Switzerland

Keywords Intradural spinal tumor · Degenerative disc disease · Neurinoma · Meningioma · Treatment strategies · COMI · Spine Tango · Outcome

Introduction

Back/neck pain is common in industrialized countries with a prevalence of up to 90 % [1]. It is the most frequently reported musculoskeletal problem and the third most common symptom of any kind making it one of the most common causes of work incapacity [1]. Successful treatment is hence important not only to minimize individual suffering but also for socioeconomic reasons. Underlying pathologies are mostly of degenerative origin and include spinal canal stenosis, facet joint arthrosis, degenerative disc disease, degenerative spondylolisthesis and degenerative scoliosis. Surgical treatment of back/neck pain typically includes endoscopic or microsurgical discectomy, spinal canal decompression and fusion with or without instrumentation, but even in well-selected patients, surgical treatment may fail in up to 32 % of patients [2].

Intradural spinal tumors are rare with an annual incidence of 2–4/1,00,000 and are mostly associated with neurological deficits and radicular and nocturnal pain [3, 4]. Back/neck pain is not commonly described as a concomitant symptom. Intradural spinal tumors can be classified as intramedullary or extramedullary. Intramedullary tumors account for around 20 % of intradural spinal tumors [4] and mainly comprise ependymomas (60 %) and astrocytomas (30 %). Extramedullary tumors account for up to 80 % of intradural tumors and mostly comprise schwannomas (30 %) and meningiomas (25 %) [4]. The treatment of choice in patients with intradural spinal tumor is open microsurgery to establish a histopathological diagnosis and to decompress the neural structures. In patients with benign tumors like meningioma (WHO I) or schwannoma (WHO I) a complete tumor resection should be aimed for to reduce the risk of tumor recurrence [5, 6].

In patients with intradural spinal tumor, back/neck pain is not commonly described as a concomitant symptom, such that in patients with both tumor and degenerative spine disease, any back/neck pain is typically attributed to the degeneration rather than the tumor.

The aim of the present retrospective investigation was to analyze the impact of microsurgery of intradural spinal tumor on back/neck pain in the presence of degenerative spinal disease in the same or adjacent spinal segments.

Patients and methods

Patient demographics

Fifty-eight patients undergoing 58 consecutive surgical procedures for intradural spinal tumor were included in this retrospective study. Detailed analysis of their preoperative symptoms, tumor characteristics and surgical procedures

was carried out. Fifty-seven of the 58 procedures were performed to address the tumor alone using various approaches. Twenty-seven of the 58 patients were diagnosed with intradural spinal tumor, back/neck pain as a leading symptom and degenerative spine disease in the same or adjacent segments based on neuroradiological findings. These 27 patients comprised the group under investigation in the present study.

There were 15 male (56 %) and 12 female (44 %) patients with a mean age of 57.4 ± 17.8 years. Three patients (11 %) had a history of spine surgery: one patient had undergone discectomy; another, tumor surgery; and the final one, single segment stabilization and fusion for degenerative spine disease prior to surgical tumor removal.

Pre- and early postoperative management

Patients were seen in the outpatients department and underwent clinical, neurological and electrophysiological examination as well as neuroradiological examination (MRI, X-rays). Patients were admitted to the hospital 1 day before surgery for neurological examination and final anaesthesiological check and clearance for surgery. After surgery patients were observed on an intermediate care unit for 24 h and afterwards transferred to the ward. Mobilization under physiotherapeutic observation started 1–2 days after surgery. Patients were hospitalized for 1 week before being discharged or transferred to a rehabilitation clinic, in the case of major neurological deficits.

Surgery and intraoperative multimodal neuromonitoring

Surgery was performed under general anesthesia and patients were put in a prone position after application of the neuromonitoring electrodes. The tumor segment was localized using fluoroscopy. Before skin incision patients received antibiotic prophylaxis. The surgical approach included hemilaminectomy or laminectomy and laminoplasty. To reduce the tumor volume a SöringTM ultrasound tissue ablation machine was used to minimize manipulation of the neural structures. Dura opening was always in the midline exposing both ends of the tumor and using sutures to keep the dura opening wide open. Part of the tumor mass was sent for histological examination. In all patients gross total resection of the tumor was intended.

Multimodal intraoperative monitoring was performed by two board-certified neurologists (third and fourth authors) with at least 14 years of neuromonitoring experience, using the protocol described by Sutter et al. [7, 8].

In the case of deterioration of the neurophysiological parameters, the surgery was modified to prevent severe neurological deficit. This included the elevation of blood

pressure, stopping of tumor removal, application of steroids and/or use of a different surgical approach.

Follow-up

Patients were followed up by board certified neurologists or neurosurgeons at the clinic's outpatient department at 6 weeks (6.3 ± 1.3 weeks), 4 months (4.8 ± 3.4 months) and 1 year (13.1 ± 7.6 months) after surgery. Further follow-up was carried out depending on clinical symptoms, histopathological findings and the existence of recurrent tumor. The follow-up at 6 weeks was a clinical follow-up only, whereas those at 3 months and 1 year also comprised clinical and neuroradiological (MRI) examinations.

Clinical symptoms, surgery characteristics and complications were evaluated from the surgeons' and the patients' perspectives using the Spine Tango Registry [9] and the Core Outcomes Measures Index (COMI) [10].

The Spine Tango registry and its accompanying documentation forms were first introduced in November 2002 by EuroSpine, in cooperation with the M.E. Müller Center at the University of Berne in Switzerland [9]. The aim was to standardize outcome measures in patients undergoing spinal surgery [11]. The Spine Tango registry Surgery and Follow-up forms contain questions regarding the surgical procedure, intraoperative and postoperative complications, underlying pathology, and neurological and clinical outcome from the perspective of the treating surgeon.

The COMI is a short, multidimensional outcome instrument, with excellent psychometric properties, used in monitoring the outcome of spinal surgery from the patient's perspective. The items that it contains were first described in the literature in 1998 [12] for use in individuals with back pain. The COMI has one question each on back pain intensity, leg/buttock pain intensity, function, symptom-specific well being, general quality of life, work disability and social disability, and is scored as a 0–10 index. The questionnaire has been found to be feasible to implement on a prospective basis in routine practice [10], and as responsive as many longer spine outcome questionnaires [13]. The shortness of the COMI and its multidimensional nature make it an attractive option to comprehensively assess all patients within a given Spine Centre and hence avoid selection bias in reporting outcomes [10].

The Spine Tango Surgical form was completed by the surgeon over the time period from admission to discharge and the Spine Tango Follow-up form was completed at all postoperative follow-up examinations. The COMI questionnaire was completed by the patient preoperatively and at 3, 12 and 24 months, postoperatively.

Table 1 Patient demographic and clinical data

Variables	Mean $\pm$ SD, or absolute number (%)
Number of patients/surgical procedures	58
Study group: patients with intradural spinal tumor, DSD and back pain	27
Number of surgical procedures	27
Sex (male/female)	15/12 (56/44 %)
Mean age in years	57.4 ± 17.8
Mean duration of hospitalization in days	9.7 ± 3.1
Localization of tumor	10 CS (37 %), 7 ThS (26 %), 10 LS (37 %)
Histopathological diagnosis	
Ependymoma (WHO I)	5 (19 %)
Ependymoma (WHO II)	2 (7 %)
Astrocytoma (WHO I)	1 (4 %)
Schwannoma (WHO I)	9 (33 %)
Meningioma (WHO I)	6 (22 %)
Other (lymphoma, metastasis, hemangiopericytoma)	4 (15 %)
Mean tumor volume (mm^3)	22,324 (1,437–1,04,720)
Previous surgery	3 (11 %)

Imaging and classification of spinal degeneration

Preoperative MRI imaging was performed using different MRI scanners (all Siemens, Philips or GE scanners with a magnetic field strength of at least 1.5 T). Postoperative imaging was performed using a Philips Achieva 3 T MRI scanner. Imaging was analyzed independently and blinded to the clinical outcome using standardized software (picture archiving and communication system, PACS) by board certified radiologists.

Spinal degeneration was classified based on the preoperative imaging examinations. Spondylarthritis was classified according to the Fujiwara classification [22], disc degeneration was classified according to the Pfirrmann classification [23] and vertebral body marrow changes were classified according to the Modic classification [24]. Severe degeneration of a spinal segment was stated if type 3 or 4 of Fujiwara, type 1–3 of Modic and Grad 4 or 5 of Pfirrmann classification was present.

Statistical analysis

The statistical analysis was performed using Microsoft Excel (Version 2003) and SPSS software (Version 16.0). Tumor volume was calculated based on the diameter method ($\text{Tumor volume} = 4/3 \times \text{Pi} \times 1/2x \times 1/2y \times 1/2z$), where x , y and z are the maximum diameters in the three axis.

Table 2 Surgical procedures and preoperative neuroradiological findings

Variables	Mean $\pm$ SD, or absolute number (%)
Segments per surgical procedure	1.44
Gross total tumor resection	18 (67 %)
Subtotal tumor resection	7 (26 %)
Extended biopsy	2 (7 %)
Tumor volume (mm ³)	22,324 (1,437–1,04,720)
Mean diameter (mm)	39.2 $\pm$ 44.4
Same/adjacent segment degeneration	27 (100 %)
Spinal canal stenosis	5 (19 %)
Spondylarthrosis	24 (89 %)
Discopathy/osteochondrosis	24 (89 %)

Results

Surgical procedures, tumor characteristics and pathology

Overall, 58 patients underwent surgery for intradural spinal tumor between 03/2006 and 03/2011. Of these, 27 fulfilled the inclusion criteria for the present study; there were 15 men (56 %) and 12 women (44 %) with a mean age of 57.4 ± 17.8 years (Table 1). The tumors were located in the cervical spine in 10 patients (36 %), the thoracic spine in 7 patients (25 %) and the lumbar spine in 11 patients (39 %). The mean tumor volume was 22,324 mm³ (1,437–1,04,720 mm³) and the mean tumor size, $14 \times 12 \times 29$ mm³ (Table 1).

Tumor surgery included a mean of 1.44 spinal segments per patient. The largest tumor extended over four segments. In all patients a gross total resection was aimed for. This was achieved in 18 patients (67 %). Subtotal tumor resection was possible in seven (26 %) patients (two ependymoma, three meningioma, astrocytoma, metastasis). In two patients (7 %) tumor removal (vascular neoplasia, ependymoma) was stopped after deterioration of neurophysiological parameters at the stage of an extended biopsy (Table 2).

The neurohistopathological workup revealed five patients (19 %) with ependymoma (WHO I), two patients (7 %) with ependymoma (WHO II), one patient (4 %) with astrocytoma (WHO I), nine patients (33 %) with schwannoma and six patients (22 %) with meningioma. There were four patients with histopathological diagnoses of lymphoma, metastasis or hemangiopericytoma (Table 1).

All 27 patients suffered from concomitant degenerative spine disease in the spinal segments of or adjacent to the tumor as described above. There were 5 patients (19 %)

Table 3 Clinical and neurological symptoms before and 1 year after tumor surgery

	Preoperative	Postoperative (1 year follow-up)
Back/neck pain	27 (100 %)	9 (33 %)
Radicular pain	12 (44 %)	1 (4 %)
Sensory deficit	17 (63.0 %)	9 (33 %)
Paresthesia	13 (48 %)	5 (19 %)
Motor deficit	14 (52 %)	5 (19 %)

Table 4 Core Measures Outcome Index (COMI) data

	BP (COMI) Mean $\pm$ SD	EP (COMI) Mean $\pm$ SD	QL (COMI) Mean $\pm$ SD	COMI Mean $\pm$ SD
Preop	3.8 $\pm$ 3.2	5.3 $\pm$ 3.2	3.6 $\pm$ 0.9	6.5 $\pm$ 2.3
3 months FU	2.0 $\pm$ 2.2	2.6 $\pm$ 2.8	2.2 $\pm$ 1.1	3.2 $\pm$ 2.6
1 year FU	2.2 $\pm$ 1.1	2.4 $\pm$ 2.7	2.1 $\pm$ 1.1	2.8 $\pm$ 2.6

COMI and Spine Tango Registry data; complete study group

with spinal canal stenosis, 24 patients (89 %) with spondylarthrosis and 24 patients (89 %) with discopathy and osteochondrosis.

The two patients with malignant disease (lymphoma, metastasis) received further treatment after surgery. The patient with cervical spine metastasis underwent radiotherapy and the patient with thoracic spine lymphoma underwent chemotherapy.

Early postoperative course and complications

The early postoperative course was uneventful in most patients. In the overall group of 58 patients there was one case of postoperative hematoma and one case of CSF leakage; however, within the study subgroup of 27 patients there were no surgical complications and no cases of symptomatic postoperative hematoma, infection or CSF leakage. During hospitalization 2/27 patients (7 %) suffered from cardiac arrhythmias, 1 patient (4 %) from pneumonia, 1 patient (4 %) from postoperative symptomatic transitory psychotic syndrome and 2 patients (7 %) from transitory bladder dysfunction. There were no newly developed neurological deficits with the study population.

Preoperative symptoms, follow-up of symptomatology and COMI data (Tables 3, 4)

Before surgery back/neck pain was present as a leading symptom in all 27 study patients (100 %); radicular pain was present in 12 patients (44 %), sensory deficits in 17 patients (63 %), paresthesia in 13 patients (48 %) and motor deficits in 14 patients (52 %). Tumor surgery

Fig. 1 COMI data of Back Pain (18 patients) and No Back Pain (9 patients) groups showing COMI scores for back pain and quality of life preoperatively and at 3- and 12-month follow-ups. (*P* values: + = 0.035, x = 0.001, ++ = 0.034, * = 0.001)

improved the symptomatology, and at 1-year follow-up only nine patients (33 %) complained of persisting back/neck pain, one patient (4 %) of radicular pain, nine patients (33 %) of sensory deficits, five patients (19 %) of paresthesia and five patients (19 %) of motor deficits (Table 3). No patients developed new neurological symptoms.

Evaluation of the COMI questionnaires showed similar results to those reported by the patients during postoperative examination (see above). In the complete study group the mean preoperative COMI score for back pain was reduced from 3.8 ± 3.2 before surgery to 2.0 ± 2.2 at the 3-month follow-up and 2.2 ± 1.1 at the 1-year follow-up. The mean COMI score for leg/arm pain reduced from 5.3 ± 3.2 before surgery to 2.6 ± 2.8 at the 3-month follow-up and 2.4 ± 2.7 at the 1-year follow-up. Mean COMI Quality of Life score improved from 3.6 ± 0.9 to 2.2 ± 1.1 and 2.1 ± 1.1 . The mean COMI score was 6.5 ± 2.3 before surgery and 3.2 ± 2.6 and 2.8 ± 2.6 at the 3-month and 1-year follow-ups, respectively (Table 4).

For further evaluation of the data, we divided the study population into two groups—one with postoperative persisting back/neck pain (P-group) according to what patients reported during postoperative consultation and one without (no-BP-group). Both groups showed statistically significant improvement of the mean COMI scores for quality of life (Fig. 1). Regarding back/neck pain the self-rated COMI showed the same results as what patients reported during

follow-up. The mean COMI score for back pain was reduced significantly in the no-BP-group from 4.2 ± 3.6 before surgery to 1.7 ± 1.7 and 1.7 ± 1.7 at the 3-month and 1-year follow-ups, respectively (Fig. 1). In the BP-group the score remained around the preoperative level of 3.0 ± 2.2 (2.7 ± 2.8 and 3.4 ± 3.2 at the 3-month and 1-year follow-ups, respectively).

Evaluation of various patient characteristics in the BP-group and no-BP-group revealed no significant group differences with respect to male/female ratio, mean tumor volume, tumor pathology and extent of surgery.

Tumor control, neuroradiological follow-up and necessity of treatment for degenerative spine disease

Patients included in the present study group had follow-up examinations over a mean period of 31 months. MRI studies were carried out at the 3-month and 1-year follow-ups and annually thereafter. Altogether, 23/27 patients (85 %) were in complete remission regarding tumor growth. Three patients were foreseen for radio/chemotherapy due to progression of the tumor remnant (metastasis, astrocytoma, lymphoma) and one patient is scheduled for reoperation due to tumor regrowth (ependymoma).

Over the follow-up period after tumor surgery, only two patients (7 %) required further treatment for back/neck

Fig. 2 Illustrative case of 73-year-old patient with a cervical Schwannoma level C6/7 and cervical multi-level discopathy and spondylarthrosis. Preoperative MRI study: **a–c** preoperative MRI

study (T1 contrast sagittal), **d–f** postoperative MRI study (T1 contrast enhanced axial, sagittal, f T2 sagittal)

pain and degenerative spine disease. One was treated with an epidural infiltration and one underwent surgery (stabilization and fusion) for single segment lumbar degeneration. Both patients were in the group with complaints of persisting back/neck pain after tumor surgery. For the other seven patients in this group, the persisting back/neck pain had only a minimal effect on their quality of life and they underwent non-invasive treatment such as medication or physiotherapy.

Illustrative cases

A 73-year-old patient presented with neck pain, headache, non-specific vertigo, left sided radiculopathy and mild left sided motor weakness (M4/5) of the upper limbs. Tumor surgery through a laminoplasty of C6 was performed and histopathological diagnosis revealed a Schwannoma (WHO I). Postoperative imaging studies showed no tumor remnant or recurrence (Fig. 2) and the patient was in remission for 35 months. The neurological deficits recovered completely. The neck pain improved to the extent that no invasive

therapy was necessary despite the severe degeneration of the cervical spine.

A 60-year-old patient presented with severe back pain, and mild sensory and motor (M4/5) deficits of the lower limbs. Laminectomy at L3 and intradural tumor removal were performed. The histopathological diagnosis was a Schwannoma (WHO I). Postoperative MR imaging showed no tumor remnant or recurrence (Fig. 3) and the patient was in remission for 19 months. Postoperative clinical examination showed complete remission of back pain and neurological symptoms. So far the concomitant degeneration of the adjacent lumbar segments at the levels L2/3 and L4/5 has been asymptomatic.

Discussion

The present study analyzed the characteristics of intradural tumor surgery in patients with coexisting severe degenerative spinal disease and back/neck pain as a leading symptom. It demonstrated that surgery for intradural spinal tumors alone—despite the presence of severe degenerative

Fig. 3 Illustrative case of 60-year-old patient with lumbar Schwannoma level L3/4, lumbar spinal canal stenosis level L4/5 and discopathy and spondylarthrosis level L2/3. **a–c** Preoperative MRI

study (T1 contrast enhanced axial, sagittal, T2 sagittal), **d–f** postoperative MRI study (T1 contrast enhanced axial, sagittal, T2 sagittal)

spinal disease—avoids overtreatment of patients and that undertreatment seems to be rare.

Comparison of the results with previously published data shows that laminectomy/laminoplasty and microsurgical intradural tumor removal assisted by multimodal intraoperative neuromonitoring is a safe and very efficient treatment option concerning neurological symptoms and complications as well as tumor control. A remission rate of 85 % and a rate of surgical complications of 0 % are at least as good as in previous publications [5, 6, 14–19].

Surgical treatment of degenerative spinal disease in patients with intradural spinal tumor is controversial. Coexisting spinal canal stenosis, disk herniation, osteochondrosis or facet joint arthrosis can be treated during the same intervention. However, preoperative evaluation of the cause of symptoms, especially back pain, is difficult such that the surgical overtreatment of patients is also a distinct possibility. All of our study patients showed back/neck pain as a leading preoperative symptom and severe spinal degeneration as a comorbidity. In such situations one might

typically attribute the back pain to the degeneration rather than to the tumor itself. However, our results showed that with intradural tumor surgery only, back/neck pain was resolved in 67 % of the patients and of the remaining 33 %, only two patients (7 %) needed further invasive therapy for degenerative spinal disease. Hence, performing only intradural tumor surgery led to a maximum of 7 % of patients being undertreated and 26 % of patients possibly having a less-good outcome than if they had been treated for degenerative spinal disease as well. On the other hand, we would have overtreated 67 % of patients if a surgical treatment for the degenerative spinal disease had automatically been performed in addition to that for the tumor removal.

There is only one publication addressing this differential diagnostic dilemma. Börm et al. described a series of 10 patients that were diagnosed with and treated unsuccessfully for degenerative spinal disease and in whom, after more detailed diagnostics, spinal tumors were found [25]. Most of the patients in the series were treated with tumor surgery

alone as well. They concluded that with patients with intraspinal tumors comprising 0.5 % of their patients treated for spinal degeneration, further diagnostics are necessary if symptoms do not match with imaging findings.

Other previously published series have discussed the problem of progression of degeneration after tumor surgery [20]. It was found that there was no short-term difference between laminectomy or laminoplasty as far as progression of deformity following tumor surgery was concerned [20]. However, there is some evidence that patients undergoing a more than two-segment laminectomy for tumor surgery have a high risk of developing instability and requiring stabilization and fusion [21].

There are some limitations of the study that need to be mentioned. The present study is a retrospective study of mainly prospectively collected data and patients were followed only for a mean time of 13 months. Another limitation is that two patients underwent further therapy with radio- or chemotherapy, which could potentially interfere with symptoms of back/neck pain. Those two patients were without back/neck pain after surgery.

The results of the present study and evidence from the literature show that in patients with degenerative spinal disease and intradural spinal tumor with back/neck pain as a leading symptom, tumor surgery alone is a safe and effective treatment modality. The data imply that intradural spinal tumor may be the main cause of back/neck pain more often than has hitherto been appreciated. Our results show that overtreatment for concomitant degenerative disease seems to be a higher risk than undertreatment but we could not identify factors predicting the necessity for further treatment for degenerative spinal disease. Undertreatment is nonetheless a rare but existing problem. It is essential to identify the factors that might predict the need for further treatment in order to provide patients who have multiple pathologies with an individual treatment option that addresses only their specific symptomatic pathology.

Conclusions

Tumor surgery for intradural spinal tumors improves back/neck pain in the majority of patients with coexisting severe degenerative spinal disease. Intradural spinal tumors seem to be the only cause of back pain more often than is appreciated. In these patients surgical overtreatment seems to be a greater risk than undertreatment. Therefore, elaborate clinical and radiological examinations should be performed preoperatively and the indication for stabilization/fusion should be discussed carefully in patients foreseen for first time intradural tumor surgery.

Conflict of interest None.

References

1. Bressler HB, Keyes WJ, Rochon PA et al (1999) The prevalence of low back pain in the elderly: a systematic review of the literature. *Spine* 24:1813–1819
2. Celestin J, Edwards RR, Jamison RN (2009) Pretreatment psychosocial variables as predictors of outcomes following lumbar surgery and spinal cord stimulation: a systematic review and literature synthesis. *Pain Med* 10(4):639–653
3. Abul-Kasim K, Thurnher MM, McKeever P et al (2008) Intradural spinal tumors: current classification and MRI features. *Neuroradiology* 50:301–314
4. Hermann DM, Barth A, Porchet F et al (2008) Nocturnal positional lumboschialgia—presenting symptom of lumbar spinal tumours. *J Neurol* 255:1836–1837
5. Gezen F, Kahraman S, Canakci Z et al (2000) Review of 36 cases of spinal cord meningeoma. *Spine* 25(6):727–731
6. Conti P, Pansini G, Moutachy H et al (2004) Spinal neurinomas: retrospective analysis and long-term outcome of 179 consecutively operated cases and review of the literature. *Surg Neurol* 61:35–44
7. Sutter MA, Eggspuehler A, Grob D et al (2007) Multimodal intraoperative monitoring (MIOM) during 409 lumbosacral surgical procedures in 409 patients. *Eur Spine J* 16(Suppl 2):221–228
8. Sutter MA, Eggspuehler A et al (2007) The validity of multimodal intraoperative monitoring (MIOM) in surgery of 109 spine and spinal cord tumors. *Eur Spine J* 16(Suppl 2):197–208
9. Aebi M, Grob D (2004) SSE spine tango: a european spine registry promoted by the Spine Society of Europe (SSE). *Eur Spine J* 13(8):661–662
10. Mannion AF, Porchet F, Kleinstück FS et al (2009) The quality of spine surgery from the patient's perspective. Part 1: the Core Outcome Measures Index in clinical practice. *Eur Spine J* 18(Suppl 3):367–373
11. Deyo RA, Battie M, Beurskens AJHM, Bombardier C, Croft P, Koes B, Malmivaara A, Roland M, Von Korf M, Waddell G (1998) Outcome measures for low back pain research: a proposal for standardized use. *Spine* 23(18):2003–2013
12. Deyo RA, Battie M, Beurskens AJHM, Bombardier C, Croft P, Koes B, Malmivaara A, Roland M, Von Korf M, Waddell G (1998) Outcome measures for low back pain research. A proposal for standardized use. *Spine* 23(18):2003–2013
13. Mannion AF, Elfering A, Staerkle R, Junge A, Grob D, Semmer NK, Jacobshagen N, Dvorak J, Boos N (2005) Outcome assessment in low back pain: how low can you go? *Eur Spine J* 14(10):1014–1026
14. Kivelev J, Niemelä M, Hernesniemi J (2012) Outcome after microsurgery in 14 patients with spinal cavernomas and review of the literature. *J Neurosurg Spine* 13:524–534
15. Wager M, Lapierre F, Blanc JL et al (2000) Cauda equine tumors: a French multicenter retrospective review of 231 adult cases and review of the literature. *Neurosurg Rev* 23:119–129
16. Song KW, Shin SI, Lee JY et al (2009) Surgical results of intradural extramedullary tumors. *Clin Orthop Surg* 1:74–80
17. Ahn DK, Park HS, Choi DJ et al (2009) The surgical treatment for spinal intradural extramedullary tumors. *Clin Orthop Surg* 1:165–172
18. Safavi-Abbasi S, Senoglu M, Theodore N et al (2008) Microsurgical management of spinal schwannomas: evaluation of 128 cases. *J Neurosurg Spine* 9:40–47
19. Jenkinson MD, Simpson C, Nicholas RS et al (2006) Outcome predictors and complications in the management of intradural spinal tumours. *Eur Spine J* 15:203–210
20. McGirt MJ, Garcés-Ambrossi GL, Parker SL et al (2010) Short-term progressive spinal deformity following laminoplasty versus

- laminectomy for resection of intradural spinal tumors: analysis of 238 patients. *Neurosurgery* 66:1005–1012
21. Sciubba DM, Chaichana KL, Woodworth GF et al (2008) Factors associated with cervical instability requiring fusion after cervical laminectomy for intradural tumor resection. *J Neurosurg Spine* 8:413–419
 22. Fujiwara A, Tamai K, Yamato M et al (1999) The relationship between facet joint osteoarthritis and disc degeneration of the lumbar spine: an MRI study. *Eur Spine J* 8(5):396–401
 23. Pfirrmann CW, Metzdorf A, Zanetti M (2001) Magnetic resonance classification of lumbar intervertebral disc degeneration. *Spine (Phila Pa 1976)* 26(17):1873–1878
 24. Modic MT, Steinberg PM, Ross JS (1988) Degenerative disk disease: assessment of changes in vertebral body marrow with MR imaging. *Radiology* 166(1 Pt 1):193–199
 25. Börm W, Gleixner M, Klasen J (2004) Spinal tumors in coexisting degenerative spine disease—a differential diagnostic problem. *Eur Spine J* 13:633–638