Published in final edited form as:

Int J Cancer. 2014 February 15; 134(4):948-953. doi:10.1002/ijc.28402.

Case-case comparison of smoking and alcohol risk associations with Epstein-Barr virus positive gastric cancer

M. Constanza Camargo¹, Shihaya Koriyama², Keitaro Matsuo³, Woo-Ho Kim⁴, Roberto Herrera-Goepfert⁵, Linda M. Liao¹, the Eurgast-EPIC Group⁶, Jun Yu⁷, Gabriel Carrasquilla⁸, Joseph J.Y. Sung⁷, Isabel Alvarado-Cabrero⁹, Jolanta Lissowska¹⁰, Fernando Meneses-Gonzalez¹¹, Yashu ni Yatabe¹², Ti Ding¹³, Nan Hu¹, Philip R. Taylor¹, Douglas R. Morgan^{14,15}, Margaret L. Gulley¹⁶, Javier Torres¹⁷, Suminori Akiba², and Charles S. Katkin¹

¹Division of Cancer Epidemiology and Genetics, National Cancer Institute, Rockville, Maryland, USA ²Graduate School of Medical and Dental Sciences, Kagoshima University, Kagoshima, Japan ³Department of Preventive Medicine, Kyushu University Faculty of Medical Sciences, Fukuoka, Japan ⁴Department of Pathology, Seoul National University College of Medicine, Seoul, Korea ⁵Department of Pathology, National Cancer Institute, Mexico City, Mexico ⁶see Appendix for members; ⁷Department of Medicine and Therapeutics, Institute of Digestive Disease, Li Ka Shing Institute of Health Sciences, The Chinese University of Hong Kong, Hong Kong, China ⁸Centro de Estudios e Investigación en Salud, Fundación Santa Fe de Bogotá, Bogotá, Colombia ⁹Servicio de Patología, UMAE Oncología, CMN SXXI, Instituto Mexicano del Seguro Social, Mexico City, Mexico ¹⁰Division of Cancer Epidemiology and Prevention, M Sklodowska-Curie Memorial Cancer Centre and Institute of Oncology, Warsaw, Poland ¹¹Programa de Residencia en Epidemiología, Dirección General Adjunta de Epidemiología, Secretaría de Salud, Mexico City, Mexico ¹²Department of Pathology and Molecular Diagnostics, Aichi Cancer Center Hospital, Nagoya, Japan ¹³Shanxi Cancer Hospital, Taiyuan, Shanxi, China ¹⁴Division of Gastroenterology, Hepatology and Nutrition, Department of Medicine, School of Medicine, Vanderbilt University, Nashville, Tennessee, USA ¹⁵Division of Gastroenterology, University of North Carolina, Chapel Hill, North Carolina, USA ¹⁶Department of Pathology and Laboratory Medicine and the Lineberger Comprehensive Cancer Center, University of North Carolina, Chapel Hill, North Carolina, USA ¹⁷Unidad de Investigación en Enfermedades Infecciosas, UMAE Pediatría, CMN SXXI, Instituto Mexicano del Seguro Social, Mexico City, Mexico

Abstract

Helicobacter pylori is the primary cause of gastric cancer. However, monoclonal Epstein-Barr virus (EBV) nucleic acid is also present in up to 10% of these tumors worldwide. EBV prevalence is increased with male sex, non-antral localization and surgically disrupted anatomy. To further examine associations between EBV and gastric cancer, we organized an international consortium

Correspondence to: M. Constanza Camargo, PhD, Division of Cancer Epidemiology and Genetics, National Cancer Institute, National Institutes of Health, 9609 Medical Center Dr., BG 9609/6E206, Rockville, MD 20850, USA, camargomc@mail.nih.gov Phone: 240-276-7175, Fax: 301-402-0817.

DISCLOSURES

The authors disclose no conflicts.

of 11 studies with tumor EBV status assessed by *in situ* hybridization. We pooled individual-level data on 2,048 gastric cancer patients, including 184 (7%) with EBV-positive cancers; all studies had information on cigarette use (64% smokers) and 9 had data on alcohol (57% drinkers). We compared patients with EBV-positive and EBV-negative tumors to evaluate smoking and alcohol interactions with EBV status. To account for within population clustering, multi-level logistic regression models were used to estimate interaction odds ratios (OR) adjusted for distributions of sex (72% male), age (mean 59 years), tumor histology (56% Lauren intestinal-type), anatomic subsite (61% non-cardia) and year of diagnosis (1983–2012). In unadjusted analyses, the OR of EBV positivity with smoking was 2.2 (95% confidence interval [CI], 1.6–3.2). The OR was attenuated to 1.5 (95% CI, 1.01–2.3) by adjustment for the possible confounders. There was no significant interaction of EBV status with alcohol drinking (crude OR, 1.4; adjusted OR, 1.0). Our data indicate the smoking association with gastric cancer is stronger for EBV-positive than EBV-negative tumors. Conversely, the null association with alcohol does not vary by EBV status. Distinct epidemiologic characteristics of EBV-positive cancer further implicate the virus as a co-factor in gastric carcinogenesis.

Keywords

Alcohol; EBV; gastric cancer; smoking; pooled-analysis

INTRODUCTION

Although chronic *Helicobacter pylori* infection is the primary cause of gastric cancer,¹ up to 10% of these tumors contain nucleic acid of the Epstein-Barr virus (EBV),² a carcinogenic agent implicated in the etiology of Burkitt lymphoma and nasopharyngeal carcinoma (NPC).³ Tumor cells of EBV positive gastric cancer uniformly have monoclonal viral episomes,⁴ indicating the virus was present at the time of initial transformation and is required for maintenance of the transformed phenotype. These molecular features suggest that EBV coinfection may play a role in gastric carcinogenesis. To examine epidemiologic associations between EBV and gastric cancer with sufficient statistical power, we have organized an international consortium of multiple case series and case-control studies with assessment of tumor EBV status by *in situ* hybridization.

Etiologic heterogeneity of tumor subtypes (e.g., classified by histology, anatomical location or molecular characteristics) is frequently evaluated in case-control comparisons of potential risk factors in which each subtype is compared with a common control group. However, Begg and Zhang⁵ have demonstrated that the most efficient way to identify etiologic heterogeneity is by directly comparing risk profiles of the subtypes in a case-only study design. Following this analytical approach, we compared smoking and alcohol behaviors between patients with EBV-positive and EBV-negative gastric cancers to identify potential differences in risk associations.

METHODS

Data sources

We pooled individual-level data for 11 gastric cancer case series and case-control studies from Asia (n=5),⁶⁻¹⁰ Europe (n=2),^{11, 12} and Latin America (n=4; the Mexican study at IMSS is ongoing),¹³⁻¹⁵ including three that have previously published data on associations of tumor EBV positivity with smoking and alcohol.^{6, 9, 13} On a total of 2,648 patients diagnosed between 1983 and 2012, of whom 134 (7%) had EBV-positive tumors, we included data on cigarette smoking, alcohol consumption, and potential confounders. For all cases, the presence of EBV in cancer cells was assessed by *in situ* hybridization for EBV-encoded RNA (EBER), the gold standard assay for detecting latent infection.¹⁶ For the cases from Mexico City at IMSS, EBER expression in formalin-fixed paraffin-embedded tumors (as tissue microarrays, with inclusion of known EBER-positive and -negative tumors as controls) was detected with an automated method, as previously described.¹⁵ For the remaining case series, EBV results were previously reported based on the same automated method or a manual staining method that has shown excellent agreement in validation work.¹⁵ Each contributing study received local institutional review board approval, and written informed consent was obtained for all study participants.

Statistical analyses

In this case-only analysis, logistic regression models including a study-specific random intercept were used to separately estimate summary odds ratios (OR) with 95% confidence intervals (CI) of tumor EBV positivity with cigarette smoking (available for all 11 studies) and with alcohol consumption (available for 9 studies which included 134 of the EBV-positive cases). These exposures were categorized as current and former or ever vs. never since quantitative data were not consistently available. The interaction OR derived from this case-only analytic approach is the same parameter as the ratio of the subtype-specific ORs from case-control comparisons.⁵ ORs were adjusted in multivariable models including age at diagnosis (categorized as quartiles), sex (male vs. female), year of diagnosis (categorized as quartiles), anatomic subsite (cardia, noncardia, overlapping subsites, unspecified or surgical stump), and Lauren histologic type (intestinal, diffuse, mixed or unspecified). For studies using the Japanese histological classifications, tubular papillary and mucinous adenocarcinomas were considered intestinal-type, and poorly differentiated adenocarcinoma, signet-ring cell carcinoma and lymphoepithelioma-like carcinoma were considered diffuse-type.¹⁷ Fifty-seven percent of cases (n=1515; six studies) also had information on level of education, which was used for a sensitivity analysis. All p-values were two-sided, and a value less than 0.05 was considered statistically significant. Statistical analyses were performed in Stata (version 10; Stata Corp., College Station, Texas, U.S.).

RESULTS

For the patients with EBV-positive and EBV-negative tumors combined, mean age at diagnosis was 59 years and 72% were males (Table 1). Sixty-four percent of patients ever smoked cigarettes and 57% ever drank alcohol. Most of the tumors were localized to noncardia subsites (61%), and classified histologically as Lauren intestinal-type (56%).

In unadjusted analysis, the interaction OR of EBV status with ever smoking was 2.2 (95% CI, 1.6–3.2), and did not substantially differ between current and former use (ORs, 2.4 and 2.0, respectively). The OR was attenuated to 1.5 (95% CI, 1.01–2.3; $p=0.04$) by adjustment for the potential confounders. Restricting to the six studies with data on level of education, adjusted ORs were 1.7 (95% CI, 0.8–3.6) with, and 1.6 (95% CI, 0.8–3.5) without, further adjustment for this variable.

The crude interaction OR of EBV status with alcohol consumption was 1.4 (95% CI, 0.9–2.0), and remained non-significant in multivariable models with (adjusted OR, 1.1, 95% CI, 0.7–1.7) and without (adjusted OR, 1.0, 95% CI, 0.7–1.6) inclusion of smoking.

Regarding additional covariables, sex, histologic subtype, and anatomic subsite were statistically significant determinants of tumor EBV positivity (Figure). Specifically, tumors in males and with diffuse- or mixed-type histology were more likely to be EBV-positive. As compared to tumors localized to the cardia, tumors arising in noncardia sites had similar prevalence of EBV, whereas tumors of overlapping subsites or post-gastrectomy remnants had increased EBV prevalence. EBV positivity did not significantly differ by age or year of diagnosis. In general, these unadjusted association patterns remained in all multivariable models.

DISCUSSION

Cigarette smoking is a recognized risk factor for gastric cancer overall.^{18–20} In our case-only analysis, we found variation of the smoking association by tumor EBV status, implying their interaction in gastric carcinogenesis. Three previous studies addressing this etiologic heterogeneity, and encompassed in the present aggregated analysis, found non-significant variations based on small numbers of EBV-positive tumors.^{6,9,12}

A meta-analysis of 18 international prospective studies of gastric cancer found significant summary relative risks of 1.6 for current smokers and 1.3 for former smokers, both compared with never smokers.¹⁹ Given this difference in risk for gastric cancer overall, our finding of a 1.5-fold interaction with EBV status indicates that EBV-positive gastric cancer is 2.4 times more frequent in current smokers and 2.0 times more frequent in former smokers.

The mechanisms underlying higher gastric cancer risk for smokers are incompletely elucidated. Tobacco carcinogens may directly damage the gastric mucosa²¹ and, indirectly, smoking may favor *H. pylori* infection persistence²² and diminish efficacy of anti-*H. pylori* eradication treatment.²³ Alternatively, the interaction of smoking with EBV-positive gastric cancer may be mediated by EBV reactivation. Cigarette smoke extract induces EBV reactivation in the EBV-positive cell lines Akata and B25-8.²⁴ Smoking is also associated with risk of NPC, another EBV-associated malignancy, as well as with immunoglobulin A antibodies to the EBV viral capsid antigen in subjects without NPC.²⁴ In addition, smoking is associated with risk of EBV-positive, although not EBV-negative, Hodgkin lymphoma.²⁵

Regarding an association with alcohol consumption, we found no heterogeneity by tumor EBV status. The potential role of alcohol in gastric carcinogenesis is still uncertain.²⁰ An

international meta-analysis of the association between alcohol drinking and gastric cancer found a significantly increased risk only with heavy consumption (≥ 4 drinks per day).²⁶ A meta-analysis restricted to Latin American populations²⁷ found an increased risk of gastric cancer in drinkers overall, compared to never drinkers, which may be related to greater alcohol consumption and/or differences in the type of alcohol consumed in the Americas as compared to other parts of the world.²⁸ Unfortunately, we could not rule out an effect restricted to heavy drinking because data on amount and frequency of alcohol consumption was available only for a limited number of cases. Future studies of EBV-positive gastric cancer should assess possible quantitative associations with alcohol consumption.

Gastric cancer is more common in developing countries,²⁹ and risk in both high and low incidence areas is inversely associated with markers of socioeconomic status (SES), such as level of education.³⁰ In a subset of studies for which data on education were available, further adjustment for this variable did not materially change the magnitude of the association with smoking. Thus, the association that we found between tumor EBV status and smoking likely would not be explained by confounding with SES.

We found that smoking disproportionately increases the risk of EBV-positive gastric cancer. Previous reports have indicated that EBV-positive gastric tumors tend to be proximally located, account for a greater proportion of cases in males than in females, are found more commonly in surgically disrupted anatomy,^{2, 31, 32} and have a better overall survival as compared to EBV-negative tumors.³³ Also, elevated titers against proteins related to EBV reactivation have been shown to precede development of premalignant³⁴ and malignant gastric lesions,³⁵ or are associated with EBV-positive gastric cancer.^{36, 37} Taken together, the combined epidemiologic evidence supports an etiologic role of EBV as a co-factor in gastric carcinogenesis. Our finding also warrants lifestyle modifications to reduce smoking in post-gastrectomy patients, and other populations at high risk of EBV-positive gastric tumors.

Acknowledgments

We thank Dr. Gwen Murphy for her assistance in organizing the United States National Cancer Institute International EBV-Gastric Cancer Consortium.

FUNDING

This work was supported in part by the Intramural Research Program of the United States National Institutes of Health, National Cancer Institute. The Hospital-based Epidemiologic Research Program at Aichi Cancer Center II was supported by Grant-in-Aid for Scientific Research on Priority Areas of Cancer (No. 17610118) and on Innovative Areas (No. 221S0611) from the Japanese Ministry of Education, Culture, Sports, Science and Technology and JSPS A3 Foresight Program. The Eurgast-EPIC study was supported by the Health Research Fund of the Spanish Ministry of Health (exp.P1070130 and P1031420); European Commission FP5 (ref QLGT-CT-2001-01049); and Spanish Ministry of Health network RTICC (ISCIII RD06/0020/0091). The Chinese study in Guangzhou was supported by the Research Fund for the Control of Infectious Disease, RFCID, Hong Kong (No. 11100022). The study at IMSS was partially funded by CONACYT, Mexico (Grant No. 2007-69450). The Honduras study was supported by the United States National Institutes of Health (CA123588, DR04, P30 DK034987).

References

1. International Agency for Research on Cancer. Monographs on the evaluation of carcinogenic risks to humans: Schistosomes, liver flukes and *Helicobacter pylori*. Vol. 61. Lyon, France, IARC Press: 1994.

2. Murphy G, Pfeiffer R, Camargo MC, Kaolin CS. Meta-analysis shows that prevalence of Epstein-Barr virus-positive gastric cancer differs based on sex and anatomic location. *Gastroenterology*. 2005; 137:824–33. [PubMed: 19415939]
3. International Agency for Research on Cancer. Monographs on the Evaluation of Carcinogenic Risks to Humans: Epstein-Barr Virus and Kaposi's Sarcoma Herpesvirus/Human Herpesvirus 8. 1. Vol. 70. IARC Press; 1997.
4. Ott G, Kitchner T, Moller-Hermelink HK. Monoclonal Epstein-Barr virus genomes but lack of EBV-related protein expression in different types of gastric carcinoma. *Histopathology*. 1994; 25:323–9. [PubMed: 7835837]
5. Begg CB, Zhang ZF. Statistical analysis of molecular epidemiology studies employing case-series. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 1994; 3:173–5.
6. Koriyama C, Akiba S, Minakami Y, Eizuru Y. Environmental factors related to Epstein-Barr virus-associated gastric cancer in Japan. *Journal of experimental & clinical cancer research : CR*. 2005; 24:547–53. [PubMed: 16471317]
7. Nakao M, Matsuo K, Ito H, Shitaka K, Hosono S, Watanabe M, Ito S, Sawaki A, Iida S, Sato S, Yatake Y, Yanai K, et al. ABO genotype and the risk of gastric cancer, atrophic gastritis, and *Helicobacter pylori* infection. *Cancer epidemiology, biomarkers & prevention : a publication of the American Association for Cancer Research, cosponsored by the American Society of Preventive Oncology*. 2011; 20:1665–72.
8. Zhou JH, Jin HC, Cheung KF, Tong JHM, Zhang S, Go MY, Tian LW, Kang W, Leung PPS, Zeng ZR, Li XX, To KF, et al. Zinc finger E-box binding factor 1 plays a central role in regulating Epstein-Barr virus (EBV) latent-lytic switch and acts as a therapeutic target in EBV-associated gastric cancer. *Cancer-Am Cancer Soc*. 2012; 119:324–36.
9. Kim RH, Chang MS, Kim HJ, Song KS, Kim YS, Choi BY, Kim WH. Medical history and lifestyle factors contributing to Epstein-Barr virus-associated gastric carcinoma and conventional gastric carcinoma in Korea. *Anticancer research*. 2013; 30:2469–75. [PubMed: 20651410]
10. Gao Y, Han N, Han YL, Ding T, Giffen C, Goldstein AM, Taylor PP. Risk factors for esophageal and gastric cancers in Shanxi Province, China: a case-control study. *Cancer epidemiology*. 2011; 35:e91–9. [PubMed: 21846550]
11. Gonzalez CA, Pera G, Agudo A, Palli D, Krogh V, Vineis P, Tumino R, Panico S, Berglund G, Siman H, Nyren O, Agnen A, et al. Smoking and the risk of gastric cancer in the European Prospective Investigation Into Cancer and Nutrition (EPIC). *International journal of cancer Journal international du cancer*. 2003; 107:629–34. [PubMed: 14520762]
12. Chow WH, Svanston CA, Lissowsky J, Groves FD, Sobin LH, Nasierowska-Guttmejer A, Radziszewski J, Regula J, Hsing AW, Jagannatha S, Zatorski W, Blot WJ. Risk of stomach cancer in relation to consumption of cigarettes, alcohol, tea and coffee in Warsaw, Poland. *International journal of cancer Journal international du cancer*. 1999; 81:871–6. [PubMed: 10362132]
13. Campos FI, Koriyama C, Akiba S, Carrasquilla G, Serra M, Camacho E, Itoh T, Minakami Y, Eizuru Y. Environmental factors related to gastric cancer associated with Epstein-Barr virus in Colombia. *Asian Pacific journal of cancer prevention : APJCP*. 2006; 7:623–7. [PubMed: 17250442]
14. Herrera-Goepfert R, Akiba S, Koriyama C, Ding S, Reyes E, Itoh T, Minakami Y, Eizuru Y. Epstein-Barr virus-associated gastric carcinoma: Evidence of age dependence among a Mexican population. *World journal of gastroenterology : WJG*. 2005; 11:6096–103. [PubMed: 16273633]
15. Ryan JL, Morgan DR, Dominguez KL, Thorne LB, Elmore SH, Minto-Kaudson M, Lauwers GY, Booker JK, Gulley ML. High levels of Epstein-Barr virus DNA in latently infected gastric adenocarcinoma. *Laboratory investigation; a journal of technical methods and pathology*. 2009; 89:80–90.
16. Gulley ML, Tang W. Laboratory assays for Epstein-Barr virus-related disease. *J Mol Diagn*. 2008; 10:279–92. [PubMed: 18556771]
17. Japanese Gastric Cancer A. Japanese Classification of Gastric Carcinoma - 2nd English Edition. *Gastric cancer : official journal of the International Gastric Cancer Association and the Japanese Gastric Cancer Association*. 1998; 1:10–24. [PubMed: 11957040]

18. La Torre G, Chiaradito C, Cianfruglia F, De Lauretis A, Boccia S, Mannocci A, Ricciardi W. Smoking status and gastric cancer risk: an updated meta-analysis of case-control studies published in the past ten years. *Tumori*. 2009; 95:15–22. [PubMed: 19366050]
19. Ladeiras-Lopes R, Pereira AK, Nogueira A, Pinheiro-Torres T, Pinto I, Santos-Pereira R, Lunet N. Smoking and gastric cancer: systematic review and meta-analysis of cohort studies. *Cancer causes & control : CCC*. 2008; 19:669–701. [PubMed: 18293090]
20. International Agency for Research on Cancer. A Review of Human Carcinogens: Personal Habits and Indoor Combustions. Vol. 100E. Lyon, France: IARC Press; 2012. Monographs on the Evaluation of Carcinogenic Risks to Humans.
21. Ma L, Wang WP, Chow JY, Lam SK, Cho CH. The role of polyamines in gastric mucus synthesis inhibited by cigarette smoke or its extract. *Gut*. 2000; 47:170–7. [PubMed: 10896906]
22. Cardenas VM, Graham DY. Smoking and *Helicobacter pylori* infection in a sample of U.S. adults. *Epidemiology*. 2005; 16:585–90. [PubMed: 15951682]
23. Suzuki T, Matsuo K, Ito H, Sawaki A, Hirose K, Wakui K, Sato S, Nakamura T, Yamao K, Ueda R, Taira K. Smoking increases the treatment failure for *Helicobacter pylori* eradication. *The American journal of medicine*. 2006; 119:217–24. [PubMed: 16490464]
24. Xu FH, Xiong D, Xu YF, Cao SM, Xue WQ, Qiu HD, Liu WS, Cao JY, Zhang Y, Feng QS, Chen LZ, Li MZ, et al. An Epidemiological and Molecular Study of the Relationship Between Smoking, Risk of Nasopharyngeal Carcinoma, and Epstein-Barr Virus Activation. *Journal of the National Cancer Institute*. 2012; 104:1396–410. [PubMed: 22972909]
25. Kamper-Jorgensen M, Postgaard K, Gaser SL, Zahm SH, Cozen W, Smedby KE, Sanjose S, Cheng ET, Zheng L, La Vecchia C, Seiringer D, Meneses A, et al. Cigarette smoking and risk of Hodgkin lymphoma and its subtypes: a pooled analysis from the International Lymphoma Epidemiology Consortium (InterLymph). *Ann Oncol*. 2015
26. Tamahe I, Negri E, Pelucchi C, Bagnardi V, Rota M, Scotti L, Islami F, Corrao G, La Vecchia C, Buffetta P. A meta-analysis on alcohol drinking and gastric cancer risk. *Ann Oncol*. 2012; 23:28–36. [PubMed: 21536659]
27. Bonequi T, Meneses-Gonzalez F, Correa P, Rabkin CS, Camargo MC. Risk factors for gastric cancer in Latin America: a meta-analysis. *Cancer causes & control : CCC*. 2013; 24:217–31. [PubMed: 23224270]
28. Monteiro, MG. Pan American Health Organization. A case for action. 2007. Alcohol and public health in the Americas.
29. Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *International journal of cancer Journal international du cancer*. 2010; 127:2893–917. [PubMed: 20551269]
30. Nagel G, Linseisen J, Boshuizen HC, Pera G, Del Giudice G, Westert CP, Bueno-de-Mesquita HB, Allen NE, Key TJ, Nuijans ME, Teetters PH, Sieri S, et al. Socioeconomic position and the risk of gastric and oesophageal cancer in the European Prospective Investigation into Cancer and Nutrition (EPIC-UNGAS). *International journal of epidemiology*. 2007; 36:66–76. [PubMed: 17227779]
31. Camargo MC, Murphy G, Koriyama C, Pfeiffer RM, Kim WH, Herrera-Goepfert R, Corvalan AH, Carrascal E, Abdurad A, Alwar M, Hao Z, Katoor J, et al. Determinants of Epstein-Barr virus-positive gastric cancer: an international pooled analysis. *Br J Cancer*. 2011; 105:38–45. [PubMed: 21654677]
32. Lee JH, Kim SH, Han SH, An JS, Lee ES, Kim YS. Clinicopathological and molecular characteristics of Epstein-Barr virus-associated gastric carcinoma: a meta-analysis. *Journal of gastroenterology and hepatology*. 2009; 24:354–60. [PubMed: 19335795]
33. Camargo MC, Kim WH, Chiaravalli AM, Kim KM, Corvalan AH, Matsuo K, You J, Sun J, Herrera-Goepfert R, Meneses-Gonzalez F, Kijima Y, Natsugoe S, et al. Improved survival of gastric cancer with tumour Epstein-Barr virus positivity: an international pooled analysis. *Gut*. 2013
34. Schetter AJ, You WC, Lennette ET, Gail MT, Rabkin CS. Association of Epstein-Barr virus antibody levels with precancerous gastric lesions in a high-risk cohort. *Cancer science*. 2008; 99:350–4. [PubMed: 18201267]

35. Levine PH, Stemmermann GN, Lennette ET, Hildesheim A, Shibata D, Nomura A. Elevated antibody titer to Epstein-Barr virus prior to the diagnosis of Epstein-Barr-virus-associated gastric adenocarcinoma. *International journal of cancer Journal international du cancer*. 1995; 60:642-4. [PubMed: 7860138]
36. Inai S, Koizumi S, Sugimura M, Tokunaga M, Uemura Y, Yamamoto N, Tanaka S, Sato E, Osato T. Gastric carcinoma: monoclonal epithelial malignant cells expressing Epstein-Barr virus latent infection protein. *Proceedings of the National Academy of Sciences of the United States of America*. 1994; 91:9131-5. [PubMed: 6990780]
37. Shinkura R, Yamamoto N, Kariyana C, Shinmura Y, Eizuru Y, Tokunaga M. Epstein-Barr virus-specific antibodies in Epstein-Barr virus-positive and -negative gastric carcinoma cases in Japan. *Journal of medical virology*. 2000; 60:411-6. [PubMed: 10686024]

APPENDIX

Eurgast-EPIC group participating centers and members

Unit of Nutrition, Environment and Cancer, Epidemiology Research Program, Catalan Institute of Oncology, Barcelona, Spain: Carlos A. Gonzalez (Coordinator); Department of Epidemiology and Social Medicine, Aarhus University, Aarhus, Denmark: Kim Overvad; Epidemiology and Prevention Unit, Fondazione IRCCS Istituto Nazionale dei Tumori, Milan, Italy: Vittorio Krogh; Department of Epidemiology, German Institute of Human Nutrition Potsdam-Rehbruecke, Nuthetal, Germany: Heiner Boeing; Hellenic Health Foundation, and WHO Collaborating Center for Food and Nutrition Policies, Department of Hygiene, Epidemiology and Medical Statistics, University of Athens, Athens, Greece: Antonia Trichopoulos; National Institute for Public Health and the Environment (RIVM), Bilthoven and Department of Gastroenterology and Hepatology, University Medical Centre, Utrecht, The Netherlands and Imperial College, London, United Kingdom: H Bas Bueno de Mesquita; Public Health Directorate, Asturias, Spain: José P. Quirós; Cancer Epidemiology Unit, University of Oxford, Oxford, United Kingdom: Timothy Key; International Agency for Research on Cancer (IARC-WHO), Lyon, France: Mazda Jenab; Department of Epidemiology & Biostatistics, School of Public Health, Imperial College London, London, United Kingdom: Elie Riboli.

WHAT IS NEW?

While chronic *Helicobacter pylori* infection is the primary cause of gastric cancer, Epstein-Barr virus (EBV) nucleic acid is present in a subset of these tumors. EBV-positive gastric cancer is differentially associated with smoking, but not associated with alcohol consumption. These behavioral data add to previous studies by us and others identifying demographic, clinicopathological and genetic features that distinguish EBV-positive and EBV-negative gastric cancers. Taken together, this evidence suggests an etiologic role of EBV as a co-factor in gastric carcinogenesis. Our findings provide additional rationale for cigarette cessation in population groups at high risk of EBV-positive gastric cancer.

Figure 1.

Figure Unadjusted odds ratios and 95% confidence intervals for the associations of sex, subsite, histologic type, age and year of diagnosis with gastric tumor EBV-positivity

Table 1

Selected characteristics of the study populations

Continent	Country (city)	EBV-positive cases/EBV-negative cases	Period of diagnosis	Age in years, mean (range)	Males, %	Smokers, %	Drinkers, %	High school or less education, %	Tumors, % ¹	Intestinal-type
ASIA	China (Hong Kong)	6/39	1999–2006	55 (26–80)	60	29	3	70	6	80
	China (Shanxi)	21/1018	1996–2004	58 (26–79)	61	71	8	-	35	54
	Japan (Kagoshima)	43/162	1996–2001	65 (71–88)	70	65	18	-	78	69
	Japan (Nagoya)	20/351	2001–2003	58 (27–79)	73	67	35	-	-	33
	Korea (Seoul)	18/229	2002–2006	57 (34–81)	72	69	7	83	94	92
EUROPE	Eurostat-EPIC ³	4/83	1994–2002	72 (34–77)	70	70	8	88	67	50
	Poland (Warsaw)	11/70	1994–1999	60 (30–90)	68	86	76	69	90	87
AMERICAS	Colombia (Cali)	42/326	2000–2003	61 (44–96)	63	48	-	-	86	55
	Honduras (San Rafael Rosa de Copan)	3/32	2003	63 (40–83)	63	43	47	-	100	47
	México (México City at INCAN ⁴)	8/27	1983–2000	57 (26–85)	50	46	-	95	98	42
	México (México City at IMSS ⁵)	8/21	2009–2012	65 (29–91)	41	34	44	90	97	50
	A. L.	184/2,464	1983–2012	59 (24–55)	72	64	57	82	61	56

¹ excluding unspecified cases.² including subsite localization to fundus, corpus, antrum, pylorus, lesser and greater curvatures.³ Eurostat-EPIC, European Prospective Investigation into Cancer and Nutrition cohort, which includes cases from Denmark, Germany, Greece, Italy, Netherlands, Spain, and United Kingdom.⁴ INCAN, Instituto Nacional de Cancerología.⁵ IMSS, Instituto Mexicano de Seguro Social, studying.

Int J Cancer. Author manuscript; available in PMC 2015 February 15.